

U.S. Department of the Interior
National Park Service

**Bird Monitoring Protocol for Agate Fossil Beds National Monument,
Nebraska and Tallgrass Prairie National Preserve, Kansas**

Prepared by:

David G. Peitz (Ecologist)¹
Steven G. Fancy (National Monitoring Coordinator)²
Lisa P. Thomas (Program Coordinator)³
Gareth A. Rowell (Data Manager)¹
Mike D. DeBacker (Ecologist-Program Coordinator)¹

¹Prairie Cluster Prototype Monitoring Program
Wilson's Creek National Battlefield
6424 West Farm Road 182
Republic, MO 65738

²National Park Service
1201 Oak Ridge Drive, Suite 200
Fort Collins, CO 80525

and

³Southern Colorado Plateau I&M
Northern Arizona University
P.O. Box 5765
Flagstaff, AZ 86011

Prepared for:

Prairie Cluster Prototype Monitoring Program,
National Park Service, U.S. Department of the Interior

Version 1.00 (July 18, 2003)
Revised Version 1.01 (June 18, 2004)

Important Note: This sampling protocol consists of this Protocol Narrative and the following Standard Operating Procedures (SOPs):

SOP 1:	Preparations and Equipment Setup Prior to the Field Season
SOP 2:	Training Observers
SOP 3:	Using the Global Positioning System
SOP 4:	Establishing and Marking Sampling Plots
SOP 5:	Conducting the Variable Circular Plot Count
SOP 6:	Documenting Habitat Variables
SOP 7:	Data Management
SOP 8:	Data Analysis
SOP 9:	Reporting
SOP 10:	Procedures and Equipment Storage After the Field Season
SOP 11:	Revising the Protocol

Table of Contents:

I. Background and Objectives

- Issue being Addressed and Rationale for Monitoring Grassland Bird Populations
- Historical Development of Bird Monitoring in Prairie Parks
- Measurable Objectives

II. Sampling Design

- Rationale for Selecting this Sampling Design over Others
- Site Selection
- Population being Monitored
- Sampling Frequency and Replication

III. Field Methods

- Field Season Preparations, Field Schedule and Equipment Setup
- Sampling Methods
- Conducting the Variable Circular Plot Count
- Collecting Habitat Data

IV. Data Management

- Overview of Database Design
- Data Entry
- Data Verification and Editing
- Metadata Procedures
- Data Archival Procedures

V. Analysis and Reporting

- Bird Data Analysis
- Habitat Data Analysis
- Reporting

VI. Personnel Requirements and Training

- Roles and Responsibilities
- Qualifications and Training

VII. Operational Requirements

- Annual Workload and Field Schedule
- Facility and Equipment needs

Startup Costs and Budget Considerations
 Procedures for Revising the Protocol

VIII. References

Appendix A. Field data forms

Appendix B. Bird Species

Appendix C. Vegetation Guilds – by park

Appendix D. Common Tree Species

I. Background and Objectives

Issue being Addressed and Rationale for Monitoring Grassland Bird Populations

Birds are an important component of park ecosystems, and their high body temperature, rapid metabolism, and high ecological position in most food webs make them a good indicator of the effects of local and regional changes in ecosystems. It has been suggested that management activities aimed at preserving habitat for bird populations, such as for neotropical migrants, can have the added benefit of preserving entire ecosystems and their attendant ecosystem services (Karr 1991, Maurer 1993). Moreover, birds have a tremendous following among the public, and many parks provide information on the status and trends of birds in the park through their interpretive program.

Native Great Plains grasslands and their constituent avian fauna once covered vast areas of the North American continent. During the last century, large portions of grassland landscapes were plowed for cropland or converted to livestock pasture (29% of shortgrass, 41% of mixed-grass, and 99% of tallgrass prairie; Knopf and Sampson 1997). Remaining grasslands have been altered through continued fragmentation and isolation, interruption of driving ecological processes such as periodic wildfire, and loss of significant faunal species, including bison (*Bos bison*), elk (*Cervus elaphus*), and wolves (*Canis lupus*).

While not to the extent of large native ungulates and mammalian predators, many grassland avian species have also demonstrated declining abundance as prairie habitat loss continues. Data collected during the U.S. Geological Survey's annual North American Breeding Bird Surveys (BBS) between 1966 and 1999 indicates that 70% of 29 grassland bird species show evidence of population declines (Sauer et al. 2000). Many prairie species such as the grasshopper sparrow (*Ammodramus savannarum*), eastern meadowlark (*Sturnella magna*), horned lark (*Eremophila alpestris*), bobolink (*Dolichonyx oryzivorus*), lark bunting (*Calamospiza melanocrys*), and dickcissel (*Spiza americana*) have declined at alarming rates (Sauer et al. 2003). The destruction and fragmentation of prairie landscapes, as well as the structural degradation resulting from fire suppression and changes in grazing regimes of remaining prairie habitats contribute to these declines.

Accumulation of litter and standing dead material are significant components in the structure of a tallgrass prairie. In the absence of fire and grazing, dead material accumulates, decreasing availability of nutrients and reducing solar radiation reaching the ground resulting in a homogeneous canopy dominated by big bluestem (*Andropogon gerardii*) and indiagrass (*Sorghastrum nutans*), with eventual invasion by woody plants where precipitation permits (Vinton and Collins 1997). Fire tends to remove accumulated dead material more evenly in grasslands than grazing that produces a heterogeneous mix of live and dead plant material. The season, frequency and spatial extent of fire and grazing interact to influence litter and standing

dead material accumulation, creating a mosaic in vegetation structure across the landscape (Glenn et al. 1992). The accumulation of litter and standing dead material and the resulting vegetation structure vary along an east (tallgrass) to west (shortgrass) gradient with mixed-grass prairies falling somewhere in between. Net annual production and the accumulation of dead material are significantly greater in eastern tallgrass prairies than western shortgrass prairies. Shortgrass prairies also develop a more heterogeneous canopy than tallgrass prairie, with bunchgrasses, such as blue grama (*Bouteloua gracilis*), and shrubs dominating (Lane 1995, Vinton and Collins 1997). Consequently, the accumulation of litter and standing dead material in shortgrass prairie systems is an insignificant component of vegetation structure.

Responses of avian communities to changes in habitat structure have received much attention in recent years (Cody 1981, 1985, Zimmerman 1997, Fitzgerald and Pashley 2000, Tappe et al. in press). Much of this work has been carried out in forested systems, which are structurally more diverse than grassland systems. However, work done by Zimmerman (1982, 1988, 1993) on the Konza Prairie Research Natural Area in the Flint Hills of Kansas has demonstrated how avian species respond to fire and grazing in a tallgrass prairie system. Zimmerman (1997) categorized three types of bird species inhabiting tallgrass prairie: grass/forb-dependent, woody-dependent and habitat-independent and recorded their responses to grazing and fire. Woody-dependent species are generally restricted to riparian corridors where woods are protected from fire. In prairies invaded by woody species, restoration of fire and grazing lead to a decline in woody species through removal of their habitat. Grass/forb-dependent species in tallgrass prairie generally respond well to moderate grazing that increases structural diversity. On the other hand, annual spring burns to promote warm season grass production have a negative influence on structural development by removing cool season grasses, forbs, litter and standing dead material, thus reducing habitat quality for grass/forb-dependent species. While some elements of habitat-avian community relationships are clearly understood for prairie systems, such as the response following removal of woody plants, others are not. The influence of habitat alterations on invertebrate communities and the birds that feed upon them is one component that is not fully understood.

It is against a backdrop of vanishing or altered prairie ecosystems and declining grassland bird populations that we propose monitoring avian communities on National Park Service lands. Long-term trends in the community composition and abundance of breeding bird populations provide one measure for assessing the ecological integrity and sustainability of prairie systems. Long-term patterns in community composition and species abundance in relation to changes in the structural diversity of vegetation resulting from fire and grazing will improve our understanding of the effects of various management actions.

Historical Development of Bird Monitoring in Prairie Parks

Trends in the composition and abundance of grassland bird populations have been proposed as a long-term indicator of prairie ecosystem integrity, which is defined as the capability to support and maintain a balanced, integrated, adaptive community of organisms having a community composition, diversity, and functional organization comparable to that of natural habitat of the region (Karr and Dudley 1981). During 1998 and 1999, as part of the design phase of the Prairie Cluster Prototype Long-term Ecological Monitoring Program (PC-LTEM) led by the U.S. Geological Survey, bird inventories and pilot monitoring work was conducted in eight prairie parks by Dr. Abby Powell of the USGS to identify the species present at each park and assess the feasibility of using grassland birds as indicators of ecosystem

integrity (Powell 2000). Work was conducted at Agate Fossil Beds National Monument (AGFO), Badlands National Park (BADL), Homestead National Monument of America (HOME), Pipestone National Monument (PIPE), Scotts Bluff National Monument (SCBL), Tallgrass Prairie National Preserve (TAPR), Theodore Roosevelt National Park (THRO), and Wilson's Creek National Battlefield (WICR). Powell (2000) concluded that Agate Fossil Beds National Monument and Tallgrass Prairie National Preserve were of sufficient size to support healthy grassland bird communities, but recommend against avian community monitoring in other PC-LTEM parks (HOME, PIPE, SCBL and WICR) because of their small size. Following Powell's recommendation, the PC-LTEM Program initiated a pilot project in the spring of 2001 to monitor avian communities at Agate Fossil Beds National Monument and Tallgrass Prairie National Preserve. Grassland bird monitoring may be of particular importance at Tallgrass Prairie National Preserve, where current management includes annual prescribed fire and heavy cattle grazing. As the park General Management Plan is implemented, greater structural diversity and improved avian habitat may be observed.

Measurable Objectives

There are two primary objectives for the monitoring described in this protocol:

1. Identify significant temporal changes in composition and abundance of bird communities at Agate Fossil Beds National Monument and Tallgrass Prairie National Preserve during the breeding season.
2. Improve our understanding of breeding bird – habitat relationships and the effects of management actions such as grazing and prescribed fire regimes on bird populations by correlating changes in bird community composition and abundance with changes in habitat variables.

II. Sampling Design

Rationale for Selecting this Sampling Design over Others

Hundreds of different sampling approaches have been used to quantify status or trends in bird populations and dozens of different monitoring programs are currently in place throughout North America to determine local, regional, or national trends in bird numbers. Most common survey methods allow simultaneous collection of information about species that share a common life history or habitat, but no single method will adequately sample the diversity of either habitats that birds occupy or life history groups such as seabirds, songbirds, raptors and shorebirds.

The sampling design described in this protocol involves a series of sampling stations laid out on a systematic grid that are sampled during 5 minute counts using the variable circular plot (VCP) methodology (Reynolds et al. 1980, Scott et al. 1986, Buckland et al. 1993, 2001, Fancy 1997, Nelson and Fancy 1999). Variable circular plots and line transect sampling are two types of distance sampling. Distance sampling has been used for more than 30 years to estimate animal abundance and is the best omnibus method currently available for determining relative abundance or trends for birds in most sampling situations. In practice, the method documented in this protocol is basically the same as point counts, such as those used in the Breeding Bird Survey (BBS), except that for each bird heard or seen during the count, its horizontal distance from the observer is estimated. In the case of line transect sampling, the observer walks down a transect and records either the perpendicular distance to each bird heard or seen, or else records the sighting angle and sighting distance instead of the perpendicular distance. Line transects are usually more efficient than VCP counts where they can be conducted because the observer

collects data continuously while walking along the transect, whereas during VCP counts the observer only records birds detected during 3-10 minute sampling periods from stations located along a transect or on a systematic grid. However, VCP counts are the preferred approach in patchy habitats where correlations between bird data and vegetation or other habitat information are sought and in dense, rugged or hazardous terrain where you need to watch your footing as you traverse the landscape. Another advantage of VCP counts is that data can be directly compared to historical point count data such as those obtained during BBS counts and can contribute to ongoing programs such as the National Point Count Database.

An important benefit of using the variable circular plot method is the ability to accommodate a wide range of bird species, each of which possesses a different singing style and each of which may occur in a variety of acoustically-different habitats (BCRIB 1999). VCP counts operate by essentially allowing the habitat to determine the size of the area being surveyed. The maximum detectable distance to a bird may change between different habitats, but the radius of the survey will also change. For example, surveys of grassland birds usually cover a larger area per point because of the absence of a screen of trees, and because bird species may be flushed at greater distances in open habitats (BCRIB 1999).

In addition to the rationale for VCP counts given above, this protocol uses variable circular plot counts arranged in a systematic grid to address the two objectives for bird monitoring in prairie parks for the following reasons:

- A primary objective is to associate changes in bird community composition and abundance with long-term changes in vegetation that occur as a result of natural succession or various management actions (e.g., prescribed fire, grazing). We cannot (prior to the first year of the long-term sampling scheme) stratify the park into areas that will be subjected to different management regimes (e.g., grazed versus ungrazed) because the location and timing of future management will be based on feedback from adaptive management and other factors and cannot be predicted. Thus, the approach is to sample birds on a systematic grid that will allow inferences to be made to the entire park, and to later post-stratify the data to investigate changes in bird community composition and numbers as a result of different management regimes.
- Unadjusted point counts, such as those used in the Breeding Bird Survey, do not account for the number of birds missed during the counting interval. The number of birds that are counted at a sampling station is a combination of the number of birds that are actually there, and the proportion of them detected. Use of unadjusted point counts to calculate trends in abundance assumes that the proportion of birds detected is the same among species, observers and counting conditions, which is usually not the case. Without a measure of detectability, counts of birds are an unreliable measure of differences in the actual number of birds present (see Burnham 1981, Barker and Sauer 1995, Nelson and Fancy 1999). To obtain credible long-term trend data, and to allow comparisons among bird species and habitat types where detectability is expected to differ, we will incorporate distance measures to improve the quality of the data.
- By recording distances and by keeping track of whether a bird was detected during the first 3 minutes of the count or between minutes 3-5, we will be able to make direct comparisons between our data and those collected by others using the BBS or Ralph et al. (1995) methods.
- Our approach allows data from repeated surveys of the same area or areas with similar habitat characteristics to be combined to increase sample sizes. By combining surveys, it is

possible to develop detection functions for uncommon species for which few detections are recorded during any single survey.

Note: Occasionally, there is detectability issues in bird sampling that VCP and other estimation procedures cannot address. For example, there may be unobservable portions of the population (such as cryptic and silent females or nocturnal birds) that are not detected at all during counting, or it may be impossible to estimate detectability at the appropriate scale, for example when habitat-specific detectability exists in a rare species. In structurally complex habitats (e.g., forests in the northwest U.S. with very tall trees and multiple shrub and tree layers that make it impossible to see birds directly above the observer but high up in the canopy) it may be difficult or impossible to detect birds at close horizontal distances to the observer. Thus, even with a measure of detectability factored into estimates derived from counts of birds, such estimates may still be an unreliable measure of differences in the actual number of birds present in some situations. However, it is important to note that although data may be less reliable for species encountered at low rates, it is still valuable to collect the data, and pooling of data over time may allow certain limited analyses. Interpretation of survey data requires sensitivity to these extra-statistical limitations of the estimation procedures.

Site Selection

Sampling locations or ‘plots’ are selected as described in SOP #4, “Establishing and Marking Sampling Plots.” Briefly, permanent sampling locations were selected by overlaying systematic grids of 400 x 400-m cells (originating from a random starting point) on park maps. Systematic sampling across the park will allow us to make park-wide inferences concerning the avian community. The orientation of each grid was rotated 45 degrees from North to prevent sampling sites from being influenced by man-made features (roads, fences, etc.) oriented along cardinal directions. In each park, the riparian corridor was identified as a separate stratum, with sampling extending 125 m on either side of the stream channel (Niobrara River, AGFO; Palmer and Fox Creeks, TAPR). The riparian stratum makes up 15.6% of the total park area (965 ha) at Agate Fossil Beds National Monument, and 5.3% of the total park area (4398 ha) at Tallgrass Prairie National Preserve. Within the riparian stratum, plots were located at 250-m intervals along the extent of the stream. Any plots from the overall park grid that fell within the riparian stratum were discarded. A total of 242 and 40 plots were established in upland prairie areas at Tallgrass Prairie National Preserve and Agate Fossil Beds National Monument, respectively. An additional 18 riparian plots were established at Tallgrass Prairie National Preserve and 14 at Agate Fossil Beds National Monument. Sampling locations are not physically marked on the ground; rather UTM coordinates are maintained and located in the field with a GPS unit. Refer to SOP #3 “Using GPS” and SOP #4 “Establishing and Marking Sampling Plots” for navigation between sampling plots and plot establishment instructions.

This systematic approach to selecting sampling sites allows flexibility to choose the appropriate reference frame to answer different monitoring questions. When making park-wide inferences, results from each stratum will be weighted by area and combined to give an overall park mean and variance. At the same time, more intensive sampling in the riparian corridor will ensure an adequate sample to describe habitat relationships specific to this less common, but important stratum. The systematic grid will also allow us to limit the reference frame appropriately when asking more specific monitoring questions (e.g. only those sampling points within particular management units would be used to compare the avian response to different fire or grazing regimes).

Population being Monitored

Sampling will be limited to the bird-breeding season (early May through mid-June), and sampling will include those species that may potentially breed in the park (see Appendix B). Thus, the population being sampled includes breeding bird species within the park boundary that are present during the time that the survey is being conducted.

Sampling Frequency and Replication

All plots will be sampled annually at Agate Fossil Beds National Monument. To limit the effort needed to complete annual breeding bird surveys at Tallgrass Prairie National Preserve, an interpenetrating, split panel revisit plan [1-0,1-4] is utilized in which plots in one panel will be sampled in all years ($n = 41$); and plots in five additional panels will be visited on a rotating basis once every five years ($n = 40$ or 41) (McDonald 2003). Panel membership was assigned to all plots in the systematic sample by numbering plots sequentially along transects aligned northwest to southeast, starting with the most northeastern plot. Assignment to panels A through F was done in a repeated fashion to the sequenced numbers (i.e. number 1 = A, 2 = B, 3 = C, 4 = D, 5 = E, 6 = F, 7 = A, 8 = B and so on). To determine membership in the annual panel and the sequence of the remaining panels, the six letters (A – F) were randomly sequenced. The result was E, A, D, B, F and C. Therefore, panel E will be visited annually, panel A in years 1 and 6, panel D in years 2 and 7, B in years 3 and 8, and so on.

Along with two panels, annual sampling will include the 18 riparian sites at Tallgrass Prairie National Preserve resulting in an annual work load of approximately 100 sites per year, with 50 percent of the annual sampling effort in upland prairie allocated to the annually sampled panel.

Schematic for the revisit design for upland prairie

	Sample Occasion									
	1	2	3	4	5	6	7	8	9	10
Panel E	x	x	x	x	x	x	x	x	x	x
A	x					x				
D		x					x			
B			x					x		
F				x					x	
C					x					x




Figure 0.01.1. Sampling plot memberships in design panels for Tallgrass Prairie National Preserve.

III. Field Methods

Field Season Preparations, Field Schedule and Equipment Setup

Prior to the field season each year, usually beginning in March or April, the observer(s) should review this entire protocol, including all of the SOPs. The observer should pay special attention to the tasks described in SOP #1 “Before the Field Season” and SOP #2 “Training Observers.” Review of bird identification by sight and sound (see SOP #2) is particularly important each year, as the misidentification of a species is perhaps the most serious error one can make during a bird count with greater consequences than errors in estimating distances or double-counting a bird. All of the equipment and supplies listed in SOP #1 should be organized and made ready for the field season, and copies of the field data forms in Appendix A should be made, 25% of which should be on write-in-the-rain paper.

Staff workloads and unpredictable weather (e.g. delays in the onset of spring, periods of rain, spring storms) necessitate maintaining some flexibility in scheduling the sequence and duration of sampling trips. Sampling dates should be scheduled and logistics organized prior to the start of each field season. Twelve to fifteen VCP counts should be scheduled for completion each field day. Field trips up to ten days in length will be necessary to complete bird surveys within the breeding season. In years when both Tallgrass Prairie National Preserve and Agate Fossil Beds National Monument are being surveyed, it is recommended that surveys be

completed at Tallgrass Prairie National Preserve first, to accommodate differences in vegetation phenology.

Sampling Methods

The protocol comprises two data sets collected concurrently; bird abundance (VCP) and habitat measures. The bird crew should arrive at the park the afternoon prior to the first day of sampling to familiarize themselves with the area and the birds present. Each night, the group discusses the next day's plans and establishes the route for visiting bird plots. The birding crew arrives at the first plot each day before sunrise and begins sampling as soon as it is light enough to do so. The habitat crew follows the route of the bird observer, being careful to avoid walking to the next plot until bird observations are completed and the observer has moved on. Habitat sampling on a plot commences after bird observations are complete. Loud voices, motion, or walking through a plot may disrupt the birds and bias survey results. The bird observer completes counts around four hours after local sunrise and then joins the habitat crew. Each day, the habitat crew attempts to complete habitat work on all plots sampled for birds that morning. However, this is not always possible and some plots may need to be sampled at a later date. No habitat sampling should be scheduled after June 30. If for any reason a habitat subplot extends beyond park boundaries, the plot is dropped from sampling. Refer to SOP #5 "Conducting the Variable Circular Plot Count" for conducting bird counts and SOP #6 "Documenting Habitat Variables" for doing habitat work.

Before leaving the field each day, data sheets are checked for completeness and readability. All information pertinent to the plots sampled that day is recorded to avoid repeating or skipping sampling plots. The project manager is responsible for the safekeeping and organization of the data sheets, and ensuring that data get entered into the database.

Conducting the Variable Circular Plot Count

Variable Circular Plot Counts are a point count methodology that incorporates a measure of detectability into population estimates (Buckland et al. 2001). Details of how to conduct VCP counts during the breeding season and for filling in data forms are given in SOP #5 "Conducting the Variable Circular Plot Count" and are summarized here. All birds seen or heard at each plot are recorded during a 5 minute sampling period. Bird observations are separated into two time segments: those detected during the first 3 minutes of the count (to allow comparisons with BBS data), and any new birds detected during the final 2 minutes of the count. All birds, regardless of distance detected from the observer are counted and recorded. For most species, each individual bird will be recorded as a separate observation. For species that usually occur in clusters or flocks, the appropriate unit to record is the cluster or flock size, and not the individual bird.

When we conduct a VCP count, we are attempting to get an "instantaneous count" of birds present. Birds flushed from the plot when approached by the observer should be recorded and the count started as soon as the observer is at plot center. The method takes into account the fact that birds close to the observer have a higher probability of being detected (if they are not flushed) than birds far from the observer and that different species have different detection functions (i.e., the probability of detecting a bird at different distances from the observer). An important assumption of the method is that a bird exactly at the center of the plot has a probability of $p = 1$ of being detected, and that there is a high probability of detecting birds within the first 5-10 meters of the plot center. The most important birds to detect are those very close to the observer (within the first 5-10 meters), and it is highly desirable that estimated

distances be within 1-2 meters of actual distances for any bird within 20 m of the observer. However, all birds seen or heard should be recorded with an estimate of distance from the observer. Distances should be estimated to the nearest meter in the field (during subsequent data analysis, distances will be grouped into intervals to lessen the effect of errors in estimating distances); do not round distances off to the nearest 5 or 10 m. Birds far from the center of the plot usually have little effect on the resulting distance function; during data analysis, 5-10% of the furthest detection distances for each species will be truncated before a detection function is calculated.

Once a count is completed at a plot and the data sheet filled out the observer uses a GPS unit to navigate to the next plot to sample. Twelve to fifteen plots can be sampled each morning during a period when it is light enough to observe birds to four hours post sunrise.

Collecting Habitat Data

Habitat data are collected in a 50-m radius plot centered on each sampling point and in four nested 5-m radius and 1.78-m radius subplots. SOP #6 “Documenting Habitat Variables” gives a detailed description of how habitat data are collected; a brief summary is presented here.

Habitat data are collected to meet several objectives. Large-scale plot attributes such as vegetation type, slope, and water cover are relevant to bird community composition and also place the plot in a landscape context. Canopy structure and horizontal vegetation profile data (collected at 5-m radius scale) characterize habitat structure available to birds and also link bird data to park vegetation types. Similarly, ground cover and foliar cover of vegetation guilds (collected at the 1.78-m radius scale) allow us to integrate bird monitoring with more in-depth vegetation data collected as part of plant community monitoring. Integrating these monitoring projects will allow us to provide feedback to management on how grazing and fire regimes influence vegetation, which in turn affects bird community composition and abundance.

The collection of habitat data starts each morning after the first VCP count has been completed. The habitat crew samples VCP plots in the same order as the bird observer to avoid disturbing birds on plots where the bird count has not been conducted. The bird observer flags the center point of each VCP, making it easier for the habitat crew to find. The habitat crew navigates to the plots using a compass, a field map that includes compass bearings for the grid, and directions from the bird observer. Once the habitat crew arrives at a plot, they set up subplot one (plot center) and complete all habitat measures for this subplot and for the 50-m radius plot, before locating and completing subplots two, three and four.

IV. Data Management

Overview of Database Design

Microsoft Access 97 is the primary software environment for managing bird and habitat data. ESRI ArcInfo 8 serves as a tool for validation of spatial data residing in Access. Data products will be posted at the NPS I&M website:

<http://science.nature.nps.gov/im/monitor/protocoldb.cfm>. Metadata for grassland bird monitoring data will be available at NPS I&M application server: <http://science.nature.nps.gov/im/apps.htm>. QA/QC guidelines in this document are based on recommendations of DeBacker et al. (2002) and S. Fancy at <http://science.nature.nps.gov/im/monitor> and citations therein.

The general data model for grassland bird monitoring consists of two sets of tables. One set manages bird abundance data the other associated habitat data. Abundance and habitat data

are linked in time and space by way of standardized event and location tables. For data management purposes, it is probably easier to manage bird and habitat data separately. When the two are correctly linked, they can be integrated transparently for analysis. One issue that becomes readily apparent when managing the two data sets is that time-scale for bird observation events is in minutes and hours, whereas time-scale for associated habitat data is in days and weeks. While the location linkage is unaffected, this has consequences for the structure of the event-relationships. For example, tbl_SamplingPeriod is used rather than tbl_SamplingEvents in the database to accommodate the differences in time-scale between bird and habitat data collections.

The primary table for storing bird abundance contains information about bird species and bird distance from observer as well as flock size, age, sex, whether the observation was a flyover and species code (AOUcode). Supporting tables include water and climatic information, observers and the period of observation. A locations table provides detailed location information associated with each sampling point.

A habitat subplot table serves as the star table about which all other habitat information is linked. Small look-up tables are linked to this table by way of multiple junction tables. This facilitates many-to-many relationships in Microsoft Access. Small attribute tables provide the values for pick-lists on data-entry forms, thereby reducing possible error during data entry (see Data Verification and Editing below).

Data Entry

A number of features have been designed into the database to minimize errors that occur when field data is transcribed to the database for storage and analysis. Forms are used as portals for data entry into the database. Standardized identifiers (e.g. sample event or plot) are selected from list of easily interpreted codes. Species and habitat data is entered into fields linked to appropriate tables. Look-up tables contain project specific data and prohibit entry of data into a field if a corresponding value is not included in the look-up table. Consequently, only valid names or measures may be entered and spelling mistakes are eliminated. Species or habitat measures are selected using a pick list or by typing the beginning of the name. While typing, the form searches for a similar entry, typically locating the desired species after typing only a few characters.

If a species name or habitat measure is not accepted during data entry, it may be the result of a name change or it may be a new species or habitat measure. To check the current nomenclature for a data variable, a nomenclature update form is attached to the Access data file via a button. If the species or habitat measure name has changed, the form will indicate the current accepted name. If the species or habitat measure is new to the nomenclature update form, a form is provided to update the nomenclature table with the new attribute data.

Data Verification and Editing

Data verification immediately follows data entry and involves checking the accuracy of computerized records against the original source, usually paper field records. While the goal of data entry is to achieve 100% correct entries, this is rarely accomplished. To minimize transcription errors, our policy is to verify 100% of records to their original source by staff familiar with project design and field implementation. Further, 10% of records are reviewed a second time by the Project Manager and the results of that comparison reported with the data. If errors are found in the Project Manager's review, then the entire data set is verified again. Once

the computerized data are verified as accurately reflecting the original field data, the paper forms are archived and the electronic version is used for all subsequent data activities.

Although data may be correctly transcribed from the original field forms, they may not be accurate or logical. For example, a canopy height of 1125 m instead of 11.25 m or a temperature of 95°C instead of 9.5°C is illogical and almost certainly incorrect, whether or not it was properly transcribed from field forms. The process of reviewing computerized data for range and logic errors is the validation stage. Certain components of data validation are built into data entry forms (e.g. range limits). Data validation can also be extended into the design and structure of the database. As much as possible, values for data-entry forms have been limited to valid entries stored in the look-up tables.

Additional data validation can be accomplished during verification, if the operator is sufficiently knowledgeable about the data. The Project Manager will validate the data after verification is complete. Validation procedures seek to identify generic errors (e.g. missing, mismatched or duplicate records) as well as errors specific to particular projects. For example, validation of bird data includes database query and comparison of data among years. One query detects records with a location ID from a park and a period ID from a different park. Another query counts the number of subplots per plot (typically there are 4) to assure that all subplots were entered. Finally, data are compared to previous years to identify gross differences. For example, *Sturnella neglecta* or *S. magna* may be recorded this year, but *Sturnella spp.* the previous. During the entry, verification and validation phases, the Project Manager is responsible for the data. The Project Manager must assure consistency between field forms and the database by noting how and why any changes were made to the data on the original field forms. In general, changes made to the field forms should not be made via erasure, but rather through marginal notes or attached explanations. Once validation is complete, the data set is turned over to the Data Manager for archiving and storage.

Spatial validation of database sample coordinates can be accomplished using ArcGIS (ESRI, Inc.). Because Tweety is maintained as an Access database, it can be imported directly into ArcCatalog (ArcGIS, ESRI, Inc.) as an OLE DB object. Fields “StartUTMX” and “StartUTMY” of the locations table can then be used to validate the UTM coordinate values for point locations stored in Access against the original GPS coordinates.

Metadata Procedures

Preliminary notes for metadata were developed using the Electronic Metadata Guide (developed by G. Lienkaemper, USGS FRESC, Corvallis, OR - see Appendix A of SOP # 7). Metadata are then entered into the National Park Service Inventory and Monitoring Dataset Catalog (Ver. 2002.1). Then an extended report from Dataset Catalog is generated. The Dataset Catalog report tool can be used to generate output in the text-based FGDC-compliant metadata format. The output file should be validated and parsed into html format using the USGS metaparser command tool with the following syntax “mp -e efile -h hfile infile.”

Data Archival Procedure

Data sets are rarely static. They often change through additions, corrections, and improvements made following the archival of a data set. There are three main caveats to this process:

- Only make changes that improve or update the data while maintaining data integrity.
- Once archived, document any changes made to the data set.

- Be prepared to recover from mistakes made during editing.

Any editing of archived data is accomplished jointly by the Project Manager and Data Manager. Every change must be documented in the edit log and accompanied by an explanation that includes pre- and post-edit data descriptions. The reader is referred to Tessler & Gregson (1997) for a complete description of prescribed data editing procedures and an example edit log.

Prior to any major changes of a data set a copy is stored with the appropriate version number. This allows for the tracking of changes over time. With proper controls and communication, versioning ensures that only the most current version is used in any analysis. Versioning of archived data sets is handled by adding a three digit number to the file name, with the first version being numbered 001. Each additional version is assigned a sequentially higher number. Frequent users of the data are notified of the updates, and provided with a copy of the most recent archived version.

Once the data are archived, any changes made to the data must be documented in an edit log. At this point forward, original field forms are not altered. Field forms can be reconciled to the database through the use of the edit log. Secure data archiving is essential for protecting data files from corruption. Once a data set has passed all QA/QC procedures specified in the protocol, a formal entry is made in the I&M Data Set Catalog. Subsequently, an electronic version of the data set is maintained in a read-only format on the program server. Backup copies of the data are maintained at the Wilson's Creek visitor center, and an additional digital copy is forwarded to the NPS Inventory and Monitoring Program Archive. Tape backups of the bird project databases are made twice weekly. Two backup copies are maintained at the PC-LTEM building and two additional copies at the Wilson's Creek National Battlefield Visitor Center, Republic Missouri. Once-a-month, one tape copy is placed into permanent archive.

Currently, data are available for research and management applications on request, for database versions where all QA/QC has been completed and the data have been archived. Data can also be transferred using e-mail (most Access files are smaller than 10 Mbytes or can be compressed into .zip files) or ftp where available. As mentioned in the introduction, the plan is to make grassland bird monitoring data available for download directly from the NPS I&M Monitoring webpage. In addition, metadata will become available directly from the NPS I&M metadata server. Data requests should be directed to:

Gareth Rowell, Data Manager
Wilson's Creek National Battlefield
6424 W. Farm Road 182
Republic, MO 65738-9514
gareth_rowell@nps.gov

V. Analysis and Reporting

A critical component of any long-term monitoring protocol is a consistent and systematic way of analyzing and reporting on information (data) collected. That being said, it is also important that the data analysis and reporting requirements be flexible enough to accommodate new and varied analytical methods and reporting styles as they become available. Data visualization, geostatistics and time-series analysis are but three analysis methods that may be employed with this protocol as the size and complexity of the database increases through time.

SOP # 8 “Data Analysis” gives step-by-step details on how to: 1) determine species abundance, bird species richness, diversity and evenness, 2) estimate abundance of common species, adjusted for detectability using Distance 4.0 analysis, 3) calculate descriptive statistics for habitat variables and 4) examine the relationships between bird species abundance and habitat conditions by way of correlation analysis. All calculations are made at the plot level so means and standard errors can be calculated by habitats, across habitats or average for park wide inferences. These parameters should be analyzed each time a survey is completed.

The proposed sampling design at Tallgrass Prairie National Preserve allows for a wide variety of powerful analyses while restricting annual field efforts to manageable amounts of time. We will be able to make design-based estimates of total abundance (or density) in stratum annually and at the end of year 5 (and every 5 years thereafter). We will also be able to construct park-wide design-based estimates annually and after every 5th year. The park-wide estimates will involve weighting the 2 strata by their areas, so the riparian corridor will be given little weight. Because of the annual visits (i.e., the [1-0] panel), we will also be able to use a linear model (repeated measures) to estimate trend and compare abundance (or density) between pastures. For this, a univariate repeated measures analysis that treats the pastures as “treatments” in on the “main plots”, and points within pastures as the “subjects”. In this repeated measures analysis, much of the power for comparing pastures will come from the [1-0] panel, but points in the [1-4] split will also contribute. This analysis is very flexible and could not only estimate trends and compare pastures, but could also estimate variance components, temporal variance, and spatial variance. For pure comparative purposes, equal numbers of points in each pasture is convenient, but given the differences in pasture sizes and other objectives, the proportional allocation of our grid sample is appropriate.

Bird Data Analysis

Species richness is defined as the total number of bird species observed on a plot. Community diversity, calculated using the Shannon Diversity Index (Shannon, 1949), measures the equitability of species across samples. The disparity among species abundance is calculated using Pielou’s “J” (Pielou, 1969). To compensate for an observers inability to detect all individuals present during a survey, Distance 4.0 is used to obtain more accurate and robust estimates of population sizes for common species (Buckland et al., 2001). Relative occurrence (individual / plot visit) of each bird species is compared between surveys each time a new one is conducted to determine changes in the status of a population.

When using Distance 4.0 to estimate population sizes for species of greatest importance to the parks, the formula recommended by Buckland et al. (1993) may require additional plots be sampled within a park. Ideally, 60-80 detections per observer are needed to compute a distance function for each species (Buckland et al. 1993). However, it is possible to pool data from multiple surveys over a number of years to obtain adequate sample sizes for development of these detection functions, and it is not necessary that all distance data be collected in a single survey. Also, if the detectability of a particular species is high and factors affecting detection rate are not complex, it is possible to obtain adequate results with fewer than 60 detections.

Habitat Data Analysis

Summary reports for habitat attributes at the 50 m plot level are required for “permanent” and “semi-permanent” features. For permanent features, the values for slope, aspect and topographic position are measured, reported and assigned to a permanent locations table within

the database only once. Semi-permanent plot features, those not expected to change much including percent cover of upland prairie, riparian woodland, road cover and water cover are recorded each time a survey is conducted, with these changes reported.

The bulk of habitat data is collected at the 5 m subplot level with the first subplot located at plot center and three remaining ones located at a distance of 50 m from center plot and 120° offset from one another. Permanent subplot features include azimuth from plot center to a subplot as well as slope and aspect of each subplot. Like permanent attributes measured on plots, permanent features for subplots are measured only once and stored in a table within the database. Subplot features that are reported on each time a survey is conducted include tree tallies, canopy height, canopy coverage, basal area, ground and foliar coverage, horizontal vegetation coverage and structural diversity. Values reported for each subplot feature are either averaged or used in calculations to establish single values representing each plot, see SOP # 8 “Data Analysis” for details.

Reporting

In an effort to disseminate findings in a timely manner, reports should be completed by March 31 of the year following data collection. More extensive summary reports should be completed every five to ten years depending on how fast habitat conditions are changing and how critical summary information is to setting management goals influencing bird populations within a park. Summary reports may be used in place of annual reports for the year in which the last data is collected. Refer to SOP # 9 “Reporting” for details on report structure and style. Also see, “Status Report 2001-2003: Grassland Bird Monitoring at Agate Fossil Beds National Monument, Nebraska and Tallgrass Prairie National Preserve, Kansas” (Peitz and Rowell, draft) for content included in a report and reporting style.

Tables in a report should be placed within the text or immediately following the literature cited section (see Appendix B). Tables should be numbered in sequence regardless of where they are located. Table headers are placed at the top of a table. Horizontal lines are used to separate the table heading from column headings, column headings from the table and to signify the end of the table. Vertical lines should not appear on a table.

Figures (including pictures) should be placed within the text of a report or immediately following tables behind the literature cited sections (see Figure 6.01.1. Spatial arrangement of vegetation subplots). Figures should be numbered in sequence regardless of where they are located. Figure captions are placed below the figure if it is included in the text or on a separate sheet of paper preceding the figure if included after the literature cited section. Both tables and figures should contain information not presented in the body of the text. Also, tables and figures should not duplicate information already presented in the body of the text.

VI. Personnel Requirements and Training

Roles and Responsibilities

The project manager is the lead ecologist for implementing this monitoring protocol, and is supervised by the Program Coordinator for the PC-LTEM Program. Because of the need for a high level of training and consistency in implementing the protocol, the project manager will usually be the person conducting the VCP counts in the field. The data management aspect of the monitoring effort is the shared responsibility of the project manager and the data manager. Typically, the project manager is responsible for data collection, data entry, data verification and

validation, as well as data summary, analysis and reporting. The data manager is responsible for data archiving, data security, dissemination and database design. The data manager, in collaboration with the project manager, also develops data entry forms and other database features as part of quality assurance and automates report generation. The data manager is ultimately responsible that adequate QA/QC procedures are built into the database management system and appropriate data handling procedures followed.

Qualifications and Training

The most essential component for the collection of credible, high-quality data on birds is a competent observer. This cannot be overemphasized. Various studies have shown that observer bias is one of the most noteworthy bias factors in trend analysis of songbird populations (Scott et al. 1986, Sauer et al. 1994, Kendall et al. 1996).

As well as being able to visually identify birds, field observers must be proficient at identifying species by their songs and calls. Therefore, recordings of birds in the study area, especially for the less common or unexpected species should be provided for surveyors. Observers should be tested frequently on their ability to identify bird calls. Good hearing ability is essential because many birds, particularly in forested habitats, are detected by sound only. Differences in hearing ability between observers may strongly affect survey results (Ramsey and Scott 1981).

The quality of the observer will determine the quality of the data. Time should also be invested in training personnel to estimate the distance from themselves to singing birds within different habitats. This will require training in the field (see SOP #2, "Training Observers") to become proficient with the use of a laser rangefinder and to gain experience with estimating distances to different species in different habitat types.

The project manager and a staff botanist will assist in training the bird crew to conduct habitat monitoring. Observers doing habitat work should familiarize themselves with the PC-LTEM standard cover classes and vegetation guilds described in SOP #6. Prior to the field season, observers should practice estimating cover of different vegetation guilds and ground cover categories within 1.78-m practice plots. If habitat monitoring cannot be practiced prior to the field season, a staff botanist will accompany the bird crew on the first field trip of the year.

VII. Operational Requirements

Annual Workload and Field Schedule

Breeding bird surveys will begin no sooner than the first full week of May and extend no later than the second full week of June, a period that coincides with the peak-breeding activity of most birds. Inclement weather and personnel workloads will preclude the scheduling of sampling events to specific annual dates. Bird surveys may be scheduled for the third week of June if spring green-up is unusually late. Habitat monitoring should be completed no later than June 30, in order for habitat measures to accurately reflect conditions experienced by breeding birds. Monitoring efforts will require a three to four person crew. Approximately five to seven field days are required to complete bird sampling at Agate Fossil Beds National Monument, and 10 – 12 days at Tallgrass Prairie National Preserve. Twelve to fifteen VCP counts should be scheduled for completion each field day.

Facility and Equipment Needs

The nature of bird survey work does not require special facilities beyond normal office space and equipment storage needs. Table 1.01.1 is a list of field equipment needs for one crew. If two or more crews work simultaneously, equipment requirements will increase accordingly.

Startup Costs and Budget

Personnel expenses for field work are based on a crew of three to four people: an ecologist or biologist to conduct the bird counts, a botanist to initially work with or train the crew to conduct habitat sampling, and two seasonal biological science technicians. Five to seven field days should be planned for Agate Fossil Beds National Monument and approximately 10 – 12 days for Tallgrass Prairie National Preserve. Field costs will vary somewhat from year to year depending on the skill level and size of the crew. Data management personnel expenses include staff time of biological science technicians, the project manager and data manager. Startup cost for equipment include the purchase of equipment and supplies listed in Table 1.01.1, as well as maintenance and or replacement of equipment shared among multiple projects (e.g. GPS units, cameras).

Estimated Costs	TAPR	AGFO
Field Work (Salary)	10,500	6,000
Field Work (Travel)	3,200	1,200
Data Management/Reporting (Salary)	8,100	2,600
Startup Equipment Costs	1,200	1,200
Annual equipment/supplies	1,400	1,000
Total	24,400	12,000

Procedure for Revising the Protocol

Over time, revisions to both the Protocol Narrative and to specific Standard Operating Procedures (SOPs) are to be expected. Careful documentation of changes to the protocol and a library of previous protocol versions are essential for maintaining consistency in data collection and for appropriate treatment of the data during data summary and analysis. The MicroSoft Access® database for each monitoring component contains a field that identifies which version of the protocol was being used when the data were collected.

The rationale for dividing a sampling protocol into a Protocol Narrative with supporting SOPs is based on the following:

- The Protocol Narrative is a general overview of the protocol that gives the history and justification for doing the work and an overview of the sampling methods, but that does not provide all of the methodological details. The Protocol Narrative will only be revised if major changes are made to the protocol.
- The SOPs, in contrast, are very specific step-by-step instructions for performing a given task. They are expected to be revised more frequently than the protocol narrative.
- When a SOP is revised, in most cases, it is not necessary to revise the Protocol Narrative to reflect the specific changes made to the SOP.
- All versions of the Protocol Narrative and SOPs will be archived in a Protocol Library.

The steps for changing the protocol (either the Protocol Narrative or the SOPs) are outlined in SOP #11, “Revising the Protocol.” Each SOP contains a Revision History Log that should be filled out each time a SOP is revised to explain why the change was made, and to

assign a new Version Number to the revised SOP. The new version of the SOP and/or Protocol Narrative should then be archived in the PC-LTEM Protocol Library under the appropriate folder.

VIII. References

- Barker, R. J., and J. R. Sauer. 1995. Statistical aspects of point count sampling. Pages 125-130 in C. J. Ralph, J. R. Sauer, and S. Droege, eds. *Monitoring Bird Populations by Point Counts*, USDA Forest Service, Pacific Southwest Research Station, General Technical Report PSW-GTR-149.
- BCRIB (British Columbia Resource Inventory Branch). 1999. Inventory methods for forest and grassland songbirds: standards for components of British Columbia's biodiversity No. 15. Resource Inventory Branch, British Columbia Ministry of Environment, Lands and Parks. Available from <http://www.for.gov.bc.ca/RIC/Pubs/teBioDiv/>
- Buckland, S. T., D. R. Anderson, K. P. Burnham, and J. L. Laake. 1993. *Distance sampling: Estimating abundance of biological populations*. Chapman and Hall, New York. 446 pp.
- Buckland, S. T., D. R. Anderson, K. P. Burnham, J. L. Laake, D. L. Borchers, and L. Thomas. 2001. *Introduction to distance sampling: estimating abundance of biological populations*. Oxford University Press. 432 pp.
- Burnham, K. P. 1981. Summarizing remarks: environmental influences. *Studies in Avian Biology* 6:324-325.
- Cody, M. L. 1981. Habitat selection in birds: the roles of vegetation structure, competitors, and productivity. *BioScience* 31:107-111.
- Cody, M. L. 1985. *Habitat selection in birds*. Academic Press, New York, New York.
- DeBacker, M.D., B. Witcher, and J.R. Boetsch. 2002. *Prairie Cluster Prototype Long-term Ecological Monitoring Program. Data Management Plan*. Prairie Cluster Prototype LTEM Program, National Park Service. Wilson's Creek National Battlefield. Republic, Missouri. 26pp.
- Fancy, S. G. 1997. A new approach for analyzing bird densities from variable circular-plot counts. *Pacific Science* 51:107-114.
- Fitzgerald, J. A. and D. N. Pashley. 2000. *Partners in Flight bird conservation plan for the Ozark/Ouachitas (physiographic area 19)*. Partners in Flight Midwest Region, Brentwood, Missouri. 78pp.
- Glenn, S.M., S.L. Collins, and D.J. Gibson. 1992. Disturbance in tallgrass prairie: local versus regional effects on community heterogeneity. *Landscape Ecology* 7:243-252.
- Graetz J. L., R. A. Garrott, and S. R. Craven. 1995. *Faunal survey of Agate Fossil Beds National Monument*. Report to Midwest Region, National Park Service, U.S. Department of the Interior, Omaha, Nebraska. 54 p.
- Karr, J. R. and D. R. Dudley. 1981. Ecological perspective on water quality goals. *Environmental Management* 5:55-68.
- Karr, J. R. 1991. Biological integrity: a long-neglected aspect of water resource management. *Ecological Applications* 1:66-84.
- Kendall, W. L., B. G. Peterjohn, and J. R. Sauer. 1996. First-time observer effects in the North American Breeding Bird Survey. *Auk* 113:823-829.
- Kepler, C. B. and J. M. Scott. 1981. Reducing bird count variability by training observers. *Studies in Avian Biology* 6:366-371.

- Knopf, F.L. and F.B. Samson. 1997. Conservation of grassland vertebrates. Pages 273-289 in F.L. Knopf and F.B. Samson, eds. *Ecology and Conservation of Great Plains Vertebrates*. Springer-Verlag, New York, New York.
- Lane, D. L. 1995. Above-ground net primary production across a precipitation gradient in the central grassland region. M.S. thesis, Colorado State University, Fort Collins.
- Lichtenberg, J. S. and A. N. Powell. 1999. Avian inventory of Tallgrass Prairie National Preserve, Kansas. Report to Tallgrass Prairie National Preserve, National Park Service, U.S. Department of the Interior, Strong City, Kansas. 38 p.
- Maurer, B.A. 1993. Biological diversity, ecological integrity, and neotropical migrants: New perspectives for wildlife managers. Pages 24-31 in D.M. Finch and P.W. Stangel, editors. *Status and management of neotropical migratory birds*. U.S. Forest Service General Technical Report RM-229.
- McDonald, Trent L. 2003. Review of environmental monitoring methods: survey designs. *Environmental Monitoring and Assessment* 85: 277-292.
- Nelson, J. T. and S. G. Fancy. 1999. A test of the variable circular-plot method when exact density of a bird population was known. *Pacific Conservation Biology* 5:139-143.
- Peitz, D.G. and G.A. Rowell. Draft. Status Report 2001-2003: Grassland Bird Monitoring at Agate Fossil Beds National Monument, Nebraska and Tallgrass Prairie National Preserve, Kansas. Prairie Cluster Prototype Long-term Ecological Monitoring Program, National Park Service, U.S. Department of the Interior, Republic, Missouri
- Powell, A. N. 2000. Grassland bird inventory of seven prairie parks. Final report to the Great Plains Prairie Cluster Long-term Ecological Monitoring Program, Wilson's Creek National Battlefield, National Park Service, U.S. Department of the Interior, Republic, Missouri. 47 p.
- Pielou, E.C. 1969. *An introduction to mathematical ecology*. John Wiley and Sons, New York, New York. 286pp.
- Ralph, C. J., S. Droege, and J. R. Sauer. 1995. Managing and monitoring birds using point counts: standards and applications. Pages 161-168 in C. J. Ralph, J. R. Sauer, and S. Droege, eds. *Monitoring Bird Populations by Point Counts*, USDA Forest Service, Pacific Southwest Research Station, General Technical Report PSW-GTR-149.
- Ramsey, F. L. and J. M. Scott. 1981. Tests of hearing ability. *Studies in Avian Biology* 6:341-345.
- Reynolds, R. T., J. M. Scott, and R. A. Nussbaum. 1980. A variable circular-plot method for estimating bird numbers. *Condor* 82:309-313.
- Sauer, J. R., B. G. Peterjohn, and W. A. Link. 1994. Observer differences in the North American Breeding Bird Survey. *Auk* 111:50-62.
- Sauer, J. R., J. E. Hines, I. Thomas, J. Fallon, and G. Gough. 2000. The North American breeding bird survey, results and analysis 1966 – 1999. Version 98.1, USGS Patuxent Wildlife Research Center, Laurel, Maryland. Available at <http://www.mbr-pwrc.usgs.gov/bbs/bbs.html>
- Sauer, J. R., J. E. Hines, and J. Fallon. 2003. The North American breeding bird survey, results and analysis 1966 - 2002. Version 2003.1, USGS Patuxent Wildlife Research Center, Laurel, Maryland. Available at <http://www.mbr-pwrc.usgs.gov/bbs/bbs.html>
- Scott, J. M., S. Mountainspring, F. L. Ramsey, and C. B. Kepler. 1986. Forest bird communities of the Hawaiian Islands: Their dynamics, ecology and conservation. *Studies in Avian Biology* 9:1-431.

- Shannon, C.E. 1949. The mathematical theory of communication. University of Illinois Press, Urbana, Illinois. 177 pp.
- Tappe, P. A., R. E. Thill, M. A. Melchoirs, and T. B. Wigley. (in press). Breeding bird community composition and structure in four watersheds under different management scenarios in the Ouachita Mountains of Arkansas. *in* J.M. Guldin, ed. Symposium on ecosystem management in the Ouachita and Ozark Mountains. USDA Forest Service General Technical Report, Southern Forest Experiment Station, Ashville, North Carolina.
- Tessler, S. and J. Gregson. 1997. Draft Data Management Protocol. National Park Service. Document available at: www.nature.nps.gov/im/dmproto/joe40001.htm.
- Vinton, M. A. and S. L. Collins. 1997. Landscape gradients and habitat structure in native grasslands of the central Great Plains. Pages 3 – 19 *in* F.L Knopf and F.B. Samson, eds. Ecology and Conservation of Great Plains Vertebrates, Springer-Verlag New York Inc., New York, New York.
- Zimmerman, J. L. 1982. Nesting success of dickcissels (*Spiza americana*) in preferred and less preferred habitats. Auk 99:292-298.
- Zimmerman, J. L. 1988. Breeding season habitat selection by the Henslow's sparrow (*Ammodramus henslowii*) in Kansas. Wilson Bulletin 100:17-24.
- Zimmerman, J. L. 1993. The birds of Konza: avian ecology of the tallgrass prairie. University Press of Kansas. 176 pp.
- Zimmerman, J. L. 1997. Avian community responses to fire, grazing and drought in tallgrass prairie. Pages 167-180 *in* F.L Knopf and F.B. Samson, eds. Ecology and Conservation of Great Plains Vertebrates, Springer-Verlag New York Inc., New York, New York.

Appendix A. Field data forms for recording (a) bird detections, (b) habitat variables, and (c) location data for new sampling stations. These forms should be copied onto write-in-the-rain paper and taken into the field. The database has data entry screens that mimic these forms.

- (a) Field Data Form – Variable Circular Plot Counts
Use one of these forms each time that a 5 minute count is conducted.
- (b) Habitat Data Form
Use one of these forms once each year at each plot that is sampled.
- (c) New Location Data Form
This form should be filled out each time that a new plot location is established. The data from this form is entered into the Locations table of the database.

[illegible]

Variable Circular Plot Count Data Form: Categories, Definitions and Descriptions.

Codes (Beaufort scale) used to record wind strength during bird counts.

Wind Code	Explanation
0	Calm, smoke rises vertically (< 2 km/h)
1	Smoke drifts (2-5 km/h)
2	Light breeze felt on face, leaves rustle (6-12 km/h)
3	Leaves and twigs in constant motion (13-19 km/h)
4	Small branches move, raises loose paper, dust rises (20-29 km/h)
5	Fresh breeze, small trees sway (30-39 km/h)
6	Strong breeze, large branches moving, wind whistling (40-50 km/h)

Codes used to record precipitation codes during bird counts.

Rain Code	Explanation
0	No rain
1	Mist or fog
2	Light drizzle
3	Light rain
4	Heavy rain; difficult to hear birds
5	Snow

Codes used to record level of background noise as it effects observer's ability to hear birds.

Noise Code	Explanation
0	quiet; normal background noises; no interference
1	low noise; might be missing some high-pitched songs/calls of distant birds
2	medium noise; detection radius is probably substantially reduced
3	high noise; probably detecting only the loudest/closest birds

Codes used to record the type of detection (Detection Type, DT) of birds during a count.

DT Code	Explanation
1	heard first, but not seen (i.e., detected initially by sound) during the 5-min count
2	seen first (regardless of whether it was later heard or not) during the 5-min count
4	heard first, but then seen (a DT of 1 can be changed to a 4) during the 5-min count
8	heard between plots or not within the 5-minute sampling period
9	seen between plots or not within the 5-minute sampling period

Field Data Form - PRAIRIE BIRD HABITAT MONITORING DATA FORM

Park Code: _____ Date (mm/dd/yyyy) _____ Observer(s) Initials: _____

Management Unit: _____ Plot: _____ Riparian Corridor: Yes No

PLOT ATTRIBUTES: (50m radius)

Slope (°): _____ Var: high med low Aspect (°): _____ Var: high med low

Topographic position:

level lower-slope mid-slope upper-slope escarpment/face ledge crest depression draw

Vegetation Type:	Cover Class:	Vegetation Type:	Cover Class:	Road Cover:	Cover Class:
Upland Prairie		Other: _____		Paved	
Riparian Woodland (TAPR)		Other: _____		Pasture	
Other: _____				Water Cover:	
Other: _____				Stream	
Other: _____				Pond	

Plot Notes: _____

SUBPLOT: _____ Random Azimuth (°) _____ Slope(°): _____ Aspect (°) _____

TREE TALLY (by dbh size class)

Species	<1.0 cm	1.1 – 2.5 cm	2.6 – 8.0 cm	8.1 – 15.0 cm	15.1 – 23.0 cm	23.1 – 38.0 cm	>38.0 cm

CANOPY HEIGHT: (m) CANOPY COVER: (dots covered out of 96) BASAL AREA: (cruz-all tree count -- use 10 factor)

Hardwood: _____ Hardwood: 1__ 2__ 3__ 4__ Hardwood: _____

Conifer: _____ Conifer: 1__ 2__ 3__ 4__ Conifer: _____

Total: 1__ 2__ 3__ 4__

GROUND COVER:	Cover Class:	FOLIAR COVER (by guilds):	Cover Class:
Deciduous Litter		Warm Season Grass	
Conifer Litter		Cool Season Grass	
Grass Litter		Forbs	
Bare Soil		Moss and Lichen	
Rock		Shrubs & Vines	
Woody Debris		Tree Seedlings	
Unvegetated		Total Foliar Cover (< 1.5 m tall)	

HORIZONTAL VEGETATION PROFILE BOARD:

5-m distance: 0.50 m: _____ 0.75 m: _____ 1.0 m: _____ 1.25 m: _____ 1.5 m: _____ 1.75 m: _____ 2.0 m: _____

15-m distance: 0.50 m: _____ 0.75 m: _____ 1.0 m: _____ 1.25 m: _____ 1.5 m: _____ 1.75 m: _____ 2.0 m: _____

VEGETATION PROFILE (D = Deciduous, H = Herbaceous, C = Conifer)

0–1 m	1–2 m	2–3 m	3–4 m	4–5 m	5–6 m	6–7 m	7–7.5 m

Subplot Notes: _____

Prairie Bird Habitat Data Form: Categories, Definitions and Descriptions.

Vegetation Type

TAPR		AGFO	
	Prairie		Upland prairie
	Riparian woodland		Upland disturbance
	Brome fields		Western wheatgrass prairie (river terraces)
	Upland disturbance		Annual-dominated floodplain disturbance
			Willow shrubland
			Wetland (cattail-iris marsh and rush-sedge wetland)

Cover Class Definitions and Cut-point Descriptions

Cover class	Definition	Upper cutpoints for 50 m radius plot	Upper cutpoints for 5 m radius plot	Upper cutpoints for 1.78 m radius plot
0	None present			
1	Trace to 1 %	A circle of radius 5.00 m	A circle of radius 0.50 m	A circle of radius 0.18 m
2	1 – 5 %	A circle of radius 11.18 m	A circle of radius 1.12 m	A circle of radius 0.40 m
3	5 – 25 %	A circle of radius 25.00 m	A circle of radius 2.50 m	A circle of radius 0.89 m
4	25 – 50 %	A circle of radius 35.36 m	A circle of radius 3.54 m	A circle of radius 1.26 m
5	50 – 75 %	A circle of radius 43.30 m	A circle of radius 4.33 m	A circle of radius 1.54 m
6	75 – 95 %	A circle of radius 48.73 m	A circle of radius 4.87 m	A circle of radius 1.73 m
7	95 – 100 %	Full plot radius	Full plot radius	Full plot radius

New Location Data Form (use this form to record coordinates for new sampling locations)

Park: _____ Project: _____ LocationID: _____

Description: _____

UTM Coordinates:**Ending Coordinates for Transects:**

UTMX: _____

UTM Zone: _____

UTMX end: _____

UTMY: _____

Error (m): _____

UTMY end: _____

Latitude: _____

Datum: NAD83

Latitude end: _____

Longitude: _____

Longitude end: _____

Elevation: _____

Vertical Error (m): _____

Aspect: N NE E SE S SW W NW

How was accuracy determined? _____

Slope (deg): _____

New Location Data Form (use this form to record coordinates for new sampling locations)

Park: _____ Project: _____ LocationID: _____

Description: _____

UTM Coordinates:**Ending Coordinates for Transects:**

UTMX: _____

UTM Zone: _____

UTMX end: _____

UTMY: _____

Error (m): _____

UTMY end: _____

Latitude: _____

Datum: NAD83

Latitude end: _____

Longitude: _____

Longitude end: _____

Elevation: _____

Vertical Error (m): _____

Aspect: N NE E SE S SW W NW

How was accuracy determined? _____

Slope (deg): _____

Appendix B. List of avian species recorded during breeding seasons (May-June) at Agate Fossil Beds National Monument, Nebraska and Tallgrass Prairie National Preserve, Kansas. Sources of data includes Graetz et al. 1995 (a), Powell 2000 (b), and Lichtenberg and Powell 1999 (c).

Common name / Code	Scientific name	TSN	AGFO		TAPR
Great Blue Heron / GBHE	<i>Ardea herodias</i>	174773	Xa	Xb	Xc
Green (Green-backed) Heron / GRHE	<i>Butorides virescens</i>	174793			X
Turkey Vulture / TUVU	<i>Cathartes aura</i>	175265	X		X
Canada Goose / CAGO	<i>Branta canadensis</i>	174999			X
Wood Duck / WODU	<i>Aix sponsa</i>	175122	X	X	X
Mallard / MALL	<i>Anas platyrhynchos</i>	175063	X	X	X
Blue-winged Teal / BWTE	<i>Anas discors</i>	175086	X		
Northern Harrier / NOHA	<i>Circus cyaneus</i>	175430	X		
Red-tailed Hawk / RTHA	<i>Buteo jamaicensis</i>	175350	X	X	X
Swainson's Hawk / SWHA	<i>Buteo swainsoni</i>	175367	X		X
Golden Eagle / GOEA	<i>Aquila chrysaetos</i>	175407	X		
Prairie Falcon / PRFA	<i>Falco mexicanus</i>	175603	X		
American Kestrel / AMKE	<i>Falco sparverius</i>	175622	X	X	X
Wild Turkey / WITU	<i>Meleagris gallopavo</i>	176136	X	X	X
Sharp-tailed Grouse / STGR	<i>Tympanuchus phasianellus</i>	175841	X		
Greater Prairie-chicken / GRCH	<i>Tympanuchus cupido</i>	175834		X	X
Northern Bobwhite / NOBO	<i>Colinus virginianus</i>	175863		X	X
Ring-necked Pheasant / RPHE	<i>Phasianus colchicus</i>	175905	X	X	
Virginia Rail / VIRA	<i>Rallus limicola</i>	176221	X		
Sora / SORA	<i>Porzana Carolina</i>	176242	X		
American Avocet / AMAV	<i>Recurvirostra americana</i>	176721	X		
Killdeer / KILL	<i>Charadrius vociferus</i>	176520	X	X	X
Upland Sandpiper / UPSA	<i>Bartramia longicauda</i>	176610	X	X	X
Wilson's Phalarope / WIPH	<i>Phalaropus tricolor</i>	176736	X		
Common Snipe / COSN	<i>Gallinago gallinago</i>	176700	X		
Black Tern / BLTE	<i>Chlidonias niger</i>	176959	X		
Rock Dove / RODO	<i>Columba livia</i>	177071		X	X
Mourning Dove / MODO	<i>Zenaidura macroura</i>	177125	X	X	X
Black-billed Cuckoo / BBCU	<i>Coccyzus erythrophthalmus</i>	177834			X
Yellow-billed Cuckoo / YBCU	<i>Coccyzus americanus</i>	177831			X
Great Horned Owl / GHOW	<i>Bubo virginianus</i>	177884	X		X
Short-eared Owl / SEOW	<i>Asio flammeus</i>	177935		X	
Barn Owl / BNOW	<i>Tyto alba</i>	177851	X		
Common Nighthawk / CONI	<i>Chordeiles minor</i>	177979	X	X	X
Chimney Swift / CHSW	<i>Chaetura pelagica</i>	178001			X
Ruby-throated Hummingbird / RTHU	<i>Archilochus colubris</i>	178032			X
Belted Kingfisher / BEKI	<i>Ceryle alcyon</i>	178119	X		X
Northern Flicker / NOFL	<i>Colaptes auratus</i>	178154		X	X
Pileated Woodpecker / PIWO	<i>Dryocopus pileatus</i>	178166			X
Red-headed Woodpecker / RHWO	<i>Melanerpes erythrocephalus</i>	178186			X
Red-bellied Woodpecker / RBWO	<i>Melanerpes carolinus</i>	178195			X
Downy Woodpecker / DOWO	<i>Picoides pubescens</i>	178259	X		X
Hairy Woodpecker / HAWO	<i>Picoides villosus</i>	178262			X
Eastern Wood-pewee / EAWP	<i>Contopus virens</i>	178359			X
Western Wood-pewee / WEWP	<i>Contopus sordidulus</i>	178360	X		
Yellow-bellied Flycatcher / YBFL	<i>Empidonax flaviventris</i>	178338			X
Acadian Flycatcher / ACFL	<i>Empidonax virescens</i>	178339			X
Great Crested Flycatcher / GCFL	<i>Myiarchus crinitus</i>	178309			X
Scissor-tailed Flycatcher / STFL	<i>Tyrannus forficatus</i>	178293			X
Eastern Phoebe / EAPH	<i>Sayornis phoebe</i>	178329			X
Say's Phoebe / SAPH	<i>Sayornis saya</i>	178333	X	X	

Appendix B. List of avian species, cont'd.

Common name / Code	Scientific name	TSN	AGFO		TAPR
Cassin's Kingbird / CAKI	Tyrannus vociferans	178288		X	
Western Kingbird / WEKI	Tyrannus verticalis	178287	X	X	X
Eastern Kingbird / EAKI	Tyrannus tyrannus	178279	X	X	X
Loggerhead Shrike / LOSH	Lanius ludovicianus	178515	X	X	X
White-eyed Vireo / WEVI	Vireo griseus	178991			X
Yellow-throated Vireo / YTVI	Vireo flavifrons	179009			X
Warbling Vireo / WAVI	Vireo gilvus	179023			X
Red-eyed Vireo / REVI	Vireo olivaceus	179021			X
Blue Jay / BLJA	Cyanocitta cristata	179680			X
American Crow / AMCR	Corvus brachyrhynchos	179731	X	X	X
Horned Lark / HOLA	Eremophila alpestris	554256	X	X	X
Northern Rough-winged Swallow / NRWS	Stelgidopteryx serripennis	178443	X	X	X
Bank Swallow / BANS	Riparia riparia	178436	X	X	X
Cliff Swallow / CLSW	Hirundo pyrrhonota	178453	X	X	
Barn Swallow / BARS	Hirundo rustica	178448	X	X	X
Black-capped Chickadee / BCCH	Parus atricapillus	178699			X
(Eastern) Tufted Titmouse / ETTI	Parus bicolor	178738			X
White-breasted Nuthatch / WBNU	Sitta carolinensis	178775			X
House Wren / HOWR	Troglodytes aedon	178541	X		X
Bewick's Wren / BEWR	Thryomanes bewickii	178562			X
Carolina Wren / CARW	Thryothorus ludovicianus	178581			X
Rock Wren / ROWR	Salpinctes obsoletus	178614	X	X	
Marsh Wren / MAWR	Cistothorus palustris	178608	X		
Blue-gray Gnatcatcher / BGGN	Poliophtila caerulea	179853			X
American Robin / AMRO	Turdus migratorius	179759	X		X
Swainson's Thrush / SWTH	Catharus ustulatus	179788			X
Eastern Bluebird / EABL	Sialia sialis	179801			X
Mountain Bluebird / MOBL	Sialia currucoides	179811	X		
Gray Catbird / GRCA	Dumetella carolinensis	178625	X		X
Northern Mockingbird / NOMO	Mimus polyglottos	178620			X
Brown Thrasher / BRTH	Toxostoma rufum	178627	X	X	X
European Starling / EUST	Sturnus vulgaris	179637		X	X
Tennessee Warbler / TEWA	Vermivora peregrina	178855			X
Northern Parula / NOPA	Parula americana	178868			X
Yellow Warbler / YWAR	Dendroica petechia	178878	X		X
Common Yellowthroat / COYE	Geothlypis trichas	178944	X	X	X
Yellow-breasted Chat / YBCH	Icteria virens	178964			X
Summer Tanager / SUTA	Piranga rubra	179888			X
Eastern (Rufous-sided) Towhee / EATO	Pipilo erythrophthalmus	179276			X
Spotted (Rufous-sided) Towhee / SPTO	Pipilo maculatus	554380	X	X	
Chipping Sparrow / CHSP	Spizella passerina	179435			X
Field Sparrow / FISP	Spizella pusilla	179443			X
Clay-colored Sparrow / CCSP	Spizella pallida	179439		X	
Vesper Sparrow / VESP	Poocetes gramineus	179366		X	X
Lark Sparrow / LASP	Chondestes grammacus	179371	X	X	X
Lark Bunting / LARB	Calamospiza melanocorys	179312	X	X	
Grasshopper Sparrow / GRSP	Ammodramus savannarum	179333	X	X	X
Chestnut-collared Longspur / CCLO	Calcarius ornatus	179530		X	
Northern Cardinal / NOCA	Cardinalis cardinalis	179124			X
Rose-breasted Grosbeak / RBGR	Pheucticus ludovicianus	179139			X
Black-headed Grosbeak / BHGR	Pheucticus melanocephalus	179140		X	
Blue Grosbeak / BLGR	Guiraca caerulea	179145	X	X	X
Indigo Bunting / INBU	Passerina cyanea	179150			X
Lazuli Bunting / LAZB	Passerina amoena	179151	X	X	
Dickcissel / DICK	Spiza americana	179165	X		X
Bobolink / BOBO	Dolichonyx oryzivorus	179032	X		
Eastern Meadowlark / EAME	Sturnella magna	179034			X
Western Meadowlark / WEME	Sturnella neglecta	179039	X		X

Appendix B. List of avian species, cont'd.

Common name / Code	Scientific name	TSN	AGFO	TAPR
Yellow-headed Blackbird / YHBL	Xanthocephalus xanthocephalus	179043	X	
Red-winged Blackbird / RWBL	Agelaius phoeniceus	179045	X	X
Great-tailed Grackle / GTGR	Quiscalus mexicanus	179109		X
Common Grackle / COGR	Quiscalus quiscula	179104	X	X
Brown-headed Cowbird / BHCO	Molothrus ater	179112	X	X
Orchard Oriole / OROR	Icterus spurius	179064	X	X
Baltimore (Northern) Oriole / BAOR	Icterus galbula	179083		X
American Goldfinch / AMGO	Carduelis tristis	179236	X	X
House Sparrow / HOSP	Passer domesticus	179628		X

Appendix C. Plant species and community guilds by park.

Agate Fossil Beds National Monument, Nebraska.

Scientific Name	Common Name	Plant Community Guild
<i>Allium textile</i>	Onion	Forb
<i>Ambrosia psilostachya</i>	Western ragweed	Forb
<i>Ambrosia trifida</i>	Giant ragweed	Forb
<i>Amphiachyris dracunculoides</i>	Broomweed	Forb
<i>Andropogon hallii</i>	Sand bluestem	warm season grass
<i>Arabis holboellii</i>	Rock-cress	Forb
<i>Arenaria hookeri</i>	Sandwort	Forb
<i>Aristida purpurea</i>	Purple three-awn	warm season grass
<i>Artemisia campestris</i>	Wormwood, sage, mugwort	Forb
<i>Artemisia dracunculus</i>	Wormwood, sage, mugwort	Forb
<i>Artemisia frigida</i>	Prairie-sagewort	Forb
<i>Asclepias pumila</i>	Milkweed	Forb
<i>Asclepias speciosa</i>	Showy milkweed	Forb
<i>Asclepias viridiflora</i>	Green milkweed	Forb
<i>Aster ericoides</i>	Squarrose white wild aster	Forb
<i>Astragalus ceramicus</i>	painted milkvetch	Forb
<i>Astragalus crassicaupus</i>	Ground-plum, prairie plum	Forb
<i>Astragalus gracilis</i>	Milk-vetch	Forb
<i>Astragalus laxmannii</i> var. <i>robustior</i>	Prairie milkvetch	Forb
<i>Astragalus lotiflorus</i>	Lotus milk-vetch	Forb
<i>Astragalus missourienses</i>	Nuttall milk-vetch	Forb
<i>Astragalus sericoleucus</i>	Milk-vetch	Forb
<i>Astragalus spatulatus</i>	Milk-vetch	Forb
<i>Astragalus</i> spp		Forb
<i>Bouteloua curtipendula</i>	Side-oats grama-grass	warm season grass
<i>Bouteloua gracilis</i>	Blue grama-grass	warm season grass
<i>Bromus japonicus</i>	Japanese brome	cool season grass
<i>Bromus tectorum</i>	Downy brome	cool season grass
<i>Calamovilfa longifolia</i>	Sand-reed	warm season grass
<i>Calylophus serrulatus</i>	Evening-primrose	Forb
<i>Camelina microcarpa</i>	Small-seed false flax	Forb
<i>Carex filifolia</i>	Sedge	grass-like (cool season grass)
<i>Castilleja sessiliflora</i>	Downy paintbrush	Forb
<i>Chamaesyce serpens</i>	Round-leaved spurge	Forb
<i>Chenopodium berlandieri</i>	Pitseed goosefoot	Forb
<i>Chenopodium pratericola</i>	Narrow-leaf goosefoot	Forb
<i>Cirsium canescens</i>	Thistle	Forb
<i>Cirsium flodmanii</i>	Prairie thistle	Forb
<i>Cleome serrulata</i>	Stinking clover	Forb
<i>Comandra umbellata</i>	Bastard toad-flax	Forb
<i>Conyza canadensis</i>	Horseweed	Forb
<i>Corispermum hyssopifolium</i>	Bugseed	Forb
<i>Coryphantha vivipara</i>	Pincushion cactus	Forb
<i>Croton texensis</i>	Texas croton	Forb
<i>Cryptantha cana</i>	Borage	Forb
<i>Cryptantha celosioides</i>	Borage	Forb
<i>Cymopterus acaulis</i>	Wild parsley	Forb
<i>Dalea candida</i> var. <i>oligophylla</i>	White prairie clover	Forb
<i>Dalea purpurea</i>	Purple prairie clover	Forb

Appendix C. Agate Fossil Beds National Monument, Nebraska, cont'd.

Scientific Name	Common Name	Plant Community Guilds
<i>Descurainia pinnata</i>	Tansy-mustard	Forb
<i>Descurainia sophia</i>	Herb-sophia, tansy-mustard	Forb
<i>Ellisia nyctelea</i>	Water-pod	Forb
<i>Elymus elymoides</i>	Wild rye	cool season grass
<i>Elymus lanceolatus</i> ssp. <i>Lanceolatus</i>	Wild rye	cool season grass
<i>Equisetum laevigatum</i>	Smooth scouring rushes	Forb
<i>Erigeron bellidiastrum</i>	Fleabane	Forb
<i>Erigeron pumilus</i>	Fleabane	Forb
<i>Eriogonum annuum</i>	Annual eriogonum	Forb
<i>Eriogonum cernuum</i>	Nodding wild buckwheat	Forb
<i>Eriogonum flavum</i>	Yellow wild buckwheat	Forb
<i>Erysimum asperum</i>	Western wallflower	Forb
<i>Euphorbia glyptosperma</i>	Ridge-seeded spurge	Forb
<i>Euphorbia missurica</i>	Prairie spurge	Forb
<i>Euphorbia robusta</i>	Spurge	Forb
<i>Gaura coccinea</i>	Scarlet gaura	Forb
<i>Gaura parviflora</i>	Gaura	Forb
<i>Gutierrezia sarothrae</i>	Matchbrush	Forb
<i>Hedeoma drummondii</i>	Drummond false pennyroyal	Forb
<i>Hedeoma hispidum</i>	Mint	Forb
<i>Helianthus annuus</i>	Common sunflower	Forb
<i>Helianthus petiolaris</i>	Plains sunflower	Forb
<i>Heterotheca villosa</i>	Golden aster	Forb
<i>Hymenopappus filifolius</i> var. <i>polycephalus</i>	Aster	Forb
<i>Iva xanthifolia</i>	Big marsh-elder	Forb
<i>Kochia scoparia</i>	Summer-cypress	Forb
<i>Koeleria macrantha</i>	Junegrass	cool season grass
<i>Lactuca Serriola</i>	Prickly lettuce	Forb
<i>Lactuca tatarica</i> var. <i>pulchella</i>	Blue lettuce	Forb
<i>Lappula occidentalis</i> var. <i>occidentalis</i>	Western stickseed	Forb
<i>Lathyrus polymorphus</i>	Vetchling, wild pea	Forb
<i>Lepidium densiflorum</i>	Prairie-pepperweed	Forb
<i>Lesquerella arenosa</i>	Great Plains bladderpod	Forb
<i>Lesquerella ludoviciana</i>	Mustard	Forb
<i>Liatris punctata</i>	Blazing star, gay feather	Forb
<i>Linum puberulum</i>	Plains flax	Forb
<i>Linum rigidum</i>	Stiffstem yellow flax	Forb
<i>Lithospermum incisum</i>	Narrow-leaved puccoon	Forb
<i>Lupinus plattensis</i>	Lupine	Forb
<i>Lupinus pusillus</i>	Lupine	Forb
<i>Lygodesmia juncea</i>	Skeleton-weed	Forb
<i>Machaeranthera grindelioides</i>	Rayless aster	Forb
<i>Machaeranthera pinnatifida</i> ssp. <i>Pinnatifida</i>	Aster	Forb
<i>Melilotus</i> ssp.	<i>M. albus</i> , <i>M. officinalis</i>	Forb
<i>Mirabilis hirsuta</i>	Hairy umbrella-wort	Forb
<i>Mirabilis linearis</i>	Narrow-leaved umbrella-wort	Forb
<i>Muhlenbergia pungens</i>	Muhly	warm season grass
<i>Musineon tenuifolium</i>	Parsley	Forb
<i>Oenothera albicaulis</i>	Evening-primrose	Forb
<i>Oenothera cespitosa</i>	Evening-primrose	Forb

Appendix C. Agate Fossil Beds National Monument, Nebraska, cont'd.

Scientific Name	Common Name	Plant Community Guilds
<i>Oenothera nuttallii</i>	White-stemmed evening-primrose	Forb
<i>Oenothera villosa</i>	Evening-primrose	Forb
<i>Opuntia fragilis</i>	Little prickly pear	Forb
<i>Opuntia macrorhiza</i>	Plains prickly pear	Forb
<i>Opuntia polyacantha</i>	Plains prickly pear	Forb
<i>Orobanche fasciculata</i>	Clustered broom-rape	Forb
<i>Oryzopsis hymenoides</i>	Indian ricegrass	cool season grass
<i>Oxytropis sericea</i>	White locoweed	Forb
<i>Paronychia depressa</i>	Whitlow-wort	Forb
<i>Paronychia sessiliflora</i>	Nailwort	Forb
<i>Pascopyrum smithii</i>	Western wheatgrass	cool season grass
<i>Penstemon angustifolius</i>	Beard-tongue	Forb
<i>Penstemon eriantherus</i>	Beard-tongue	Forb
<i>Phacelia hastata</i>	Phacelia	Forb
<i>Phlox andicola</i>	Phlox, Sweet William	Forb
<i>Phlox hoodii</i>	Phlox, Sweet William	Forb
<i>Physalis hispida</i>	Plains-sandhill ground cherry	Forb
<i>Plantago patagonica</i>	Wooly plantain	Forb
<i>Poa pratensis</i>	Kentucky bluegrass	cool season grass
<i>Poa secunda</i>	Bluegrass	cool season grass
<i>Polygonum ramosissimum</i>	Smartweed	Forb
<i>Psoralea esculenta</i>	Breadroot scurf-pea	Forb
<i>Psoralidium lanceolatum</i>	Pea, Bean	Forb
<i>Psoralidium tenuiflorum</i>	Gray scurf-pea	Forb
<i>Rhus trilobata</i>	Squaw-bush	woody (shrubs and vines)
<i>Ribes americanum</i>	Gooseberry	woody (shrubs and vines)
<i>Rumex venosus</i>	Veiny dock	Forb
<i>Salix exigua</i>	Sandbar willow	woody (shrub and vines)
<i>Salsola</i> spp.	Russian thistle (<i>S. kali</i> or <i>S. collina</i>)	Forb
<i>Schizachyrium scoparium</i>	Little bluestem	warm season grass
<i>Senecio canus</i>	Gray groundsel	Forb
<i>Senecio riddellii</i>	Groundsel, ragwort	Forb
<i>Silene drummondii</i>	Drummond's campion	Forb
<i>Sisymbrium altissimum</i>	Tumbling-mustard	Forb
<i>Sphaeralcea coccinea</i>	Scarlet mallow	Forb
<i>Sporobolus cryptandrus</i>	Sand dropseed	warm season grass
<i>Stipa comata</i>	Needle-and-thread grass	cool season grass
<i>Symphoricarpos occidentalis</i>	Western snowberry	woody (shrubs and vines)
<i>Taraxacum officinale</i>	Common dandelion	Forb
<i>Tetrameuris acaulis</i>	Bitterweed	Forb
<i>Thermopsis rhombifolia</i>	Buckbean	Forb
<i>Townsendia grandiflora</i>	Easter daisy	Forb
<i>Tradescantia occidentalis</i>	Prairie spiderwort	Forb
<i>Tragopogon dubius</i>	Fistulous goat's beard	Forb
<i>Viola nuttallii</i>	Yellow prairie violet	Forb
<i>Vulpia octoflora</i>	Six-weeks fescue	cool season grass
<i>Yucca glauca</i>	Soap plant	woody (shrubs and vines)

Appendix C. cont”d.

Tallgrass Prairie National Preserve, Kansas

ScientificName	Common Name	Plant Community Guild
<i>Acalypha ostryifolia</i>	Rough-pod copperleaf	Forb
<i>Acalypha</i> spp		Forb
<i>Acalypha virginica</i>	Virginia copperleaf	Forb
<i>Achillea millefolium</i>	Common yarrow	Forb
<i>Aegilops cylindrica</i>	Jointed goatgrass	cool season grass
<i>Aesculus glabra</i>	Western buckeye	woody (shrub or small tree)
<i>Agalinis aspera</i>	Tall false foxglove	Forb
<i>Agalinis densiflora</i>	Fineleaf gerardia	Forb
<i>Agalinis</i> spp.-TAPR		Forb
<i>Agalinis tenuifolia</i>	Slender gerardia	Forb
<i>Agastache nepetoides</i>	Catnip giant hyssop	Forb
<i>Ageratina altissima</i>	Tall ageratina	Forb
<i>Agrimonia parviflora</i>	Many-flowered agrimony	Forb
<i>Agrostis hyemalis</i>	Ticklegrass	cool season grass
<i>Alisma subcordatum</i>	Water plantain	Forb
<i>Alliaria petiolata</i>	Garlic-mustard	Forb
<i>Allium canadense</i>	Onion	Forb
<i>Alopecurus carolinianus</i>	Carolina foxtail	cool season grass
<i>Amaranthus blitoides</i>	Prostrate pigweed	Forb
<i>Amaranthus retroflexus</i>	Rough pigweed	Forb
<i>Amaranthus rudis</i>	Water hemp	Forb
<i>Amaranthus spinosus</i>	Spiny pigweed	Forb
<i>Ambrosia artemisiifolia</i>	Common ragweed	Forb
<i>Ambrosia psilostachya</i>	Western ragweed	Forb
<i>Ambrosia trifida</i>	Giant ragweed	Forb
<i>Ammannia coccinea</i>	Purple ammannia	Forb
<i>Amorpha canescens</i>	Lead-plant	woody (shrubs and vines)
<i>Amorpha fruticosa</i>	False indigo	woody (shrubs and vines)
<i>Amphiachyris dracunculoides</i>	Broomweed	Forb
<i>Andropogon gerardii</i>	Big bluestem	warm season grass
<i>Androsace occidentalis</i>	Western androsace	Forb
<i>Anemone caroliniana</i>	Carolina anemone	Forb
<i>Antennaria neglecta</i>	Field pussytoes	Forb
<i>Apocynum cannabinum</i>	Hemp dogbane	Forb
<i>Aquilegia canadensis</i>	Wild columbine	Forb
<i>Arctium minus</i>	Common burdock	Forb
<i>Arenaria serpyllifolia</i>	Thyme-leaved sandwort	Forb
<i>Arisaema dracontium</i>	Dragonroot	Forb
<i>Aristida oligantha</i>	Prairie three-awn	warm season grass
<i>Arnoglossum atriplicifolium</i>	Pale indian plantain	Forb
<i>Arnoglossum plantagineum</i>	Tuberous Indian plantain	Forb
<i>Artemisia ludoviciana</i>	White sage	Forb
<i>Asclepias amplexicaulis</i>	Clasping milkweed	Forb
<i>Asclepias asperula</i>	Antelope horns	Forb
<i>Asclepias</i> spp.		Forb

Appendix C. Tallgrass Prairie National Preserve, Kansas, cont'd.

ScientificName	Common Name	Plant Community Guild
<i>Asclepias stenophylla</i>	Narrow-leaved milkweed	Forb
<i>Asclepias sullivantii</i>	Smooth milkweed	Forb
<i>Asclepias syriaca</i>	Common milkweed	Forb
<i>Asclepias tuberosa</i>	Butterfly-weed	Forb
<i>Asclepias verticillata</i>	Whorled milkweed	Forb
<i>Asclepias viridiflora</i>	Green milkweed	Forb
<i>Asclepias viridis</i>	Ozark milkweed	Forb
<i>Aster drummondii</i>	Hairy heart-leaved wild aster	Forb
<i>Aster ericoides</i>	Squarrose white wild aster	Forb
<i>Aster laevis</i>	Smooth wild aster	Forb
<i>Aster oblongifolius</i>	Aromatic wild aster	Forb
<i>Aster sericeus</i>	Western silvery wild aster	Forb
<i>Astragalus Crassicaarpus</i>	Ground-plum, prairie plum	Forb
<i>Astragalus lotiflorus</i>	Lotus milk-vetch	Forb
<i>Baptisia australis</i>	blue wild indigo	Forb
<i>Baptisia bracteata</i> var. <i>leucophaea</i>	Plains wild indigo	Forb
<i>Bassia scoparia</i>	Summer cypress	Forb
<i>Bidens aristosa</i>	Coreopsis beggar-ticks	Forb
<i>Bidens cernua</i>	Nodding beggar-ticks	Forb
<i>Bidens frondosa</i>	Beggar-ticks	Forb
<i>Bothriochloa bladii</i>	Caucasian bluestem	warm season grass
<i>Bothriochloa saccharoides</i>	Silver bluestem	warm season grass
<i>Botrychium virginianum</i>	Rattlesnake fern	Forb
<i>Bouteloua curtipendula</i>	Side-oats grama-grass	warm season grass
<i>Bouteloua gracilis</i>	Blue grama-grass	warm season grass
<i>Bouteloua hirsuta</i>	Hairy grama-grass	warm season grass
<i>Brickellia eupatorioides</i>	Aster	Forb
<i>Bromus inermis</i>	Smooth brome	cool season grass
<i>Bromus japonicus</i>	Japanese brome	cool season grass
<i>Bromus tectorum</i>	Downy brome	cool season grass
<i>Buchloe dactyloides</i>	Buffalo-grass	cool season grass
<i>Buglossoides arvensis</i>	Corn gromwell	Forb
<i>Cacalia plantaginea</i>	Tuberous Indian plantain	Forb
<i>Callirhoe alcaeoides</i>	Pale poppy-mallow	Forb
<i>Callirhoe involucrata</i>	Purple poppy mallow	Forb
<i>Callitriche heterophylla</i>	Water starwort	Forb
<i>Calylophus serrulatus</i>	Evening-primrose	Forb
<i>Calystegia macounii</i>	Macoun's bindweed	Forb
<i>Camelina microcarpa</i>	Small-seeded false flax	Forb
<i>Capsella bursa-pastoris</i>	Shepherd's purse	Forb
<i>Carduus nutans</i>	Musk thistle	Forb
<i>Carex amphibola</i>	Eastern narrowleaf sedge	grass-like (cool season grass)
<i>Carex blanda</i>	Sedge	grass-like (cool season grass)
<i>Carex brevior</i>	Sedge	grass-like (cool season grass)
<i>Carex davisii</i>	Davis sedge	grass-like (cool season grass)
<i>Carex emoryi</i>	Emory sedge	grass-like (cool season grass)
<i>Carex frankii</i>	Frank's sedge	grass-like (cool season grass)
<i>Carex gravida</i>	Heavy sedge	grass-like (cool season grass)
<i>Carex hystericina</i>	Bottlebrush sedge	grass-like (cool season grass)
<i>Carex meadii</i>	Sedge	grass-like (cool season grass)

Appendix C. Tallgrass Prairie National Preserve, Kansas, cont'd.

ScientificName	Common Name	Plant Community Guild
Carex mulilenbergii	Muhlenberg sedge	grass-like (cool season grass)
Carex spp.		grass-like (cool season grass)
Carex vulpinoidea	Fox sedge	grass-like (cool season grass)
Ceanothus americanus	New Jersey tea, redroot	woody (shrubs and vines)
Ceanothus herbaceus	Prairie-redroot	woody (shrubs and vines)
Celastrus scandens	American bittersweet	vine
Cenchrus longispinus	Sandbur	warm season grass
Cephalanthus occidentalis	Buttonbush	Shrub
Cerastium brachypodum	Mouse-ear chickweed	Forb
Ceratophyllum demersum	Hornwort	Forb
Chaerophyllum procumbens	Spreading chervil	Forb
Chamaecrista fasciculata	Showy partridge pea	Forb
Chamaesyce glyptosperma	Ridge-seed spurge	Forb
Chamaesyce missurica	Missouri spurge	Forb
Chamaesyce nutans	Eyebane	Forb
Chamaesyce prostrata	Spurge	Forb
Chenopodium album	Lamb's quarters, pigweed	Forb
Chenopodium berlandieri	Pitseed goosefoot	Forb
Chenopodium simplex	Mapleleaf goosefoot	Forb
Chenopodium spp.		Forb
Chloris verticillata	Windmill finger-grass	Cool season grass
Cirsium altissimum	Tall thistle	Forb
Cirsium undulatum	Wavy-leaved thistle	Forb
Clematis pitcheri	Pitcher's clematis	vine
Clematis terniflora	Sweet autumn virginbower	woody (shrubs and vines)
Commelina erecta	Erect dayflower	Forb
Conium maculatum	Poison hemlock	Forb
Convolvulus arvensis	Field-bindweed	Forb
Conyza Canadensis	Horseweed	Forb
Coreopsis grandiflora	Bigflower coreopsis	Forb
Cornus drummondii	Rough-leaved dogwood	woody (shrubs and vines)
Corydalis micrantha	Slender corydalis	annual-biennial
Crataegus mollis	Downy hawthorn	woody (shrub or small tree)
Croton capitatus	Woolly croton	Forb
Croton monanthogynus	Prairie-tea, one-seed croton	Forb
Croton spp.		Forb
Croton texensis	Texas croton	Forb
Cucurbita foetidissima	Buffalo-gourd	Forb
Cynanchum laeve	Honey vine	Forb
Cyperus esculentus	Yellow nutsedge	cool season grass
Cyperus lupulinus	Houghton flatsedge	cool season grass
Cyperus odoratus	Fragrant flatsedge	cool season grass
Cyperus setigerus	Roughedge flatsedge	cool season grass
Cyperus spp.		grass-like (cool season grass)
Cyperus strigosus	False nutgrass	cool season grass
Cystopteris tennesseensis	Tennessee bladder fern	Forb
Dalea aurea	Golden prairie clover	Forb
Dalea candida	White prairie clover	Forb
Dalea multiflora	roundhead prairieclover	Forb
Dalea purpurea	Purple prairie clover	Forb
Dalea spp.		Forb

Appendix C. Tallgrass Prairie National Preserve, Kansas, cont'd.

ScientificName	Common Name	Plant Community Guild
<i>Datura stramonium</i>	Jimson weed	Forb
<i>Delphinium carolinianum</i> spp. <i>Virescens</i>	Prairie larkspur	Forb
<i>Descurainia pinnata</i>	Tansy-mustard	Forb
<i>Desmanthus illinoensis</i>	Bundleflower	Forb
<i>Desmodium cuspidatum</i>	Big tick-trefoil	Forb
<i>Desmodium illinoense</i>	Prairie tick-trefoil	Forb
<i>Desmodium sessilifolium</i>	Sessile tickclover	Forb
<i>Desmodium</i> spp.		Forb
<i>Dichanthelium acuminatum</i>	Panic grass	cool season grass
<i>Dichanthelium oligosanthos</i>	Panic grass	cool season grass
<i>Dichanthelium</i> spp.		cool season grass
<i>Digitaria cognata</i>	Carolina crab grass	cool season grass
<i>Digitaria sanguinalis</i>	Hairy crabgrass	warm season grass
<i>Draba branchycarpa</i>	Mustard	Forb
<i>Draba cuneifolia</i>	Mustard	Forb
<i>Draba reptans</i>	Mustard	Forb
<i>Dyssodia papposa</i>	Fetid marigold	Forb
<i>Echinacea angustifolia</i>	Prairie coneflower	Forb
<i>Echinochloa crus-galli</i>	Barnyard grass	warm season grass
<i>Echinochloa muricata</i>	Rough barnyard grass	warm season grass
<i>Eclipta prostrata</i>	Yerba-de-tajo	Forb
<i>Eleocharis compressa</i>	Spike-rush	grass-like (cool season grass)
<i>Eleocharis macrostachya</i>	Longstem spike sedge	cool season grass
<i>Eleocharis obtusa</i>	Blunt spike sedge	cool season grass
<i>Eleocharis</i> spp.		grass-like (cool season grass)
<i>Ellisia nyctelea</i>	Water-pod	Forb
<i>Elymus canadensis</i>	Canada wild rye	cool season grass
<i>Elymus</i> spp.		cool season grass
<i>Elymus virginicus</i>	Virginia wild rye	warm season grass
<i>Equisetum laevigatum</i>	Smooth horsetail	warm season grass
<i>Eragrostis cilianensis</i>	Stinkgrass	warm season grass
<i>Eragrostis spectabilis</i>	Purple lovegrass	warm season grass
<i>Erigeron philadelphicus</i>	Philadelphia daisy	Forb
<i>Erigeron</i> spp.		Forb
<i>Erigeron strigosus</i>	Rough fleabane	Forb
<i>Eriochloa contracta</i>	Prairie cupgrass	warm season grass
<i>Erysimum repandum</i>	Bushy wallflower	Forb
<i>Erythronium mesochoreum</i>	Midland fawn lily	Forb
<i>Escobaria missouriensis</i> var. <i>missouriensis</i>	Missouri coryphanthe	Forb
<i>Euonymus atropurpurea</i>	Wahoo	shrub
<i>Eupatorium altissimum</i>	Tall Joe-Pye weed	Forb
<i>Euphorbia corollata</i>	Flowering spurge	Forb
<i>Euphorbia cyathophora</i>	Fire-on-the-mountain	Forb
<i>Euphorbia dentata</i>	Toothed spurge	Forb
<i>Euphorbia marginata</i>	Snow-on-the-mountain	Forb
<i>Euphorbia spathulata</i>	Prairie spurge	Forb
<i>Euthamia gymnospermoides</i>	Great Plains flat-topped goldenrod	Forb
<i>Evolvulus nuttallianus</i>	Evolvulus	Forb
<i>Festuca subverticillata</i>	Nodding fescue	cool season grass
<i>Fuirena simplex</i>	Umbrella grass	cool season grass

Appendix C. Tallgrass Prairie National Preserve, Kansas, cont'd.

ScientificName	Common Name	Plant Community Guild
Galium aparine	Cleavers	Forb
Galium circaezans	Forest bedstraw, wild licorice	Forb
Galium triflorum	Sweet-scented bedstraw	Forb
Gaura longiflora	Large flowered gaura	Forb
Gaura mollis	Velvety gaura	Forb
Geranium carolinianum	Carolina crane's-bill	Forb
Geum canadense	White avens	Forb
Glandularia canadensis	Rose verberna	Forb
Glyceria striata	Fowl mannagrass	cool season grass
Glycyrrhiza lepidota	Wild licorice	Forb
Grindelia squarrosa	Curly-top gumweed	Forb
Hedeoma hispidum	Mint	Forb
Hedyotis nigricans var. nigricans	Madder	Forb
Helianthus annuus	Common sunflower	Forb
Helianthus maximiliani	Maximilian sunflower	Forb
Helianthus pauciflorus	Stiff sunflower	Forb
Helianthus salicifolius	Willow-leaved sunflower	Forb
Helianthus spp.		Forb
Helianthus tuberosus	Jerusalem artichoke	Forb
Heliopsis helianthoides	False sunflower	Forb
Heterotheca canescens	Golden aster	Forb
Hieracium longipilum	Long-haired hawkweed	Forb
Holcus lanatus	Common velvet-grass	cool season grass
Hordeum pusillum	Little barley	cool season grass
Hybanthus verticillatus	Nodding green violet	Forb
Hymenopappus scabiosaeus	Carolina woollywhite	Forb
Hypericum perforatum	Common ST. Johns-wort	Forb
Ipomoea hederacea	Ivyleaf morning-glory	Forb
Iva annua	Annual sumpweed	Forb
Juncus dudleyi	Dudley rush	cool season grass
Juncus interior	Rush	grass-like (cool season grass)
Juncus spp.		grass-like (cool season grass)
Juncus tenuis	Path rush	cool season grass
Juncus torreyi coville	Torrey's rush	cool season grass
Justicia americana	Water willow	Forb
Kochia scoparia	Summer-cypress	Forb
Koeleria macrantha	Junegrass	cool season grass
Krigia cespitosa	Sunflower	Forb
Kummerowia stipulacea	Korean clover	Forb
Lactuca ludoviciana	Western wild lettuce	Forb
Lactuca Serriola	Prickly lettuce	Forb
Lactuca spp.		Forb
Lamium amplexicaule	Henbit	Forb
Laportea canadensis	Nettle	Forb
Leersia oryzoides	Rice cut-grass	warm season grass
Lemna minor	Duckweed	Forb
Lepidium densiflorum	Prairie-pepperweed	Forb
Lespedeza capitata	Bush-clover	Forb
Lespedeza violacea	Violet lespedeza	Forb
Lespedeza virginica	Slender bush lespedeza	Forb
Leucospora multifida	Leucospora	Forb

Appendix C. Tallgrass Prairie National Preserve, Kansas, cont'd.

ScientificName	Common Name	Plant Community Guild
<i>Liatris aspera</i>	Lacerate blazing star	Forb
<i>Liatris mucronata</i>	Gayfeather	Forb
<i>Liatris punctata</i>	Blazing star, gay feather	Forb
<i>Liatris</i> spp.–TAPR	<i>L. mucronata</i> , <i>L. punctata</i>	Forb
<i>Linum sulcatum</i>	Grooved yellow flax	Forb
<i>Lithospermum Incisum</i>	Narrow-leaved puccoon	Forb
<i>Lobelia cardinalis</i>	Cardinal-flower	Forb
<i>Lobelia siphilitica</i>	Blue lobelia	Forb
<i>Lomatium foeniculaceum</i>	Yellow wild parsley	Forb
<i>Ludwigia peploides</i>	Floating evening primrose	Forb
<i>Lycopus americanus</i>	American bugleweed	Forb
<i>Lythrum alatum</i>	Winged loosestrife	Forb
<i>Medicago lupulina</i>	Black medick	Forb
<i>Melilotus albus</i>	White sweet clover	Forb
<i>Melilotus officinalis</i>	Sweet clover	Forb
<i>Menispermum canadense</i>	Moonseed	woody (shrubs and vines)
<i>Mentha arvensis</i>	Field mint	Forb
<i>Mentzelia oligosperma</i>	Stickleaf mentzelia	Forb
<i>Mimosa nuttallii</i>	Catclaw sensitive brier	Forb
<i>Mirabilis albida</i>	White four o'clock	Forb
<i>Mirabilis linearis</i>	Narrow-leaf Four o'clock	Forb
<i>Mirabilis nyctaginea</i>	Heart-leaved umbrella-wort	Forb
<i>Mirabilis</i> spp. – TAPR	<i>M. albida</i> , <i>M. linearis</i>	Forb
<i>Monarda fistulosa</i>	Wild bergamot	Forb
<i>Monolepis nuttalliana</i>	Poverty weed	Forb
<i>Muhlenbergia frondosa</i>	Muhly	warm season grass
<i>Muhlenbergia cuspidata</i>	Muhly	cool season grass
<i>Myagrum perfoliatum</i>	Bird's-eye cress	Forb
<i>Myosotis verna</i>	Early scorpion grass	Forb
<i>Myosurus minimus</i>	Tiny mousetail	Forb
<i>Myriophyllum heterophyllum</i>	Water milfoil	Forb
<i>Nepeta cataria</i>	Catnip	Forb
<i>Nothocalais cuspidata</i>	Wavyleaf agoseris	Forb
<i>Nothoscordum bivalve</i>	Lily	Forb
<i>Oenothera biennis</i>	Common evening-primrose	Forb
<i>Oenothera macrocarpa</i>	Wing-fruit evening-primrose	Forb
<i>Oenothera speciosa</i>	White evening-primrose	Forb
<i>Oligoneuron rigidum</i>	Rigid goldenrod	Forb
<i>Onosmodium molle</i>	Western false gromwell	Forb
<i>Opuntia macrorhiza</i>	Plains prickly pear	Forb
<i>Osmorhiza longistylis</i>	Anise root	Forb
<i>Oxalis dillenii</i>	Southern yellow wood-sorrel	Forb
<i>Oxalis</i> spp.	wood-sorrel	Forb
<i>Oxalis violacea</i>	Violet wood-sorrel	Forb
<i>Packera plattensis</i>	Prairie ragwort	Forb
<i>Panicum capillare</i>	Witch-grass	warm season grass
<i>Panicum dichotomiflorum</i>	Fall panicum	warm season grass
<i>Panicum obtusum</i>	Vine-mesquite	warm season grass
<i>Panicum virgatum</i>	Switchgrass	warm season grass
<i>Parietaria pensylvanica</i>	Pennsylvania pellitory	Forb
<i>Parthenocissus quinquefolia</i>	Virginia-creeper, woodbine	woody (shrubs and vines)

Appendix C. Tallgrass Prairie National Preserve, Kansas, cont'd.

ScientificName	Common Name	Plant Community Guild
Pascopyrum smithii	Western wheatgrass	cool season grass
Paspalum pubiflorum	Hairyseen paspalum	warm season grass
Paspalum spp.-TAPR		warm season grass
Pedimelum argophyllum	Silver-leaf scurf pea	Forb
Pedimelum esculentum	Breadroot scurf-pea	Forb
Pedimelum tenuiflorum	Wild alfalfa	Forb
Pellaea atropurpurea	Purple cliff-brake	Forb
Pennisetum glaucum	Yellow foxtail	warm season grass
Penstemon cobaea	cobaea beardtongue	Forb
Penstemon grandiflorus	Large beard-tongue	Forb
Penstemon tubiflorus	Tube beard-tongue	Forb
Penthorum sedoides	Ditch stonecrop	Forb
Phlox divaricata	Forest phlox	Forb
Phlox spp.		Forb
Phryma leptostachya	Lopseed	Forb
Phyla lanceolata	Northern frog fruit	Forb
Physalis heterophylla	Clammy ground cherry	Forb
Physalis longifolia	Longflower ground cherry	Forb
Physalis pumila	Prairie ground cherry	Forb
Physalis spp.		Forb
Physalis virginiana	Virginia ground cherry	Forb
Phytolacca americana	Pokeweed, pokeberry	Forb
Plantago patagonica	Wooly plantain	Forb
Plantago rhodosperma	Plantain	Forb
Plantago rugelii	American plantain	Forb
Plantago spp.		Forb
Plantago virginica	Pale-seeded plantain	Forb
Poa arida	Plains bluegrass	cool season grass
Poa pratensis	Kentucky bluegrass	cool season grass
Poa spp.		cool season grass
Polanisia dodecandra	Clammy-weed	Forb
Polygala verticillata	Whorled milkwort	Forb
Polygonatum biflorum	Solomon's seal	Forb
Polygonum lapathifolium	Pale smartweed	Forb
Polygonum pensylvanicum	Pink smartweed	Forb
Polygonum persicaria	Lady's-thumb	Forb
Polygonum punctatum	Dotted smartweed	Forb
Polygonum ramosissimum	Smartweed	Forb
Polygonum scandens	False buckwheat	Forb
Polygonum virginianum	Jumpseed	Forb
Polytaenia nuttallii	Nuttall's prairie parsley	Forb
Portulaca oleracea	Common purslane	Forb
Potamogeton foliosus	Leafy pondweed	Forb
Potamogeton nodosus	Longleaf pondweed	Forb
Prenanthes aspera	Rough rattlesnake root	Forb
Prunella vulgaris	Selfheal	Forb
Prunus americana	American wild plum	Tree/shrub
Prunus virginiana	Choke cherry	Tree/shrub
Psoralea argophylla	Silvery scurf-pea	Forb
Psoralea esculenta	Breadroot scurf-pea	Forb

Appendix C. Tallgrass Prairie National Preserve, Kansas, cont'd.

ScientificName	Common Name	Plant Community Guild
<i>Psoraleum tenuiflorum</i>	Gray scurf-pea	Forb
<i>Ranunculus sceleratus</i>	Cursed crowfoot	Forb
<i>Ratibida columnifera</i>	Columnar coneflower	Forb
<i>Rhus aromatica</i>	Fragrant sumac	shrub
<i>Rhus glabra</i>	Smooth sumac	woody (shrubs and vines)
<i>Ribes aureum</i>	Buffalo currant	shrub
<i>Ribes missouriense</i>	Missouri gooseberry	woody (shrubs and vines)
<i>Rorippa nasturtium</i>	Watercress	Forb
<i>Rosa arkansana</i>	Dwarf prairie rose	woody (shrubs and vines)
<i>Rosa multiflora</i>	Multiflora rose	shrub
<i>Ruellia humilis</i>	Fringeleaf ruellia	Forb
<i>Ruellia strepens</i>	Acanthus	Forb
<i>Rumex altissimus</i>	Pale dock	Forb
<i>Rumex crispus</i>	Curly dock	Forb
<i>Rumex</i> spp.	Dock	Forb
<i>Sagittaria brevirostra</i>	Arrowhead	Forb
<i>Salvia azurea</i>	Sage	Forb
<i>Salvia reflexa</i>	Lance-leaved sage	Forb
<i>Sambucus canadensis</i>	Common elder	woody (shrubs and vines)
<i>Sambucus nigra</i>	Common elderberry	Tree/shrub
<i>Sanicula canadensis</i>	Black snakeroot	Forb
<i>Schizachyrium scoparium</i>	Little bluestem	warm season grass
<i>Schoenoplectus tabernaemontani</i>	Soft-stem bulrush	cool season grass
<i>Schrankia nuttallii</i>	Sensitive brier	Forb
<i>Scirpus atrovirens</i>	Darkgreen bulrush	cool season grass
<i>Scirpus pallidus</i>	Pale bulrush	cool season grass
<i>Scirpus pendulus</i>	Rusty bulrush	cool season grass
<i>Scrophularia marilandica</i>	Maryland figwort	Forb
<i>Scutellaria parvula</i>	Little skullcap	Forb
<i>Senecio platensis</i>	Platte groundsel	Forb
<i>Senna marilandica</i>	Wild senna	Forb
<i>Setaria</i> spp.		warm season grass
<i>Setaria viridis</i>	Green foxtail-grass	warm season grass
<i>Sicyos angulatus</i>	Bur cucumber	Forb
<i>Sida spinosa</i>	Prickly spinosa	Forb
<i>Silene antirrhina</i>	Catchfly, campion	Forb
<i>Silene stellata</i>	Starry campion	Forb
<i>Silphium integrifolium</i>	Showy rosinweed	Forb
<i>Silphium laciniatum</i>	Compass-plant	Forb
<i>Sisyrinchium campestre</i>	Blue-eyed grass	Forb
<i>Smilax tamnoides</i>	Catbrier	woody (shrubs and vines)
<i>Solanum carolinense</i>	Horse-nettle	Forb
<i>Solanum elaeagnifolium</i>	Silver-leaf nightshade	Forb
<i>Solanum rostratum</i>	Buffalo-bur	Forb
<i>Solidago canadensis</i>	Common goldenrod	Forb
<i>Solidago Canadensis</i> var. <i>gilvocanescens</i>	Common goldenrod	Forb
<i>Solidago missouriensis</i>	Missouri goldenrod	Forb
<i>Solidago petiolaris</i>	Downy goldenrod	Forb
<i>Solidago rigida</i>	Stiff goldenrod	Forb
<i>Solidago</i> spp.		Forb

Appendix C. Tallgrass Prairie National Preserve, Kansas, cont'd.

ScientificName	Common Name	Plant Community Guild
<i>Sonchus asper</i>	Prickly sow thistle	Forb
<i>Sorghastrum nutans</i>	Indian grass	warm season grass
<i>Sorghum halepense</i>	Johnson grass	warm season grass
<i>Spartina pectinata</i>	Prairie cordgrass	warm season grass
<i>Spermolepis inermis</i>	Western spermolepis	Forb
<i>Spiranthes cernua</i>	Nodding ladies' tresses	Forb
<i>Spiranthes vernalis</i>	Twisted lady's tresses	Forb
<i>Sporobolus asper</i>	Tall dropseed	warm season grass
<i>Sporobolus compositus</i>	Rough dropseed	warm season grass
<i>Sporobolus neglectus</i>	Dropseed	warm season grass
<i>Sporobolus spp.</i>		warm season grass
<i>Stellaria media</i>	Common chickweed	Forb
<i>Stenosiphon linifolius</i>	Stenosiphon	Forb
<i>Strophostyles helvula</i>	Annual woolly bean	Forb
<i>Strophostyles leiosperma</i>	Small-flowered woolly bean	Forb
<i>Symphoricarpos orbiculatus</i>	Coralberry	woody (shrubs and vines)
<i>Symphyotrichum drummondii</i>	Drummond aster	Forb
<i>Symphyotrichum ericoides</i>	White aster	Forb
<i>Symphyotrichum lanceolatum</i>	Simple panicled aster	Forb
<i>Symphyotrichum oblongifolium</i>	Aromatic aster	Forb
<i>Symphyotrichum sericeum</i>	Silky aster	Forb
<i>Taraxacum officinale</i>	Common dandelion	Forb
<i>Teucrium canadense</i>	American germander	Forb
<i>Thalictrum dasycarpum</i>	Purple meadow rue	Forb
<i>Thelesperma filifolium</i>	Greenthread	Forb
<i>Thlaspi arvense</i>	Field penny-cress	Forb
<i>Toxicodendron radicans</i>	Common poison-ivy	woody (shrubs and vines)
<i>Tradescantia bracteata</i>	Sticky spiderwort	Forb
<i>Tradescantia ohiensis</i>	Ohio spiderwort	Forb
<i>Tragia betonicifolia</i>	Many-flowered noseburn	Forb
<i>Tragia ramosa</i>	branched noseburn	Forb
<i>Tragopogon dubius</i>	Goats beard	Forb
<i>Tribulus terrestris</i>	Puncture vine	Forb
<i>Trichostema brachiatum</i>	Blue curls	Forb
<i>Tridens flavus</i>	Purpletop	warm season grass
<i>Trifolium repens</i>	White clover	Forb
<i>Triodanis leptocarpa</i>	Narrow-leaved triodanis	Forb
<i>Triodanis perfoliata</i>	Round-leaved triodanis	Forb
<i>Tripsacum dactyloides</i>	Eastern gammagrass	warm season grass
<i>Typha latifolia</i>	Common cattail	Forb
<i>Ulmus pumila</i>	Siberian elm	Tree/shrub
<i>Urtica dioica</i> ssp. <i>Gracilis</i>	Nettle, stinging nettle	Forb
<i>Verbascum blattaria</i>	Moth mullein	Forb
<i>Verbascum thapsus</i>	Common mullein	Forb
<i>Verbena bracteata</i>	Prostrate vervain	Forb
<i>Verbena hastata</i>	Blue vervain	Forb
<i>Verbena simplex</i>	Narrow-leaved vervain	Forb
<i>Verbena stricta</i>	Hoary vervain	Forb
<i>Verbesina alternifolia</i>	Wingstem	Forb

Appendix C. Tallgrass Prairie National Preserve, Kansas, cont'd.

ScientificName	Common Name	Plant Community Guild
Vernonia baldwinii	Western ironweed	Forb
Veronica anagallis-aquatica	Water speedwell	Forb
Veronica arvensis	Corn speedwell	Forb
Veronica peregrina	Purslane speedwell	Forb
Vicia americana	American vetch	Forb
Viola bicolor	Violet	Forb
Viola nephrophylla	Blue prairie violet	Forb
Viola pedatifida	Wood violet	Forb
Viola pratincola	Violet	Forb
Viola sororia	Downy Blue violet	Forb
Viola spp.		Forb
Vitis riparia	Frost grape	woody (shrubs and vines)
Vulpia octoflora	Six-weeks fescue	cool season grass
Xanthium strumarium	Common cocklebur	Forb
Yucca glauca	Yucca	Forb
Zanthoxylum americanum	Common prickly ash	woody (shrubs and vines)
Zigadenus nuttallii	Nuttall's deathcamas	Forb
Zizia aurea	Common golden alexanders	Forb

Appendix D. Common tree species found on Tallgrass Prairie National Preserve, Kansas.

Scientific Name	Common Name	Plant Community Guild
Acer negundo	Box elder	Tree
Acer saccharinum	Silver maple	Tree
Carya cordiformis	Bitternut Hickory	Tree
Carya illinoensis	Pecan	Tree
Catalpa bignonioides	Catalpa	Tree
Celtis occidentalis	Hackberry	Tree
Cercis canadensis	Redbud	Tree
Cornus drummondii	Rough-leaved dogwood	Tree
Fraxinus pennsylvanica	Green ash	Tree
Gleditsia triacanthos	Honey locust	Tree
Gymnocladus dioica	Kentucky coffee tree	Tree
Juglans nigra	Black Walnut	Tree
Juniperus virginiana	Red cedar	Tree
Maclura pomifera	Osage-orange	Tree
Morus alba	White mulberry	Tree
Platanus occidentalis	Sycamore	Tree
Populus deltoides	Eastern cottonwood	Tree
Quercus macrocarpa	Bur oak	Tree
Robinia pseud-acacia	Black locust	Tree
Salix amygdaloides	Peachleaf willow	Tree
Salix caroliniana	Carolina willow	Tree
Salix nigra	Black willow	Tree
Ulmus americana	American elm	Tree