Macroinvertebrates *Resource Brief*

Importance

A macroinvertebrate is any animal without a backbone that is large enough to be seen by the naked eye. Macroinvertebrates that are aquatic (water-dwelling) or benthic (occupying the lowest level of a body of water) include aquatic insects, crayfish, and mussels. Crayfish, and aquatic insects (including stoneflies, mayflies, and caddisflies), consume organic matter that falls into rivers and streams, while mussels filter nutrients and sediment from the water. All are a food source for other animals. Fish and other stream-dependent creatures like water turtles and salamanders consume aquatic insects; crayfish are eaten by fish, birds, raccoons, and others; and mussels provide food and habitat to other animals.

Macroinvertebrates are good indicators of watershed health and water quality because they live in the water for all or most of their life, have limited mobility, and differ in their tolerance to amounts and types of pollution.



Photo: NPS/Nortrup. Illustration: University of Maryland, Integration & Application Network

Above: A fine-meshed D-shaped net is used to scoop up macroinvertebrates from promising stream habitat. Left: Presence of pollution-sensitive species like this mayfly (Ephemeroptera), along with stoneflies and caddisflies can indicate healthy stream condition.

Monitoring

Macroinvertebrate monitoring is part of a broader effort by the National Capital Region Network (NCRN) Inventory & Monitoring (I&M) program to assess the condition of streams and watersheds.

Long-term macroinvertebrate monitoring at thirty-seven park sites throughout the NCRN began in 2008 and follows a set, six-year rotation. Each spring 5-8 sites are visited. At Manassas monitoring is done on Youngs Branch. Macroinvertebrate monitoring is co-located with fish monitoring and stream physical habitat analysis.

The objectives of this combined monitoring are to:

- determine current conditions and track long-term trends in stream condition,
- determine trends in species composition and functional groups of fish and benthic invertebrates

At each site, monitoring teams work within a 75 meter stream segment. Organisms are dislodged from available riffles and other productive stream habitats and captured in a fine-meshed, D-shaped net (see photo). A subsample of 100 individuals are preserved and identified to genus if possible, using stereoscopes (low-magnification microscopes). Chironomidae (chironomids or non-biting midges) are slide-mounted and identified using compound microscopes.

BIBI Scores

The species and number of macroinvertebrates present in a stream segment are used to calculate a Benthic Index of Biotic Integrity (BIBI) score. Scoring takes into account factors such as:

- number and diversity of taxonomic groups
- the proportion of individuals belonging to specific taxonomic groups
- ability of individuals present to survive short- and long-term exposure to stressors from pollution, changes in stream flow, or habitat degradation
- mode of feeding including scrapers, predators, collectors, filterers, and shredders
- habit tendencies including burrowers, climbers, clingers, and sprawlers

Scores also take into account Manassass's location in the Eastern Piedmont physiographic province. BIBI scores range from 1 to 5 and are ranked as follows: 1.0-1.9 (very poor), 2.0-2.9 (poor), 3.0-3.9 (fair), and 4.0-5.0 (good). For results, see reverse.

Manassas National Battlefield Park

National Park Service
U.S. Department of the Interior

Results

Youngs Branch was sampled in 2010 and in 2004. BIBI scores indicate an improvement in stream condition from very poor to fair.

During 2010 sampling of Youngs Branch, dead individuals of the exotic Asian clam species, *Corbicula fluminea* were found.

Youngs Branch

2010 BIBI = 3 (fair)

2004 BIBI = 1.25 (very poor)

Number of taxa found in 2010: 37

Most common taxa in 2010: *Cricotopus* (14), *Tanytarsus* (11), *Simulium* (9), *Stenelmis* (9), and *Perlesta* (8)

Discussion

The change from very poor to fair BIBI scores for Youngs Branch stems from the increased presence of pollution- and disturbance-sensitive macroinvertebrates of the"clinger" variety and a reduction in the amount of pollution-tolerant Chironomids.

Taxa Explained

Cricotopus – genus of non-biting midges in the subfamily Orthocladiinae of the bloodworm family Chironomidae

Perlesta – genus of stonefly

Simulium- a genus of blackflies

Stenelmis - genus of beetles in the family Elmidae

Tanytarsus - a large genus of non-biting midges in the subfamily Orthocladiinae of the bloodworm family Chironomidae

References:

National Capital Region Network Biological Stream Survey Macroinvertebrate Data (2008-2010). Versar, Inc. National Capital Region Inventory and Monitoring Program, Washington, DC. Generic Dataset-2175665. https://irma.nps.gov/App/Reference/Profile/2175665

NCRN Macroinvertebrate webpage. http://science.nature.nps.gov/im/units/ncrn/monitoring_macroinverts.cfm

Norris, M. E., and G. Sanders. 2009. National Capital Region Network biological stream survey protocol: Physical habitat, fish, and aquatic macroinvertebrate vital signs. Natural Resource Report NPS/NCRN/NRR—2009/116. National Park Service, Fort Collins, Colorado. http://science.nature.nps.gov/im/monitor/protocols/NCRN_Biological_Stream_Survey_Protoco&SOPs_Ver_2.0.pdf