Monocacy National Battlefield

National Park Service
U.S. Department of the Interior

Macroinvertebrates *Resource Brief*

Importance

A macroinvertebrate is any animal without a backbone that is large enough to be seen by the naked eye. Macroinvertebrates that are aquatic (water-dwelling) or benthic (occupying the lowest level of a body of water) include aquatic insects, crayfish, and mussels. Crayfish, and aquatic insects (including stoneflies, mayflies, and caddisflies), consume organic matter that falls into rivers and streams, while mussels filter nutrients and sediment from the water. All are a food source for other animals. Fish and other stream-dependent creatures like water turtles and salamanders consume aquatic insects; crayfish are eaten by fish, birds, raccoons, and others; and mussels provide food and habitat to other animals.

Macroinvertebrates are good indicators of watershed health and water quality because they live in the water for all or most of their life, have limited mobility, and differ in their tolerance to amounts and types of pollution.



Photo: NPS/Nortrup. Illustration: University of Maryland, Integration & Application Network

Above: A fine-meshed D-shaped net is used to scoop up macroinvertebrates from promising stream habitat. Left: Presence of pollution-sensitive species like this mayfly (Ephemeroptera), along with stoneflies and caddisflies can indicate healthy stream condition.

Monitoring

Macroinvertebrate monitoring is part of a broader effort by the National Capital Region Network (NCRN) Inventory & Monitoring (I&M) program to assess the condition of streams and watersheds.

Long-term macroinvertebrate monitoring at thirty-seven park sites throughout the NCRN began in 2008 and follows a set, six-year rotation. Each spring 5-8 sites are visited. At Monocacy monitoring is done on Gambrill Creek and Bush Creek. Macroinvertebrate monitoring is co-located with fish monitoring and stream physical habitat analysis.

The objectives of this combined monitoring are to:

- determine current conditions and track long-term trends in stream condition,
- determine trends in species composition and functional groups of fish and benthic invertebrates

At each site, monitoring teams work within a 75 meter stream segment. Organisms are dislodged from available riffles and other productive stream habitats and captured in a fine-meshed, D-shaped net (see photo). A subsample of 100 individuals are preserved and identified to genus if possible, using stereoscopes (low-magnification microscopes). Chironomidae (chironomids or non-biting midges) are slide-mounted and identified using compound microscopes.

BIBI Scores

The species and number of macroinvertebrates present in a stream segment are used to calculate a Benthic Index of Biotic Integrity (BIBI) score. Scoring takes into account factors such as:

- number and diversity of taxonomic groups
- the proportion of individuals belonging to specific taxonomic groups
- ability of macroinvertebrates present to survive short- and long-term exposure to stressors from pollution, changes in stream flow, or habitat degradation
- mode of feeding including scrapers, predators, collectors, filterers, and shredders
- habit tendencies including burrowers, climbers, clingers, and sprawlers

Scores also take into account Monocacy's location in the Eastern Piedmont physiographic province. BIBI scores range from 1 to 5 and are ranked as follows: 1.0-1.9 (very poor), 2.0-2.9 (poor), 3.0-3.9 (fair), and 4.0-5.0 (good). For results, see reverse.

Published May 2012. Revised June 2015.

Monocacy National Battlefield

National Park Service
U.S. Department of the Interior

Results

Gambrill Creek and Bush Creek were sampled in 2010. BIBI scores indicate poor stream conditions.

No exotic crayfish or mussel species were found during 2010 sampling.

Gambrill Creek (MONO-217-N-2010)

2010 BIBI = 1.67 (very poor)

Number of taxa found: 33

Most common taxa: *Tvetenia* (22), *Parametriocnemus* (15), *Nais* (13), *Gammarus* (13), and *Micropsectra* (11)

Bush Creek (MONO-316-N-2010)

2010 BIBI = 2.33 (poor)

Number of taxa found: 25

Most common taxa: *Nais* (34), *Baetis* (25), *Orthocladius* (9), *Cricotopus* (8), and *Polypedilum* (7)

Discussion

Poor BIBI scores for Monocacy's streams stem from the relative absence of pollution- and disturbance-sensitive macroinvertebrates such as Ephemeroptera (mayflies), Plecoptera (stoneflies), and Trichoptera (caddisflies). Sites also had moderate amounts of pollution-tolerant Chironomid taxa.

Taxa Explained

Baetis- genus of mayflies in the family Baetidae

Cricotopus – genus of non-biting midges in the subfamily Orthocladiinae of the bloodworm family Chironomidae




Illustration: Univ. of MD, Integration & Application Network

Gammarus - genus of amphipod crustaceans

Micrispectra - genus of non-biting midges

Nais - subgroup of pollution-tolerant Oligochaetes (worms)

Orthocladius- same description as Cricotopus

Parametriocnemus - genus of non-biting midges in the subfamily Orthocladiinae of the bloodworm family Chironomidae

Polypedilum - a genus of non-biting midges in the subfamily Chironominae of the bloodworm family Chironomidae

Tvetenia- a genus of non-biting midges of the bloodworm family Chironomidae

References:

National Capital Region Network Biological Stream Survey Macroinvertebrate Data (2008-2010). Versar, Inc. National Capital Region Inventory and Monitoring Program, Washington, DC. Generic Dataset-2175665. https://irma.nps.gov/App/Reference/Profile/2175665

NCRN Macroinvertebrate webpage. http://science.nature.nps.gov/im/units/ncrn/monitoring/stream_survey/index.cfm

Norris, M. E., and G. Sanders. 2009. National Capital Region Network biological stream survey protocol: Physical habitat, fish, and aquatic macroinvertebrate vital signs. Natural Resource Report NPS/NCRN/NRR—2009/116. National Park Service, Fort Collins, Colorado. https://irma.nps.gov/App/Reference/Profile/662917