




Landbird Monitoring in the Northern Colorado Plateau Network

2011 Field Season

Natural Resource Technical Report NPS/NCPN/NRTR—2012/596



ON THE COVER

Clockwise from top: lazuli bunting (Bill Schmoker), black-chinned sparrow (Gerry Dewaghe), Canyonlands National Park (Jeff Birek). Used with permission.

Landbird Monitoring in the Northern Colorado Plateau Network

2011 Field Season

Natural Resource Technical Report NPS/NCPN/NRTR—2012/596

Prepared by

Jora Fogg
Rocky Mountain Bird Observatory
P.O. Box 1232
Brighton, CO 80603

Editing and Design

Alice Wondrak Biel
Northern Colorado Plateau Network
National Park Service
P.O. Box 848
Moab, UT 84532

June 2012

U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science
Fort Collins, Colorado

The National Park Service, Natural Resource Stewardship and Science office in Fort Collins, Colorado, publishes a range of reports that address natural resource topics of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Technical Report Series is used to disseminate results of scientific studies in the physical, biological, and social sciences for both the advancement of science and the achievement of the National Park Service mission. The series provides contributors with a forum for displaying comprehensive data that are often deleted from journals because of page limitations.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner. This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available from the Northern Colorado Plateau Network website, <http://www.nature.nps.gov/im/units/ncpn>, as well as at the Natural Resource Publications Management web site, <http://www.nature.nps.gov/publications/nrpm>.

Please cite this publication as:

Fogg, J. A. 2012. Landbird monitoring in the Northern Colorado Plateau Network: 2011 field season. Natural Resource Technical Report NPS/NCPN/NRTR—2012/596. National Park Service, Fort Collins, Colorado.

Contents

Figures v

Tables vii

Executive Summary ix

Author Information xi

Acronyms xiii

Acknowledgements xv

1 Introduction 1

 1.1 Program history 1

 1.2 Reasons for monitoring 1

 1.3 Monitoring objectives 1

2 Methods 3

 2.1 Study area 3

 2.2 Field personnel 3

 2.3 Site selection 3

 2.4 Sampling design 7

 2.5 Data analysis 7

3 Results 9

 3.1 Low-elevation riparian 9

 3.2 Pinyon-juniper 20

 3.2 Sage shrubland 32

 3.3 Pipe Spring National Monument 32

4 Discussion 43

 4.1 Status and trends 43

 4.2 Knowledge-sharing 43

 4.3 Pipe Spring National Monument 44

5 Literature Cited 45

Appendices 47

 Appendix A. Total bird species observed during surveys in the Northern Colorado Plateau Network by habitat (2011) and year (2005–2011). 47

 Appendix B. Priority species observed on transects in the Northern Colorado Plateau Network, 2005–2011 52

Figures

Figure 2-1. Transect locations by habitat, Utah units..... 4

Figure 2-2. Transect locations by habitat, Colorado units..... 5

Figure 2-3. Transect locations, Fossil Butte National Monument. 6

Figure 2-4. Point-count and area-inventory locations, Pipe Spring National Monument. 6

Figure 3-1. Estimated densities and population trend of rock wren in NCPN low-elevation riparian habitat, 2005–2011 11

Figure 3-2. Estimated densities and population trend of Bewick’s wren in NCPN pinyon-juniper habitat, 2005–2011 21

Figure 3-3. Estimated densities and population trend of black-throated gray warbler in NCPN pinyon-juniper habitat, 2005–2011 21

Figure 3-4. Estimated densities and population trend of white-throated swift in NCPN pinyon-juniper habitat, 2005–2011 22

Figure 3-5. Estimated densities and population trend of gray vireo in NCPN pinyon-juniper habitat, 2005–2011 22

Figure 3-6. Estimated densities and population trend of plumbeous vireo in NCPN pinyon-juniper habitat, 2005–2011 23

Figure 3-7. Estimated densities and population trend of Virginia’s warbler in NCPN pinyon-juniper habitat, 2005–2011 23

Figure 3-8. Estimated densities and population trend of black-billed magpie in NCPN sage shrubland habitat, 2005–2011 33

Figure 3-9. Estimated densities and population trend of violet-green swallow in NCPN sage shrubland habitat, 2005–2011 33

Figure 3-10. Estimated densities and population trend of dusky flycatcher in NCPN sage shrubland habitat, 2005–2011 34

Figure 3-11. Estimated densities and population trend of sage thrasher in NCPN sage shrubland habitat, 2005–2011 34

Figure 3-12. Estimated densities and population trend of Say’s phoebe in NCPN sage shrubland habitat, 2005–2011 35

Tables

Table 3-1. Bird sampling periods and effort in each NCPN habitat, 2011.....	9
Table 3-2. Bird totals and averages by NCPN habitat, 2011.....	9
Table 3-3. Estimated densities per km ² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in low-elevation riparian habitat, 2005–2011.....	12
Table 3-4. Estimated densities per km ² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (% CV), and sample sizes (n) of breeding birds in pinyon-juniper habitat, 2005–2011.	24
Table 3-5. Estimated densities per km ² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (% CV), and sample sizes (n) of breeding birds in sage shrubland habitat, 2005–2011.	36

Executive Summary

In 2011, the Rocky Mountain Bird Observatory (RMBO), in cooperation with the National Park Service, completed the seventh year of a habitat-based landbird monitoring program in national parks of the Northern Colorado Plateau Network (NCPN). This program is designed to provide rigorous population-trend data on most diurnal, regularly occurring, breeding landbird species throughout the network. This information is useful for land managers and supports the National Park Service's goal of developing long-term monitoring of biological indicators for network parks. The program is also consistent with goals emphasized by the U.S. North American Bird Conservation Initiative Committee (USNABCIMS 2007).

In 2011, RMBO staff surveyed 45 transect locations within 11 NCPN units, placing 15 transect locations in each of the three habitats of interest: low-elevation riparian, pinyon-juniper, and sage shrubland. To increase sample size, all transect locations were surveyed twice during the breeding season. Using a modified monitoring design, staff also conducted four point counts two times each, as well as four area searches at Pipe Spring National Monument.

This year, RMBO completed 100% of the assigned surveys with 94 transect visits between May 7 and July 7. A total of 9,626 birds of 115 species were recorded. The monitoring crew detected 2,997 birds of 85 species in low-elevation riparian, 2,850 birds of 76 species in pinyon-juniper, and 3,493 birds of 88 species in sage shrubland habitat. At Pipe Spring National Monument, observers detected a total of 286 birds of 36 species. The crew recorded one new species (bobolink) at Dinosaur National Monument.

Density estimates were generated for species with more than 80 independent detections from 2005 to 2011. This year, density estimates were calculated for 57 species in at least one habitat. The data yielded robust density estimates (coefficient of variation (CV) < 50%) for 40 species in 2011. The RMBO staff expects to reach the target of detecting a population change of at least 3% within 30 years for these 40 species. The crew recorded 44 bird species of conservation or management concern throughout the network, and calculated density estimates for 24 of those species.

Because seven years of data have been collected, the minimum number of detections used to estimate densities was increased from 60 to 80, following recommendations by Thomas and others (2010). Making this change initially decreased the number of species with density estimates but will provide more reliable estimates over time.

A second year of trend analyses was conducted for those 25 of 44 species of conservation or management concern. Separate analyses were performed for each habitat. Significant trends were detected for 12 species. The results for several species differed from those obtained in 2010. For instance, black-billed magpie in sage shrubland habitat went from a negative trend in 2010 to a quadratic trend in 2011. Bewick's wren continued to decline in pinyon-juniper. Long-term monitoring will be necessary for interpreting population status and trends for avian species of the NCPN.

Author Information

ROCKY MOUNTAIN BIRD OBSERVATORY (RMBO)

Mission: To conserve birds and their habitats

Vision: Native bird populations are sustained in healthy ecosystems

Core Values

1. Science provides the foundation for effective bird conservation.
2. Education is critical to the success of bird conservation.
3. Stewardship of birds and their habitats is a shared responsibility.

RMBO accomplishes its mission by:

- Monitoring long-term bird population trends to provide a scientific foundation for conservation action.
- Researching bird ecology and population response to anthropogenic and natural processes to evaluate and adjust management and conservation strategies using the best available science.
- Educating people of all ages through active, experiential programs that create an awareness and appreciation for birds.
- Fostering good stewardship on private and public lands through voluntary, cooperative partnerships that create win-win situations for wildlife and people.
- Partnering with state and federal natural resource agencies, private citizens, schools, universities, and other non-governmental organizations to build synergy and consensus for bird conservation.
- Sharing the latest information on bird populations, land management and conservation practices to create informed publics.
- Delivering bird conservation at biologically relevant scales by working across political and jurisdictional boundaries in western North America.



Acronyms

AICc	Akaike's Information Criterion corrected
CV	coefficient of variation
LR	low-elevation riparian
NCPN	Northern Colorado Plateau Network
NM	national monument
NPS	National Park Service
PIF	Partners in Flight
PISP	Pipe Spring National Monument
PJ	pinyon-juniper
RMBO	Rocky Mountain Bird Observatory
SA	sage shrubland

Acknowledgements

The National Park Service funded this project through a cooperative agreement with the Rocky Mountain Bird Observatory (RMBO). This report fulfills requirements in RMBO's contracts with the National Park Service, Northern Colorado Plateau Network agreement number J2118100015/RMBCP-16.

We thank Dusty Perkins of the Northern Colorado Plateau Network for logistical assistance before, during, and after the field season. We thank the superintendents, resource managers, and biologists in the individual parks for providing us with research permits, allowing us access into the backcountry, and assisting with logistics. Thank you to our 2011 crew of field biologists—Glenn Giroir, Scott Schuette, and Leah Smith—who spent many weeks in the field, sometimes under difficult conditions, conducting transects and collecting data. We thank our IT staff for their work on the RMBO database and data entry and management system, Jennifer Blakesley for providing her expertise in statistical analysis, Nick Van Lanen for conducting the trend analyses in SAS, and RMBO staff for their careful review of this report.

1 Introduction

1.1 Program history

In 2011, the Rocky Mountain Bird Observatory (RMBO), in cooperation with its partner, the National Park Service (NPS), completed the seventh year of a habitat-based landbird monitoring program in national parks of the Northern Colorado Plateau Network (NCPN). This program is designed to provide rigorous population trend data on most diurnal, regularly occurring, breeding landbird species in 11 national parks in Colorado, Utah, and Wyoming. We also sampled Pipe Spring National Monument (PISP), Arizona for the third year, using a modified monitoring design. In addition to monitoring landbird populations, this program generates information useful for managing birds, such as annual density estimates, habitat associations, and spatial distribution. It also supports the NCPN's efforts to develop long-term natural resource monitoring programs for its park units. Modeled after our Colorado habitat-based monitoring program (Leukering et al. 2000), the NCPN monitoring effort is consistent with goals emphasized by the U.S. North American Bird Conservation Initiative Committee (USNABCIMS 2007).

1.2 Reasons for monitoring

Monitoring is an essential component of wildlife management and conservation science (Witmer 2005; Marsh and Trenham 2008). Common goals of population monitoring are to estimate the population status of target species and detect changes in populations over time (Thompson et al. 1998; Sauer and Knutson 2008). Effective monitoring programs can identify species that are at risk due to small or declining populations (Dreitz et al. 2006), provide an understanding of how management actions affect populations (Alexander et al. 2008; Lyons et al. 2008), evaluate population responses to landscape alteration and climate change (Baron et al. 2008; Lindenmayer and Likens 2009), and provide basic information on species distributions.

The apparent large-scale declines of avian populations at regional and continental levels, and the loss, fragmentation, and degradation of native habitats, highlight the need for extensive and rigorous landbird moni-

toring programs (Rich et al. 2004; USNAB-CIC 2009). Population monitoring helps to achieve the intent of legislation such as the Migratory Bird Treaty Act (1918), National Environmental Policy Act (1969), Endangered Species Act (1973), National Forest Management Act (1976), and various state laws (Manley et al. 1993; Sauer 1993).

1.3 Monitoring objectives

The NCPN/RMBO landbird monitoring program uses the Partners in Flight (PIF) plan (Rich et al. 2004) as a guideline for bird conservation. PIF is a partnership of federal and state agencies, industry, non-governmental organizations, and many others, with the goal of conserving North American birds. In 1991, PIF began developing a formal species-assessment process that could provide consistent scientific evaluations of conservation status across all bird species in North America and identify the most important focus areas for the conservation of each species. This process applies quantitative rule sets to data on population size, distribution, trends, threats, and regional abundance of birds to rank species in terms of biological vulnerability and regional status. The process results in global and regional conservation assessments of each bird species that can be used to objectively assign regional and continental conservation priorities among birds.

The NCPN/RMBO landbird monitoring program is designed to provide population status and trend information for regularly occurring breeding landbird species within low-elevation riparian, pinyon-juniper, and sage shrubland habitats, and to provide “early-warning” information for all monitored species through a habitat-based approach to data collection. We calculated an 80% probability of detecting a minimum population change rate of 3.0% per year in 30 years, with a Type I error rate of 10% to provide long-term status and trend estimates for landbirds in those habitats. We anticipate that these data will prompt additional research to determine possible reasons for observed changes and enable better-informed management decisions.

RMBO maintains a high-quality online database of raw and summarized data that is accessible to collaborators and the public.

We will use these data to generate decision support tools, such as population estimate models, to help guide conservation efforts

and provide a better measure of conservation success.

2 Methods

2.1 Study area

In 2005, the NPS selected three habitats in which to implement landbird monitoring in the NCPN: low-elevation riparian (LR), pinyon-juniper (PJ), and sage shrubland (SA). A panel of NPS resource managers selected these habitats because they represent distinct bird communities and are each associated with park management questions. During the spring and summer of 2005, RMBO staff established 45 transect locations (15 in each habitat, Figures 2-1, 2-2, 2-3). In 2009, four point-count locations and area inventories were added at Pipe Spring National Monument (Figure 2-4).

2.1.1 Low-elevation riparian

Low-elevation riparian (LR) habitat comprises mostly scattered stands of Fremont cottonwood (*Populus fremontii*) and box-elder (*Acer negundo*) along perennial streams, sometimes within deeply-cut canyons. Tamarisk (*Tamarix* spp.), also known as saltcedar, is an exotic species that has invaded much of the LR habitat of the western United States. While the NPS is working to eradicate tamarisk in many park units, it is still fairly common in this habitat type.

2.1.2 Pinyon-juniper

Pinyon-juniper (PJ) habitat typically occurs at elevations just above semidesert shrubland habitat, typically above 1,500 m. PJ is present on most of the ridges and mesas, and is the most extensive habitat, in the NCPN. Pinyon pine (*Pinus edulis*) and juniper (*Juniperus* spp.) are the dominant species in this habitat, and depending on the site may have a significant sage component. Their relative abundance and composition vary significantly in this habitat type.

2.1.3 Sage shrubland

The sagebrush shrubland (sage shrubland; SA) community occurs extensively on the Colorado Plateau. The stands of sage surveyed in the NCPN are generally narrow “fingers” of pure sage, and the point-count stations are often near forests. The most

common species of sagebrush in the NCPN are big sagebrush (*Artemisia tridentata*) and mountain sagebrush (*Artemisia frigida*).

2.1.4 Pipe Spring National Monument

There are four point-count stations in a mixture of pinyon-juniper, low-elevation riparian, and semidesert shrubland habitats in Pipe Spring National Monument (NM). Semidesert shrubland habitats are dry landscapes that contain shrubs but lack a co-dominant grass component. Dominant shrubs may include sagebrush, greasewood, barberry, and saltbush. The ground-cover layer is typically dominated by bare ground and rock, with limited forbs and grasses present.

2.2 Field personnel

In 2011, the RMBO field staff consisted of three experienced biologists with excellent aural and visual bird-identification skills. All three had conducted surveys for RMBO in past years. Biologists completed a five-day training program at the beginning of the season to ensure full understanding of the field protocol.

2.3 Site selection

The NPS and RMBO selected survey sites for LR, PJ, and SA during winter 2005. For LR survey sites, the options were limited to crossable streams, excluding the Colorado, Green, Gunnison, and Virgin rivers. Because there is a limited amount of riparian habitat, survey locations were manually selected; RMBO staff ground-proofed riparian stands and established transects in 2005. During ground-proofing, a few of the stands were found not to fit the selection criteria. In these cases, replacement stands were chosen following the same conditions. These same locations have been surveyed every season since 2005.

For PJ and SA habitat, GIS and the Southwest Regional Re-GAP Analysis Project (Lowry et al. 2005) were used to randomly select sites from a pool of habitat “stands” that were large enough to accommodate transects. Areas with >50% slope were excluded from the list of potential sites to ensure that selected stands could be accessed safely on foot.




Figure 2-1. Transect locations by habitat, Utah units.




Figure 2-2. Transect locations by habitat, Colorado units.




Figure 2-3. Transect locations, Fossil Butte National Monument.




Figure 2-4. Point-count and area-inventory locations, Pipe Spring National Monument.

Point-count and area-search locations at Pipe Spring NM were placed in each distinct habitat represented in the monument as previously described, spaced at least 250 m apart. Area-search start locations were selected in order to efficiently cover all areas of the monument.

2.4 Sampling design

RMBO staff sampled landbird populations in each habitat following the protocol established by Leukering (rev. 2005) and modified by Hanni and others (2011). All transects were surveyed between one half-hour before sunrise and 11AM. Up to 15 five-minute point counts were conducted at stations located at 250-m intervals along each transect. In order to increase the sample size, the crew surveyed each of the 45 transect locations twice during the summer, with each visit occurring on a separate day. At each point, all birds detected during the five-minute point count were recorded. For every bird detected during a point count, the crew recorded species, sex, horizontal distance from the observer, the minute each bird was detected, and the type of detection (e.g., call, song or visual).

Observers measured horizontal distances to each bird using laser rangefinders. When it was not possible to measure distance to a bird, observers estimated distance by measuring to some nearby object. Observers also recorded birds flying over but not using the immediate surrounding landscape. For distribution mapping purposes, observers recorded the presence of all rare or difficult-to-detect species encountered while traveling between points (e.g., woodpeckers, owls, raptors). All non-independent detections of birds were considered to be part of a “cluster” (i.e., flocks or pairs of conspecific birds together in close proximity) rather than separate independent observations.

At the start and end of each transect, observers recorded time, temperature, percent cloud cover, and precipitation type, and estimated the Beaufort scale wind-speed category. The crew measured distances between points using hand-held GPS units and recorded all GPS data in Universal Transverse Mercator (UTM) North American Datum 1983 (NAD 83). At each point, the crew recorded UTM coordinates, habitat informa-

tion (within a 50-m radius), and distance from a road (if within 100 m). Habitat information included the structural stage as well as types, relative abundance, percent coverage, and mean height of trees, shrubs, and groundcover. If there was a distinct tree subcanopy present, the tree species was/were recorded. All of these data were recorded prior to beginning each point count.

At Pipe Spring NM, observers used the protocol described above for the point-count portion of surveys. This year, monitoring was further refined to include diurnal and nocturnal area searches, because the small size of the monument makes it feasible to record species that may be missed by traditional point-count methods, as well as breeding data. A total of four 20-minute area searches of the entire monument were conducted this year. Two diurnal-area searches were conducted upon completion of the four point counts on separate dates, and two nocturnal-area searches covering the same areas were conducted beginning at dusk and concluding after 20 minutes. The observer continuously walked the monument, tallying all individual birds heard and seen along with information on any breeding or flocking behavior. The observer also recorded the UTM coordinates of start locations, start and end times, and weather information.

2.5 Data analysis

2.5.1 Distance analysis

RMBO staff used the analysis software Distance 6.0 (Thomas et al. 2010) to estimate detection probabilities using the point-count data. Analysis of distance data is accomplished by fitting a detection function to the distribution of recorded distances. The distribution of distances can be a function of characteristics of the object being recorded (e.g., a bird’s size, color, movement, volume, or frequency of song), the sampling landscape (e.g., density of vegetation), and observer ability. Because the detection function is unique to each species in each habitat, Distance analysis avoids some serious problems inherent in traditional analyses of point-count data (e.g., unquantifiable differences in detectability among habitats, species, and years). Distance analysis relies on three assumptions, all of which are reasonably well

met by this program: (1) all birds at the point (distance = 0 m) are detected, (2) distances of birds close to the point are measured accurately, and (3) birds do not move in response to the observer's presence.

RMBO staff estimated densities of species with at least 80 independent detections within a habitat across all years. Birds flying over and not using the immediate surrounding landscape, and birds detected between points, were excluded from analyses. For each species, the staff fit a half normal key function, uniform function, and hazard rate key function with cosine series expansions and a hazard rate key function with simple polynomial series expansion to the distribution of distances (Buckland et al. 2001). We used Akaike's Information Criterion corrected (AICc) for small sample size and model-selection theory to select the most parsimonious detection function for each species (Burnham and Anderson 2002). We estimated variance in Distance 6.0 using bootstrapping of transects within strata rather than empirical estimates. Empirical methods tend to underestimate variance from small sample sizes.

Unless otherwise specified, all bird species names listed in this report are from the American Ornithologists' Union Check-list of North American Birds (AOU 2007).

2.5.2 Trend detection

Using weighted regression and Information-Theoretic model selection (Burnham and Anderson 2002), RMBO staff modeled observed trends in populations of all species of conservation concern (Appendix B) in each habitat using data from all years (2005–2011). For each species, staff modeled four different functions using Proc REG in program SAS (SAS Institute 2007): no trend (intercept-only model), linear trend, quadratic trend, and log-linear (pseudo-threshold) trend. For small sample sizes, AICc was used to select the most parsimonious trend model for each species (Burnham and Anderson 2002). Input data were density estimates and their variances, with the inverse of the coefficient of variation (CV) used as a variable weight (giving more weight to more precise estimates). Confidence limits of the beta estimate (β_{trend}) were used to determine significance of the trend. Trend detection was also performed in 2010, but the number of detections used in analysis changed in 2011 due to new truncations used in Distance analyses.

3 Results

In 2011, the seventh year of RMBO land-bird monitoring in the NCPN, 1,233 point counts were conducted along 45 transects in three habitats between May 7 and July 7, 2011 (Table 3-1). All transect locations were surveyed twice. Ninety-one percent of the point counts were completed (1,233 out of 1,350). Some point counts were not completed because of weather (wind, rain or heat: 76 points), poor GPS reception along the canyons of the low-elevation riparian transects (17 points), and rivers that could not be safely crossed (24 points).

Excluding between-point detections of low-density species, observers recorded 9,626 birds of 115 species (Table 3-2, Appendix A). We detected 2,997 birds of 85 species in low-elevation riparian, 2,850 birds of 76 species in pinyon-juniper, and 3,493 birds of 88 species in sage shrubland habitat. One new species, a female bobolink, was recorded at Dinosaur NM on May 24. This bird was likely a migrant.

A total of 286 birds of 36 species were detected during surveys in Pipe Spring NM. Observers recorded 119 birds of 28 species during point counts. During the nocturnal- and diurnal-area searches of the monument, 170 birds and five additional species that had not been previously detected were recorded: black phoebe (confirmed nesting), Bewick's wren, lazuli bunting, common poorwill, and great horned owl.

This year, density estimates were calculated for 57 species in at least one habitat. These 57 species represent 50% of species detected on transects in the NCPN during 2011, and 91% of birds observed on transects during 2011. The habitat-stratified data yielded robust density estimates (CV <50%) for 40 species.

RMBO recorded 44 bird species that are of conservation and management concern (priority species) throughout the NCPN (Appendix B). Density estimates were calculated for 24 of these species.

3.2 Low-elevation riparian

All 15 LR transect locations were surveyed twice this year. Density estimates were calcu-

Table 3-1. Bird sampling periods and effort in each NCPN habitat, 2011.

Habitat	Dates sampled	Transects*	Point counts
Low-elevation riparian	May 10–June 30	15	408
Pinyon-juniper	May 07–July 1	15	390
Sage shrubland	May 14–July 7	15	435
All habitats	May 07– July 7	45	1,233

*Technicians attempted to survey all transect locations twice.

Table 3-2. Bird totals and averages by NCPN habitat, 2011.

Habitat	Birds detected	Birds per transect (avg)	Species detected	Species per transect (avg)
Low-elevation riparian	2,997	206	85	30
Pinyon-juniper	2,850	195	76	27
Sage shrubland	3,493	241	88	27
All habitats	9,340	214	115	28

lated for 37 species, of which 14 were priority species, from 2005 to 2011. Density estimates for 2011 could not be calculated for seven of the 37 species because of small sample size. The pooled data for 2005–2011 yielded robust density estimates (CV <50%) for 24 species and a moderately robust estimate (CV = 50–75%) for six additional species (Table 3-3). RMBO expects to reach the target of detecting population changes of at least 3% within 30 years for these 30 species, which represent 44% of species detected and 91% of birds detected in LR habitat.

Estimated densities recorded in LR were highest for the following 10 species in 2011 (listed in order from highest to lowest density):

1. Blue-gray gnatcatcher
2. Black-chinned hummingbird
3. Spotted towhee
4. Yellow warbler
5. Lazuli bunting
6. Ash-throated flycatcher
7. Yellow-rumped warbler
8. House finch
9. Lesser goldfinch
10. Chipping sparrow

For the following 18 species, estimated densities were higher in LR than in the other two habitats sampled in 2011 (listed in order from highest to lowest density):

1. Blue-gray gnatcatcher
2. Black-chinned hummingbird
3. Spotted towhee
4. Yellow warbler
5. Lazuli bunting
6. Ash-throated flycatcher
7. Yellow-rumped warbler
8. House finch
9. Lesser goldfinch
10. White-throated swift
11. House wren
12. Song sparrow
13. Brown-headed cowbird
14. Yellow-breasted chat
15. Say's phoebe
16. Warbling vireo
17. Common raven
18. Western wood-pewee

3.1.1 Trend detection

Rock wren ($\beta_{\text{trend}} = -0.46$; 95% CI = -0.22, -0.71) showed evidence of significant population change over the last seven years because the confidence limits around the beta estimate do not contain zero. The best approximating model for rock wren was a quadratic trend (Figure 3-1). The best model for all other species of concern with density estimates in LR habitat was the intercept-only (constant) model, indicating no significant change in population estimates over time.




Figure 3-1. Estimated densities and population trend of rock wren in NCPN low-elevation riparian habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.

Table 3-3. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in low-elevation riparian habitat, 2005–2011.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
American robin	2005	5.33	2.57	11.06	46	15
	2006	7.38	3.72	14.65	43	23
	2007	7.41	3.90	14.10	40	24
	2008	8.29	4.48	15.31	38	26
	2009	--	--	--	--	4
	2010	5.46	2.01	14.88	66	16
	2011	--	--	--	--	9
Ash-throated flycatcher	2005	15.13	10.85	21.10	20	153
	2006	23.51	17.86	30.96	17	239
	2007	20.01	15.09	26.54	17	207
	2008	18.37	13.35	25.27	19	191
	2009	17.20	12.43	23.80	20	141
	2010	21.58	15.54	29.97	20	192
	2011	23.86	17.66	32.22	18	225
Bewick's wren	2005	7.02	3.63	13.55	42	44
	2006	15.52	9.08	26.52	33	100
	2007	8.54	4.32	16.86	43	56
	2008	8.43	4.52	15.71	39	57
	2009	15.37	8.83	26.77	35	85
	2010	3.58	1.91	6.72	40	20
	2011	2.28	1.17	4.45	42	14
Black phoebe	2005	5.00	1.59	15.73	76	10
	2006	4.45	1.37	14.44	81	10
	2007	8.58	2.62	28.12	82	19
	2008	--	--	--	--	9
	2009	8.35	3.17	21.99	64	20
	2010	--	--	--	--	2
	2011	--	--	--	--	2
Black-chinned hummingbird	2005	149.79	94.18	238.25	29	28
	2006	185.35	107.92	318.34	34	35
	2007	113.93	68.48	189.55	32	22
	2008	116.51	70.30	193.11	31	23
	2009	119.37	69.05	206.35	34	20
	2010	74.48	43.88	126.40	33	13
	2011	97.83	58.78	162.81	32	18

Table 3-3. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in low-elevation riparian habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Black-headed grosbeak	2005	3.74	1.22	11.46	76	11
	2006	7.14	4.03	12.64	35	22
	2007	3.98	1.76	8.99	52	12
	2008	5.36	2.80	10.26	41	17
	2009	6.65	2.53	17.49	64	17
	2010	--	--	--	--	9
	2011	4.27	2.13	8.58	44	12
Black-throated gray warbler	2005	7.10	3.76	13.42	40	38
	2006	14.09	8.55	23.21	31	76
	2007	9.93	6.66	14.81	25	54
	2008	19.70	10.26	37.84	41	106
	2009	10.21	4.42	23.57	54	45
	2010	10.99	6.62	18.24	32	52
	2011	14.13	7.97	25.04	36	70
Black-throated sparrow	2005	5.63	2.62	12.12	49	49
	2006	5.98	3.21	11.12	39	51
	2007	6.68	2.60	17.17	62	62
	2008	7.46	3.68	15.12	45	67
	2009	20.13	9.17	44.18	50	138
	2010	8.15	4.22	15.77	42	60
	2011	11.52	6.21	21.37	39	96
Blue-gray gnatcatcher	2005	57.05	42.17	77.17	18	112
	2006	74.57	51.21	108.59	22	149
	2007	55.82	40.41	77.11	19	115
	2008	66.76	44.46	100.23	24	136
	2009	144.63	103.63	201.87	20	230
	2010	93.20	68.64	126.55	18	166
	2011	105.95	82.97	135.30	14	196
Brown-headed cowbird	2005	--	--	--	--	7
	2006	9.37	4.82	18.24	42	24
	2007	6.51	2.84	14.88	53	17
	2008	5.44	2.62	11.30	46	14
	2009	7.76	2.96	20.34	63	15
	2010	--	--	--	--	5
	2011	9.93	4.09	24.12	58	20

Table 3-3. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in low-elevation riparian habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Bullock's oriole	2005	--	--	--	--	3
	2006	8.66	5.19	14.47	32	22
	2007	6.19	2.72	14.10	53	16
	2008	--	--	--	--	9
	2009	5.89	2.04	17.01	70	12
	2010	--	--	--	--	9
	2011	--	--	--	--	9
Bushtit	2005	36.32	20.12	65.59	36	17
	2006	61.16	37.77	99.03	30	55
	2007	8.30	3.70	18.62	52	11
	2008	--	--	--	--	5
	2009	12.89	5.99	27.74	48	14
	2010	--	--	--	--	0
	2011	--	--	--	--	0
Canyon wren	2005	1.87	1.06	3.28	35	30
	2006	4.04	2.41	6.77	32	63
	2007	2.25	1.23	4.13	38	37
	2008	1.66	0.69	3.97	57	27
	2009	1.65	0.78	3.46	47	23
	2010	--	--	--	--	1
	2011	--	--	--	--	9
Chipping sparrow	2005	--	--	--	--	6
	2006	--	--	--	--	8
	2007	15.51	7.67	31.34	45	29
	2008	14.88	7.22	30.66	46	28
	2009	18.91	9.12	39.21	46	24
	2010	16.69	7.62	36.55	50	28
	2011	16.99	8.82	32.70	40	21
Common raven	2005	2.65	1.48	4.74	36	16
	2006	1.82	0.89	3.71	45	15
	2007	1.98	0.97	4.03	45	14
	2008	4.58	2.16	9.72	48	37
	2009	--	--	--	--	8
	2010	4.05	1.98	8.28	45	27
	2011	2.60	1.53	4.41	33	19

Table 3-3. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in low-elevation riparian habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Common yellowthroat	2005	--	--	--	--	9
	2006	13.45	5.81	31.15	54	19
	2007	14.40	5.09	40.75	69	19
	2008	--	--	--	--	4
	2009	15.14	5.72	40.04	64	18
	2010	8.17	2.61	25.63	78	10
	2011	--	--	--	--	9
Gray vireo	2005	1.09	0.60	2.00	38	17
	2006	2.33	1.15	4.76	45	36
	2007	1.14	0.49	2.66	54	18
	2008	1.54	0.83	2.88	39	24
	2009	2.61	1.58	4.32	31	35
	2010	--	--	--	--	4
	2011	--	--	--	--	9
House finch	2005	18.47	12.80	26.66	22	109
	2006	26.63	19.29	36.76	19	178
	2007	27.88	19.28	40.31	22	189
	2008	28.26	20.35	39.25	19	192
	2009	31.75	18.63	54.12	32	168
	2010	18.60	12.67	27.31	23	112
	2011	22.77	16.98	30.54	18	130
House wren	2005	11.62	5.01	26.94	54	40
	2006	11.44	4.66	28.05	58	42
	2007	8.29	3.39	20.27	58	30
	2008	12.64	5.08	31.41	59	47
	2009	13.10	5.27	32.54	59	38
	2010	9.13	2.39	34.97	95	27
	2011	13.01	5.14	32.94	59	45
Juniper titmouse	2005	8.24	4.12	16.50	44	28
	2006	7.11	3.89	13.01	38	24
	2007	10.69	6.54	17.48	30	42
	2008	10.87	6.37	18.55	33	41
	2009	10.15	4.69	21.93	49	28
	2010	5.06	2.57	9.96	43	17
	2011	6.00	3.21	11.23	39	22

Table 3-3. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in low-elevation riparian habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Lazuli bunting	2005	53.78	25.90	111.71	45	151
	2006	47.95	22.74	101.11	45	132
	2007	54.71	30.19	99.13	36	159
	2008	52.33	23.57	116.17	49	148
	2009	62.24	22.60	171.42	64	149
	2010	58.74	28.01	123.18	45	144
	2011	38.18	23.76	61.35	29	101
Lesser goldfinch	2005	12.53	5.64	27.84	51	25
	2006	23.18	12.02	44.69	41	46
	2007	42.76	22.72	80.48	40	91
	2008	29.51	12.14	71.71	57	54
	2009	40.50	20.71	79.20	42	75
	2010	14.92	7.39	30.13	45	31
	2011	17.70	8.93	35.09	43	33
Mourning dove	2005	9.58	5.39	17.04	36	73
	2006	16.27	10.83	24.43	25	114
	2007	18.58	12.28	28.12	25	140
	2008	10.96	6.00	20.03	38	91
	2009	5.38	3.27	8.86	31	35
	2010	15.03	9.55	23.65	28	96
	2011	10.69	6.42	17.81	32	73
Plumbeous vireo	2005	12.57	7.36	21.48	33	50
	2006	20.05	11.42	35.20	35	81
	2007	15.23	8.86	26.19	34	62
	2008	10.61	6.42	17.54	31	44
	2009	18.41	10.76	31.50	33	65
	2010	19.52	11.64	32.75	32	71
	2011	8.53	4.95	14.69	34	32
Rock wren	2005	6.26	3.58	10.95	34	75
	2006	10.44	7.00	15.56	24	104
	2007	9.85	7.15	13.58	20	135
	2008	9.48	5.06	17.79	39	80
	2009	7.62	5.28	10.99	22	93
	2010	5.06	3.24	7.91	27	57
	2011	3.61	2.39	5.46	24	97

Table 3-3. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in low-elevation riparian habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Say's phoebe	2005	3.41	2.22	5.26	27	41
	2006	5.10	3.00	8.68	33	61
	2007	2.99	1.91	4.67	28	37
	2008	4.23	2.25	7.97	40	51
	2009	5.28	2.73	10.23	42	51
	2010	2.41	1.29	4.52	40	25
	2011	5.75	3.35	9.85	34	64
Song sparrow	2005	19.90	7.54	52.56	64	45
	2006	22.14	9.41	52.07	56	56
	2007	18.00	8.28	39.11	50	49
	2008	16.15	8.01	32.57	45	42
	2009	14.27	5.25	38.84	67	28
	2010	3.44	0.86	13.74	101	8
	2011	11.40	4.71	27.58	58	27
Spotted towhee	2005	48.32	36.48	64.02	17	225
	2006	86.99	66.67	113.51	16	413
	2007	57.69	41.58	80.06	20	274
	2008	42.90	32.53	56.58	17	208
	2009	76.65	56.22	104.51	19	305
	2010	67.93	50.03	92.23	18	281
	2011	72.80	56.18	94.34	16	316
Violet-green swallow	2005	55.02	34.00	89.03	30	82
	2006	193.15	122.24	305.21	28	198
	2007	115.76	74.53	179.81	27	122
	2008	128.42	72.37	227.87	36	108
	2009	200.53	122.79	327.48	30	138
	2010	59.18	31.43	111.41	40	87
	2011	15.27	9.55	24.41	29	52
Virginia's warbler	2005	7.94	4.30	14.64	38	23
	2006	13.90	5.79	33.41	57	41
	2007	9.90	5.70	17.18	34	31
	2008	17.96	10.95	29.48	31	58
	2009	--	--	--	--	2
	2010	8.89	4.77	16.58	39	23
	2011	4.59	2.42	8.70	40	13

Table 3-3. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in low-elevation riparian habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Warbling vireo	2005	5.38	2.25	12.88	57	17
	2006	11.26	4.89	25.95	54	37
	2007	7.75	3.89	15.44	44	27
	2008	8.76	5.46	14.06	29	30
	2009	6.87	3.31	14.24	46	18
	2010	6.44	1.80	23.09	90	20
	2011	4.33	1.88	9.99	54	13
Western scrub-jay	2005	4.84	2.41	9.74	44	18
	2006	4.77	2.34	9.73	45	25
	2007	4.94	2.35	10.37	47	22
	2008	4.25	1.53	11.78	68	22
	2009	2.42	1.00	5.83	57	11
	2010	3.55	1.49	8.46	56	14
	2011	3.95	2.07	7.54	41	20
Western wood-pewee	2005	3.83	2.14	6.87	36	27
	2006	4.58	2.28	9.23	44	33
	2007	4.62	2.17	9.87	48	34
	2008	4.45	1.70	11.66	64	31
	2009	4.53	2.09	9.80	49	28
	2010	2.19	0.90	5.32	58	14
	2011	2.40	1.26	4.56	40	16
White-throated swift	2005	30.65	19.94	47.13	27	103
	2006	50.83	32.29	80.03	28	127
	2007	63.27	40.59	98.62	27	190
	2008	36.61	21.29	62.96	34	86
	2009	15.56	9.10	26.61	33	73
	2010	11.22	5.87	21.43	40	50
	2011	15.25	8.44	27.56	37	46
Yellow-breasted chat	2005	6.80	2.04	22.66	84	37
	2006	8.42	3.07	23.12	67	51
	2007	6.57	2.17	19.89	75	42
	2008	5.48	2.58	11.61	48	32
	2009	8.22	4.21	16.02	42	43
	2010	9.52	4.76	19.05	44	50
	2011	7.90	3.38	18.50	55	45

Table 3-3. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in low-elevation riparian habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Yellow-rumped warbler	2005	--	--	--	--	4
	2006	--	--	--	--	2
	2007	--	--	--	--	3
	2008	--	--	--	--	8
	2009	--	--	--	--	3
	2010	27.57	15.51	49.00	36	43
	2011	23.31	8.98	60.54	62	26
Yellow warbler	2005	62.28	31.43	123.43	41	131
	2006	64.04	31.12	131.79	44	135
	2007	52.70	24.41	113.76	47	113
	2008	67.68	34.14	134.15	41	142
	2009	129.16	61.39	271.76	45	220
	2010	98.19	54.07	178.30	36	179
	2011	68.22	36.32	128.14	38	127

Dashes indicate the sample size was insufficient for estimating density. Priority species are bolded and shaded.

3.2 Pinyon-juniper

All 15 PJ transect locations were surveyed twice this year. Density estimates were calculated for 36 species, 15 of which were priority species, from 2005 to 2011. Density estimates could not be calculated for three of the 36 species specifically for 2011. The pooled data for 2005–2011 yielded robust density estimates (CV <50%) for 21 species and a moderately robust estimate (CV = 50–75%) for 9 additional species (Table 3-4). RMBO expects to reach the target of detecting population changes of at least 3% within 30 years for these 30 species, which represent 46% of species and 90% of individuals detected in PJ habitat.

Estimated densities recorded in PJ were highest for the following 10 species in 2011:

1. Blue-gray gnatcatcher
2. Black-throated gray warbler
3. Chipping sparrow
4. Spotted towhee
5. Juniper titmouse
6. Gray flycatcher
7. House finch
8. Plumbeous vireo
9. Virginia's warbler
10. Violet-green swallow

For the following 15 species, estimated densities were higher in PJ than in the other two habitats sampled in 2011 (listed in order from highest to lowest density):

1. Black-throated gray warbler
2. Chipping sparrow
3. Gray flycatcher
4. Plumbeous vireo
5. Virginia's warbler
6. Gray vireo
7. Mourning dove
8. Mountain chickadee

9. Western scrub-jay
10. Bewick's wren
11. White-breasted nuthatch
12. Pinyon jay
13. Hermit thrush
14. Grace's warbler
15. Western tanager

3.2.1 Trend detection

Significant population trends were detected for six species in PJ habitat. Bewick's wren ($\beta_{\text{trend}} = -2.75$; 95% CI = -3.68, -1.81), black-throated gray warbler ($\beta_{\text{trend}} = -8.78$; 95% CI = -16.07, -1.49), and white-throated swift ($\beta_{\text{trend}} = -2.64$; 95% CI = -4.61, -0.66) showed evidence of population change over the last seven years with a decreasing linear trend (Figure 3-2, 3-3, 3-4). Gray vireo ($\beta_{\text{trend}} = 6.88$; 95% CI = 0.48, 13.29) showed evidence of population change over the last seven years with an increasing log-linear trend (Figure 3-5). Plumbeous vireo ($\beta_{\text{trend}} = 0.39$; 95% CI = 0.15, 0.64) and Virginia's warbler ($\beta_{\text{trend}} = 0.61$; 95% CI = 0.33, 0.89) also showed evidence of significant population change with a quadratic trend (Figures 3-6, 3-7). The best approximating model for all other species of concern with density estimates in PJ habitat was the intercept-only (constant) model, which indicated no significant trend.




Figure 3-2. Estimated densities and population trend of Bewick's wren in NCPN pinyon-juniper habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.




Figure 3-3. Estimated densities and population trend of black-throated gray warbler in NCPN pinyon-juniper habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.




Figure 3-4. Estimated densities and population trend of white-throated swift in NCPN pinyon-juniper habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.




Figure 3-5. Estimated densities and population trend of gray vireo in NCPN pinyon-juniper habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.




Figure 3-6. Estimated densities and population trend of plumbeous vireo in NCPN pinyon-juniper habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.




Figure 3-7. Estimated densities and population trend of Virginia's warbler in NCPN pinyon-juniper habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.

Table 3-4. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (% CV), and sample sizes (n) of breeding birds in pinyon-juniper habitat, 2005–2011.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
American robin	2005	1.82	1.00	3.32	38	26
	2006	2.67	1.17	6.07	53	37
	2007	2.65	0.86	8.21	77	35
	2008	1.91	0.61	5.97	78	25
	2009	1.13	0.37	3.45	76	14
	2010	0.87	0.32	2.33	65	12
	2011	2.78	1.34	5.75	46	33
Ash-throated flycatcher	2005	9.94	8.38	11.80	10	121
	2006	16.44	11.33	23.87	23	125
	2007	8.78	7.33	10.53	11	117
	2008	12.03	9.92	14.58	12	116
	2009	9.65	6.94	13.43	20	104
	2010	5.54	4.17	7.37	17	82
	2011	4.79	3.92	5.85	12	83
Bewick's wren	2005	20.15	15.49	26.22	16	134
	2006	20.09	12.60	32.04	29	134
	2007	16.67	9.89	28.09	32	112
	2008	15.84	8.62	29.11	38	100
	2009	13.41	9.36	19.21	22	79
	2010	3.40	1.86	6.21	38	22
	2011	4.36	2.52	7.57	34	26
Black-throated gray warbler	2005	99.28	57.75	170.65	34	240
	2006	95.88	68.53	134.14	20	314
	2007	94.92	65.53	137.50	22	305
	2008	66.56	46.98	94.31	21	301
	2009	111.81	72.61	172.18	27	313
	2010	53.11	39.28	71.80	18	236
	2011	46.28	32.53	65.85	22	262
Black-throated sparrow	2005	5.38	2.66	10.90	45	45
	2006	8.87	4.16	18.88	48	70
	2007	12.27	5.14	29.32	56	99
	2008	9.58	3.60	25.49	64	73
	2009	11.33	5.89	21.79	41	80
	2010	8.45	4.69	15.23	37	67
	2011	10.53	5.21	21.29	44	74

Table 3-4. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in pinyon-juniper habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Blue-gray gnatcatcher	2005	74.20	51.39	107.12	22	125
	2006	56.26	34.91	90.68	30	92
	2007	76.65	53.48	109.85	22	125
	2008	64.52	42.29	98.44	26	98
	2009	148.69	115.81	190.91	15	215
	2010	85.27	60.81	119.58	21	136
	2011	91.63	71.01	118.25	16	134
Brown-headed cowbird	2005	4.49	2.16	9.30	46	16
	2006	4.08	1.82	9.14	51	13
	2007	4.69	1.93	11.39	57	16
	2008	3.12	1.25	7.79	60	10
	2009	5.63	2.92	10.86	41	17
	2010	--	--	--	--	7
	2011	--	--	--	--	9
Bushtit	2005	96.86	55.49	169.05	34	32
	2006	62.00	39.50	97.32	27	25
	2007	36.37	19.18	68.96	40	14
	2008	--	--	--	--	7
	2009	91.55	54.39	154.10	32	36
	2010	--	--	--	--	0
	2011	--	--	--	--	1
Canyon wren	2005	0.42	0.15	1.20	57	14
	2006	0.61	0.22	1.74	56	22
	2007	0.70	0.21	2.33	67	24
	2008	--	--	--	--	5
	2009	0.42	0.15	1.17	68	15
	2010	--	--	--	--	3
	2011	--	--	--	--	6
Chipping sparrow	2005	21.46	13.02	35.38	31	70
	2006	16.80	10.51	26.87	29	53
	2007	24.37	14.49	40.99	32	75
	2008	29.23	18.53	46.13	28	88
	2009	53.25	29.70	95.48	36	134
	2010	31.19	18.19	53.48	34	92
	2011	36.11	25.37	51.40	22	102

Table 3-4. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in pinyon-juniper habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Common raven	2005	0.69	0.42	1.12	30	35
	2006	0.30	0.17	0.53	36	19
	2007	1.08	0.67	1.76	30	68
	2008	0.64	0.38	1.06	32	38
	2009	--	--	--	--	27
	2010	1.07	0.65	1.75	31	66
	2011	1.12	0.71	1.79	29	59
Dusky flycatcher	2005	--	--	--	--	9
	2006	13.65	5.91	31.53	54	34
	2007	10.93	4.24	28.20	62	28
	2008	18.59	5.54	62.44	84	38
	2009	5.42	1.73	17.02	78	13
	2010	6.23	2.13	18.21	72	15
	2011	5.29	2.07	13.49	61	12
Grace's warbler	2005	--	--	--	--	6
	2006	1.40	0.35	5.55	99	12
	2007	2.39	0.58	9.79	103	19
	2008	--	--	--	--	8
	2009	2.51	0.62	10.23	102	17
	2010	1.33	0.34	5.20	98	11
	2011	2.05	0.57	7.39	91	17
Gray flycatcher	2005	24.85	16.00	38.58	27	116
	2006	26.12	16.86	40.48	27	86
	2007	24.84	13.67	45.13	37	85
	2008	37.37	22.48	62.13	31	101
	2009	34.64	21.71	55.26	29	179
	2010	23.82	17.08	33.21	20	123
	2011	15.94	10.10	25.15	28	113
Gray vireo	2005	6.42	4.08	10.11	28	67
	2006	5.74	3.11	10.61	39	59
	2007	6.72	4.06	11.14	31	70
	2008	8.91	5.30	14.98	32	86
	2009	13.15	9.63	17.94	19	125
	2010	10.60	7.41	15.16	22	108
	2011	8.84	5.67	13.78	27	81

Table 3-4. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in pinyon-juniper habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Hermit thrush	2005	--	--	--	--	8
	2006	0.70	0.15	3.15	113	13
	2007	--	--	--	--	1
	2008	--	--	--	--	0
	2009	0.55	0.15	1.97	90	11
	2010	0.70	0.18	2.70	97	12
	2011	2.19	0.61	7.85	88	41
House finch	2005	18.44	10.97	31.02	32	94
	2006	10.52	6.10	18.17	34	69
	2007	25.45	14.20	45.60	36	146
	2008	17.09	11.53	25.34	24	128
	2009	21.67	13.12	35.78	31	136
	2010	11.37	7.86	16.46	23	101
	2011	14.10	8.89	22.34	28	148
Juniper titmouse	2005	29.71	16.56	53.30	36	79
	2006	25.99	15.37	43.95	32	76
	2007	28.46	14.97	54.08	40	82
	2008	23.82	13.07	43.41	37	89
	2009	46.65	32.86	66.23	21	199
	2010	41.45	27.38	62.74	25	141
	2011	22.41	13.25	37.92	33	112
Lark sparrow	2005	2.49	1.08	5.75	54	20
	2006	2.21	0.54	9.00	102	17
	2007	2.11	0.79	5.66	65	19
	2008	1.57	0.62	3.98	61	13
	2009	1.96	0.85	4.53	54	17
	2010	1.97	0.60	6.42	81	17
	2011	3.50	1.44	8.51	58	27
Mountain bluebird	2005	9.27	3.62	23.73	62	63
	2006	4.98	1.54	16.12	81	31
	2007	5.99	2.41	14.88	59	36
	2008	5.18	2.13	12.60	58	34
	2009	6.84	2.33	20.09	72	33
	2010	6.71	2.40	18.80	69	40
	2011	5.46	2.03	14.67	65	26

Table 3-4. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in pinyon-juniper habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Mountain chickadee	2005	4.95	2.03	12.08	58	19
	2006	--	--	--	--	9
	2007	2.75	1.10	6.87	60	10
	2008	--	--	--	--	9
	2009	3.85	1.36	10.91	69	13
	2010	--	--	--	--	4
	2011	8.36	3.08	22.65	66	26
Mourning dove	2005	17.34	8.62	34.88	44	139
	2006	19.30	11.34	32.88	33	154
	2007	23.83	13.02	43.61	38	167
	2008	17.35	9.95	30.27	35	158
	2009	9.02	5.80	14.03	27	96
	2010	14.33	8.97	22.89	29	154
	2011	8.36	5.17	13.53	30	114
Pinyon jay	2005	1.85	1.04	3.32	36	40
	2006	4.35	2.15	8.79	45	66
	2007	6.21	3.49	11.07	36	93
	2008	1.57	0.65	3.78	56	22
	2009	2.25	1.27	4.00	36	41
	2010	6.99	3.87	12.63	37	103
	2011	2.88	1.33	6.26	49	36
Plumbeous vireo	2005	7.98	4.51	14.14	36	62
	2006	5.98	3.19	11.21	40	45
	2007	5.85	3.36	10.20	35	44
	2008	6.00	3.35	10.74	36	43
	2009	5.95	2.45	14.46	58	42
	2010	10.91	6.33	18.83	34	83
	2011	11.05	7.35	16.61	25	76
Rock wren	2005	4.27	2.81	6.48	26	84
	2006	4.13	2.57	6.64	29	80
	2007	4.38	2.86	6.71	26	85
	2008	5.61	3.29	9.56	33	101
	2009	5.45	3.25	9.11	32	93
	2010	2.06	1.24	3.44	32	39
	2011	4.98	3.29	7.51	25	84

Table 3-4. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in pinyon-juniper habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Say's phoebe	2005	1.70	0.92	3.14	32	28
	2006	--	--	--	--	9
	2007	--	--	--	--	8
	2008	0.68	0.34	1.36	36	11
	2009	0.89	0.48	1.63	31	14
	2010	1.12	0.47	2.70	47	19
	2011	1.09	0.48	2.45	43	17
Spotted towhee	2005	14.68	9.08	23.74	30	80
	2006	12.96	7.15	23.48	37	69
	2007	20.21	11.01	37.11	38	109
	2008	9.19	4.86	17.35	40	46
	2009	19.95	10.96	36.31	37	97
	2010	13.94	7.54	25.75	39	72
	2011	24.44	13.40	44.58	37	112
Vesper sparrow	2005	--	--	--	--	9
	2006	1.31	0.44	3.90	74	15
	2007	1.27	0.39	4.06	80	15
	2008	--	--	--	--	6
	2009	1.35	0.36	5.07	94	13
	2010	1.81	0.58	5.62	77	20
	2011	2.35	0.66	8.43	90	25
Violet-green swallow	2005	9.21	4.74	17.91	42	32
	2006	13.62	6.49	28.59	47	48
	2007	17.64	8.32	37.38	48	58
	2008	16.24	7.83	33.67	46	40
	2009	11.76	6.74	20.53	35	31
	2010	10.61	4.66	24.15	52	29
	2011	9.04	4.94	16.54	38	21
Virginia's warbler	2005	8.93	4.32	18.46	46	54
	2006	6.01	2.22	16.26	66	35
	2007	5.31	1.95	14.42	67	30
	2008	4.26	1.92	9.46	51	24
	2009	3.99	1.65	9.65	58	22
	2010	9.41	4.62	19.13	45	53
	2011	10.22	3.97	26.26	62	52

Table 3-4. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in pinyon-juniper habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Western meadowlark	2005	2.34	0.72	7.60	81	40
	2006	1.18	0.29	4.79	102	20
	2007	1.35	0.29	6.23	116	21
	2008	0.86	0.23	3.25	95	14
	2009	1.15	0.29	4.57	100	17
	2010	--	--	--	--	4
	2011	1.98	0.69	5.70	71	34
Western scrub-jay	2005	5.31	2.88	9.79	38	31
	2006	6.90	3.55	13.44	42	32
	2007	6.41	3.56	11.56	37	30
	2008	7.40	3.60	15.21	46	33
	2009	3.38	1.48	7.76	54	15
	2010	6.05	3.16	11.61	41	27
	2011	5.17	3.05	8.75	33	22
Western tanager	2005	1.17	0.50	2.76	55	17
	2006	1.74	0.94	3.21	38	25
	2007	2.16	1.00	4.67	49	30
	2008	1.32	0.44	3.96	75	17
	2009	--	--	--	--	8
	2010	1.21	0.41	3.58	73	17
	2011	1.59	0.65	3.91	59	21
White-breasted nuthatch	2005	3.86	1.36	10.99	70	15
	2006	4.16	1.43	12.10	72	14
	2007	7.18	2.02	25.51	89	22
	2008	--	--	--	--	9
	2009	--	--	--	--	8
	2010	3.20	0.81	12.70	99	11
	2011	3.15	1.06	9.37	73	11
White-throated swift	2005	18.12	7.12	46.10	62	64
	2006	25.15	7.49	84.46	85	82
	2007	25.81	10.10	65.96	62	94
	2008	22.76	6.63	78.20	87	66
	2009	9.96	3.05	32.60	82	35
	2010	7.71	2.54	23.43	75	25
	2011	9.02	4.08	19.91	51	27

Table 3-4. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in pinyon-juniper habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Yellow-rumped warbler	2005	1.65	0.48	5.65	86	11
	2006	3.90	1.10	13.82	89	27
	2007	1.84	0.45	7.46	102	11
	2008	2.41	0.64	9.11	95	15
	2009	--	--	--	--	1
	2010	--	--	--	--	8
	2011	4.24	2.01	8.96	47	24

Dashes indicate the sample size was insufficient for estimating density. Priority species are bolded and shaded.

3.2 Sage shrubland

We surveyed all 15 SA transect locations twice this year. We calculated density estimates for 33 species, 14 of which were priority species, from 2005 to 2011. We were unable to calculate density estimates for three of the 33 species specifically for 2011. The pooled data for 2005–2011 yielded robust density estimates (CV <50%) for 21 species and a moderately robust estimate (CV = 50–75%) for eight additional species (Table 3-5). We expect to reach our target of detecting a population change of at least 3% within 30 years for these 29 species, which represent 35% of species and 90% of individuals detected in SA habitat.

Estimated densities recorded in SA were highest for the following 10 species in 2011:

1. Brewer's sparrow
2. Vesper sparrow
3. Violet-green swallow
4. Green-tailed towhee
5. Blue-gray gnatcatcher
6. Spotted towhee
7. Chipping sparrow
8. Lark sparrow
9. Mountain bluebird
10. Dusky flycatcher

For the following 16 species, estimated densities were higher in SA than in the other two habitats sampled in 2011 (listed in order from highest to lowest density):

1. Brewer's sparrow
2. Vesper sparrow
3. Violet-green swallow
4. Green-tailed towhee
5. Lark sparrow
6. Western meadowlark
7. Mountain bluebird
8. Dusky flycatcher

9. Sage thrasher
10. Rock wren
11. Brewer's blackbird
12. American robin
13. Horned lark
14. Black-billed magpie
15. Northern flicker
16. Sage sparrow

3.3.1 Trend detection

Significant population trends were detected for five species in sage shrubland habitat. Black-billed magpie ($\beta_{\text{trend}} = 0.15$; 95% CI = 0.06, 0.24) and violet-green swallow ($\beta_{\text{trend}} = 1.76$, 95% CI = 0.56, 2.97) showed evidence of significant population change over the last seven years with quadratic trend (Figures 3-8, 3-9). Dusky flycatcher ($\beta_{\text{trend}} = 4.09$; 95% CI = 1.38, 6.80) and sage thrasher ($\beta_{\text{trend}} = 4.01$, 95% CI = 1.06, 6.95) each showed an increasing log-linear trend (Figures 3-10, 3-11), and Say's phoebe ($\beta_{\text{trend}} = 0.18$; 95% CI = 0.08, 0.28) showed an increasing linear trend (Figure 3-12). The best approximating model for all other species of concern with density estimates in SA habitat was the intercept-only (constant) model, which indicates no significant trend.

3.3 Pipe Spring National Monument

We detected a total of 286 birds of 36 species at Pipe Spring NM in 2011 (Appendix A). To date, we have detected 644 birds of 48 species. Likely due to the late migration season and the early date of the first survey visit (May 8), two species not previously detected in the park were found in 2011: black-throated gray warbler and sage thrasher.




Figure 3-8. Estimated densities and population trend of black-billed magpie in NCPN sage shrubland habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.




Figure 3-9. Estimated densities and population trend of violet-green swallow in NCPN sage shrubland habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.




Figure 3-10. Estimated densities and population trend of dusky flycatcher in NCPN sage shrubland habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.




Figure 3-11. Estimated densities and population trend of sage thrasher in NCPN sage shrubland habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.




Figure 3-12. Estimated densities and population trend of Say's phoebe in NCPN sage shrubland habitat, 2005–2011. Error bars represent 90% confidence intervals. The dashed line represents the best estimate of observed population trend for the NCPN data.

Table 3-5. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (% CV), and sample sizes (n) of breeding birds in sage shrubland habitat, 2005–2011.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
American robin	2005	2.14	1.22	3.74	35	36
	2006	4.07	2.11	7.87	42	71
	2007	2.82	1.42	5.60	44	50
	2008	2.17	1.27	3.70	33	36
	2009	1.72	0.89	3.33	42	28
	2010	2.64	1.43	4.91	39	46
	2011	3.27	1.89	5.66	34	58
Ash-throated flycatcher	2005	1.04	0.46	2.35	52	12
	2006	--	--	--	--	9
	2007	1.38	0.46	4.14	74	14
	2008	0.93	0.38	2.27	58	10
	2009	--	--	--	--	2
	2010	1.07	0.47	2.45	53	11
	2011	0.92	0.39	2.15	54	10
Black-billed magpie	2005	3.77	1.74	8.16	49	62
	2006	3.60	2.27	5.71	29	85
	2007	2.16	1.34	3.47	29	45
	2008	1.85	1.03	3.33	37	40
	2009	1.29	0.56	2.95	54	29
	2010	1.81	1.00	3.27	37	44
	2011	2.29	1.18	4.43	42	50
Black-throated gray warbler	2005	2.71	0.74	9.83	91	25
	2006	1.76	0.46	6.65	95	17
	2007	1.48	0.43	5.11	87	14
	2008	2.52	0.50	12.69	126	24
	2009	4.18	1.47	11.82	70	38
	2010	1.74	0.61	4.99	71	17
	2011	2.33	0.55	9.85	106	22
Black-throated sparrow	2005	0.78	0.21	2.88	92	10
	2006	2.05	0.61	6.86	84	29
	2007	1.33	0.41	4.35	81	18
	2008	--	--	--	--	3
	2009	1.86	0.46	7.44	101	23
	2010	--	--	--	--	8
	2011	--	--	--	--	7

Table 3-5. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in sage shrubland habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Blue-gray gnatcatcher	2005	9.16	4.53	18.54	45	27
	2006	5.32	3.08	9.19	34	16
	2007	3.98	1.90	8.35	46	11
	2008	14.21	8.54	23.64	32	40
	2009	13.20	6.31	27.63	47	34
	2010	13.89	8.46	22.81	31	41
	2011	10.49	5.70	19.29	38	30
Brewer's blackbird	2005	--	--	--	--	5
	2006	6.35	3.11	12.93	45	40
	2007	2.55	1.07	6.09	55	17
	2008	--	--	--	--	8
	2009	4.56	1.61	12.87	69	25
	2010	2.64	1.38	5.04	41	18
	2011	3.57	1.29	9.88	68	22
Brewer's sparrow	2005	104.18	70.25	154.51	24	437
	2006	81.00	49.73	131.94	30	473
	2007	68.03	44.31	104.45	26	405
	2008	57.74	38.18	87.33	25	351
	2009	102.54	66.59	157.88	25	521
	2010	73.33	54.49	98.68	18	504
	2011	63.16	35.75	111.59	34	490
Broad-tailed hummingbird	2005	14.60	5.34	39.90	67	19
	2006	8.75	2.84	26.99	76	12
	2007	10.45	3.50	31.24	74	15
	2008	20.43	6.11	68.32	83	23
	2009	11.62	3.45	39.10	84	13
	2010	--	--	--	--	9
	2011	--	--	--	--	4
Brown-headed cowbird	2005	--	--	--	--	4
	2006	4.67	2.39	9.11	42	26
	2007	1.90	0.92	3.94	46	12
	2008	2.56	1.33	4.90	41	15
	2009	--	--	--	--	8
	2010	--	--	--	--	6
	2011	1.93	1.09	3.42	35	12

Table 3-5. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in sage shrubland habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Chipping sparrow	2005	5.67	3.07	10.45	38	29
	2006	9.15	5.11	16.37	36	50
	2007	9.38	4.38	20.10	49	46
	2008	7.53	3.24	17.49	54	38
	2009	7.72	3.80	15.67	45	38
	2010	18.76	10.14	34.71	39	93
	2011	8.99	4.96	16.32	37	44
Common raven	2005	--	--	--	--	7
	2006	1.00	0.47	2.13	48	19
	2007	1.52	0.96	2.39	28	40
	2008	1.00	0.60	1.67	32	25
	2009	0.49	0.22	1.08	50	11
	2010	1.23	0.77	1.97	29	32
	2011	1.55	0.98	2.44	28	40
Dusky flycatcher	2005	3.04	1.12	8.26	66	30
	2006	4.70	2.14	10.29	50	48
	2007	5.06	2.19	11.71	54	53
	2008	6.58	2.84	15.24	54	63
	2009	7.88	3.25	19.08	57	71
	2010	6.17	3.13	12.15	43	62
	2011	6.35	3.67	11.01	34	64
Gray flycatcher	2005	1.85	1.02	3.36	37	23
	2006	--	--	--	--	8
	2007	--	--	--	--	7
	2008	1.13	0.33	3.87	86	13
	2009	2.13	0.90	5.04	56	23
	2010	1.88	0.94	3.77	44	23
	2011	1.79	0.81	3.92	50	21
Green-tailed towhee	2005	34.51	20.83	57.17	31	266
	2006	41.66	20.15	86.14	46	387
	2007	40.31	19.58	82.98	45	283
	2008	32.75	16.93	63.36	41	227
	2009	45.26	22.15	92.48	44	242
	2010	21.42	11.96	38.37	36	183
	2011	22.72	13.83	37.30	30	273

Table 3-5. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in sage shrubland habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Horned lark	2005	2.81	1.33	5.98	48	27
	2006	3.15	1.65	6.01	41	31
	2007	3.88	1.95	7.74	44	38
	2008	2.01	0.68	5.93	73	18
	2009	4.75	2.26	10.01	47	42
	2010	2.98	1.46	6.08	45	29
	2011	2.93	1.39	6.19	47	28
House finch	2005	--	--	--	--	3
	2006	4.46	2.73	7.28	30	52
	2007	2.41	0.97	6.01	60	29
	2008	2.63	1.43	4.84	38	31
	2009	4.43	1.53	12.85	72	44
	2010	1.59	0.83	3.08	41	19
	2011	1.64	0.83	3.22	43	18
Lark sparrow	2005	8.10	3.80	17.25	48	83
	2006	8.88	4.57	17.25	42	91
	2007	5.42	2.50	11.76	50	61
	2008	2.38	1.03	5.48	54	24
	2009	11.82	5.02	27.83	54	111
	2010	7.44	3.74	14.81	44	72
	2011	8.60	4.48	16.52	41	97
Lazuli bunting	2005	--	--	--	--	8
	2006	--	--	--	--	6
	2007	--	--	--	--	5
	2008	--	--	--	--	0
	2009	9.33	3.30	26.40	70	74
	2010	4.97	2.33	10.61	48	43
	2011	2.38	0.98	5.77	58	20
Mountain bluebird	2005	6.18	3.73	10.24	31	60
	2006	7.99	5.52	11.57	23	79
	2007	8.25	4.99	13.65	31	81
	2008	6.95	4.85	9.95	22	65
	2009	6.04	3.16	11.52	41	53
	2010	7.77	5.43	11.11	22	75
	2011	6.57	4.29	10.06	26	62

Table 3-5. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in sage shrubland habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Mourning dove	2005	2.44	1.37	4.34	36	55
	2006	4.41	3.03	6.42	23	101
	2007	2.70	1.45	5.01	39	63
	2008	3.10	1.51	6.36	45	69
	2009	3.11	1.65	5.85	40	61
	2010	3.74	1.99	7.01	39	89
	2011	2.42	1.56	3.74	27	56
Northern flicker	2005	1.00	0.59	1.68	32	18
	2006	1.47	0.83	2.62	36	27
	2007	0.78	0.40	1.52	42	14
	2008	0.58	0.34	0.98	32	10
	2009	1.28	0.66	2.47	42	21
	2010	1.38	0.84	2.25	30	25
	2011	1.68	1.02	2.76	31	30
Rock wren	2005	3.39	2.13	5.40	29	83
	2006	6.31	3.82	10.43	31	153
	2007	2.92	1.43	5.96	45	71
	2008	3.64	2.12	6.24	34	83
	2009	3.09	1.78	5.37	35	68
	2010	2.58	1.45	4.60	36	62
	2011	5.18	3.02	8.90	34	122
Sage sparrow	2005	2.96	1.09	8.05	66	36
	2006	1.97	0.71	5.49	68	25
	2007	2.46	0.83	7.24	73	31
	2008	1.74	0.46	6.61	96	20
	2009	2.80	1.01	7.74	68	27
	2010	--	--	--	--	4
	2011	1.40	0.53	3.69	64	18
Sage thrasher	2005	3.34	1.57	7.09	47	72
	2006	3.16	1.46	6.83	49	70
	2007	2.99	1.48	6.01	44	69
	2008	6.26	2.97	13.19	45	131
	2009	5.40	2.46	11.85	49	109
	2010	6.27	2.62	14.96	54	138
	2011	5.33	2.67	10.61	42	118

Table 3-5. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in sage shrubland habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Say's phoebe	2005	0.65	0.31	1.36	47	10
	2006	0.82	0.47	1.41	34	15
	2007	--	--	--	--	7
	2008	0.89	0.33	2.41	66	15
	2009	1.10	0.43	2.85	62	16
	2010	1.08	0.47	2.53	55	19
	2011	1.82	1.06	3.13	34	31
Spotted towhee	2005	6.01	2.96	12.22	45	55
	2006	6.29	3.56	11.12	36	61
	2007	5.21	2.86	9.51	38	50
	2008	7.64	4.27	13.68	36	68
	2009	4.90	2.54	9.45	41	41
	2010	8.40	4.46	15.81	40	75
	2011	10.27	5.83	18.08	35	95
Vesper sparrow	2005	32.87	21.86	49.44	25	252
	2006	27.28	17.29	43.05	28	324
	2007	47.27	25.93	86.18	37	382
	2008	51.41	24.45	108.08	46	346
	2009	51.33	33.12	79.54	27	396
	2010	47.74	36.38	62.64	17	461
	2011	38.90	28.81	52.54	18	473
Violet-green swallow	2005	5.40	2.75	10.62	42	14
	2006	8.52	4.05	17.91	47	26
	2007	4.04	2.11	7.74	41	13
	2008	6.29	3.33	11.91	39	13
	2009	6.57	3.04	14.19	49	19
	2010	5.18	0.70	38.31	171	15
	2011	37.18	15.38	89.90	56	22
Virginia's warbler	2005	1.42	0.61	3.34	55	23
	2006	1.80	0.86	3.80	47	30
	2007	1.08	0.28	4.10	96	17
	2008	2.57	0.96	6.91	66	40
	2009	1.43	0.55	3.71	63	21
	2010	1.55	0.64	3.77	58	25
	2011	1.32	0.57	3.08	55	21

Table 3-5. Estimated densities per km² (D), lower and upper 90% confidence limits on D (LCL, UCL), percent coefficient of variation of estimates (%CV), and sample sizes (n) of breeding birds in sage shrubland habitat, 2005–2011, cont.

Species	Year	D	Confidence limits		%CV	n
			LCL	UCL		
Warbling vireo	2005	1.27	0.44	3.62	70	17
	2006	0.68	0.28	1.66	58	10
	2007	1.44	0.54	3.87	65	22
	2008	--	--	--	--	2
	2009	--	--	--	--	3
	2010	--	--	--	--	9
	2011	1.11	0.43	2.87	62	16
Western meadowlark	2005	5.72	2.99	10.96	41	152
	2006	5.64	3.35	9.51	32	166
	2007	5.73	2.93	11.22	42	135
	2008	10.51	5.00	21.86	46	178
	2009	6.95	3.60	13.41	41	209
	2010	7.57	4.31	13.29	35	169
	2011	7.76	4.09	14.72	39	246
Western tanager	2005	0.48	0.16	1.45	74	13
	2006	0.69	0.20	2.47	89	17
	2007	0.90	0.31	2.60	71	25
	2008	0.73	0.23	2.30	78	19
	2009	0.79	0.27	2.26	71	20
	2010	0.52	0.18	1.49	70	13
	2011	--	--	--	--	4

Dashes indicate the sample size was insufficient for estimating density. Priority species are bolded and shaded.

4 Discussion

The density estimates shown in Tables 3-3, 3-4, and 3-5 allow for comparison across seven years and, more importantly, provide new, more robust, estimates for data collected in 2005–2010 (because pooling data across all years increased overall sample sizes). For most species, each year of additional data will improve our ability to accurately estimate densities of the species that occupy the NCPN. The density estimates presented in this report replace the estimates provided in the 2005–2010 reports. Sample sizes (n) reported in the tables indicate the number of detections used in analyses, after truncation.

The minimum post-truncation sample size was increased from 60 to 80 detections this year. This method will initially reduce the number of species for which density can be estimated, but will eventually provide more reliable estimates. In low-elevation riparian habitat, density estimates for western tanager (n = 72) were not included. In pinyon-juniper habitat, density estimates were not included for Brewer's sparrow (n = 62) or green-tailed towhee (n = 77), but enough detections were accrued to obtain density estimates for a new species, hermit thrush (n = 86). In sage shrubland habitat, density estimates for yellow-rumped warbler (n = 79) were not included for the first time since 2009. However, sufficient sample sizes for several species (including yellow-rumped warbler in SA) will likely allow their inclusion in next season's report.

4.1 Status and trends

In support of the project objective, which is to determine the population status and trends of breeding landbird species in LR, PJ, and SA habitats, the second year of landbird population trend results for several species of conservation concern in the NCPN are presented here. The results this year used slightly different density estimates to populate trends than those used in 2010, because increased truncation distances for some species resulted in smaller sample sizes. In other cases, the smaller sample sizes were related to the increase, from 60 to 80, in the minimum number of detections used for analysis this year.

The trend results indicate that many sensitive species appear to have stable populations within certain habitats. As expected, the number of species showing significant trends increased in 2011—particularly, quadratic trends, which are

significant though not necessarily positive or negative.

The Bewick's wren (BEWR) population continued to have a decreasing population trend. Potential explanations of BEWR declines include loss of suitable nesting habitat (cavities) and competition with other cavity nesters. Additional investigation of BEWR populations is recommended, as RMBO has documented a decline of more than 50% in the last seven years. Black-billed magpie (BBMA) had evidence of a negative population trend in 2010, but showed a quadratic population trend with the addition of the 2011 data. With continued data collection, species such as BBMA could reveal trends in population variability over time. Other species for which RMBO has detected significant decreasing trends are black-throated gray warbler and white-throated swift, in pinyon-juniper. Because staff did not detect significant declines in these species in 2010, RMBO is exploring whether these are true declines, whether observer detectability is varying among years, or if these species have cyclical populations. Long-term monitoring will continue to be necessary for interpreting population status and trends for avian species of the NCPN.

4.2 Knowledge-sharing

In 2007, the NABCI monitoring subcommittee outlined recommendations for improving monitoring programs (USNABCIMS 2007). The first recommendation was that bird-monitoring programs should integrate an adaptive management approach into the monitoring process to incorporate management and conservation priorities. It is hoped that trends identified in this early warning program will lead to additional research and projects that explore fluctuations in NCPN landbird populations.

The second recommendation was to coordinate landbird monitoring among organizations and across spatial scales to make monitoring more efficient and effective. RMBO continues to work with a variety of federal, state, and local agencies throughout 13 different states, monitoring landbirds through a spatially balanced study design that uses Bird Conservation Regions (BCRs) as the sampling frame. RMBO expects to post-stratify the NCPN surveys in this framework by habitat and compare the results to those presented here. Furthermore, RMBO can use BCR-based data from lands outside the network for further comparisons. Data collected from Colorado

and Wyoming from 2008 to 2011 are currently available for such comparisons. In future years, RMBO hopes to add the Utah portions of BCR 9 and BCR 16. Comparisons of densities in these areas may help guide the National Park Service in making key management decisions and assessing the effectiveness of past and existing management practices.

RMBO's Avian Data Center (ADC) meets a third NABCI recommendation by providing decision-support tools, such as distribution maps, species counts, monitoring reports, and species information to land managers and the public. In 2012, RMBO will release an updated version of the ADC. This will include species accounts of the 44 priority species (Appendix B) we detected during counts in 2011, as well as information on other species found in the NCPN. In addition, RMBO is a partner of the Avian Knowledge Network (AKN), whose goals are to (1) compile bird-monitoring data from various contributor organizations and organize them into one format and (2) make those data available to land managers, scientists, and the public for decisionmaking, research, and educational purposes.

4.3 Pipe Spring National Monument

In 2011, the monitoring protocol at PISP was further refined to include diurnal and nocturnal area searches. These area searches provided additional information on species occurrences, as well as migratory and breeding behavior. For example, the presence of black phoebe in the park was first confirmed last year when observers located a black phoebe nest during the first area search. Staff counted 170 birds and 5 additional species by adding area searches in 2011: Bewick's wren, Black phoebe, common poorwill, great horned owl, and lazuli bunting. The monument's small size allows these area searches to be repeated by volunteers who come to observe birds or conduct other surveys whose results can be used to inform park management actions or habitat enhancement decisions.

5 Literature Cited

- Alexander, J. D., J. L. Stevens, G. R. Geupel, and T. C. Will. 2008. Decision support tools: Bridging the gap between science and management. Pages 283–291 *in* Proceedings of the Fourth International Partners in Flight Conference: Tundra to Tropics. February 13–16, 2008, McAllen, Texas.
- American Ornithologists' Union (AOU). 2007. Checklist of North American birds, seventh edition. American Ornithologists' Union, Washington, D.C..
- Baron, J. S., S. H. Julius, J. M. West, L. A. Joyce, G. Blate, C. H. Peterson, M. Palmer, B. D. Keller, P. Kareiva, J. M. Scott, and B. Griffith. 2008. Some guidelines for helping natural resources adapt to climate change. *International Human Dimensions Programme on Global Environmental Change Update* 2:46–52.
- Buckland, S. T., D. R. Anderson, K. P. Burnham, J. L. Laake, D. L. Borchers, and L. Thomas. 2001. Introduction to distance sampling: Estimating abundance of biological populations. Oxford, U.K.: Oxford University Press.
- Burnham, K. P., and D. R. Anderson. 2002. Model selection and multimodel inference: A practical information-theoretic approach. New York: Springer-Verlag.
- Dreitz, V. J., P. M. Lukacs, and F. L. Knopf. 2006. Monitoring low density avian populations: An example using Mountain Plovers. *Condor* 108:700–706.
- Hanni, D. J., C. M. White, R. A. Sparks., J. A. Blakesley, J. J. Birek, N. J. Van Lanen, and J. A. Fogg. 2011. Field protocol for spatially balanced sampling of landbird populations. Unpublished report. Rocky Mountain Bird Observatory, Brighton, Colorado.
- Leukering, T., M. Carter, A. Panjabi, D. Faulkner, and R. Levad. 2005. Point transect protocol (rev). Unpublished document. Rocky Mountain Bird Observatory, Brighton, Colorado.
- Lindenmayer, D. B., and G. E. Likens. 2009. Adaptive monitoring: A new paradigm for long-term research and monitoring. *Trends in Ecology and Evolution* 24:482–486.
- Lowry, J. H., Jr., R. D. Ramsey, K. Boykin, D. Bradford, P. Comer, S. Falzarano, W. Kepner, J. Kirby, L. Langs, J. Prior-Magee, G. Manis, L. O'Brien, T. Sajwaj, K. A. Thomas, W. Rieth, S. Schrader, D. Schrupp, K. Schulz, B. Thompson, C. Velasquez, C. Wallace, E. Waller, and B. Wolk. 2005. Southwest Regional Gap Analysis Project: Final report on land cover mapping methods. Logan, Utah. <http://earth.gis.usu.edu/swgap/swregap_landcover_report.pdf>. Accessed January 28, 2011.
- Lyons, J. E., M. C. Runge, H. P. Laskowski, and W. L. Kendall. 2008. Monitoring in the context of structured decision-making and adaptive management. *Journal of Wildlife Management* 72:1683–1692.
- Manley, P. N., W. M. Block, F. R. Thompson, G. S. Butcher, C. Paige, L. H. Suring, D. S. Winn, D. Roth, C. J. Ralph, E. Morris, C. H. Flather, and K. Byford. 1993. Guidelines for monitoring populations of Neotropical migratory birds on National Forest system lands. USDA Forest Service Monitoring Task Group Report, Washington, D. C.
- Marsh, D. M., and P. C. Trenham. 2008. Current trends in plant and animal population monitoring. *Conservation Biology* 22:647–655.
- Partners in Flight. 2005. Species assessment database. <<http://www.rmbo.org/pif/pifdb.html>>. Accessed November 1, 2010.
- Rich, T. D., C. J. Beardmore, H. Berlanga, P. J. Blancher, M. S. W. Bradstreet, G. S. Butcher, D. W. Demarest, E. H. Dunn, W. C. Hunter, E. E. Iñigo-Elias, J. A. Kennedy, A. M. Martell, A. O. Panjabi, D. N. Pashley, K. V. Rosenberg, C. M. Rustay, J. S. Wendt, and T. C. Will. 2004. Partners in Flight North American landbird conservation plan. Cornell Lab of Ornithology, Ithaca, New York.

- Sauer, J. R. 1993. Monitoring goals and programs of the U.S. Fish and Wildlife Service. Pages 245–251 *in* Conference proceedings, Status and management of neotropical migratory birds: September 21–25, 1992, Estes Park, Colorado, D. M. Finch and P. W. Stangel, eds. Gen. Tech. Rep. RM-229. U.S. Forest Service Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado.
- Sauer, J. R., and M. G. Knutson. 2008. Objectives and metrics for wildlife monitoring. *Journal of Wildlife Management* 72:1663–1664.
- Thomas, L., S. T. Buckland, E. A. Rexstad, J. L. Laake, S. Strindberg, S. L. Hedley, J. R. B. Bishop, T. A. Marques, and K. P. Burnham. 2010. Distance software: Design and analysis of distance sampling surveys for estimating population size. *Journal of Applied Ecology* 47:5–14.
- Thompson, W. L., G. C. White, and C. Gowan. 1998. Monitoring vertebrate populations. San Diego, Calif.: Academic Press.
- U.S. Fish and Wildlife Service. 2008. Birds of conservation concern 2008. Arlington, Virginia. <<http://www.fws.gov/migratorybirds/NewReportsPublications/SpecialTopics/BCC2008/BCC2008.pdf>>. Accessed November 1, 2010.
- U.S. North American Bird Conservation Initiative Committee. 2009. The state of the birds, United States of America, 2009. U.S. Fish and Wildlife Service, Washington, D.C. <http://www.stateofthebirds.org/pdf_files/State%20of%20the%20Birds_FINAL.pdf>. Accessed December 1, 2010.
- U.S. North American Bird Conservation Initiative Monitoring Subcommittee (USNABCIMS). 2007. Opportunities for improving avian monitoring. Division of Migratory Bird Management, U.S. Fish and Wildlife Service, Arlington, Virginia. <<http://www.nabci-us.org/>>. Accessed November 11, 2010.
- Utah Division of Wildlife Resources. 2005. Utah comprehensive wildlife conservation strategy. Salt Lake City, Utah. <http://wildlife.utah.gov/cwcs/10-01-21_utah_cwcs_strategy.pdf>. Accessed November 1, 2010.
- Witmer, G. W. 2005. Wildlife population monitoring: Some practical considerations. *Wildlife Research* 32:259–263.

Appendix A. Total bird species observed during surveys in the Northern Colorado Plateau Network by habitat (2011) and year (2005–2011).

Species	Individuals observed per habitat, 2011				Individuals observed per year (all habitats), 2005–2011							
	LR	PJ	SA	PISP	2005	2006	2007	2008	2009	2010	2011	Total
American crow					10	1	2	4	2	2		21
American dipper					1	1	1	2	1			6
American goldfinch	10				10	4	9	7	8	34	10	82
American kestrel			17		10	10	11	9	9	13	17	79
American robin	30	39	70	1	100	154	144	129	64	100	140	831
Ash-throated flycatcher	257	109	10	3	300	412	357	357	442	384	379	2,631
Bank swallow							2	1		1		4
Barn swallow			2	2	1	2	3	1	2		4	13
Belted kingfisher							1					1
Bewick's wren	17	33		1	237	306	197	205	219	50	51	1,265
Black phoebe	3			2	10	15	28	12	32	10	5	112
Black-billed magpie		7	84		77	108	53	52	52	74	91	507
Black-capped chickadee	2		1		1	4	4		2	1	3	15
Black-chinned hummingbird	48	19	2		50	67	52	52	104	53	69	447
Black-chinned sparrow		3			9	6	4		12	4	3	38
Black-headed grosbeak	16	18	7		27	54	37	37	46	38	41	280
Black-throated gray warbler	99	379	23	5	393	551	455	566	627	420	506	3,518
Black-throated sparrow	123	93	7	12	113	171	193	151	334	183	235	1,380
Blue grosbeak	6				14	32	6	6	35	3	6	102
Blue-gray gnatcatcher	218	180	36		340	302	291	332	493	424	434	2,616
Blue-winged teal					1							1
Bobolink	1										1	1
Brewer's blackbird	1		33		10	48	23	10	30	29	34	184
Brewer's sparrow	6	4	647	1	584	635	593	535	710	652	658	4,367
Broad-tailed hummingbird	11	1	10	1	68	30	46	75	29	78	23	349
Brown creeper			1								1	1
Brown-headed cowbird	24	23	22	11	39	70	50	47	56	27	80	369
Bullock's oriole	12	2	8	9	7	30	19	13	20	34	31	154
Bushtit		1			67	93	33	24	61	11	1	290
California gull					3	3						6
Canada goose	2	4	17		6	6	20	9	5	8	23	77

Appendix A. Total bird species observed during surveys in the Northern Colorado Plateau Network by habitat (2011) and year (2005–2011), cont.

Species	Individuals observed per habitat, 2011				Individuals observed per year (all habitats), 2005–2011							Total
	LR	PJ	SA	PISP	2005	2006	2007	2008	2009	2010	2011	
Canyon wren	12	6			52	103	70	36	58	7	18	344
Cassin's finch		9	4		4	38	14	13	2	4	13	88
Cassin's kingbird	1			6		1			19		7	27
Cedar waxwing							1	1	1			3
Chipping sparrow	43	149	76		153	152	186	213	245	272	268	1,489
Chukar						3	3		2	3		11
Clark's nutcracker		6	2		16	24	26	17	24	7	8	122
Cliff swallow	1		15		50	36	15	6		5	16	128
Common merganser					1	2	7	4	1			15
Common nighthawk		2	1		2	1	1	2	6	14	3	29
Common poorwill				1	1	1	1			2	1	6
Common raven	64	75	55	5	116	134	176	135	131	192	199	1,083
Common yellowthroat	15	2	3		13	30	27	9	31	16	20	146
Cooper's hawk	2		1		15	17	14	5	8	15	3	77
Cordilleran flycatcher		1			4	1	2	3	3	3	1	17
Dark-eyed junco		1	7		38	18	45	35	4	3	8	151
Downy woodpecker	6				5	11	3	7	15	1	6	48
Dusky flycatcher	3	19	77		51	111	98	136	108	98	99	701
Dusky grouse					1		1					2
Eastern kingbird	1										1	1
Eared grebe									1			1
Eurasian collared-dove	1	1		8			1			19	10	29
European starling			12	15	9	16	8	10		8	27	78
Evening grosbeak	1	1					1				2	3
Ferruginous hawk										1		1
Gadwall			2		1	1	3				2	7
Gambel's quail				14	9	15		1	7	15	14	61
Golden eagle					7	10	4	6	1			28
Grace's warbler		18	15		15	31	39	31	26	19	33	194
Gray flycatcher	8	128	22		160	116	108	142	256	227	158	1,167
Gray vireo	10	108	3		136	128	104	150	252	164	121	1,055
Great blue heron			3		9	3	3	3	1	6	3	28
Great horned owl				1	2			1		1		4

Appendix A. Total bird species observed during surveys in the Northern Colorado Plateau Network by habitat (2011) and year (2005–2011), cont.

Species	Individuals observed per habitat, 2011				Individuals observed per year (all habitats), 2005–2011							
	LR	PJ	SA	PISP	2005	2006	2007	2008	2009	2010	2011	Total
Greater sage-grouse					7	2	1					10
Green-tailed towhee	1	17	351		346	462	364	302	330	220	369	2,393
Green-winged teal						1						1
Hairy woodpecker	7	4	3		13	21	21	16	26	21	14	132
Hammond's flycatcher	1						9	9	19		1	38
Hermit thrush		43	1		15	21	2		24	14	44	120
Horned lark			29		31	39	43	23	47	32	29	244
House finch	151	175	20	22	234	336	423	416	506	357	368	2,640
House sparrow				16						1	16	17
House wren	57	11	20		59	61	39	55	57	48	88	407
Indigo bunting						1			6			7
Juniper titmouse	28	121	2		130	112	135	156	251	192	151	1,127
Killdeer					4	3	2	2	5	1		17
Lark bunting	1										1	1
Lark sparrow	13	27	101	5	131	146	101	50	168	130	146	872
Lazuli bunting	155	11	21	5	183	155	223	186	275	247	192	1,461
Lesser goldfinch	47	5	1	7	38	67	106	75	159	50	60	555
Lincoln's sparrow					1		2	2	1			4
Loggerhead shrike					3	1		1		4		9
Long-eared owl						2						2
Lucy's warbler	19				1		11	20	39	7	19	97
Macgillivray's warbler					3	9	3	4	7	11		37
Mallard	2		4	1	2	15	2	11	3	2	7	42
Mountain bluebird	10	32	77		149	134	157	111	109	144	119	923
Mountain chickadee		27	4		34	22	24	14	25	8	31	158
Mourning dove	91	151	72	16	411	536	499	393	270	428	330	2,867
Northern flicker	9	18	37		48	57	58	22	50	72	64	371
Northern goshawk								1				1
Northern harrier	1		1		2	4	5	6	2	8	2	29
Northern mockingbird	10	7	2	5		25	11	17	37	32	24	146
Northern rough-winged swallow	6		1		6	6	2	6	16	8	7	51
Northern shoveler							2					2
Olive-sided flycatcher		2			16	4	4	2	18		2	46

Appendix A. Total bird species observed during surveys in the Northern Colorado Plateau Network by habitat (2011) and year (2005–2011), cont.

Species	Individuals observed per habitat, 2011				Individuals observed per year (all habitats), 2005–2011							Total
	LR	PJ	SA	PISP	2005	2006	2007	2008	2009	2010	2011	
Orange-crowned warbler	1		2		2	2	2	2		2	3	13
Osprey		1			1			1			1	3
Peregrine falcon	2	3			4	5	7	10	3	7	5	41
Phainopepla									3			3
Pine siskin	11	22	3	25	4	15	4	15	15	18	61	132
Pinyon jay	3	49	7		75	106	123	50	79	163	59	655
Plumbeous vireo	57	91	10		147	165	138	131	148	216	158	1,103
Prairie falcon	2				2			2	1	1	2	8
Pygmy nuthatch		9	15		10	7	12	5	10	7	24	75
Red crossbill		1			1	11	4		19	6	1	42
Red-breasted nuthatch	1		2		11	5	7	1	2	1	3	30
Red-naped sapsucker						2		1	2			5
Red-tailed hawk	2	3	2		12	17	10	9	12	5	7	72
Red-winged blackbird	4		3			4	1	1	1	2	7	16
Ring-billed gull							2					2
Rock pigeon		1			2	5	3			2	1	13
Rock wren	131	94	135	3	315	403	348	303	417	256	363	2,405
Ruby-crowned kinglet	2	4	1		4	4	3	9	11	10	7	48
Sage sparrow			18		43	32	31	21	37	10	18	192
Sage thrasher			130	1	92	93	88	155	128	150	131	837
Sandhill crane						1		1	1			3
Savannah sparrow						1	1					2
Say's phoebe	72	19	32	5	101	97	63	95	121	91	128	696
Scott's oriole							1	5	6			12
Sharp-shinned hawk						1	1	1				3
Short-eared owl						3			1			4
Song sparrow	34	2	9		60	75	69	61	33	10	45	353
Spotted sandpiper					1	2	7		3	8		21
Spotted towhee	377	128	101	5	428	609	499	405	528	543	611	3,623
Steller's jay			3		5	5	16	9	5	9	3	52
Summer tanager									1			1
Swainson's thrush									1			1
Townsend's solitaire	2	6	4		5	6	12	6	18	12	12	71

Appendix A. Total bird species observed during surveys in the Northern Colorado Plateau Network by habitat (2011) and year (2005–2011), cont.

Species	Individuals observed per habitat, 2011				Individuals observed per year (all habitats), 2005–2011							
	LR	PJ	SA	PISP	2005	2006	2007	2008	2009	2010	2011	Total
Tree swallow	28	4	3		3	20	20	25	2	2	35	107
Turkey vulture	2	7	3		19	23	16	8	18	18	12	114
Veery						1						1
Vesper sparrow		25	583		379	492	484	473	477	599	608	3,512
Violet-green swallow	106	29	28	5	194	320	265	204	276	183	168	1,610
Virginia's warbler	20	56	21		121	109	102	154	46	120	97	749
Warbling vireo	16	2	16		44	58	61	51	37	32	34	317
Western bluebird	5	3			15	14	15	17	22	9	8	100
Western grebe						1						1
Western kingbird	2			13	2	4	1	6	24	24	15	76
Western meadowlark	7	45	6		227	239	182	213	276	227	58	1,422
Western screech-owl			1		1						1	2
Western scrub-jay	36	34	4		86	87	71	78	55	80	74	531
Western tanager	13	22	1		44	69	72	59	45	55	36	380
Western wood-pewee	20		2		40	44	64	42	56	28	22	296
White-breasted nuthatch	1	12	259		27	27	30	17	22	27	272	422
White-crowned sparrow	2		7	33		6	4	7	2	3	42	64
White-faced ibis		60	4		1						64	65
White-throated swift	106		5		271	286	376	200	236	183	111	1,663
Wild turkey	3	1			2	7	1	4	7	6	4	31
Williamson's sapsucker					1		1	1				3
Willow flycatcher					1	3	3		1			8
Wilson's warbler	1		3			4		3	3	6	4	20
Yellow warbler	178	1	9	18	157	175	155	181	273	252	206	1,399
Yellow-billed cuckoo						1						1
Yellow-breasted chat	57		5		52	57	54	48	72	89	62	434
Yellow-rumped warbler	30	24	14	3	26	65	32	49	14	69	71	326

Habitats: LR=Low-Elevation Riparian; PJ=Pinyon-Juniper; SA=Sage Shrubland; PISP=Pipe Spring National Monument. Priority species are bolded and shaded.

Appendix B. Priority species observed on transects in the Northern Colorado Plateau Network, 2005–2011.

Species	UDWR ¹	Management designations					Number of individuals observed per habitat, 2005–2011			
		USFWS ²		Partners in Flight ³			LR	PJ	SA	PISP
		BCR 10	BCR 16	Region 6	BCR 10	BCR 16				
American dipper					RS		6			
Bewick's wren				BCC			454	779	26	6
Black-billed magpie						RS	4	18	486	
Black-chinned sparrow						CC	16	21	1	
Black-throated gray warbler	Tier III					RC	558	2,761	198	5
Black-throated sparrow						RC	624	582	123	58
Bobolink	Tier II						1			
Brewer's sparrow	Tier III	X	X		CC,RC	CC,RC	29	73	4,280	2
Broad-tailed hummingbird	Tier III					RS	96	99	155	1
Canyon wren						RC	236	102	4	2
Cassin's finch		X	X	BCC	RC,CS,RS	RC	7	53	32	
Clark's nutcracker					CS,RS	CS,RS	2	58	62	
Common nighthawk						RC	2	23	4	
Cordilleran flycatcher						RS	13	4		
Dusky flycatcher					CS,RS		31	195	476	
Dusky grouse					CC,RC	CC			2	
Ferruginous hawk	Tier II	X	X	BCC	RC	RC			1	
Gambel's quail	Tier III						1	22	2	36
Golden eagle			X	BCC		RC	3	12	13	
Grace's warbler			X			CC,RC	3	111	80	
Gray vireo	Tier III		X	BCC		CC,RC,RS	172	829	51	3
Greater sage-grouse	Tier II				CC,RC,CS,RS	CC,RC			10	
Green-tailed towhee						CS,RS	22	91	2,291	
Hammond's flycatcher					RS		2	32	5	
Juniper titmouse			X			RC,RS	223	863	35	7
Lazuli bunting					RS		1,205	68	185	5
Loggerhead shrike		X		BCC	RC	RC	1	4	4	
Lucy's warbler	Tier III						101			
Mountain bluebird						RC,CS,RS	47	315	583	
Northern goshawk	Tier I				RC,RS				1	

Appendix B. Priority species observed on transects in the Northern Colorado Plateau Network, 2005–2011, cont.

Species	UDWR ¹	Management designations					Number of individuals observed per habitat, 2005–2011			
		USFWS ²		Partners in Flight ³		Region 6	LR	PJ	SA	PISP
		BCR 10	BCR 16	BCR 10	BCR 16					
Northern harrier							6			
Olive-sided flycatcher		X			CC		1	10	35	
Osprey	Tier III							2	1	
Peregrine falcon	Tier III	X				BCC	22	13	3	3
Pine siskin							40	46	36	25
Pinyon jay			X		CC	BCC	19	558	82	
Plumbeous vireo					RS		514	495	94	
Prairie falcon					RC	BCC	4		5	
Pygmy nuthatch					RC			33	42	
Red crossbill				RS				12	32	
Red-naped sapsucker				CS,RS			4		1	
Rock wren					RS		824	791	781	14
Sage sparrow	Tier III	X			RC	BCC	3		189	
Sage thrasher	Tier III	X				BCC		1	836	1
Say's phoebe					RS		405	140	136	16
Short-eared owl	Tier II			CC		BCC			4	
Townsend's solitaire				RS			5	43	23	
Veery			X						1	
Violet-green swallow					RS		1,093	342	178	10
Virginia's warbler	Tier III				CC,RC,RS		256	306	189	
Warbling vireo					RS		212	18	88	
Western bluebird					RS		20	34	53	
White-throated swift				CC	CC,RS		995	651	96	7
Williamson's sapsucker	Tier III	X		CS,RS	CS,RS			1	2	
Willow flycatcher	Tier I	X	X	CC,RS	CC,RC	BCC	7	1		
Yellow-billed cuckoo	Tier I	X	X				1			

Habitats: LR=Low-Elevation Riparian; PL=Pinyon-Juniper; SA=Sage Shrubland; PISP=Pipe Spring National Monument.

¹ UDWR=Utah Division of Wildlife Resources, Tier I,II,III= Utah Comprehensive Wildlife Conservation Strategy Tier I,II,III Species of Greatest Conservation Need (Utah Division of Wildlife Resources 2005)

² USFWS=United States Fish and Wildlife Service, BCR10=Bird Conservation Region 10 (Northern Rockies), BCR 16=Bird Conservation Region 16 (Southern Rockies/Colorado Plateau) BCC=Bird of Conservation Concern for Region 6 (Mountain-Prairie Region) (U.S. Fish and Wildlife Service 2008).

³ Partners In Flight, BCR10=Bird Conservation Region 10, BCR 16=Bird Conservation Region 16, CC=Continental Concern Species, RC=Regional Concern Species, CS=Continental Stewardship Species, RS = Regional Stewardship Species (Partners in Flight 2005).

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS 960/115251, June 2012

National Park Service
U.S. Department of the Interior



Natural Resource Stewardship and Science

1201 Oak Ridge Drive, Suite 150
Fort Collins, Colorado 80525

www.nature.nps.gov

EXPERIENCE YOUR AMERICA™