

D-1172

File:

NPS General

Proceedings of the Second Biennial Conference on Research in Colorado Plateau National Parks



Transactions and Proceedings Series NPS/NRNAU/NRTP-95/11
United States Department of the Interior
National Park Service **PLEASE RETURN TO:**

Color Scans

ON MICROFILM

TECHNICAL INFORMATION CENTER
DENVER SERVICE CENTER
NATIONAL PARK SERVICE

1/31/2003

The National Park Service publishes scientific studies of significant natural resources in units of the National Park System. These studies are of scholarly quality and may include any discipline of the biological, physical, or social sciences.

Editorial Staff

National Biological Service

Paul A. Opler	Managing Editor
Paul A. Vohs	Subject Editor

National Park Service

Donna L. O'Leary	Publications Coordinator
Jerry D. Cox	Technical Editor
Martha W. Nichols	Editorial Assistant

Cover photo: Grand Canyon National Park,
Arizona (© Al Richmond)

This report may be cited as follows:

van Riper, C., III, editor. 1995. Proceedings of the Second Biennial Conference on Research in Colorado Plateau National Parks. 25-28 October 1993. National Park Service Transactions and Proceedings Series NPS/ NRNAU/ NRTP-95/11.

Copies of this report are available from the Station Leader, Colorado Plateau Research Station, Northern Arizona University, P.O. Box 5614, Flagstaff, AZ 86011-5614.

ISSN 0270-8655

Proceedings of the Second Biennial Conference on Research in Colorado Plateau National Parks

Editor

Charles van Riper III

National Biological Service

and

Department of Biological Sciences

P.O. Box 5614

Northern Arizona University

Flagstaff, Arizona 86011

Second Biennial Conference of Research on the Colorado Plateau

Northern Arizona University

Flagstaff

25-28 October 1993

Sponsored and organized by

National Biological Service

Colorado Plateau Research Station

Cosponsored by

Northern Arizona University

Transactions and Proceedings Series NPS/NRNAU/NRTP-95/11

United States Department of the Interior

National Park Service

•1995•

Contributors

W. Sylvester Allred

Northern Arizona University
Department of Biology
P.O. Box 5640
Flagstaff, Arizona 86011

Janet L. Brown

U.S. Geological Survey
Denver Federal Center
P.O. Box 25046, MS 939
Denver, Colorado 80225

Brian L. Cluer

National Park Service
National Natural Resource Center
1201 Oak Ridge Drive, Suite 250
Fort Collins, Colorado 80525

Brian J. Cooper

Department of Geology
Sam Houston State University
Huntsville, Texas 77341

Vidal Davila, Jr.

National Park Service
Great Basin National Park
Baker, Nevada 89311

Leland R. Dexter

Department of Geography
Northern Arizona University
P.O. Box 15016
Flagstaff, Arizona 86011

Charles A. Drost

National Biological Service
Colorado Plateau Research Station
Northern Arizona University
Box 5614
Flagstaff, Arizona 86011

Lisa Floyd-Hanna

Department of Environmental Studies
Prescott College
220 Grove Avenue
Prescott, Arizona 86301

James F. Fowler

State Fair Community College
Department of Natural Science
Sedalia, Missouri 65301

Phil R. Geib

Navajo Nation Archaeological Department
Northern Arizona University
P.O. Box 6013
Flagstaff, Arizona 86011

Tim B. Graham

National Biological Service
University of California
DES, Wichson Hall
Davis, California 95616

David D. Hanna

Department of Environmental Studies
Prescott College
220 Grove Avenue
Prescott, Arizona 86301

Ronald L. Hartman

Department of Botany
University of Wyoming
Laramie, Wyoming 82071

Matthew J. Johnson

National Biological Service
Colorado Plateau Research Station
Northern Arizona University
P.O. Box 5614
Flagstaff, Arizona 86011

Yiqun Lin

School of Forestry
Northern Arizona University
Box 15018
Flagstaff, Arizona 86011

Allan Loy

National Park Service
Mesa Verde National Park
Mesa Verde, Colorado 83110

Mark Manone

Department of Geography
Northern Arizona University
Flagstaff, Arizona 86011

Cathleen L. May

U.S.D.A. Forest Service
740 Simms
Golden, Colorado 80401

Ron W. Meyer

411 N. Pine
Gunnison, Colorado 81230

Donald Moore

Adobe Creek Animal Clinic
Fruita, Colorado 81521

Dennis I. Netoff

Department of Geography and Geology
Sam Houston State University
Huntsville, Texas 77341

Trevor Persons

27 Penobscot Street
Orono, Maine 04473

Gilbert C. Pogany

Department of Biological Sciences
Northern Arizona University
Box 5640
Flagstaff, Arizona 86011

H. Ronald Pulliam

U.S. Department of the Interior
National Biological Service
1849 C. Street NW, MS ARLSQ 725
Washington, D.C. 20240

Elaine Rodriguez

Department of Political Science
Northern Arizona University
P.O. Box 15036
Flagstaff, Arizona 86011

William H. Romme

Biology Department
Fort Lewis College
Durango, Colorado 81301

Peter G. Rowlands

National Biological Service
Colorado Plateau Research Station
Northern Arizona University
P.O. Box 5614
Flagstaff, Arizona 86011

William Schaedla

Saiyoke National Park
Moo 7 Tambol Saiyoke
Amphur Saiyoke
Kanchanaburi 71150 Thailand

Ralph Shroba

U.S. Geological Survey, MS 913
Denver Federal Center
Box 25046
Denver, Colorado 80225

Mark Sogge

National Biological Service
Colorado Plateau Research Station
P.O. Box 5614
Flagstaff, Arizona 86011

Frederic I. Solop

Department of Political Science
Northern Arizona University
P.O. Box 15036
Flagstaff, Arizona 86011

John R. Spence

National Park Service
Glen Canyon National Recreation Area
P.O. Box 1507
Page, Arizona 86040

Kimberly Spurr

Navajo Nation Archaeological Department
Northern Arizona University
Box 6013
Flagstaff, Arizona 86011

Nancy L. Stanton

Department of Zoology and Physiology
University of Wyoming
Laramie, Wyoming 82071

Charles van Riper III

National Biological Service
Colorado Plateau Research Station
P.O. Box 5614
Flagstaff, Arizona 86011

Michael Wagner

School of Forestry
Northern Arizona University
P.O. Box 15018
Flagstaff, Arizona 86011

Contents

	Page
List of Contributors	ii
Preface. <i>H. Ronald Pulliam</i>	vii
Introduction. <i>Charles van Riper III</i>	xi
Section I. Physical Resources	1
Geomorphology of the Hanging Gardens of the Colorado Plateau. <i>Cathleen L. May, James F. Fowler, Nancy L. Stanton</i>	3
Giant Sandstone Weathering Pits Near Cookie Jar Butte, Southeastern Utah. <i>Dennis I. Netoff, Brian J. Cooper, Ralph R. Shroba</i>	25
Large-scale Geologic Mapping in Great Basin National Park, Nevada. <i>Janet L. Brown, Vidal Davila, Jr.</i>	55
Using Land-based Photogrammetry to Monitor Sandbar Stability in Grand Canyon on a Daily Time Scale. <i>Leland R. Dexter, Brian L. Cluer, Mark Manone</i>	67
Section II. Cultural Resources	87
Radiocarbon Record for Archaic Occupation of the Central Colorado Plateau. <i>Phil R. Geib</i>	89
Compositional Analysis of Temper in Emery Gray Ceramics From Central Utah. <i>Kimberly Spurr</i>	137
A Model to Study Fire Effects on Cultural Resource Studies of Mesa Verde. <i>Lisa Floyd-Hanna, William H. Romme, Allan Loy, David D. Hanna</i>	171
Public Support for Protecting Grand Canyon Resources. <i>Frederic I. Solop, Elaine Rodriquez</i>	179
Section III. Natural Resources	191
A Preliminary Vegetation Classification for the Colorado Plateau. <i>John R. Spence, William H. Romme, Lisa Floyd-Hanna, Peter G. Rowlands</i>	193

Level of Endemism in Hanging Gardens of the Colorado Plateau. <i>James F. Fowler, Nancy L. Stanton, Ronald L. Hartman,</i> <i>Cathleen L. May</i>	215
Tissue Culture of Mature Ponderosa Pine: Callus Induction and Differentiation. <i>Yiqun Lin, Michael R. Wagner</i>	225
Preliminary Survey of Leopard Frogs in Glen Canyon National Recreation Area. <i>Charles A. Drost, Mark K. Sogge</i>	239
Temperature Patterns of Rattlesnakes at National Bridges National Monument, Utah. <i>Tim B. Graham, Trevor Persons, William Schaedla,</i> <i>Donald Moore</i>	255
Cowbird Concentrations at Livestock Corrals in Grand Canyon National Park. <i>Matthew J. Johnson, Mark K. Sogge</i>	275
Observations of a Great Blue Heron Colony From 1987 to 1993. <i>Tim B. Graham, Ron W. Meyer</i>	285
Abert's Squirrels of the Colorado Plateau: Their Reproductive Cycle. <i>Gilbert C. Pogany, W. Sylvester Allred</i>	293

Foreword

The papers in this volume are contributions from National Biological Service (NBS) scientists, university students, and resource managers throughout the Colorado Plateau. The focus of all studies in this volume is on providing baseline scientific information on the physical, cultural, and natural resources of the Colorado Plateau. Support for these studies came from a myriad of federal, state, and private partners concerned about the well-being of the Plateau's resources.

The rich variety of the 68 presentations given at the conference and the 16 papers included here reflects the diversity of science presently being carried out on the Colorado Plateau. I applaud the effort of the contributors who, with modest funding and a broad base of public and institutional support, have pursued important lines of work in the four states that make up this vast biogeographic region.

All across America, we face the prospect of extensive environmental changes that will continue to affect the physical, cultural, and natural resources on our federal lands. As the biological and ecological research branch of the Department of the Interior, we in the NBS are committed to providing sound scientific information that can be used by both public lands managers and private landowners to conserve and manage natural resources.


National parks represent the best remaining examples of America's natural ecosystems and, as such, are baselines for measuring future change. To protect parks and other federal lands, we must increase our efforts to inventory and monitor their biological resources, to understand the factors, both natural and human-induced, that threaten them, and to assist with the design of resource management programs that will ensure that future generations can use and enjoy them as we have.

In short, we must provide the sound science needed to conserve and manage the natural resources that sustain us, inspire us, and represent our biological and environmental heritage.

H. RONALD PULLIAM

Director
National Biological Service

Cultural Resources



A Model to Study Fire Effects on Cultural Resource Studies of Mesa Verde

Lisa Floyd-Hanna

*Environmental Studies Program
Prescott College
Prescott, Arizona 86301*

William H. Romme

*Biology Department
Fort Lewis College
Durango, Colorado 81301*

Allan Loy

*National Park Service
Mesa Verde National Park
Mesa Verde, Colorado 83110*

David D. Hanna

*Environmental Studies Program
Prescott College
Prescott, Arizona 86301*

Abstract. On 8 July 1989, lightning struck the dry terrain of Long Mesa in Mesa Verde National Park and ignited a 2-week burn that scorched 1,200 ha. As a result, park management initiated studies of the effects of high-intensity fire on cultural resources. Mesa Verde National Park was established to protect works of prehistoric humans including pit-houses and cliff dwellings of the Anasazi. Later, park management was charged with the preservation of historic buildings and wood structures of 20th-century Americans. Our research will provide the necessary background information to predict specific fire effects on cultural sites. We mapped the vegetation communities, reconstructed the prehistoric fire history of half of Mesa Verde, and will document postfire succession. These data sets, incorporated into spatially explicit layers in the park's Geographic Information System, will be used to model the risk and specific effects of fire as related to particular classes of cultural resources.

Key words: Anasazi, postfire succession, vegetation mapping.

Mesa Verde is a series of north–south mesas in sandstone and shale substrates. Mancos Shale of marine origin is topped by Point Lookout Sandstone, then covered in parts of Mesa Verde National Park by the diverse Menifee Shale. The Cliffhouse Sandstone is exposed above these formations, and it is in this final layer that the most impressive cliff dwellings were built (Griffiths 1990). Water is ephemeral throughout the Mesa system—seeps and springs are infrequently encountered. The Mancos River forms the eastern boundary of the park and provides the only perennial water source.

We focused our postfire succession study on Long Mesa, the site of the 1989 fire that burned 1,200 ha. Long Mesa, one of the western mesas, ranges in elevation from 2,180 m in the south to 2,517 m at the north end. Long Canyon floor is 2,133 m. The vegetation on Long Mesa is a mosaic of mature piñon–juniper woodlands and mountain shrub associations. Shrub associations range from oak (*Quercus gambelii*) and serviceberry (*Amelanchier utahensis*) to sagebrush (*Artemisia nova*) and bitterbrush (*Purshia tridentata*).

Piñon (*Pinus edulis*) and juniper (*Juniperus osteosperma*) woodlands dominate the lower, southern ends of the mesas, whereas mountain shrub communities dominate on the northern ends. Differential fire frequency is hypothesized to be the major factor controlling this pattern (e.g., Erdman 1970). Douglas-fir (*Pseudotsuga menziesii*) forms dense stands on the north escarpment or on steep north-facing slopes. Ponderosa pine (*Pinus ponderosa*) is found in a few limited stands in Morefield and Prater canyons and in isolated pockets elsewhere. Scattered, small deciduous forest communities include aspen (*Populus tremuloides*) and maple (*Acer* sp.).

Methods and Preliminary Results

Mapping of Vegetation and Woody Fuels

The first step in modeling the effects of fire on cultural resources was to compile a digital map to define the vegetation of Mesa Verde to accurately predict the fire potential of each plant community. Landsat TM and SPOT panchromatic scenes from May and June 1990 were used as a spatial and spectral base. An unsupervised classification of the image and initial field surveys showed that 17% of the image was in shadow and that important vegetation characteristics were obscured. To mitigate the topographic effects in the image data, a transformation was used to separate the spectral and illumination

information (Pouch and Campagna 1990). The transformed image was then reclassified. Ground-truth efforts guided by this reclassification were made with relevé stand analyses (Mueller-Dombois and Ellenburg 1974) of more than 300 sites, each located with a Trimble Basic Pathfinder Global Positioning System (GPS). Species cover and abundance ratings were clustered with TWINSpan, a multivariate clustering program (Gauch 1982). This ordination was used to guide the final supervised classification of the image data with training signatures from the mean spectral signatures of the clustered sample points. In this way, the spectral and field information were related to one another.

Woody fuels were assessed in a subsample of the sampling points ($n = 26$). Canopy fuels—dead snags and woody fuels in live trees and shrubs—were measured (Meeuwig and Budy 1981). Downed fuels were sampled on transects with the plane-intercept method (Brown 1974). These data will be used in the fire behavior model BEHAVE (Burgan and Rothermel 1984) to predict the fire potential within the major vegetation communities in Mesa Verde National Park.

Fire History and Fire Effects

Our study of fire history and fire effects at Mesa Verde Park had two objectives (Floyd-Hanna and Romme 1993*¹). First was to document the postfire succession patterns following the 1989 fire on Long Mesa. This large fire affected at least three different vegetation types—piñon-juniper, mountain shrub, and Douglas-fir. Three permanent sampling grids containing 291 sampling points were established to document postfire patterns. Percent cover of plants, litter, or bare substrates was recorded in 1991, 1992, and 1994.

The second objective was to develop and apply a method of dating past fires in Mesa Verde (for background see Arno and Sneek 1977; Romme 1982). Because few fire-scarred trees are in Mesa Verde National Park and no unequivocal fire scars were located, we developed a method to age shrubs, which resprout vigorously after fires, using annual ring counts to determine their time of origin. All dominant shrub species were sampled and aged in the historically documented 1934, 1959, 1972, and 1989 burns. Although there is some variability within an individual plant as to the date of shoot emergence, we determined that by selecting the centermost shoot and restricting ourselves to the species *Quercus gambelii* we could determine reliably the known fire date. In

¹ Asterisk indicates unpublished material.

1992 and 1994, we applied this method to sample areas of unknown fire history in the northern portion of Mesa Verde National Park.

Results of the postfire succession sampling indicate that perennial resprouting has returned the cover of the northern sector of the burn to a shrub-dominated community much like the prefire community, whereas the southern burned areas (formerly piñon-juniper) are proceeding through a herbaceous phase of succession. The effects of prefire vegetation have been significant, and previous fire history has also affected the postfire successional patterns since the 1989 fire. Noxious weed invasion of the southern end of the fire is becoming an increasing problem.

Using the technique to age shrubs, we successfully identified and mapped prehistoric fires that occurred between 1850 and the 1920's. The median fire return interval varied from 55 years in the west to greater than 130 years in the eastern portion of the sampling area.

Discussion

Effects of Fire on Cultural Resources

These data will allow us to predict the probability of occurrence and the effect that a fire might have on a given type of archaeological site, depending on its location in the landscape. Data layers, each spatially defined for points or extrapolated to areas, will be as follows (Figure):

1. Median fire interval—expressed as a probability of occurrence: high = 0.02 (from a 50-year fire return interval), medium = 0.01 (from a 100-year fire return interval), and low < 0.01 (for fire return intervals greater than 100 years).
2. Vegetation—fuels and fire intensity potential—expressed as high, medium, or low intensity. High potential occurs in Douglas-fir forests or piñon-juniper woodlands, medium potential occurs in sparse piñon-juniper woodlands, and low potential occurs in mountain shrublands and meadows.
3. Significant and vulnerable cultural resources—Although all cultural resources at Mesa Verde have intrinsic value, some are more vulnerable to fires than other. The following classification was




Figure. A fire-risk summary of Mesa Verde National Park showing fire probability–fire intensity–vulnerability of cultural resources; L = low; M = medium; H = high. This is a simplified example of three layers for illustrative purposes.

developed by a group of archaeologists, ecologists, and fire behavior specialists (Romme et al. 1993):

A. Sites with high vulnerability to fire damage:

1. Native American historic structure (e.g., sweat lodges and corrals)
2. Alcoves and cliff dwellings containing organic materials (e.g., packet rat middens, wooden beams, corn cobs)
3. Rock-art panels

4. Culturally-scarred trees
- B. Sites with moderate vulnerability to fire damage:
 1. Euro-American historic structures (e.g., museum and administration buildings)
 2. Lithic scatter and shallow hearth
- C. Sites with low vulnerability to fire damage:
 1. Deeply buried, unexcavated pueblos
 2. Lithic scatter
 3. Check dams

A risk model is being developed by the authors and other National Park Service personnel where these three categories of risk overlap each other (Figure). Although each data layer is not yet complete, we begin to see the pattern of fire risk at Mesa Verde National Park as a mosaic of superimposed risk probabilities. For example, an area with dense fuels and highly valued cultural sites such as Chapin Mesa is at higher risk than an area that has burned frequently and has few archaeological resources (the north end of the park). Many of the most vulnerable sites—and perhaps those on which fire fighting efforts and *presuppression-suppression* activities such as fuel reduction should focus—are on the southern ends of Chapin and Wetherill mesas. High risk rating occurs there in two categories. Another high risk area is at the head of Prater Canyon where culturally scarred trees and dangerous fuels exist together.

We will continue to improve on the risk assessment for Mesa Verde National Park. The mass and configuration of woody fuels in each vegetation type will be used to predict actual fire behavior under particular sets of weather conditions (e.g., using BEHAVE, a fire behavior model), and this information will add significantly to our modeling potential. Fire history will be determined for the northeastern and southern portions of the park in 1994. Completion of the inventory and mapping of cultural sites in 1994 and combining of these data with the risk model will allow accurate assessment of the vulnerability of archaeological sites to fire.

Acknowledgments

All studies were funded by the Fire Initiative established by the Rocky Mountain Region of the National Park Service.

Cited Literature²

- Arno, S. F., and K. M. Sneek. 1977. A method for determining fire history in the coniferous forests of the Mountain West. U.S. Forest Service General Technical Report INT-42. 27 pp.
- Brown, J. K. 1974. Handbook for inventorying downed woody material. U.S. Forest Service Publication General Technical Report INT-16. 24 pp.
- Burgan, R. E., and R. C. Rothermel. 1984. BEHAVE: fire behavior prediction and fuel modeling system. U.S. Forest Service General Technical Report INT-167. 126 pp.
- Erdman, J. A. 1970. Pinyon-juniper succession after natural fires on residual soils of Mesa Verde, Colorado. Brigham Young University Biological Series Vol XI (2). 58 pp.
- *Floyd-Hanna, M. L., and W. H. Romme. 1993. Fire history and fire effects in Mesa Verde National Park, National Park Service, Rocky Mountain Region.
- Griffiths, M. O. 1990. Guide to the geology of Mesa Verde National Park. Mesa Verde Museum Association, Lorraine Press, Utah. 88 pp.
- Meeuwig, R. O., and J. Budy. 1981. Point and line intersect sampling in pinyon-juniper woodland. U.S. Forest Service General Technical Report INT-104. 88 pp.
- Mueller-Dombois, D., and H. Ellenburg. 1974. Aims and methods in vegetation ecology. Wiley & Sons, New York. 547 pp.
- Pouch, G. W., and D. J. Campagna. 1990. Hyperspherical direction cosine transformation for separation of spectral and illumination information in digital scanner data. *Photogrammetric Engineering and Remote Sensing* 56:475-479.
- Romme, W. H. 1982. Fire and landscape diversity in subalpine forests of Yellowstone National Park. *Ecological Monographs* 52:199-221.
- Romme, W. H., L. Floyd-Hanna, and M. Conner. 1993. Effects of fire on cultural resources at Mesa Verde National Park. *Park Science* 13(3):28-30.

²Asterisk indicates unpublished material.