

Congaree National Park

National Park Service
U.S. Department of the Interior

Inventory & Monitoring Program
Southeast Coast Network



Resource Brief: Supplementary Anuran Community Monitoring, 2011

Vital Sign Overview

Amphibian communities of the Southeast Coast Network (SECN) are monitored because of their global importance as ecological indicators, documented population declines and extinctions, and their high level of diversity in the Southeast. With an estimated 140 amphibian species, the Southeast accounts for about half of the total number of amphibians in the U.S.

The SECN has 61 known amphibian species: 26 in Caudata (salamanders, newts, amphiumas, sirens), and 35 in Anura (frogs and toads). Because of their complex life histories, habitat requirements, anatomy, and physiology, amphibians are considered to be good indicators of changes in ecosystem conditions as they are affected by climate change, land use development and conversion, contaminants, and changes in hydrology.

Significant Findings

We detected 133 vocalizations from 12 anuran species using automated recording devices (ARDs).

Spring peeper (*Pseudacris crucifer*) (Figure 1) and Southern leopard frog (*Lithobates sphenoccephalus*) had the highest frequency of occurrence over the subset of sampling locations at CONG.

Southern leopard frog had the highest relative detection frequency of vocalizations on the ARD recordings.

Spring peeper and Southern leopard frog were also the most widely distributed amphibians over the subset of sampling locations at CONG.

No non-native species were found.



Figure 1. Spring peeper (*Pseudacris crucifer*) at Congaree National Park. Photo by Briana Smrekar.

Sampling Effort

Data were collected at a subset of 12 of the 2010 sampling locations (Figure 2) using ARDs. Locations were chosen based on three criteria; 1) 2010 VES detections, 2) species of management concern were present or were expected to be present, and 3) representative of the types of freshwater resources available at CONG.

Auditory recordings were collected at the park from 3/2/2011 through 5/27/2011.

The full monitoring protocol will next be implemented at CONG in 2014.




Figure 2. Select sampling locations for ARD placement at Congaree National Park, 2011.

Monitoring Techniques

Determining the species that are present at a site can provide a great deal of insight to the condition of that location. Surveying for an elusive species, however, can be challenging and a single visit to a site cannot determine a species' true presence or absence. Further, weather variables such as temperature, precipitation, and humidity play a crucial role in whether or not an Anuran will be active or be vocalizing during a survey. Therefore, an approach maximizing sampling effort while minimizing safety hazards and cost is necessary to ensure a site is adequately characterized.

Automated recording devices (ARDs) were used to monitor anurans. Since most vocal anurans (i.e.,

frogs and toads that vocalize) call at night, monitoring these species can be challenging. Often times, personnel need to work through the night and this not only presents several safety concerns but also can be expensive. The ARD was deployed for 77 days and programmed to record at specific intervals throughout the evening, night, and morning; times when we know anurans are most likely to call. When viewed in specialized software, Anuran calls display unique patterns that can be identified through an automated process. Each ARD collects vocalizations within a sample site of 0.5 ha (~ 1acre).

About the Southeast Coast Network

The Southeast Coast Network (SECN) includes 20 parks, 17 of which contain significant and diverse natural resources (Figure 3). In total, SECN parks encompass more than 184,000 acres of federally-managed land across North Carolina, South Carolina, Georgia, Alabama, and Florida. The parks span a wide diversity of cultural missions also, including four national seashores, two national historic sites, two national memorials, seven national monuments, two national military parks, as well as a national recreation area, national battlefield and an ecological and historic preserve. The parks range in size from slightly more than 20 to nearly 60,000 acres, and when considered with non-federal lands jointly managed with NPS, the Network encompasses more than 253,000 acres.




Figure 3. Locations of parks within the Southeast Coast Network.

About the Inventory & Monitoring Program

In 1999, the National Park Service initiated a long-term ecological monitoring program, known as "Vital Signs Monitoring," to provide the minimum infrastructure to allow more than 270 national park system units to identify and implement long-term monitoring of their highest-priority measurements of resource condition. The overarching purpose of natural resource monitoring in parks is to develop scientifically sound information on the current status and long-term trends in the composition, structure, and function of park ecosystems, and to determine how well current management practices are sustaining those ecosystems.

The NPS Vital Signs Monitoring Program addresses

five goals for all parks with significant natural resources:

- Determine the status and trends in selected indicators of the condition of park ecosystem,
- Provide early warning of abnormal conditions,
- Provide data to better understand the dynamic nature and condition of park ecosystems,
- Provide data to meet certain legal and Congressional mandates, and
- Provide a means of measuring progress towards performance goals.

For More Information

SECN Home Page: <http://science.nature.nps.gov/im/units/secn/index.cfm>

SECN Reports & Publications: <http://science.nature.nps.gov/im/units/SECN/publications.cfm>

Inventory & Monitoring Program: <http://science.nature.nps.gov/im/index.cfm>

Data Downloads via the Natural Resource Information Portal: <http://nrinfo.nps.gov/Home.mvc>

Follow us on Twitter at: <http://twitter.com/SECoastNetwNPS> or @SECoastNetwNPS