



Vegetation Mapping at Fort Pulaski National Monument

Natural Resource Report NPS/SECN/NRR—2013/718



ON THE COVER

Fort Pulaski

Photograph courtesy of Jennifer Asper, NPS

Vegetation Mapping at Fort Pulaski National Monument

Natural Resource Report NPS/SECN/NRR—2013/718

Rachel H. McManamay, Anthony C. Curtis, and Sarah Corbett Heath

National Park Service
Southeast Coast Network
101 Wheeler Street
Saint Marys, GA 31558

October 2013

U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science
Fort Collins, Colorado

The National Park Service, Natural Resource Stewardship and Science office in Fort Collins, Colorado, publishes a range of reports that address natural resource topics. These reports are of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Data Series is intended for the timely release of basic data sets and data summaries. Care has been taken to assure accuracy of raw data values, but a thorough analysis and interpretation of the data has not been completed. Consequently, the initial analyses of data in this report are provisional and subject to change.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner.

This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data. Data in this report were collected and analyzed using methods based on established, peer-reviewed protocols and were analyzed and interpreted within the guidelines of the protocols.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available from the Southeast Coast Network (<http://science.nature.nps.gov/um/units/secn>) and the Natural Resource Publications Management website (<http://www.nature.nps.gov/publications/nrpm/>). To receive this report in a format optimized for screen readers, please email irma@nps.gov.

Please cite this publication as:

McManamay, R. H, A.C. Curtis, and S.C. Heath. 2013. Vegetation mapping at Fort Pulaski National Monument. Natural Resource Report NPS/SECN/NRR—2013/718. National Park Service, Fort Collins, Colorado.

Contents

	Page
Figures.....	v
Tables.....	vii
Executive Summary.....	ix
Acknowledgments.....	xi
Introduction:.....	1
Fort Pulaski National Monument Vegetation Mapping Project	1
The Vegetation Inventory Program	1
Vegetation Inventory Program Standards.....	1
Protocols	1
Standards.....	1
National Vegetation Classification Standard.....	2
Federal Geographic Data Committee Standards.....	3
Methods.....	5
Study Area	5
Location	5
Vegetation.....	6
Climate and Weather.....	6
Soils.....	6
Project Scoping and Planning.....	12
Project Boundary Extent	12
Field Survey.....	12
Location and Plot Identifiers.....	14
Environmental Description	14
Vegetation Description	14
Other Ancillary Information Useful to Classification	14
Vegetation Classification.....	14
Final Map Classifications	15

Contents (continued)

	Page
Digital Imagery and Interpretation	16
Aerial Photography Acquisition and Processing	16
Preliminary Photo-Interpretation	19
Field Reconnaissance.....	19
Photo-Interpretation and Polygon Attribution	19
Field Verification	21
Accuracy Assessment	21
Sampling Design.....	22
Field Data Collection	24
Data Analysis	24
Results.....	27
Vegetation Classification.....	27
Digital Imagery and Interpretation	27
Vegetation Map	29
Map Layer Presentation	32
Accuracy Assessment	35
User's Accuracy (Errors of Commission).....	35
Producer's Accuracy (Errors of Omission).....	36
Literature Cited	39

Figures

	Page
Figure 1. Location of Fort Pulaski National Monument (map courtesy Jennifer Asper, Southeast Coast Network).....	5
Figure 2. The largest vegetation association found in Fort Pulaski National Monument is the Southern Atlantic Coast Salt Marsh (Photo courtesy: Jennifer Asper, National Park Service)	7
Figure 3. Example of drainage canal at Fort Pulaski National Monument.....	8
Figure 4. Cedar – Live Oak – Cabbage Palmetto Marsh Hammock that occurs on dredge spoils at Fort Pulaski National Monument.....	9
Figure 5. Chinese Tallow (<i>Triadica sebifera</i>) is one non-native of concern in Fort Pulaski National Monument.	10
Figure 6. Soils of Fort Pulaski National Monument (Spatial data obtained from SSURGO, available at http://SoilDataMart.nrcs.usda.gov).	11
Figure 7. Locations of plots sampled in Fort Pulaski National Monument used for vegetation classification by NatureServe.....	13
Figure 8. Flight paths for the 2009 color-infrared aerial photography coverage of Fort Pulaski National Monument.	18
Figure 9. Locations of accuracy assessment points sampled in Fort Pulaski National Monument.	23
Figure 10. Vegetation (NVCS 1997) and Land Use (Anderson et al. 1976) map classes for Fort Pulaski National Monument.	33
Figure 11. Distribution of wetlands (Cowardin et al. 1979) in Fort Pulaski National Monument.	34

Tables

	Page
Table 1. Example of National Vegetation Classification (NVCS) hierarchy for terrestrial vegetation (FGDC 1997).	3
Table 2. Vegetation height and cover classes used in the Fort Pulaski National Monument Vegetation Mapping Project.....	14
Table 3. Map classification codes for wetlands derived from the U.S. Fish and Wildlife Service’s Classification of Wetlands and Deepwater Habitats of the United States (Cowardin et al. 1979).....	16
Table 4. Fields used in the attribute table of the vegetation polygon layer of the Fort Pulaski National Monument Vegetation Mapping Project.	20
Table 5. Standard sample size allocations for accuracy assessment, based on map class area (Lea and Curtis, 2010).....	22
Table 6. Final vegetation associations of the NVCS used for vegetation mapping (with their assigned Map Codes) in Fort Pulaski National Monument.	28
Table 7. Map classification codes for non-vegetated areas in Fort Pulaski National Monument, following Land Use and Land Cover Level II (Anderson et al. 1976).	28
Table 8. Summary statistics for the map classes occurring in Fort Pulaski National Monument.	30

Appendices

	Page
Appendix A: Plot Sampling Form	43
Appendix B: Descriptions of Vegetation Types	47
Appendix C: Field Key to Vegetation Types Draft Key to the National Vegetation Classification (NVC) Associations that Occur or Potentially occur at Fort Pulaski and Fort Sumter National Monuments	129
Appendix D: Photo-Interpretation Guide.....	133
Appendix E: Field Reconnaissance Form.....	153
Appendix F: Accuracy Assessment Form.....	157
Appendix G: Accuracy Assessment Contingency Table	161

Executive Summary

The Southeast Coast Network (SECN) worked with the support of the National Park Service (NPS) Vegetation Inventory Program (VIP) to describe and map vegetation at Fort Pulaski National Monument (FOPU). This mapping effort was accomplished through collaboration with the Southeast regional office of NatureServe (Raleigh/Durham, NC).

The mapping area is 2,543 ha, encompassing the entire boundary of FOPU and an additional buffered area of adjacent land. NatureServe ecologists identified plant associations for FOPU and a remote sensing specialist from the SECN completed the vegetation mapping using 1:12,000-scale, color-infrared aerial photography and digital orthophotography. NatureServe collected vegetation and environmental data from 17 vegetation classification plots, and the Remote Sensing Specialist assessed 47 observation points. Two Student Conservation Association (SCA) interns hired by the SECN conducted an accuracy assessment (AA), which entailed sampling 133 additional points. Fieldwork and mapping were completed between 2005 and 2013.

A total of 25 map classes were developed to map the vegetation and general land cover of FOPU and surroundings, including the following: 12 map classes representing natural/semi-natural vegetation at the association level in the National Vegetation Classification System (NVCS), two map class representing cultural vegetation (e.g. developed) in the NVCS, and 11 map classes representing non-vegetated units (e.g. open water bodies, buildings, roads, etc.). Features were interpreted using 1:12,000 scale digital color-infrared aerial photography (flown 31 May 2009) through heads-up-digitizing in ArcGIS (Version 10.0, © 2010 Environmental Systems Research Institute, Redlands, California). Polygons were mapped to a 0.5-ha minimum mapping unit (MMU).

A geodatabase containing various feature class layers and tables show the locations of vegetation types and general land cover (vegetation map), vegetation plot samples, AA sites, project boundary extent, and aerial photographic centers. The feature class layer for the FOPU vegetation map provides 251 polygons of detailed attribute data covering 2,543 ha, with an average polygon size of 11.2 ha. Of the area mapped, 177 polygons (70.5% of all polygons) represent natural/semi-natural vegetation types in the NVCS, encompassing 1,834.6 ha (72.1%) of the total map extent.

Summary reports generated from the vegetation map layer indicate that the Southern Atlantic Coast Salt Marsh dominates the vegetated landscape (57 polygons covering 1,488 ha), with the Needlerush High Marsh (20 polygons, 91.4 ha) and the Cedar – Live Oak – Cabbage Palmetto Hammock (21 polygons, 52.9 ha) also covering a considerable portion of the park. The Southern Atlantic Coast Salt Marsh occupies the greatest area of any of the vegetation communities by far and is best developed in areas between mean high and low tides that are regularly flooded. The Needlerush High Marsh occurs within the salt marsh in areas slightly higher than where the Southern Atlantic Coast Salt Marsh is found. The Cedar – Live Oak – Cabbage Palmetto Marsh Hammock is a maritime forest community located on the highest elevations on Cockspur Island, typically on the older spoil deposits.

A thematic AA study was conducted on the eleven map classes within the boundary of FOPU representing floristic types within the NVCS. Results present an overall accuracy of 90.3% (Kappa statistic = 59%) based on data from 129 AA sites.

Products resulting from the FOPU vegetation mapping project include:

Available in this report:

- Project summary of methods and results
- Illustrated photo-interpretation guide
- Detailed descriptions of vegetation associations
- Samples of field forms

Available elsewhere¹:

- Geodatabase containing map-polygon attribute, aerial photography flight lines, plot data and Monument and project boundaries
- Ground photography of vegetation plots, observation points, and AA points in hard copy and digital formats
- All field data (plot, observation point, and AA point) stored in a Microsoft Access database
- Hard copy vegetation maps
- Hard copy orthophotographs in DOQQ format and stereo aerial photograph prints
- Metadata for all digital products

Geospatial products are in the Universal Transverse Mercator (UTM) projection, Zone 17, using the North American Datum of 1983.

¹ Products and additional information can be found at the following website:
<http://biology.usgs.gov/npsveg/>

Acknowledgments

Funding for this project was provided by the NPS VIP. The Southeast regional office of NatureServe (Raleigh/Durham, NC) completed the field survey and other tasks associated with the vegetation classification portion of the mapping project. The SECN completed the vegetation mapping and AA. Special thanks to the staff of Fort Pulaski National Monument for their cooperation with this project.

Introduction:

Fort Pulaski National Monument Vegetation Mapping Project

The FOPU Vegetation Mapping project was organized and coordinated by the SECN between 2005 and 2013, with assistance from the Southeast regional office of NatureServe for the vegetation classification portion of the project. The purpose of this project was to inventory and map existing plant associations on approximately 2,543 ha of FOPU, including a buffered area surrounding the park. The mapping information is provided in written, tabular, digital, and spatial formats that are useful to resource managers, the SECN, and others. The mapping product will serve as a useful baseline of vegetation information that will assist park natural and cultural resource stewardship. The identification and description of plant communities also provides habitat information important to understanding associated organisms (reptiles, amphibians, mammals, etc.) and providing inference to the location and abundance of species that are characteristic of each community.

The Vegetation Inventory Program

The VIP is an effort by the NPS to classify, describe, inventory, and map vegetation communities in more than 280 national park units across the United States. The primary objective of the VIP is to produce high-quality, standardized maps and associated data sets of vegetation and other land cover occurring within parks. Since vegetation species and communities are unique from park to park, the inventory of these resources assists park managers on a wide array of management issues. For more information about the VIP, visit <http://science.nature.nps.gov/im/inventory/veg/index.cfm>.

Vegetation Inventory Program Standards

The NPS and VIP established guidance and standards for all vegetation mapping projects in a series of documents:

Protocols:

- National Vegetation Classification System (TNC and ESRI 1994a, NatureServe 2003)
- Field methods and mapping procedures (TNC and ESRI 1994b)
- Statistically rigorous and consistent AA procedures (ESRI and TNC 1994, Lea and Curtis 2010)
- Guidelines for using existing vegetation data (TNC 1994)

Standards:

- The National Vegetation Classification Standard (FGDC 1997, FGDC 2008, NatureServe 2009)
- Spatial Data Transfer Standard (FGDC 1998b)
- Content Standard for Digital Geospatial Metadata (FGDC 1998a)
- United States National Map Accuracy Standards (USGS 1999)
- Integrated Taxonomic Information System (U.S. Department of Agriculture)
- Program-defined standards for map attribute accuracy and MMU

National Vegetation Classification Standard

The NVCS is the mapping standard used in SECN vegetation mapping projects. The NVCS, adopted by the Federal Geographic Data Committee (FGDC 1997), evolved from work conducted by The Nature Conservancy, NatureServe, and the Natural Heritage Program network over more than two decades. The NVCS standard supports the development and use of a consistent national vegetation classification throughout the NPS and other organizations. Such a standard facilitates the production of uniform statistics about vegetation resources across the nation and makes cooperation on vegetation management issues across jurisdictional boundaries possible (Grossman et al. 1998). The FGDC Vegetation Subcommittee works to keep the standard relevant and current. Revisions made to the upper levels of the NVCS hierarchy adopted by the Vegetation Subcommittee as Version 2 (FGDC 2008) were not used in this project since the crosswalks for vegetation associations were unavailable at the time this project began.

Vegetation classification systems attempt to recognize and describe repeating assemblages of plants in similar habitats. The NVCS is a hierarchical system that incorporates physiognomic characters and floristic data to define seven levels of terrestrial vegetation classification. The five upper levels (class, subclass, group, subgroup, and formation) are based on physiognomic features. The two lower levels (alliance and association) are distinguished by differences in floristic composition. The physiognomic units have a broad geographic perspective and the floristic units have utility in local and site-specific applications (Grossman et al. 1998). The physiognomic levels of the NVCS are based on physical, structural, and environmental characteristics identifiable from satellite imagery, aerial photography, or ground observations (Table 1). Specific criteria defining these physiognomic units are based on ecologic characteristics that vary among major vegetation groups (FGDC 1997).

The alliance and association levels form the base of the NVCS hierarchy and are determined by the most abundant or diagnostic species comprising the various layers of a homogenous vegetation community. An association is here defined as a plant community type with a consistent species composition, uniform physiognomy, and similar habitat conditions (Flahault and Schroter 1910). Species composition differentiates associations (TNC and ESRI 1994a). An alliance is "a physiognomically uniform group of plant associations sharing one or more dominant or diagnostic species which, as a rule, are found in the uppermost strata of the vegetation." (Reid and Comer 1998). NatureServe coordinates plant association data for the SECN vegetation mapping projects. Associations are added to the NVCS and older concepts are refined as new data become available.

For more information about the NVCS, see the USGS-NPS Vegetation Mapping Program standards (<http://biology.usgs.gov/npsveg/standards.html>) or Grossman et al. 1998.

Table 1. Example of National Vegetation Classification (NVCS) hierarchy for terrestrial vegetation (FGDC 1997).

Level	Criteria Delineating Level	Example
Class	Structure (height, cover) of dominant vegetation strata	Forest
Subclass	Growth form characteristics including leaf type (evergreen, deciduous) for woody plants and persistent (perennial, annual) herbaceous species	Evergreen forest
Group	Leaf morphology (broad leaf, microphyllous, xeromorphic), leaf phenology, and climatic conditions	Temperate or subpolar needle-leaved evergreen forest
Subgroup	Relative degree of human disturbance	Natural/Semi-natural temperate or subpolar needle-leaved evergreen forest
Formation	Additional physiognomic characteristics, general environmental conditions, relative landscape position, and hydrologic regimes	Rounded-crowned temperate or subpolar needle-leaved evergreen forest
Alliance	Dominant or diagnostic species of uppermost or dominant stratum	<i>Pinus taeda</i> Forest Alliance
Association	Other dominant or diagnostic species from any stratum	<i>Pinus taeda/Liquidambar styraciflua-Acer rubrum var. rubrum/Vaccinium stamineum</i> Forest

Federal Geographic Data Committee Standards

In addition to vegetation classification, the FGDC sets standards for map spatial accuracy and for metadata employed in NPS vegetation mapping projects. Standards for map products stipulate map scales of 1:24,000 or finer, and minimum polygon size of 0.5 ha (1.24 acres). Positional accuracy for vegetation maps must meet National Map Accuracy Standards, which specify horizontal errors of less than 10.2 m (33.5 ft) on the ground for 1:12,000-scale maps.

All digital vegetation products resulting from this project are accompanied by FGDC-compliant metadata. Metadata are “data about the data,” and describe the content, quality, condition, and other characteristics of the spatial dataset. Metadata are critical elements that expedite the interpretation and exchange of information among users.

Methods

Study Area

Location

FOPU is located in Chatham County, Georgia about 15 miles (24.1 km) east of Savannah (Figure 1). FOPU is located along the Savannah River, only a few miles from its junction with the Atlantic Ocean, and is comprised of two islands, Cockspur Island and McQueens Island.



Figure 1. Location of Fort Pulaski National Monument (map courtesy Jennifer Asper, Southeast Coast Network)

Vegetation

The vegetation at FOPU can generally be divided into wetland and upland communities. Approximately 90% of FOPU is classified as wetland. McQueens Island is composed of almost entirely wetland communities, a large portion being tidal marsh dominated by *Spartina alterniflora* (smooth cordgrass) (Figure 2); whereas, Cockspur Island is composed of both wetland and upland communities. The vegetation of Cockspur Island has been significantly impacted from human activities over at least the last couple hundred years. In the early nineteenth century the island would have been primarily salt marsh with some coastal hammock forest. In order to develop the island for defense, the installation of drainage canals (Figure 3) and a dike system modified the island. During this time any existing vegetation would have been removed to enhance visibility. Later, dredge spoil from the Savannah River was deposited on the island, further impacting the natural scene. Since the abandonment of the fort in the late eighteenth hundreds, a considerable portion of Cockspur Island has reverted to maritime forest. Currently, there are approximately 125 ha of upland areas which support a mosaic of maritime forest (Figure 4), shrub communities, maintained grasslands and successional spoil deposit areas.

The past history of Cockspur Island in regard to its alteration by man for fortification and spoil deposits is reflected in the number of non-native species found there. In addition, the close proximity to the port of Savannah and the long history of maritime trade along the Savannah River provides further opportunities for species not native to the area to become established. Sixty-eight species of plants have been recorded at the Monument that would be considered non-native. Many of these non-native species have been introduced in the landscape of dredge spoil and maintained grass areas for so long that they may be considered naturalized. Eighteen species have been recorded on the site that would be considered of concern to management because they tend to be invasive or disruptive to natural communities. Of these, Chinese tallowtree (*Triadica sebiferum*) (Figure 5) and China-berry (*Lonicera japonica*) are, perhaps, the non-native species of greatest concern. Although both of these species are well established on spoil deposits on Cockspur Island, they are not well established yet in the mature, maritime forest in which the park nature trail is located (courtesy [NPS.gov](https://www.nps.gov)).

Climate and Weather

Summer months at FOPU are hot and humid. The mean monthly maximum temperature in July ranges from 31.1 – 32.0° C (88.0 – 89.6° F) (1961-1990). Winter months at FOPU are typically mild with only brief cold spells. The mean monthly minimum temperature in January ranges from 4.1 – 6.0° C (39.3 – 42.8 °F) (1961-1990). Precipitation mostly comes in the form of rain from storm fronts in winter and spring months, and from thunderstorms, tropical storms, and hurricanes in the summer and fall months. FOPU receives an average precipitation amount of 1201-1400 mm/year (47.3 – 55.1 inches/year) (1961-1990) (Davey et al. 2007).

Soils

According to Soil Survey Geographic (SSURGO), there are two soils within the park boundaries of FOPU: Made Land and Tidal Marsh, Salty (Figure 6). The Made Land represents areas that have been excavated, filled, or otherwise disturbed by man. Tidal marsh represents areas of marshy soil covered by saltwater at high tide. The material is largely black to bluish-gray heavy silt clay, containing large amounts of shells and decaying organic matter. The Tidal Marsh areas are covered mainly by a thick growth of grass and/or rushes.



Figure 2. The largest vegetation association found in Fort Pulaski National Monument is the Southern Atlantic Coast Salt Marsh (Photo courtesy: Jennifer Asper, National Park Service)



Figure 3. Example of drainage canal at Fort Pulaski National Monument.



Figure 4. Cedar – Live Oak – Cabbage Palmetto Marsh Hammock that occurs on dredge spoils at Fort Pulaski National Monument.



Figure 5. Chinese Tallow (*Triadica sebifera*) is one non-native of concern in Fort Pulaski National Monument.




Figure 6. Soils of Fort Pulaski National Monument (Spatial data obtained from SSURGO, available at <http://SoilDataMart.nrcs.usda.gov>).

Project Scoping and Planning

In 2005, the SECN proposed to develop vegetation classification and maps for 16 parks (Cape Hatteras National Seashore, Cape Lookout National Seashore, Moores Creek National Battlefield, Fort Sumter National Monument, Fort Pulaski National Monument, Fort Frederica National Monument, Cumberland Island National Seashore, Fort Matanzas National Monument, Timucuan Ecological and Historic Preserve, Fort Caroline National Monument, Castillo de San Marcos National Monument, Canaveral National Seashore, Ocmulgee National Monument, Horseshoe Bend National Military Park, Kennesaw Mountain National Battlefield Park, and Chattahoochee River National Recreation Area). The SECN teamed with the Southeast regional office of NatureServe to complete this project. NatureServe's role was to collect field data and develop the local association descriptions and the field key to map classes for each of the parks. The SECN agreed to complete the photographic interpretation/image analysis, digital map and database development, AA, metadata, and final reports in-house.

Project Boundary Extent

The SECN determined that the FOPU vegetation mapping boundary extent should include the entire area within the park plus an additional buffer surrounding the park boundary of 50 meters. Therefore, the total map extent is 2,543.3 ha, with 2233.2 ha occurring within park boundaries and the remaining 310.1 ha occurring outside of park boundaries.

Field Survey

Prior to field work, NatureServe compiled a list of preliminary vegetation associations and alliances likely to occur in FOPU. Published floristic information, the plant- species list of FOPU, and expert local knowledge, including NPS staff at FOPU, were used to refine the list. The preliminary list of associations was useful for planning, estimating the amount of field work necessary, and for assigning provisional names to vegetation plots and observation points.

Plots were placed non-randomly in such a way as to sample as many of the vegetation associations as possible given the minimum number of plots for the project (C. Nordman, pers. comm., 21 Feb. 2011). Plots were deliberately placed in certain areas in order to capture the diversity of vegetation communities at the park. Plot size was variable (e.g., 10 m x 10 m, 10 m x 30 m, 20 m x 20 m, 20 m x 40 m, 20 m x 50 m) and decided at the time of sampling based on which dimension would most adequately characterize the vegetation sampled in each plot. Seventeen vegetation plots were sampled in FOPU in November 2007 (Figure 7).

Within each plot, field staff estimated and recorded an array of vegetation and environmental data using the field form featured in Appendix A. The following four categories of data were collected for vegetation plots:




Figure 7. Locations of plots sampled in Fort Pulaski National Monument used for vegetation classification by NatureServe.

Location and Plot Identifiers

The locations of the vegetation plots were not permanently marked. The bounds of each plot were marked temporarily using measuring tapes. The UTM coordinates at the center of each lot were recorded (Zone 17, NAD83) on Garmin® hand-held GPS receivers.

Environmental Description

The physical characteristics of each plot were documented in both categorical and narrative fields. These characteristics included physical site features (e.g., elevation, slope, aspect, topography), hydrology, geology, and soils. Description of the ground surface included an estimate of the cover of rocks, sand, litter, bare soil, biological soil crust, moss, and lichen. A narrative field was provided for a general description of the plot setting and the influence of disturbance and hydrology on the vegetation.

Vegetation Description

Every vascular plant was assigned to one of eight physiognomic strata. Within each stratum, the investigator recorded the height class for the most abundant species (Table 2). Percent abundance, based on a cover class percentage, for each species present in the plot was also recorded (Table 2). Field crews also recorded percent canopy cover by species.

Table 2. Vegetation height and cover classes used in the Fort Pulaski National Monument Vegetation Mapping Project.

Code	Height Scale (m)	Code	Cover Class %
1	< 0.5	1	Trace
2	0.5 - 1.0	2	0.1 - 1
3	1.0 - 2.0	3	1.0 - 2.0
4	2.0 - 5.0	4	2.0 - 5.0
5	5.0 - 10.0	5	5.0 -10.0
6	10.0- 15.0	6	10.0-25.0
7	15.0 - 20.0	7	25.0 -50.0
8	20.0 - 35.0	8	50.0 - 75.0
9	35.0 - 50.0	9	75.0 -95.0
10	> 50.0	10	> 95

Other Ancillary Information Useful to Classification

Field crews optionally recorded how well the plot represented the vegetation of the surrounding area, site disturbance history, and the overall character of the vegetation features of each plot.

Vegetation Classification

Plot data was manually entered into the NatureServe PLOTS Database Version 3.0, a Microsoft Access database designed for the NPS VIP (NatureServe 2010). The PLOTS database was designed specifically for the NPS vegetation and mapping program so that the electronic data entry fields mirror the standard field form. Data entry was facilitated by assigning each plant taxon a unique, standardized code and name based on the PLANTS database developed by the

Natural Resources Conservation Service in cooperation with the Biota of North America Program (USDA and NRCS 2009). Data was thoroughly proofed after entry to minimize errors.

Due to a relatively small sample size, cluster analysis techniques that have been traditionally used in other vegetation mapping projects to generate vegetation classifications were not used for FOPU. Instead, plots were assigned to an existing vegetation association based on a cursory overview of their best fit according to species composition (R.White, pers. comm., 25 March 2011). The vegetation associations and descriptions can be found on the NatureServe Explorer (see <http://www.natureserve.org/explorer/index.htm>) (NatureServe 2009).

Final Map Classifications

Once the associations were finalized, a dichotomous key, that was originally developed by NatureServe and later revised by the SECN (Appendix C), was updated. The full NVCS hierarchical classification and global descriptions are available in Appendix B. FOPU specific vegetation (e.g., local) descriptions were written based on FOPU plot data. In addition, the final associations were linked to map classes for use in the photo-interpretation and mapping portions of the project.

Vegetation that did not fall into the natural/semi-natural category (e.g., maintained lawn) was assigned to the Cultural Vegetation of the NVCS at the Type level (NVCS Version 2, FGDC 2008).

For non-vegetated features (e.g., open water bodies, structures, etc.), we derived map classes from Level II of the Land Use and Land Cover Classification System (Anderson et al. 1976). We slightly modified several of the land use and land cover classes so they would be more appropriate for this mapping project (e.g., ‘Mud Flat’ and ‘Tidal Wrack’).

We further classified vegetated wetlands (natural/semi-natural vegetation) to the System Level of the U.S. Fish and Wildlife Service’s Classification of Wetlands and Deepwater Habitats of the United States (Cowardin et al. 1979). The wetlands within FOPU are considered either Palustrine or Estuarine systems (Table 3).

Table 3. Map classification codes for wetlands derived from the U.S. Fish and Wildlife Service's Classification of Wetlands and Deepwater Habitats of the United States (Cowardin et al. 1979).

Map Class Code from U.S. Fish and Wildlife Wetland Classification	General Explanation of Wetland Systems
Upland	Non-Wetlands
Marine	Open ocean overlying the continental shelf and its associated high energy coastline
Estuarine	Deepwater tidal habitats and adjacent tidal wetlands in which ocean water is at least occasionally diluted by freshwater runoff from the land
Riverine	Wetlands and deepwater habitats contained within channels, except where wetlands are dominated by vegetation or where salinity exceeds 0.5‰
Lacustrine	Wetlands and deep water habitats situated in a topographic depression or dammed river channel, lacking vegetation with greater than 30% aerial coverage, total area exceeds 8 hectares (ha)
Palustrine	All non-tidal wetlands dominated by vegetation and all such wetlands that occur in tidal areas where salinity is below 0.5‰

Digital Imagery and Interpretation

Aerial Photography Acquisition and Processing

The SECN contracted with Aero-Metric Inc., through the USGS Rolla, Missouri office to provide color-infrared aerial photographs. Color-infrared imagery was chosen because healthy green vegetation is a very strong reflector of infrared radiation and appears bright red in color-infrared imagery, which helps tremendously in vegetation mapping efforts. This set of photography covered the full project boundary extent plus a buffer surrounding the boundary of 500 meters. The flight path and centers of the aerial photo mission were planned with 60% forward-lap and 30% side-lap in order to assure stereo viewing (Figure 8). Airborne GPS data was provided by Aero-Metric, Inc., as was the ground check point data. The project was referenced to the North American Datum of 1983/2007 (NAD83/07), UTM Zone 17 horizontally and to the North American Vertical Datum of 1988 vertically. The aerial photographs were taken on 31 May 2009 at a scale of 1:12,000. The calibration focal length for the photos was 152.823 mm. Contact prints of the CIR imagery were created for use in the field, and diapositives of the CIR set were created for photo-interpretation. Scanned images (13 micron) were processed to 0.3 meter (1.0 ft.) resolution and provided in a digital orthophotograph mosaic. The scanned imagery

was rectified through an aerotriangulation process. The scanned images, Airborne Global Positioning System (ABGPS) data, ground check point data, and camera calibration data were used as inputs to the Zeiss/Intergraph ImageStationAutomatic Triangulation (ISAT) softcopy program. ISAT correlated image points are aerotriangulated in a block of images to create the exposure station exterior orientations. Three ground check points were manually measured on the imagery. All of these points had vertical and horizontal coordinates. Statistics associated with the aerotriangulation bundle adjustment are as follows:

1. 43 photographs
2. 1290 image points
3. 4 Horizontal and Vertical ground control points
4. 7.1 microns image sigma
5. 1350 degrees of freedom
6. Root mean square error of residuals for 4 ground control points:
 - a. 0.068 meters for horizontal-control eastings
 - b. 0.048 meters for horizontal-control northings
 - c. 0.025 meters for vertical control
7. Root mean square error of residuals for 43 ABGPS exposure station control points:
 - a. 0.024 meters for horizontal-control eastings
 - b. 0.019 meters for horizontal-control northings
 - c. 0.069 meters for vertical control




Figure 8. Flight paths for the 2009 color-infrared aerial photography coverage of Fort Pulaski National Monument.

Preliminary Photo-Interpretation

Prior to field reconnaissance, it is advantageous for the individual(s) interpreting the imagery to become familiar with the project imagery as well as the vegetation types to be mapped. The benefit of this early mapping step is that it allows the vegetation mapper(s) to identify particular areas of interest to visit during the field reconnaissance step. For this preliminary mapping, the SECN Remote Sensing Specialist delineated vegetation polygons on the digital orthophoto mosaics to at least the Formation-level of the NVCS. In addition to vegetation, other features were mapped as well, such as non-vegetated surfaces that were classified according to the Anderson Land Use Level II mapping regime (Anderson et al. 1976). Polygons were digitized on-screen using Arc GIS (Version 10, © 2010 Environmental Systems Research Institute, Redlands, California).

Field Reconnaissance

The purpose of this step is to train the vegetation mapper(s) on the correlation of photo-signatures with vegetation associations. A photo signature is a set of diagnostic characteristics present on the imagery, such as color, tone, shape, size, etc. that represents a particular vegetation association on the imagery. During field reconnaissance it is important to explore a variety of locations that represent various images, since color-infrared (CIR) imagery is typically not consistent between photo sets. Even slight differences in sun angle, light intensity, shadow, and even the particular film batch and printing process can affect vegetation's appearance on imagery (Hershey and Befort 1995). The ability of the photo-interpreter to identify diagnostic characteristics on the imagery that can be tied directly to a vegetation association is a key component to successful and accurate mapping. For some vegetation types, a set of diagnostic characteristics may not be visible on the imagery (e.g., the understory species may be the ones that are diagnostic). In such cases, the photo-interpreter may be able to rely on an informed understanding of the ecological and environmental settings of the particular vegetation types (e.g., the environmental setting where an association is likely to occur, etc.) in order to make a mapping decision. Unfortunately, there are still instances where a photo signature cannot clearly be identified for each association. In these instances it is allowable to pool associations and map at a higher level of the hierarchy, or with less accuracy (ESRI 1994).

A field reconnaissance trip was taken to FOPU in February 2011. During this trip, the SECN Remote Sensing Specialist spent time investigating ground conditions with copies of the aerial photographs in hand. Notes were taken directly on the copies of the aerial imagery as observations were made. The Remote Sensing Specialist used a Garmin® Rino GPS unit to navigate to a set of predetermined points of interest within the vegetation polygons mapped during the preliminary mapping step. These points of interest were chosen specifically to aid in the identification of vegetation photo-signatures. Upon reaching each point, vegetation at that point was keyed to its association (as possible) by using the key developed by NatureServe and revised by the SECN. Difficulties in determining association, if any, were documented on the field sheet (Appendix E). Dominant species found in each vegetation strata were documented. Ground photographs of each point were taken.

Photo-Interpretation and Polygon Attribution

Based on the information gathered from the field reconnaissance trip, the Remote Sensing Specialist developed a photo-interpretation guide that aided the delineation of vegetation classes (Appendix D). Standard photo signature characteristics were applied during the photo-

interpretation, including texture, color, pattern, and position in the landscape (Philipson 1997). In addition to photo signature characteristics, understanding the environmental distribution of the vegetation types helped the Remote Sensing Specialist not only identify types, but also to properly place polygon boundaries. Polygons were interpreted and digitized on-screen using Arc GIS. Features were mapped to a MMU of 0.5 ha, which is the NPS VIP standard. Exceptions for mapping below the MMU were allowed for map class units with vegetation unique to the immediate surroundings. For each polygon, the appropriate map class code along with its NVCS code, alliance, association, association common name were entered into the fields of the associated attribute table (Table 4).

Table 4. Fields used in the attribute table of the vegetation polygon layer of the Fort Pulaski National Monument Vegetation Mapping Project.

Field in Attribute Table	Description	Polygon Types Field Applies to:
Cultural Type	NVCS Cultural Vegetation Type	Cultural Vegetation
Alliance	NVCS Alliance scientific name	Natural/Semi-natural vegetation
Association	NVCS Association scientific name	Natural/Semi-natural vegetation
Common_Name	NVCS Association Common Name	Natural/Semi-natural vegetation
LUC_II	Land Cover and Land Use Level II Codes (Anderson et al. 1976)	Natural/Semi-natural vegetation, Cultural Vegetation, and Non-Vegetated Areas
ELCODE	NVCS Element Code	Natural/Semi-natural vegetation
Map Code	Abbreviated code used by photo-interpreter	Natural/Semi-natural vegetation, Cultural Vegetation, and Non-Vegetated Areas
Wetland	Type of wetland	Natural/Semi-natural vegetation
Hectares	Size of polygon in hectares	Natural/Semi-natural vegetation, Cultural Vegetation, and Non-vegetated Areas

Field Verification

A second trip was taken to FOPU in June 2011 for field verification. Field verification allows the vegetation mapper(s) to gauge how accurately the photo-signatures developed during the field reconnaissance step are with respect to consistently mapping vegetation associations on the ground. Specific issues with problem classes can be resolved during this step prior to submitting the map for AA.

Methodology for the field verification is nearly identical to the field reconnaissance step. The SECN Remote Sensing Specialist navigated via GPS to a predetermined set of points. Vegetation at that point was keyed to its association by using the vegetation key developed by NatureServe (NatureServe 2008) and revised by SECN. Difficulties in determining association, if any, were documented on the field sheet.

Accuracy Assessment

The main goal of an AA is to measure the probability that a particular location on the map created from the interpreted imagery has been assigned to its correct vegetation class on the ground. A vegetation map is an abstract model of the distribution of vegetation on the ground that is made possible by making some generalizations. These generalizations lead to errors in the map product. These map errors must be quantified so that the map user and the map producer understand the limitations of the map. A formalized AA is a way to quantify these errors. In the AA, there is what is called “producer’s accuracy,” which is the probability that an AA point has been mapped correctly (also known as error of omission) and there is “user’s accuracy,” which is the probability that the map actually represents what was found on the ground (also known as error of commission). Both the producer’s accuracy and user’s accuracy can be obtained from the same set of data by using different analyses. Overall Accuracy of the map is obtained by determining the fraction of AA observations within all map classes that were mapped correctly. The Kappa index is another measure of overall accuracy, which takes into account the probability that mapped polygons will be correct due to random chance.

There are a number of factors that contribute to error on a vegetation map and some of these are listed below. It is important for the field investigator to be aware of these situations and to take actions to minimize error when at all possible.

- Locational error (when it is not possible to acquire reliable UTM coordinates, the AA point data collected may look like it is in another polygon, or if the polygons are small, narrow and the GPS receiver is not working well, it may be difficult to know which polygon is supposed to be sampled).
- Field key is difficult to use, leading the investigator to assign the point to the wrong association.
- Field key does not include all the plant associations in the park, also leading to confusion in which association to assign a point.
- Field data error-either by misidentifying diagnostic species, or by not reading the key carefully, resulting in a bad field call when the map attribute is accurate.
- AA point falls within an ecotone, which is impossible to classify but still has to be mapped to something.
- Relationship between plant associations and map units (modeling) are flawed.

- The polygon is heterogeneous, including patches of varying vegetation that are too small to map individually. The AA point may happen to fall in a part of the polygon with a different community than observed by the photo-interpreter in the bigger picture.

The formal AA consists of several parts. First, a sampling design must be developed that follows methodology which is scientifically sound as well ideally practical and cost effective. The sampling design should cover all vegetation types mapped across the park. Sample sites must be visited in order to determine the vegetation type actually present at that particular ground location. The results of the sampling are used to create a contingency table (confusion matrix) that is used to calculate the per-class and overall accuracy of the map.

Sampling Design

We used a stratified random sampling approach to select AA sites in FOPU. We included all map classes representing natural/semi-natural vegetation types as defined in the NVCS in sampling selection (FGDC 1997). Cultural vegetation (e.g. warm season lawn) and non-vegetated (e.g. Commercial and Services, etc.) polygons were excluded from the AA. Points for the AA were restricted to within park boundaries. Based on these criteria, thirteen map classes were eligible for sampling.

We determined the sample size needed for each map class by following guidelines outlined in Thematic AA Procedures (Lea and Curtis 2010) (Table 5). Based on these criteria, we calculated that a total of 133 AA points should be sampled (Figure 9).

Table 5. Standard sample size allocations for accuracy assessment, based on map class area (Lea and Curtis, 2010).

Map Class Total Area (ha)	Number of Observations per Map Class
> 50	30
8.33 to 50	0.6 per hectare
< 8.33	5

In order to account for positional error due to site location (GPS error), vegetation polygons were buffered internally by 10 m prior to generation of the AA points, wherever possible. The AA points were generated randomly in ArcGIS using the Create Random Points tool in ArcToolbox (Version 10, © 2010 Environmental Systems Research Institute, Redlands, California).




Figure 9. Locations of accuracy assessment points sampled in Fort Pulaski National Monument.

Field Data Collection

AA point data were collected at FOPU during October of 2012. We employed Field Method 3 outlined in the Thematic AA Procedures (Lea and Curtis 2010), which utilizes a qualified and trained non-expert for field observation and class assignment and allows for follow-up reviews of mismatches in the office by an expert. Two SCA interns hired and trained by the SECN conducted the AA. The interns loaded the AA points into a GPS receiver in order to navigate to the points in the field. In addition to the GPS receivers, the interns used copies of the orthophotographs overlaid with the AA points (not including any of the lines from the mapped vegetation polygons) as an additional navigation aid. Upon reaching an AA point, the interns assessed the site within an area of equal size to the minimum radius (0.25 to 0.5 ha, depending on the draft vegetation class) of the observation area following methodology outlined by Lea and Curtis (2010). At each AA point, the interns collected environmental data such as hydrology, topography, etc., where meaningful. Vegetation data collected included leaf phenology, leaf type of the dominant stratum, physiognomic class, and vegetation composition in each stratum (see Appendix F for example of AA form). A space was provided for general comments about the classification made for each plot, noting any problems that were encountered (if any). Photographs in each cardinal direction, as well as of the canopy, were taken to provide a visual documentation of each AA site. After the AA was completed, all data sheets were reviewed in office by the SECN botanist. The SECN botanist reviewed the interns' field calls, and when necessary, superseded calls that she did not agree with based on the supplemental data collected at each field site. The association provided by the SECN botanist for each AA site was considered the "final call" that was used for tabulating the results in the data analysis phase.

Data Analysis

The AA point data were manually entered into the NPS PLOTS Database Version 3.2 (The Nature Conservancy 2010). PLOTS Version 3.2 is a relational database designed to store vegetation inventory field data for taxonomy and AA, and the data were subjected to a thorough quality assessment/quality check prior to analysis. The analysis of the map accuracy included the following steps:

- Initial comparative analysis of the field and map data
- Review of all disagreements and correcting for false errors as necessary
- Final comparative analysis of the field and map data
- Individual map class analysis
- Final output of results into a contingency matrix
- Final output of the analyses and results into a spatial database for use in GIS

Initial Comparative Analysis

We used the Intersect Tool in Arc Toolbox to combine the AA point layer and the map vegetation polygon layer. The resulting layer from the intersection allowed us to compare each AA field site call (vegetation type) to the corresponding map-polygon call (map class representing vegetation type). We used Microsoft Excel 2007 (Microsoft Corporation) to compare and tabulate the field site call to the map-polygon call. We discovered that there were four AA points that were not visited either because they were inaccessible (marsh sites) or because they were overlooked. These omissions left us with 129 AA points for analysis.

Review of Disagreements

All mismatches (disagreements) were subsequently reviewed for false errors. A false error is defined as a mismatch between the AA field site call and the map-polygon call if caused by an accuracy error in the GPS field coordinates, a missing or misapplied field call, or a field site assessment of an area smaller than the MMU. This review process involves looking at the AA sites and their corresponding polygons in ArcGIS in order to locate them and view them on the aerial photographs. We also reviewed the field data sheets to gain a fuller context of the ground conditions. From this process, we determined whether an initial disagreement was either a true error or indeed a match. We determined there were five points that were initially considered mismatches, but upon review were determined to be false errors because they occurred in areas below the MMU. In each of these cases, the points were updated as matches because their matching vegetation associations occurred outside the inclusions that they occurred upon.

Final Output

We generated two outputs for the AA results: a sample contingency table and a population contingency table. The sample contingency table provides an initial summary of the AA data by displaying counts of observations, with sample data values (vegetation map classes) as rows and reference data values (vegetation types as identified on the ground) as columns. The values in the shaded cells along the diagonal represent counts for correctly classified observations, where the reference data (column) matches the mapped vegetation type (row). User's accuracy was calculated by dividing the number of samples that agreed with their corresponding map class by the total number of samples in that class. Producer's accuracy was calculated by dividing the number of samples that agreed with their corresponding map class by the total number of samples whose field call belonged to that category.

Previous NPS VIP guidance (Environmental Systems Institute et al. 1994) suggested calculating user's, producer's, and overall accuracies from the sample contingency table of observation counts. However, this practice did not account for the unequal probability sampling imposed by the stratification by map classes. While appropriate for user's accuracies, it will give inaccurate results for producer's accuracies and overall accuracy. Therefore, measures of accuracy for the AA are defined by the population contingency table. Similar to the sample contingency table, rows in the population contingency table are defined by the sample data values, and the columns are defined by the reference data values. Unlike the sample contingency table, the values in each cell are the proportion of the target area in the corresponding true and mapped vegetation classes, rather than the raw counts of observations. The population and sample contingency tables are reported together as a summary of between-class error relationships and class accuracy statistics (from Lea and Curtis 2010) (Appendix G).

We examined the overall accuracy and the accuracy for each map class on an individual basis to determine if accuracy requirements of the NPS VIP were met. Based on these results, we determined whether or not it would be necessary to merge certain map classes to gain higher accuracy. The overall accuracy generated in the population contingency table met the NPS VIP accuracy standard of 80% (Overall accuracy = $90.3\% \pm 82.7, 97.9\%$); therefore it was not necessary to merge any vegetation classes.

AA Spatial Database

For use in GIS, we produced a feature class layer of the AA site locations, along with supporting attribute tables, and incorporated them into the FOPU vegetation mapping project geodatabase.

Results

Vegetation Classification

The final classifications for FOPU resulted in 12 vegetation associations in the NVCS (see Appendix B for full association descriptions). We had 12, rather than the original thirteen vegetation associations, because after reviewing the AA we eliminated two associations and added one. We removed the Wax myrtle-Carolina Laurel-Toothache-tree Shrubland and the Atlantic Coast Interdune Swale and added the Giant Atlantic Cordgrass Marsh. In the AA, none of the nine sites sampled in what the Remote Sensing Specialist mapped as Wax myrtle-Carolina Laurel-Toothache-tree Shrubland were classified to that association by the SCA interns. The interns classified this area as either Cedar-Live Oak-Cabbage Palmetto Marsh Hammock or Cabbage Palmetto Woodland. Upon review, it was agreed that this area of vegetation was best described as either the Cedar-Live Oak-Cabbage Palmetto Marsh Hammock or the Cabbage Palmetto Woodland (depending upon the canopy cover). Similarly, none of the five sites mapped as Atlantic Coast Interdune Swale by the Remote Sensing Specialist were classified as that association by the interns. Three of the sites were classified as Cabbage Palmetto Woodland, one was classified as Cedar-Live Oak-Cabbage Palmetto Marsh Hammock, and one was classified as Tallowtree Upland Forest. We updated the vegetation map in the areas previously mapped as Atlantic Coast Interdune Swale to reflect what the interns classified these areas as during the AA. We added the Giant Atlantic Cordgrass Marsh because during the AA the interns classified two sites as this association, but the Remote Sensing Specialist did not map any area of the park to this association. Upon further review of the vegetation map, we discovered that there were additional areas of the park that should have been mapped as Giant Atlantic Cordgrass Marsh. In all, we ended up mapping about 75 ha as the Giant Atlantic Cordgrass Marsh, mainly on McQueens Island.

Digital Imagery and Interpretation

For FOPU, 12 map units were developed and directly matched to the corresponding final vegetation associations in the NVCS (Table 6). One map class representing cultural vegetation in the NVCS (e.g., Warm Season Lawn) was identified. Eleven map classes representing non-vegetated units (e.g., open water bodies, buildings, roads, etc.) were identified (Table 7). Ancillary data sources such as a digital raster graphic (DRG) and historical aerial photographs provided additional insight into ground conditions. Historical and current land management practices in the park provided details pertinent to classification issue

Table 6. Final vegetation associations of the NVCS used for vegetation mapping (with their assigned Map Codes) in Fort Pulaski National Monument.

	Association Common Name	NVCS Code	Map Code
Forest			
	Cedar - Live Oak - Cabbage Palmetto Marsh Hammock	CEGL007813	7813
	Slash Pine Plantation	CEGL007170	7170
	Tallowtree Upland Forest	CEGL007037	7037
Woodland			
	South Atlantic Coastal Shell Midden Woodland	CEGL003525	3525
	Cabbage Palmetto Woodland	CEGL003526	3526
Shrubland			
	Seaside Oxeye Tidal Shrub Flat	CEGL003924	3924
Dwarf-Shrubland			
	Salt Flat (Swampfire Type)	CEGL002278	2278
	Saltwort-Swampfire Dwarf-shrubland	CEGL003956	3956
Herbaceous			
	Needlerush High Marsh	CEGL004186	4186
	Southern Atlantic Coast Salt Marsh	CEGL004191	4191
	Atlantic Giant Cordgrass Marsh	CEGL004195	4195
	Mid-Atlantic High Salt Marsh	CEGL004197	4197

Table 7. Map classification codes for non-vegetated areas in Fort Pulaski National Monument, following Land Use and Land Cover Level II (Anderson et al. 1976).

Map Class Code	Anderson Land Use and Land Cover Level II (1976)
12	Commercial and Services
14	Transportation, Communications, and Utilities
18	Fort
50	Water
54	Bays and Estuaries
72	Beaches
73	Sandy Areas other than Beaches
74	Bare Exposed Rock
76	Transitional Areas
78	Mud Flat
79	Tidal Wrack

Vegetation Map

Table 8 provides a summary report of the spatial data layer (map) of FOPU and summarizes frequency, mean area (in hectares), and the sum of areas (in hectares) for each map class.

Collectively, the FOPU spatial database layer is composed of 251 polygons covering 2,543.3 ha, with an average polygon size of 11.2 ha. Map classes representing natural types in the NVCS apply to 177 polygons (70.5% of all polygons) covering 1,843.6 ha (72.5% of entire area) with an average polygon size of 4.4 ha. Of these natural types, the Southern Atlantic Coast Salt Marsh covers the most area (1,488.3 ha). Needlerush High Marsh also covers a considerable portion of the park (91.4 ha).

Within the category of cultural types in the NVCS, Warm Season Lawn consists of 18 polygons (7.2% of all polygons) and covers 55.5 ha (2.2% of the entire area). This particular type accounts for the maintained lawns along roadsides and around the fort. Within cultural types, we created an additional provisional category called “disturbed vegetation” in order to characterize a section of vegetation located to the southwest of the fort. This area of vegetation (2.1 ha in size divided into two polygons) does not readily fit into any of the natural types in the NVCS and appears to have undergone some sort of disturbance.

Non-vegetated areas (including developed areas) consist of 54 polygons (21.5% of all polygons) and cover 651.2 ha (25.6% of the entire area). Developed areas in the park include the NPS visitor center and maintenance buildings, the fort, and roads within the project boundary.

Wetlands consist of 146 polygons (58.2% of all polygons) and cover 2,310.6 ha (90.9% of entire area). Vegetation associations in the NVCS that are considered vegetated wetlands according to the U.S. Fish and Wildlife Service include the following: Seaside Oxeye Tidal Shrub Flat, Salt Flat (Swampfire Type), Saltwort - Swampfire Dwarf-Shrubland, Needlerush High Marsh, Southern Atlantic Coast Salt Marsh, Atlantic Giant Cordgrass Marsh, and the Mid-Atlantic High Salt Marsh.

Considerably more analyses and deductions can be attained from even these simple summary reports. For example, a query could be done to determine the average distance a certain vegetation association occurs from a road or another developed area. Additionally, more complex reports could be derived by introducing additional spatial data layers (e.g., amphibian distributions, invasive species, etc.) Such analyses provide greater insight into vegetation and its ecology.

Table 8. Summary statistics for the map classes occurring in Fort Pulaski National Monument.

Map Class	Park Boundary and Buffer			Within Park Boundary Only		
	Frequency (# Polygons)	Mean (ha)	Sum (ha)	Frequency (# Polygons)	Mean (ha)	Sum (ha)
Natural Vegetation						
Cabbage Palmetto Woodland	12	1.10	13.7	12	1.1	13.7
Cedar - Live Oak - Cabbage Palmetto Hammock	21	2.5	52.9	20	2.6	51.9
Mid-Atlantic High Salt Marsh	2	1.0	2.0	2	1.0	2.0
Needlerush High Marsh	20	4.6	91.4	20	4.6	91.4
Salt Flat (Swampfire Type)	16	3.7	58.7	16	3.7	58.7
Saltwort-Swampfire Dwarf-shrubland	4	5.1	20.2	4	5.1	20.2
Seaside Oxeye Tidal Shrub Flat	16	1.4	22.1	16	1.40	21.7
Slash Pine Plantation	2	0.97	1.9	2	0.97	1.9
South Atlantic Coastal Shell Woodland Midden	5	0.79	3.95	5	0.79	4.0
Southern Atlantic Coast Salt Marsh	57	26.0	1488.3	54	27.4	1487.2
Tallowtree Upland Forest	8	0.53	4.3	8	0.53	4.3
Giant Atlantic Cordgrass Marsh	14	5.36	75.1	14	5.36	75.1
Natural Vegetation subtotal:	177	4.4	1834.6	173	4.5	1832.1
Cultural Vegetation						
Warm Season Lawn	18	3.1	55.5	18	3.8	55.5
Disturbed Vegetation	2	1.1	2.1	2	1.1	2.1
Cultural vegetation subtotal:	20	2.1	57.6	20	2.5	57.6

Table 8. Continued.

Map Class	Park Boundary and Buffer			Within Park Boundary Only		
	Frequency (# Polygons)	Mean (ha)	Sum (ha)	Frequency (# Polygons)	Mean (ha)	Sum (ha)
Non-Vegetated Areas						
Commercial and Services	8	0.37	2.92	8	0.37	2.92
Transportation, Communications, and Utilities	4	6.53	26.1	4	6.53	26.1
Fort	2	1.05	2.05	2	1.05	2.05
Water	1	1.01	1.01	1	1.01	1.01
Bays and Estuaries	15	40.6	609.1	14	43.4	607.5
Beaches	3	0.77	2.31	3	0.77	2.31
Sandy Areas other than Beaches	6	0.29	1.76	6	0.29	1.76
Bare Exposed Rock	6	0.28	1.69	6	0.28	1.69
Transitional Areas	1	0.20	0.20	1	0.20	0.20
Mud Flat	4	0.91	3.65	4	0.91	3.65
Tidal Wrack	4	0.09	0.37	4	0.09	0.37
Non-Vegetated Areas subtotal:	54	4.7	651.2	53	8.9	649.6
Grand Total:	251	11.2	2543.3	246	6.4	2539.2

Map Layer Presentation

Figure 10 presents the map layer produced for the FOPU vegetation mapping project. The map is shown at its finest level (e.g., vegetation classes are shown at the association level). The boundary for FOPU and the mapping boundary for the project are both included. Figure 11 presents the wetlands in FOPU.




Figure 10. Vegetation (NVCS 1997) and Land Use (Anderson et al. 1976) map classes for Fort Pulaski National Monument.




Figure 11. Distribution of wetlands (Cowardin et al. 1979) in Fort Pulaski National Monument.

Accuracy Assessment

Overall accuracy is 90.3% (90% confidence interval [CI]: 82.7- 97.9%) for primary map classes representing natural/semi-natural vegetation floristic types in the NVCS. A Kappa adjustment for chance agreements resulted in a final overall accuracy of 59.0% (90% CI: 51.3-66.6%). The sample and population contingency matrices for AA results are provided in Appendix G: Accuracy Assessment Contingency Tables. The population contingency matrix shows the accuracy of each map class (along with 90% CIs), with the user's accuracy reflecting errors of inclusion (commission errors) and the producer's accuracy reflecting errors of exclusion (omission errors) present in the mapping. The width of each CI is affected by the sample size used to derive the point estimate.

Our overall AA met the accuracy requirement of 80% established by the NPS VIP. Some individual classes, however, did not meet the 80% accuracy requirement. The following sections discuss the individual map classes that did not meet the required accuracy.

User's Accuracy (Errors of Commission)

Atlantic Coast Interdune Swale (CEGL003839)

As discussed previously, the decision was made to remove this class since none of the AA points sampled were classified as this association. The user's accuracy was 0% (90% CI: 0-10%).

Cabbage Palmetto Woodland (CEGL003526)

The user's accuracy for the Cabbage Palmetto Woodland was 36% (90% CI: 8-65%). Four of the eleven sites were mapped correctly. Two of the 11 sites were mapped as Atlantic Coast Interdune Swale, but after reviewing the AA data, we felt these were two areas of Atlantic Coast Interdune Swale were not large enough to be mappable. The remaining five sites were mapped as Cedar-Live oak- Cabbage Palmetto Marsh Hammock. The Cabbage Palmetto Woodland and Cedar-Live oak- Cabbage Palmetto Marsh Hammock are fairly easy to confuse when mapping because they share similar species in the canopy (e.g. *Sabal palmetto*, *Juniperus virginiana* var. *silicicola*, *Celtis laevigata*, and *Ilex vomitoria*) and are mainly distinguished from one another by % canopy cover, which can be difficult at times to estimate (woodland has moderate tree canopy coverage of about 60%; whereas the forest has dense coverage of about 70-90%).

Mid-Atlantic High Salt Marsh (CEGL004197)

The user's accuracy for the Mid-Atlantic High Salt Marsh was 60% (90% CI: 14-100%). The sample size for this association was small (five), and two of the sites were mapped incorrectly as Salt Flat (Swampfire Type). The confusion came about because these two associations occur in close proximity to one another where they were sampled, and they can have a similar appearance to one another on the imagery, especially in areas that transition between the two types.

Needlerush High Marsh (CEGL004186)

The user's accuracy for the Needlerush High Marsh was 0% (90% CI: 0-10%) Of the five sites sampled, none were mapped correctly. These five sites were determined on the ground to be Southern Atlantic Coast Salt Marsh. Upon review, we believe the confusion likely came about because there are certain areas of Southern Atlantic Coast Salt Marsh that contain tidal wrack which can cause a darker photo signature. The darker color likely created the confusion and caused the Remote Sensing Specialist to map these areas as Needlerush High Marsh.

Furthermore, it was realized after the AA that there were other areas not mapped as Needlerush High Marsh that should have been. After the AA was complete, 91.4 ha were mapped as Needlerush High Marsh.

Salt Flat (Swampfire Type) (CEGL002278)

The user's accuracy for the Salt Flat (Swampfire Type) was 33% (90% CI: 2-65%). Of the nine sites sampled, three were correct, five were determined on the ground to be Saltwort-Swampfire Dwarf-Shrubland, and one was determined to be Southern Atlantic Coast Salt Marsh. The Salt Flat (Swampfire Type) and the Saltwort-Swampfire Dwarf-Shrubland are easy to confuse on the imagery because they both contain Swampfire and Saltwort and both occur on hypersaline tidal flats, typically side-by-side. We considered combining them to a single map class, but that did not raise the overall accuracy for the project, so we left them as separate entities.

Seaside Oxeye Tidal Shrub Flat (CEGL003924)

The user's accuracy for the Seaside Oxeye Tidal Shrub Flat was 68% (90% CI: 50-87%). Of the 22 sites sampled, 15 were correctly identified, five were determined to be Southern Atlantic Coast Salt Marsh, and two were determined to be Giant Atlantic Cordgrass Marsh. The Seaside Oxeye Tidal Shrub Flat and Southern Atlantic Coast Salt Marsh can sometimes be confused for one another on the imagery (both shades of darker pink, with smooth to moderately smooth texture). The photo signature for Giant Atlantic Cordgrass Marsh was confused with Seaside Oxeye Tidal Shrub Flat in certain locations in the park. After the AA, it was possible to distinguish the Atlantic Giant Cordgrass Marsh photo signature from the Seaside Oxeye Tidal Shrub Flat photo signature.

Wax myrtle – Carolina Laurel Cherry – Toothache-Tree Shrubland (CEGL004787)

The user's accuracy for the Wax myrtle – Carolina Laurel Cherry – Toothache-Tree Shrubland was 0% (90% CI: 0-10%). As discussed previously, the decision was made to remove this association since none of the nine sites sampled were determined on the ground to be this association.

Giant Atlantic Cordgrass Marsh (CEGL004195)

The user's accuracy for the Giant Atlantic Cordgrass Marsh was 0% (90% CI: 0-0%) because this association was not mapped at all initially. After reviewing the AA, a photo signature for this class was deciphered, and about 75 ha were mapped on McQueens Island.

Producer's Accuracy (Errors of Omission)

Atlantic Coast Interdune Swale (CEGL003839)

As discussed previously, the decision was made to remove this class. The producer's accuracy was 0% (90% CI: 0-25%) because two AA sites were identified as Atlantic Coast Interdune Swale that were mapped as Cabbage Palmetto Woodland; however, we determined these two occurrences were not large enough to be mappable.

Cabbage Palmetto Woodland (CEGL003526)

The producer's accuracy for the Cabbage Palmetto Woodland was 44% (90% CI: 12-100%). Three sites were mapped as Atlantic Coast Interdune Swale and two were mapped as Cedar-Live oak-Cabbage Palmetto Woodland.

Cedar-Live Oak-Cabbage Palmetto Marsh Hammock (CEGL007813)

The producer's accuracy for the Cedar-Live Oak-Cabbage Palmetto Marsh Hammock was 73% (90% CI: 55-91%). One site was mapped as Atlantic Coast Interdune Swale and five sites were mapped as Cabbage Palmetto Woodland.

Needlerush High Marsh (CEGL004186)

The producer's accuracy for Needlerush High Marsh was 0% (90% CI: 0-0%). One Needlerush High Marsh site was mapped as Southern Atlantic Coast Salt Marsh.

Salt Flat (Swampfire Type) (CEGL002278)

The producer's accuracy for the Salt Flat (Swampfire Type) was 50% (90% CI: 8-100%). Two sites were mapped as Mid-Atlantic Coast High Salt Marsh, one site was mapped as Saltwort-Swampfire Dwarf-shrubland, and one site was mapped as Southern Atlantic Coast Salt Marsh.

Saltwort-Swampfire Dwarf-Shrubland (CEGL003956)

The producer's accuracy for the Saltwort-Swampfire Dwarf-Shrubland was 44% (90% CI: 12-77%). Five sites were mapped as Salt Flat (Swampfire Type). As previously discussed, the Salt Flat (Swampfire Type) and the Saltwort-Swampfire Dwarf-Shrubland are easy to confuse on the imagery because they both contain the same two main species (Swampfire and Saltwort) and both occur on hypersaline tidal flats, typically side-by-side.

Southern Atlantic Coast Salt Marsh (CEGL004191)

The producer's accuracy for the Southern Atlantic Coast Salt Marsh was 70% (90% CI: 57-84%). Five sites were mapped as Needlerush High Marsh, one site was mapped as Salt Flat (Swampfire Type), and five sites were mapped as Seaside Oxeye Tidal Shrub Flat.

Giant Atlantic Cordgrass Marsh (CEGL004195)

The producer's accuracy for the Giant Atlantic Cordgrass Marsh was 0% (90% CI: 0-25%). Two sites were mapped as Seaside Oxeye Tidal Shrub Flat. The photo signature for the Giant Atlantic Cordgrass Marsh is somewhat similar to the Seaside Oxeye Tidal Shrub Flat. The Giant Atlantic Cordgrass Marsh appears reddish/pink with a rough texture. The Seaside Oxeye Shrub Flat is a deeper pink that is somewhat smoother in texture, but has some roughness to it. During the field reconnaissance phase, a photo signature for the Giant Atlantic Cordgrass Marsh was not developed; it was only after the AA that the photo signature for this association was deciphered.

Literature Cited

- Anderson, J. R., E. E. Hardy, J. T. Roach, and R. E. Witmer. 1976. A land use and land cover classification system for use with remote sensor data. Geological Survey Professional Paper 964. United States Government Printing Office. Washington, DC. 28 pp.
- Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U. S. Department of the Interior. Fish and Wildlife Service. Washington, DC. Jamestown, ND: Northern Prairie Wildlife Research Center Online. <http://www.npwrc.usgs.gov/resource/1998/classwet/classwet.htm>. (Version 04DEC98).
- Davey, C. A., K. T. Redmond, and D. B. Simeral. 2007. Weather and Climate Inventory, National Park Service, Southeast Coast Network. Natural Resource Technical Report NPS/SECN/NRTR—2007/010. National Park Service, Fort Collins, CO.
- Environmental Systems Research Institute, Inc. 2010. Arc GIS 10.0. Environmental Systems Research Institute, Inc. Redlands, CA.
- Environmental Systems Research Institute, National Center for Geographic Information and Analysis, and the Nature Conservancy. 1994. Final draft for Field Methods for Vegetation Mapping procedures. USGS/NPS Vegetation Mapping Program. Prepared for the United States Department of the Interior, Biological Resources Division and National Park Service.
- Federal Geographic Data Committee-Vegetation Subcommittee. 1997. Vegetation classification standard [online]. Available from: <http://www.fgdc.gov/standards/projects/FGDC-standards-projects/vegetation/vegclass.pdf>.
- Federal Geographic Data Committee-Vegetation Subcommittee. 1998a. Spatial data transfer standard, FGDC-STC-002 (modified version ANSI NCITS 20:19998). Web address: <http://www.fgdc.gov/standards/projects/FGDC-standards-projects/SDTS>.
- Federal Geographic Data Committee-Vegetation Subcommittee. 1998b. Content standard for digital geospatial metadata, FGDC-STD-001-1998. Web address: <http://www.fgdc.gov/standards/projects/FGDC-standards-projects/metadata/base-metadata>.
- Federal Geographic Data Committee-Vegetation Subcommittee. 2008. National Vegetation Classification Standard (Version 2). FGDC-STD-005-2008. Available online: <http://www.fgdc.gov/standards/project/FGDC-standards-projects/vegetation>.
- Flahault, C., and C. Schroter. 1910. Rapport sur la nomenclature phytogeographique. Proceedings of the Third International Botanical Congress, Brussels 1:131-164.
- Grossman, D. H., D. Faber-Langendoen, A. S. Weakley, M. Anderson, P. Bourgeron, R. Crawford, K. Goodin, S. Landaal, K. Metzler, K. D. Patterson, M. Pyne, M. Reid, and L. Sneddon. 1998. International Classification of Ecological Communities: Terrestrial Vegetation of the United States. Volume I. The National Classification System: Development, Status, and Applications. The Nature Conservancy, Arlington, VA.

- Hershey, R. R., and W. A. Befort. 1995. Aerial photo guide to New England forest cover types. General Technical Report NE-195. U.S. Forest Service, Radnor, Pennsylvania, USA.
- Lea, C. and A. C. Curtis. 2010. Thematic accuracy assessment procedures: National Park Service Vegetation Inventory, version 2.0. Natural Resources Report NPS/2010/NRR-2010/204. National Park Service, Fort Collins, Colorado.
- NatureServe. 2003. International Ecological Classification Standard: International Vegetation Classification. Natural Heritage Central Databases, NatureServe, Arlington, VA.
- NatureServe. 2008. National Vegetation Classification. Associations and Alliances of Fort Pulaski National Monument. NatureServe Central Databases. Arlington, VA. Data current as of 24 April 2008.
- NatureServe. 2009. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, VA, USA. [Available: <http://www.natureserve.org/explorer>].
- NatureServe. 2010. NPS PLOTS Database Version 3.2. NatureServe, Arlington, Virginia, USA.
- Nordman, C. 2011. Personal communication. Southeast regional office of NatureServe, Raleigh/Durham, North Carolina.
- Philipson, Warren. 1997. Manual of Photographic Interpretation, Second Edition. American Society for Photogrammetry and Remote Sensing. Bethesda, MD.
- Reid, M. and P. Comer. 1998. Vegetation Alliance Descriptions of the Western U.S. Available at <http://www.gap.uidaho.edu/bulletins/7/VADWUS.htm>.
- Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Survey Geographic (SSURGO) Database for [Charleston County, South Carolina]. Available online at <http://soildatamart.nrcs.usda.gov>. Accessed [April/11/2011].
- The Nature Conservancy. 1996. Final Draft, Methodology for Assessing the Utility of Existing Data for Vegetation Mapping. Prepared for Prepared for USDI National Biological Survey and NPS. Arlington, VA.
- The Nature Conservancy (TNC) and Environmental Systems Research Institute (ESRI). 1994a. NBS/NPS Vegetation Mapping Program: Final Draft, Standardized National Vegetation Classification System. Prepared for USDI National Biological Survey and NPS. Arlington, VA.
- _____. 1994b. NBS/NPS Vegetation Mapping Program: Final Draft, Field Methods for Vegetation Mapping. Prepared for USDI – National Biological Survey and National Park Service. Arlington, Virginia.

The Nature Conservancy and Environmental Systems Research Institute. 1994. Final Draft Methodology for Assessing the Utility of Existing Data for Vegetation Mapping. NBS/NPS Vegetation Mapping Program. Prepared for the United States Department of Interior, Biological Sciences Resource Division and National Park Service.

USDA, NRCS. 2012. The PLANTS Database (<http://plants.usda.gov>). National Plant Data Team, Greensboro, NC 27401-4901 USA.

USDA-SCS. 1971. Soil Survey of Charleston County, South Carolina. 82 p. Available online at http://soils.usda.gov/survey/online_surveys/south_carolina/charleston/sc_charleston.pdf.

U.S. Geological Survey (USGS). 1999. Map accuracy standards. Fact sheet FS-171-99 (November 1999). Web address: <http://egsc.ugs.gov/isb/pubs/factsheets/fs17199.html>.

White, R. Personal communication. March 25, 2011. Southeast regional office of NatureServe, Raleigh/Durham, North Carolina.

Appendix A: Plot Sampling Form

NPS SE COAST PROJECT: Plot Sampling Form Page 1

County: Chatham Charleston Park: Fort Sumter Fort Pulaski Plot Code: FOPU01 State: GA/SC Location org: NPS


Provisional community name Pinus elliotii var. elliotii Planted Forest
 Classified community name Pinus elliotii var. elliotii Planted Forest
 Classifier CWN Date 11/28/2007 NVC EL CODE CEGL007170
 Survey date: 11/28/2007 time 9:50 Surveyors initials: Carl Nordman Brent Rollscheid

Directions to plot (including any notes on best access route, and mile marker):
EXIT VISITOR CENTER (VC) PARKING LOT HEADING SIB THEN TURN (R) HEADING W/B.
CONTINUE W/B APPROX 1/8 MILE UNTIL YOU REACH PICNIC AREA ON (L)/SOUTH
SIDE OF ROAD.

Time taken to walk to plot: 2 minutes

PLOT CONFIGURATION (Depict the plot layout on right with respect to the surrounding area and attempt to draw in points where GPS positions were collected, witness tree, bearing of centerline, locations of pictures taken, and communities near the plot.)
 Plot representativeness - Is the surrounding area all the same? ☐ YES ☐ NO - If not, depict below
 PLOT LOCATION (Below show general plot location)

Plot dimensions: 56 m X 20 m



Est. Extent of occurrence of community:
☐ <1HA ☐ 1-10 HA ☐ 10-100HA ☐ >100HA

PIC # 32-35 Description of pictures (N,E,S & W from GPS point at middle of plot): N, E, S, W

Select one: UTM / Lat/long (If lat/long, then values are _____ N _____ W)

GPS Techniques/Equipment: Garmin V Datum (pick one) WGS84 NAD83 / NAD27 UTM Zone: 17

GPS Unit ID _____ GPS file name 039 (standard = First letter(s) of park + # of plot (e.g. CH04, CL04, MO04))

Field UTM X 0509468 m E Y 3543810 m N Elevation 4 m / ft

Averaging performed (at least 120 observations)? 870 Coordinate accuracy 8-11 m / ft DOP _____ # sats 6

NPS SE COAST PROJECT: Plot Sampling Form

Page 2

County: Chatham Charleston Park: Fort Sumter Fort Pulaski Plot Code: F0PUG1 State: GA SC Location org: NPS

ENVIRONMENTAL / SITE INFORMATION

Measured Slope _____ ° _____ % ☒ Flat 0 ° 0 % ☒ Gentle 0-5 ° 1-9 % Moderate 6-14 ° 10-25 % Somewhat steep 15-25 ° 26-49 % Steep 27-45 ° 50-100 % Very steep 45-69 ° 101-275 % Abrupt 70-100 ° 276-300 % ovrrhg/shltd > 100 ° > 300 %	Measured Aspect _____ ° (N=0 °) ☒ Flat _____ SE 113-157 ° ☒ Variable _____ S 158-202 ° N 338-22 ° SW 203-247 ° NE 23-67 ° W 248-292 ° E 68-112 ° NW 293-337 °	Topographic Position Interfluvial (Ridge/summit/crest) _____ Toeslope ☒ High Slope (upper/convex slope) _____ Low level (terrace) Midslope _____ Channel bed Lowslope (lower/footslope) _____
Cowardin System ☒ Upland _____ Palustrine Estuarine _____ Lacustrine _____ Riverine		

Landform (check most applicable)

☐ Alluvial flat ☐ Alluvial terrace ☐ Backslope Terrace ☐ Bank ☐ Bar ☐ Barrier Island ☐ Beach ☐ Beach Ridge ☐ Bench	☐ Bottomlands ☐ Carolina Bay ☐ Channel ☐ Depression ☐ Draw ☐ Dune ☐ Estuary ☐ Floodplain ☐ Gap ☐ Gravel Bar	☐ Interdune Flat ☐ Lagoon ☐ Lowland ☐ Mound ☐ Mudflat ☐ Ravine ☒ Ridge ☐ Seep ☐ Slope ☐ Slough	☐ Spit ☐ Streambed ☐ Streamhead ☐ Swale ☐ Tidal Flat ☐ Toe slope ☐ Washover Fan
---	--	--	--

Rock types present:

Hydrologic Regime:

☒ Not a wetland (Upland: XERIC: DRY - MESIC: MESIC)
☐ Intermittently flooded _____ Permanently flooded _____ Semipermanently flooded _____ Temporarily Flooded (e.g. floodplains) _____ Tidally Flooded
☐ Seasonally Flooded (e.g. seasonal ponds) _____ Saturated (e.g. bogs, perennial seeps) _____ Unknown _____ Irregularly flooded _____ Irregularly exposed

QUALITATIVE ASSESSMENT:

A) HYDROLOGY:

B) LANDSCAPE AND LANDUSE HISTORY (including disturbance history and possible threats to sustainability):

Used as picnic area

Soil Texture:

☒ Sand _____ Peat _____ Silt loam _____ Muck
☐ Sandy loam _____ Loam _____ Clay loam _____ Clay

Drainage:

☐ Rapidly drained ☒ Moderately well drained _____ Somewhat poorly drained
☐ Well drained _____ Poorly drained _____ Very poorly drained

Ground cover [Exclusive of living plants; adds to 100%]

_____ % Bedrock 80 % Litter, duff _____ % Other _____
 _____ % Large rocks (cobbles, boulders > 10cm) _____ % Wood (> 1 cm)
 _____ % Small rocks (gravel, 0.2-10 cm) _____ % Water
20 % Loose sand (0.1-2 mm) _____ % Bare soil

Leaf type:

☐ Broad-leaved _____ Succulent
☒ Needle-leaved _____ Mixed
☐ Microphyllous
☐ Graminoid
☐ Broad-leaved herbaceous
☐ Pteridophyte
☐ Extremely xeromorphic

Leaf phenology (dominant stratum)

☒ Evergreen
☐ Cold-deciduous
☐ Drought-deciduous
☐ Mixed evergreen-cold-deciduous
☐ Mixed evergreen drought deciduous
☐ Herb - Annual
☐ Herb - Perennial

Physiognomic Class

☒ Forest (closed tree canopy)
☐ Woodland (open tree canopy)
☐ Shrubland
☐ Dwarf Shrubland
☐ Herbaceous (less than 25% woody layers)
☐ Nonvascular
☐ Sparse Vegetation

Natural and Anthropogenic Disturbance

☐ logging _____ fire _____ Hydrologic
☐ erosion _____ trails/roads _____ Agriculture
☐ grazing/browsing _____ wind/ice damage _____ Old Growth
☐ pine bark beetle ☒ Exotic plants _____ Fire Suppression
☐ dogwood anthr _____ ORV _____ Redbay mortality

Disturbance and animal use comments:

- picnic area with tables and grills
- One Chinese talow seedling

Community Survey

Page 2

NPS SE COAST PROJECT: Plot Sampling Form

Page 3

County: Chatham Charleston Park: Fort Sumter Fort Pulaski Plot Code: FOPU 01 State: GA SC Location org: NPS

QUANTITATIVE VEGETATION SAMPLE

"DEFAULT" TAXONOMIC REFERENCE: K99

STRATA	STRATA HEIGHT (1 only)	COVER CLASS	DOMINANT/DIAGNOSTIC SPECIES	Height scale (use tallest one for strata)	% cover for strata = top of cover class
Emergent - T1				< .5m 1	0-5% 1
Tree Canopy T2	9	9	<i>Pinus elliotii</i> var. <i>elliotii</i>	.5-1m 2	5-10% 2
Subcanopy T3	6	2	<i>Sabal palmetto</i>	1-2m 3	10-20% 3
Tall shrub S1				2-5m 4	20-30% 4
Short shrub S2				5-10m 5	30-40% 5
Herbaceous	1	6	~~<i>Stenotaphrum</i>~~ <i>Axonopus</i>	10-15m 6	40-50% 6
Non-vascular				15-20m 7	50-60% 7
Vine/liana				20-35m 8	60-70% 8
Other notable species (indicators of distinctive conditions, e.g. high pH soil, elevation, geographic region, other particularly abundant species):				35-50m 9	70-80% 9
				>50m 10	80-90% 10
					90-100% 11

T1: Emergent | T2: Tree Canopy | T3: Subcanopy | S1: Tall Shrub (>1m; to 5m) | S2: Short Shrub (< 1m) | H: Herbaceous | N: Nonvascular | V: Vines (lianas) | E: Epiphytes

SPECIES COMPOSITION AND COVER/ABUNDANCE CLASS BY STRATUM (enter cover values for each stratum AND for Total cover)

T1	T2	T3	S1	S2	H	N	V	E	Total Cover	Name (7 letter code or full name)	Collected? Spec #?	Discarded?	Cover cls
	9									<i>Pinus elliotii</i> var. <i>elliotii</i>			1 trace
		5								<i>Sabal palmetto</i>			2 0.1-1%
					3					<i>Hydrocotyle bonariensis</i>			3 1-2%
					3					<i>Sporobolus indicus</i>			4 2-5%
					6					<i>Axonopus fissifolius</i>	✓		5 5-10%
				2						<i>Juniperus virginiana</i> var. <i>silicola</i>			6 10-25%
				4						<i>Ilex vomitoria</i>			7 25-50%
					1					<i>Sida rhombifolia</i>	✓	✓	8 50-75%
					2					<i>Dichanthelium</i>	✓		9 75-95%
				2						<i>Rubus (trailing) trivialis</i>			10 >95%
				3						<i>Smilax bonariensis</i>			
					2					<i>Eragrostis spectabilis</i>			
					2					<i>Vicia</i>	✓		
					1					<i>Celtis</i>			
					1					<i>Wahlenbergia marginata</i>	✓	✓	
				2						<i>Pyrrhopappus carolinianus</i>	✓	✓	

Page 4

SPECIES COMPOSITION AND COVER/ABUNDANCE CLASS BY STRATUM

[illegible]

Use bottom of this sheet or additional pages for other species in plot

Community Survey
Page 4

Appendix B: Descriptions of Vegetation Types

This subset of the International Ecological Classification Standard covers associations and alliances attributed to Fort Pulaski National Monument. This classification has been developed in consultation with many individuals and agencies and incorporates information from a variety of publications and other classifications. Comments and suggestions regarding the contents of this subset should be directed to Mary J. Russo, Central Ecology Data Manager, Durham, NC <mary_russo@natureserve.org> and Carl W. Nordman, Regional Ecologist, Durham, NC <carl_nordman@natureserve.org>. Some classifications were altered by SECN staff and are detailed in the descriptions below. In those instances, comments should be directed to the authors of this report.

Copyright © 2008 NatureServe, 1101 Wilson Blvd, 15th floor
Arlington, VA 22209, U.S.A. All Rights Reserved.

These vegetation data, descriptions and classifications should be cited as:

NatureServe. 2008. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. U.S.A. Data current as of 24 April 2008.

Information Warranty Disclaimer: All data are provided as is without warranty as to the currentness, completeness, or accuracy of any specific data. The absence of data in any particular geographic area does not necessarily mean that species or ecological communities of concern are not present. NatureServe hereby disclaims all warranties and conditions with regard to these data, including but not limited to all implied warranties and conditions of merchantability, fitness for a particular purpose, and non-infringement. In no event shall NatureServe be liable for any special, indirect, incidental, consequential damages, or for damages of any kind arising out of or in connection with the use of these data. Because the data in the NatureServe Central Databases are continually being updated, it is advisable to refresh data at least once a year after receipt.

Table of Contents

	Page
Forest (I.).....	49
Woodland (II.).....	63
Shrubland (III.)	70
Dwarf-Shrubland (IV.).....	87
Herbaceous Vegetation (V.).....	97
Bibliography	118
I. Forest	134
II. Woodland	138
III. Shrubland	141
IV. Dwarf-Shrubland	143
V. Herbaceous	146

Forest (I.)

Lowland temperate seasonal evergreen forest (I. A.4.N.a)

Quercus virginiana – (*Sabal palmetto*) Forest Alliance (A.55)

Alliance Concept

Summary: This alliance includes communities of barrier islands, maritime hammocks, and some more inland coastal hammocks and other fire-protected situations, which are dominated and characterized by *Quercus virginiana* (live oak), and often containing *Sabal palmetto* (cabbage palmetto) and *Juniperus virginiana* var. *silicicola* (southern redcedar) within their ranges. Habitats for associations in this alliance range from xeric and subxeric to moist. Vegetation of this alliance most typically lies just landward of maritime shrub zones; it ranges from temperate to subtropical and often has a component of deciduous broad-leaved trees as well, particularly in the north. Some examples are affected by varying intensities of salt spray; these situations display more-or-less wind- and salt spray-sculpted vegetation. Other upland examples are not affected by salt spray and correspondingly differ in composition and stature. Composition varies along a latitudinal gradient. Examples occur on sand flats, lower slopes, and on stabilized dunes that are protected from saltwater flooding but which experience light to moderate salt spray. Some more protected examples have relatively closed and diverse canopies and well-developed shrub strata; vines are often conspicuous and abundant, but the herbaceous stratum is typically sparse and low in diversity. This community occurs over moist, sandy soils, on low areas of the mainland coast, and stands are protected from the most extreme maritime influences (e.g., salt spray) but are susceptible to high winds and flooding during hurricanes. Extending south from the vicinity of Cape Fear, North Carolina, the canopy is dominated by *Quercus virginiana* (live oak) and *Pinus taeda* (loblolly pine) with some *Sabal palmetto* (cabbage palmetto). Farther south, *Pinus elliottii* var. *elliottii* (Honduras pine) replaces *Pinus taeda* (loblolly pine), and *Sabal palmetto* (cabbage palmetto) becomes more prominent. In mid-Florida, tropical species begin to dominate the understory while temperate species retain canopy dominance. South of Martin County, Florida, tropical species such as *Bursera simaruba* (gumbo limbo), *Sideroxylon foetidissimum* (false mastic), and *Ficus aurea* (Florida strangler fig) begin to dominate the forest canopy. The more tropically influenced examples may contain shrubs such as *Eugenia axillaris* (white stopper), *Myrsine floridana* (Guianese colicwood), and *Coccoloba uvifera* (seagrape) on the west coast of Florida, and *Myrcianthes fragrans* (twinberry), *Ardisia escallonioides* (island marlberry), and *Psychotria nervosa* (Seminole balsamo) on the east coast. The alliance also includes tropical/temperate maritime hammocks of the east coast of Florida, in mid-peninsula, characteristically with oak canopy and tropical subcanopy; as well as temperate maritime hammocks of the northeast and Panhandle coasts of Florida. Vegetation of this alliance may be found on xeric to mesic sites, often occurring as linear strands behind frontal dunes. The seaward edge is generally found on the leeward side of dune complexes which provide shelter from excessive salt spray and overwash; this vegetation is also found on top of relict dune ridges and other areas with xeric to mesic hydrology. While relatively protected, the vegetation frequently exhibits effects of wind-pruning and salt spray. The alliance also includes some dry hammocks, found from Florida to North Carolina; in these examples, *Quercus virginiana* (live oak) is dominant, and sometimes *Juniperus virginiana* var. *silicicola* (southern redcedar) is present in the canopy. Frequently small *Cladina* - *Cladonia*-dominated openings are present. On small hammocks in salt marshes, *Juniperus virginiana* var. *silicicola* (southern redcedar), *Baccharis*

halimifolia (eastern baccharis), and *Morella cerifera* (Wax myrtle) are characteristic. Another type of xeric hammock contains *Serenoa repens* (saw palmetto) under a *Quercus virginiana* (live oak) canopy. On Amelia Island, Florida, *Magnolia grandiflora* (southern magnolia) is characteristically in the understory, increasing after cutting or with a greater shell content in the soil. Some Mississippi vegetation included here is found on coastal sand ridges along inlets of marsh channels.

Similar Alliances:

- *Quercus virginiana* - *Carya illinoensis* Woodland Alliance (A.665)
- *Quercus virginiana* - *Celtis laevigata* Forest Alliance (A.374)
- *Quercus virginiana* - *Juniperus virginiana* - (*Sabal palmetto*) Woodland Alliance (A.479)-- related vegetation which occurs under more severe conditions.
- *Quercus virginiana* - *Quercus nigra* Saturated Forest Alliance (A.379)
- *Quercus virginiana* - *Quercus stellata* Woodland Alliance (A.668)

Similar Alliance Comments: Also compare other West Gulf Coastal Plain *Quercus virginiana* alliances.

Related Concepts:

- Cabbage Palmetto: 74 (Eyre 1980) I
- IA9a. Mid-Atlantic Barrier Island Forest (Allard 1990) I
- IA9b. South Atlantic Inland Maritime Forest (Allard 1990) I
- IA9c. Mid-Atlantic Inland Maritime Forest (Allard 1990) I
- IA9d. South Atlantic Barrier Island Forest (Allard 1990) ?
- Interdune Forest (Ambrose 1990a) ?
- Live Oak: 89 (Eyre 1980) I

Alliance Description

Environment: These forests occur on barrier islands or coastal hammocks, in situations ranging from xeric and subxeric to moist. Some examples are affected by varying intensities of salt spray; these situations display more-or-less wind- and salt spray-sculpted vegetation. Other upland examples are not affected by salt spray and correspondingly differ in composition and stature. This community occurs over moist, sandy soils, on low areas of the mainland coast, and stands are protected from the most extreme maritime influences (i.e., salt spray) but are susceptible to high winds and flooding during hurricanes. The alliance also includes tropical/temperate maritime hammocks of the east coast of Florida, in mid-peninsula, characteristically with oak canopy and tropical subcanopy; as well as temperate maritime hammocks of the northeast and Panhandle coasts of Florida. Vegetation of this alliance may be found on xeric to mesic sites, often occurring as linear strands behind frontal dunes. The seaward edge is generally found on the leeward side of dune complexes which provide shelter from excessive salt spray and overwash; this vegetation is also found on top of relict dune ridges and other areas with xeric to mesic hydrology. The alliance also includes some dry hammocks, found from Florida to North Carolina. Some Mississippi vegetation included here is found on coastal sand ridges along inlets of marsh channels.

Vegetation: These forest communities are dominated by *Quercus virginiana* (live oak), and often contain *Sabal palmetto* (cabbage palmetto) and *Juniperus virginiana* var. *silicicola* (southern redcedar) within their ranges. The northernmost examples, in Virginia and extreme northeastern North Carolina, contain *Morella pensylvanica* (northern bayberry) as a shrub. Further south, stabilized dunes where salt spray is light to moderate display wind-sculpted vegetation dominated by *Quercus virginiana* (live oak) and *Quercus hemisphaerica* (Darlington oak) with lesser amounts of *Pinus taeda* (loblolly pine) and *Juniperus virginiana* var. *silicicola* (southern redcedar); typical understory components here include *Persea borbonia* (redbay), *Carpinus caroliniana* ssp. *caroliniana* (American hornbeam), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Cornus florida* (flowering dogwood), *Osmanthus americanus* var. *americanus* (devilwood), *Ilex opaca* var. *opaca* (American holly), and *Zanthoxylum clava-herculis* (Hercules' club). Shrub species include *Ilex vomitoria* (yaupon), *Morella cerifera* (Wax myrtle), *Sabal minor* (dwarf palmetto), and *Callicarpa americana* (American beautyberry). Dominant vines are *Toxicodendron radicans* (eastern poison ivy), *Vitis rotundifolia* (muscadine), *Smilax* (greenbrier) spp., *Parthenocissus quinquefolia* (Virginia creeper), *Bignonia capreolata* (crossvine), *Berchemia scandens* (Alabama supplejack), *Ampelopsis arborea* (peppervine), and *Gelsemium sempervirens* (evening trumpetflower). Typical herbs are *Mitchella repens* (partridgeberry), *Asplenium platyneuron* var. *platyneuron* (ebony spleenwort), *Chasmanthium laxum* (slender woodoats), *Piptochaetium avenaceum* (blackseed speargrass), *Galium pilosum* (hairy bedstraw), *Dichanthelium commutatum* (variable panicgrass), *Elephantopus nudatus* (smooth elephantsfoot), and *Passiflora lutea* (yellow passionflower). Canopies in more protected examples are dominated by *Quercus virginiana* (live oak), *Quercus hemisphaerica* (Darlington oak), and *Pinus taeda* (loblolly pine) and may also contain *Quercus falcata* (southern red oak), *Carya glabra* (pignut hickory), *Quercus nigra* (water oak), and *Pinus palustris* (longleaf pine). Understory species include *Persea palustris* (swamp bay), *Magnolia virginiana* (sweetbay), *Osmanthus americanus* var. *americanus* (devilwood), *Ilex opaca* var. *opaca* (American holly), *Juniperus virginiana* var. *silicicola* (southern redcedar), and *Sassafras albidum* (sassafras). Typical shrubs in these examples include *Ilex vomitoria* (yaupon), *Morella cerifera* (Wax myrtle), *Hamamelis virginiana* (American witchhazel), and *Sabal minor* (dwarf palmetto). Vines include *Vitis rotundifolia* (muscadine), *Smilax bona-nox* (saw greenbrier), *Gelsemium sempervirens* (evening trumpetflower), and *Campsis radicans* (trumpet creeper). Common herbaceous species are *Mitchella repens* (partridgeberry) and *Asplenium platyneuron* var. *platyneuron* (ebony spleenwort). Extending south from the vicinity of Cape Fear, North Carolina, the canopy is dominated by *Quercus virginiana* (live oak) and *Pinus taeda* (loblolly pine) with some *Sabal palmetto* (cabbage palmetto). Farther south, *Pinus elliottii* var. *elliottii* (Honduras pine) replaces *Pinus taeda* (loblolly pine), and *Sabal palmetto* (cabbage palmetto) becomes more prominent. In mid-Florida, tropical species begin to dominate the understory while temperate species retain canopy dominance. South of Martin County, Florida, tropical species, such as *Bursera simaruba* (gumbo limbo), *Sideroxylon foetidissimum* (false mastic), and *Ficus aurea* (Florida strangler fig), begin to dominate the forest canopy. These more tropically influenced examples may contain shrubs such as *Eugenia axillaris* (white stopper), *Myrsine floridana* (Guianese colicwood), and *Coccoloba uvifera* (seagrape) on the west coast of Florida, and *Myrcianthes fragrans* (twinberry), *Ardisia escallonioides* (island marlberry), and *Psychotria nervosa* (Seminole balsamo) on the east coast. In some examples, small *Cladina* - *Cladonia*-dominated openings are present. Other characteristic species include *Scleria triglomerata* (whip nutrush), *Paronychia baldwinii* (Baldwin's nailwort), *Cladina evansii* (Evans' reindeer lichen),

Stipulicida setacea (pineland scalypink), and *Hypericum hypericoides* (St. Andrew's cross). Some examples (e.g., on Amelia Island, Florida), *Magnolia grandiflora* (southern magnolia) is characteristically in the understory, increasing after cutting or with a greater shell content in the soil.

Dynamics: Habitats for associations in this alliance range from xeric and subxeric to moist. Vegetation of this alliance lies just landward of maritime shrub zones; it ranges from temperate to subtropical and often has a component of deciduous broad-leaved trees as well, particularly in the north. Some examples are affected by varying intensities of salt spray; these situations display more-or-less wind- and salt spray-sculpted vegetation. Other upland examples are not affected by salt spray and correspondingly differ in composition and stature. Composition varies along a latitudinal gradient; the northernmost examples, in Virginia and extreme northeastern North Carolina, contain *Morella pensylvanica* as a shrub. Farther south, stabilized dunes where salt spray is light to moderate display variants of this vegetation as well. Some Mississippi vegetation included here is found on coastal sand ridges along inlets of marsh channels. This alliance occurs on sandy soils which are generally poorly developed and low in natural fertility and organic matter content. It is typically found on old dunes which have been stable for long enough to permit forest growth. While fire cycles are generally long (26-100 years), *Sabal palmetto* is fire-resistant and produces flammable litter. Under dry conditions, fires will burn in from adjoining pinelands and kill fire-sensitive species such as *Quercus hemisphaerica* and some tropical species. In the northern portion of this community's range, siliceous sands dominate in preference to carbonate ones; the siliceous sands are generally nutrient-poor while carbonate ones are richer. Farther south, the carbonate fraction increases. Carbonate sands begin to dominate in the Deep South, especially along the coast of Florida. Barrier island soils are derived from material carried onto the island by water and wave action and not from weathering of rock. The major nutrient input to the terrestrial vegetation is from salt spray and precipitation.

Alliance Distribution

Range: This alliance is found in the Coastal Plain from Virginia to Alabama and Florida, and possibly Mississippi.

Nations: US

Subnations: AL, FL, GA, MS?, NC, SC, VA

TNC Ecoregions: 53:C, 54:P, 55:C, 56:C, 57:C, 58:?

USFS Ecoregions: 232Bc:CCC, 232Be:CCC, 232Bf:CCC, 232Bg:CCC, 232Bh:CCC, 232Bi:CCP, 232Bj:CCC, 232Bo:CC?, 232Bp:CCP, 232Ca:CCC, 232Cb:CCC, 232Cc:CCP, 232Ce:CCC, 232Cf:CC?, 232Ch:CCC, 232Ci:CCC, 232Db:CCC, 232Dc:CCC, 232Dd:CCC, 232De:CCC, 232Ga:CCP, 232Gb:CCC, 232Gc:CCP, 232Gd:CCP

Federal Lands: DOD (Cape Canaveral, Eglin, Tyndall?); NPS (Canaveral, Cape Hatteras, Cape Lookout, Cumberland Island, Fort Caroline, Fort Matanzas, Fort Pulaski, Timucuan); USFS (Apalachicola, Croatan, De Soto, Francis Marion, Ocala, Osceola); USFWS (Back Bay, Blackbeard Island?, Bon Secour, Cape Romain, Currituck, Lake Woodruff?, Merritt Island?, Swanquarter?, Wassaw?, Wolf Island?)

(CEGL007813) Cedar - Live Oak - Cabbage Palmetto Marsh Hammock
***Juniperus virginiana* var. *silicicola* - (*Quercus virginiana*, *Sabal palmetto*) Forest**
Coastal Redcedar - (Live Oak, Cabbage Palmetto) Forest

NVC Classification

Physiognomic Class	Forest (I)
Physiognomic Subclass	Evergreen forest (I.A.)
Physiognomic Group	Temperate broad-leaved seasonal evergreen forest (I.A.4.)
Physiognomic Subgroup	Natural/Semi-natural temperate broad-leaved seasonal evergreen forest (I.A.4.N.)
Formation	Lowland temperate seasonal evergreen forest (I.A.4.N.a.)
Alliance	<i>Quercus virginiana</i> - (<i>Sabal palmetto</i>) Forest Alliance (A.55)
Alliance (English name)	Live Oak - (Cabbage Palmetto) Forest Alliance
Association	<i>Juniperus virginiana</i> var. <i>silicicola</i> - (<i>Quercus virginiana</i> , <i>Sabal palmetto</i>) Forest
Association (English name)	Coastal Redcedar - (Live Oak, Cabbage Palmetto) Forest
Association (Common name)	Cedar - Live Oak - Cabbage Palmetto Marsh Hammock
Ecological System(s):	Atlantic Coastal Plain Embayed Region Tidal Salt and Brackish Marsh (CES203.260) Central Atlantic Coastal Plain Salt and Brackish Tidal Marsh (CES203.270)

Element Concept

Global Summary: This association occupies marsh hammocks (small islands surrounded by tidal marsh) along the South Atlantic Coast from eastern North Carolina south through South Carolina and Georgia to northeastern Florida. Related vegetation along central Florida spring runs is also covered here for now. The canopy can be relatively open or completely closed. *Juniperus virginiana* var. *silicicola* (southern redcedar) typically dominates the canopy, while other species such as *Quercus virginiana* (live oak), *Sabal palmetto* (cabbage palmetto), and *Celtis laevigata* var. *laevigata* (sugarberry) may be present. This association is related to other associations in this alliance in the same geographic range but differs in being shorter in stature, coastal redcedar-dominated, with fewer strata, and floristically depauperate (often with some marsh species present).

Environmental Description

Fort Pulaski National Monument Environment: This marsh hammock association was sampled at three locations in the park. The sites are flat to gently sloping, southwest-facing lowlands. Much of the uplands where this forest occurs are on old dredge spoil. One site is on top of historic dike system located adjacent to a maintained area along the entrance road, two are old dredge spoil/fill areas that have not been recently disturbed. Soils are moderately well- to well-drained sandy loam, and may have with oyster shells on the soil surface. The unvegetated surface is made up of leaf litter (48-80% cover), wood (2-5% cover) and bare soil (15-50%). Evidence of disturbance includes the presence of trails and exotic plants. One site is located on the highest elevation and most protected portion of the island (Govus 1998).

Global Environment: This association occupies marsh hammocks (small islands surrounded by tidal marsh) along the South Atlantic Coast. In North Carolina this type is found primarily in a matrix of brackish or full salt marshes (M. Schafale pers. comm.). The community may be dependent on occasional salty inundation to keep less salt-tolerant plants out. Related vegetation found near the edge of Silver Glen Spring Run, Ocala National Forest in Florida is also covered here for now.

Vegetation Description

Fort Pulaski National Monument Vegetation: The moderately to very dense (70-90% cover) tree canopy, 10-20 m tall, is dominated by *Juniperus virginiana* var. *silicicola* (southern redcedar) along with *Sabal palmetto* (cabbage palmetto) and *Ulmus americana* (American elm) (in one stand), or codominated by *Ilex vomitoria* (yaupon) (one stand). Additional tree species include *Celtis laevigata* (sugarberry), *Quercus virginiana* (live oak), *Sabal palmetto* (cabbage palmetto), and *Melia azedarach* (Chinaberrytree) (exotic). *Tillandsia usneoides* (Spanish moss) is an epiphyte found growing in the tree canopy. The moderate (50% cover) subcanopy (10-15 m) is codominated by *Juniperus virginiana* var. *silicicola* (southern redcedar) and *Prunus caroliniana* (Carolina laurelcherry). The very sparse to moderate (5-60% cover) tall shrub layer (2-5 m) includes *Ilex vomitoria* (yaupon), *Morella cerifera* (wax myrtle), *Prunus serotina* (black cherry), *Prunus caroliniana* (Carolina laurelcherry), and *Sabal palmetto* (cabbage palmetto). The moderate (40-60% cover) short shrub layer (0.5-2 m) is dominated by *Sabal palmetto* (cabbage palmetto) and includes other species from the upper layers as well as *Rubus trivialis* (southern dewberry), *Yucca aloifolia* (aloe yucca), *Rhus copallinum* (flameleaf sumac), *Zanthoxylum clava-herculis* (Hercules' club), and *Triadica sebifera* (tallowtree) (exotic). The very sparse (<5% cover) herbaceous layer is dominated by the native grass *Oplismenus hirtellus* ssp. *setarius* (bristle basketgrass). Vines include *Ampelopsis arborea* (peppervine), *Parthenocissus quinquefolia* (Virginia creeper), *Vitis rotundifolia* (muscadine), *Campsis radicans* (trumpet creeper), *Smilax bona-nox* (saw greenbrier), and *Lonicera japonica* (Japanese honeysuckle) (exotic). Other species observed in a previous study include *Sanicula canadensis* (Canadian blacksnakeroot), *Callicarpa americana* (American beautyberry), *Dichanthelium* (rosette grass) sp., *Smilax auriculata* (earleaf greenbrier), and *Persea borbonia* (redbay) (Govus 1998).

Global Vegetation: *Juniperus virginiana* var. *silicicola* (southern redcedar) typically dominates the canopy, while other species, such as *Quercus virginiana* (live oak), *Sabal palmetto* (cabbage palmetto), and *Celtis laevigata* var. *laevigata* (sugarberry), may be present. Exotic species, such as *Triadica sebifera* (tallowtree), *Tamarix* (tamarisk) spp., and *Melia azedarach* (Chinaberrytree), can be invasive in these communities. On Ocala National Forest a plot attributed to this association is additionally characterized by *Quercus hemisphaerica* (Darlington oak), *Callicarpa americana* (American beautyberry), *Persea borbonia* (redbay), *Morella caroliniensis* (southern bayberry), *Ageratina aromatica* (lesser snakeroot), *Dichondra carolinensis* (Carolina ponysfoot), *Oplismenus hirtellus* ssp. *setarius* (bristle basketgrass), *Chasmanthium sessiliflorum* (longleaf woodoats), *Thelypteris kunthii* (Kunth's maiden fern), *Parthenocissus quinquefolia* (Virginia creeper), *Bignonia capreolata* (crossvine), *Toxicodendron radicans* (eastern poison ivy), *Smilax smallii* (lanceleaf greenbrier), and *Vitis aestivalis* (summer grape) (NatureServe Ecology unpubl. data).

Most Abundant Species

Fort Pulaski National Monument

Stratum

Lifeform

Tree (canopy & subcanopy)	Needle-leaved tree
Tree canopy	Broad-leaved deciduous tree
Tree canopy	Broad-leaved evergreen tree
Tree subcanopy	Broad-leaved evergreen tree
Shrub/sapling (tall & short)	Broad-leaved evergreen shrub
Herb (field)	Graminoid

Species

Juniperus virginiana var. *silicicola* (southern redcedar)
Celtis laevigata (sugarberry)
Ilex vomitoria (yaupon)
Prunus caroliniana (Carolina laurelcherry)
Ilex vomitoria (yaupon)
Oplismenus hirtellus ssp. *setarius* (bristle basketgrass)

Characteristic Species

Fort Pulaski National Monument: *Ampelopsis arborea* (peppervine), *Celtis laevigata* (sugarberry), *Ilex vomitoria* (yaupon), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Morella cerifera* (Wax myrtle), *Quercus virginiana* (live oak), *Rubus trivialis* (southern dewberry), *Sabal palmetto* (cabbage palmetto), *Tillandsia usneoides* (Spanish moss)

Other Noteworthy Species

Fort Pulaski National Monument

Exotic and Invasive Species

Eremochloa ophiuroides (centipede grass)
Ligustrum japonicum (Japanese privet)
Ligustrum sinense (Chinese privet)
Lonicera japonica (Japanese honeysuckle)
Melia azedarach (Chinaberrytree)
Sporobolus indicus (smut grass)
Triadica sebifera (tallowtree)

I-Rank

-
-
High/Medium
High/Medium
Medium/Low
-
High

Note

exotic
exotic
exotic
exotic
exotic
exotic
exotic

Global

Exotic and Invasive Species

Melia azedarach (Chinaberrytree)
Triadica sebifera (tallowtree)

I-Rank

Medium/Low
High

Note

invasive exotic
invasive exotic

Conservation Status Rank

Global Rank & Reasons: G3? (14-Dec-1998). This community occupies marsh hammock islands along the South Atlantic Coast of North Carolina, South Carolina, Georgia, and northern Florida. This community is restricted in occurrence, and most occurrences are small, but threats are few.

Classification

Status: Standard

Classification Confidence: 2 – Moderate

Global Comments: There are examples of this type in Onslow County, North Carolina, and in the brackish marshes of the Embayed Region of North Carolina, such as along the estuarine Pungo River, and probably Swanquarter National Wildlife Refuge as well (M. Schafale pers. comm.). This type is presumably most common in the Sea Islands region. The hydrology of this association is variable, and some examples are near the conceptual border between upland, saturated, and tidal. These communities' simpler structure may be maintained in part by

infrequent catastrophic storm events (hurricane wind and overwash events). Exotic species, such as *Triadica sebifera* (= *Sapium sebiferum*), *Tamarix* spp., and *Melia azedarach* can be invasive in these communities.

Global Similar Associations:

- *Juniperus virginiana* var. *silicicola* - *Zanthoxylum clava-herculis* - *Quercus virginiana* - (*Sabal palmetto*) / *Sageretia minutiflora* - (*Sideroxylon tenax*) Woodland (CEGL003525)--more calcareous.
- *Quercus virginiana* - (*Juniperus virginiana*) - *Zanthoxylum clava-herculis* / *Sideroxylon lanuginosum* Woodland (CEGL003523)--more open, more *Sabal palmetto*.
- *Sabal palmetto* - (*Juniperus virginiana* var. *silicicola*) Woodland (CEGL003526)

Element Distribution

Fort Pulaski National Monument Range: This community is found on the southeast side of the park entrance road before reaching the Visitor Center parking lot and near the picnic area. This is the most widespread forest type on Cockspur Island and occurs largely on the older spoil deposits to the west of the dike system and along the southwestern edge of the island within the dike system. Portions of this association also occur in low-lying areas within the dike system on the southern part of Cockspur Island, on the east side of the paved trail northeast of the Visitor Center, and on central Cockspur Island within the dike system, to the northwest of the fort (Govus 1998).

Global Range: This association is found along the South Atlantic Coast from eastern North Carolina south through South Carolina and Georgia to northeastern Florida. Related vegetation along certain central Florida spring runs is also covered here for now.

Nations: US

States/Provinces: FL, GA, NC, SC

TNC Ecoregions: 55:C, 56:C, 57:C

USFS Ecoregions: 232Bf:CCC, 232Cb:CCC, 232Ce:CCC, 232Ci:CCC

Federal Lands: NPS (Fort Pulaski, Timucuan?); USFS (Ocala); USFWS (Swanquarter?)

Element Sources

Fort Pulaski National Monument Plots: FOPU.4, FOPU.13, FOPU.16.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: C.W. Nordman, mod. J. Gramling

References: EPA 2004, Govus 1998, Keys et al. 1995, NatureServe Ecology - Southeastern U.S. unpubl. data, Peet et al. unpubl. data 2002, Schafale pers. comm., Southeastern Ecology Working Group n.d.

Planted/cultivated temperate or subpolar needle-leaved evergreen forest (I.A.8.C.x.)

Pinus elliottii Forest Alliance (A.95)

Alliance Concept

Summary: This alliance includes planted stands (plantations) of *Pinus elliottii* var. *elliottii* (Honduras pine) (typically monospecific), within and outside of its natural range. These are cultivated forests and are not considered natural or semi-natural vegetation. They are maintained as plantations for the harvest of forest products (lumber, pulpwood). Stands of this alliance are primarily in uplands, but occasional stands in wetland environments are included here as well. This alliance is also supposed to include the rare case of "exotic" stands resulting from natural regeneration outside of the natural range of *Pinus elliottii* var. *elliottii* (Honduras pine), although these are probably better accommodated as examples of *Pinus elliottii* var. *elliottii* - (*Pinus palustris*) Managed Forest (CEGL007171), a member of the *Pinus palustris* - (*Pinus elliottii*) Forest Alliance (A.123).

Classification Comments: Slash pine has been introduced into Kentucky, Virginia and eastern Texas, and now reproduces naturally in these locations.

Similar Alliances:

- *Pinus palustris* - (*Pinus elliottii*) Forest Alliance (A.123)—may include examples of abandoned plantations or silviculturally manipulated stands with a semi-natural ground layer.

Related Concepts:

- Slash Pine: 84 (Eyre 1980) I

Alliance Description

Environment: Stands of this alliance are primarily in uplands, but occasional stands in wetland environments are included here as well.

Vegetation: This alliance includes planted stands (plantations) of *Pinus elliottii* var. *elliottii* (Honduras pine) (typically monospecific), within and outside of its natural range.

Alliance Distribution

Range: This alliance is found in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, Texas, and possibly Kentucky and Virginia. In Arkansas the species is found on industrial timber land. It is reportedly introduced into Kentucky.

Nations: US

Subnations: AL, AR, FL, GA, KY?, LA, MS, NC, SC, TX, VA?

TNC Ecoregions: 40:C, 41:C, 43:C, 52:P, 53:C, 55:C, 56:P, 57:P, 58:P

USFS Ecoregions: 231:C, 232Br:CCC, 232Ca:CCC, 232Cb:CCC, 232Ce:CCC, 232Ci:CCC

Federal Lands: DOD (Fort Benning, Fort Bragg, Fort Gordon, Fort Stewart); NPS (Cape Hatteras, Fort Pulaski, Timucuan?); USFS (Angelina, Apalachicola, Conecuh, Kisatchie, Ocala, Osceola, Sabine)

(CEGL007170) Slash Pine Plantation
***Pinus elliottii* var. *elliottii* Planted Forest**
Slash Pine Planted Forest

NVC Classification

Physiognomic Class	Forest (I)
Physiognomic Subclass	Evergreen forest (I.A.)
Physiognomic Group	Temperate or subpolar needle-leaved evergreen forest (I.A.8.)
Physiognomic Subgroup	Planted/Cultivated temperate or subpolar needle-leaved evergreen forest (I.A.8.C.)
Formation	Planted/cultivated temperate or subpolar needle-leaved evergreen forest (I.A.8.C.x.)
Alliance	<i>Pinus elliottii</i> Planted Forest Alliance (A.95)
Alliance (English name)	Slash Pine Planted Forest Alliance
Association	<i>Pinus elliottii</i> var. <i>elliottii</i> Planted Forest
Association (English name)	Slash Pine Planted Forest
Association (Common name)	Slash Pine Plantation

Element Concept

Global Summary: This association includes planted stands (plantations) of *Pinus elliottii* var. *elliottii* (Honduras pine) (typically monospecific), within and outside of its natural range. These are cultivated forests and are not considered natural or semi-natural vegetation. They are maintained as plantations for the harvest of forest products (lumber, pulpwood). Stands are primarily in uplands, but occasional stands in wetland environments are included here as well. This concept is also supposed to include the rare case of "exotic" stands resulting from natural regeneration out of the natural range of *Pinus elliottii* var. *elliottii* (Honduras pine), although these are probably better accommodated as examples of *Pinus elliottii* (slash pine) var. *elliottii* - (*Pinus palustris*) Managed Forest (CEGL007171), a member of the *Pinus palustris* - (*Pinus elliottii*) Forest Alliance (A.123).

Environmental Description

Fort Pulaski National Monument Environment: Slash pine plantation was sampled once in the picnic area of the park. The site is a gently sloping area with dry-mesic, moderately well-drained sand soil. Evidence of disturbance includes the presence of tables and grills and exotic plants. This forest, located on south-central Cockspur Island, was planted by the National Park Service early on in the history of their management of Fort Pulaski (Govus 1998).

Global Environment: Stands are primarily in uplands, but occasional stands in wetland environments are included here as well.

Vegetation Description

Fort Pulaski National Monument Vegetation: The dense (80% cover) tree canopy, 15-20 m tall, is made up entirely of *Pinus elliottii* var. *elliottii* (Honduras pine). *Sabal palmetto* (cabbage palmetto) comprises the sparse (10% cover) subcanopy (10-15 m); *Prunus caroliniana* (Carolina laurelcherry) may also occur in the subcanopy (Govus 1998). The sparse (10% cover) short shrub layer (0.5-2 m) includes *Ilex vomitoria* (yaupon), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Rubus trivialis* (southern dewberry), and *Smilax bona-nox* (saw greenbrier). The moderate (50% cover) herbaceous layer is dominated by *Axonopus fissifolius* (common carpetgrass) and includes small amounts of *Eragrostis spectabilis* (purple lovegrass), *Hydrocotyle bonariensis* (largeleaf pennywort), *Pyrrhopappus carolinianus* (Carolina desert-chicory), *Phyla nodiflora* (turkey tangle fogfruit), and *Sporobolus indicus* (smut grass). It is regularly mowed.

Global Vegetation: These are planted stands dominated by *Pinus elliottii* var. *elliottii* (Honduras pine) (typically monospecific), within and outside of its natural range.

Most Abundant Species

Fort Pulaski National Monument

<u>Stratum</u>	<u>Lifeform</u>	<u>Species</u>
Tree canopy	Needle-leaved tree	<i>Pinus elliottii</i> var. <i>elliottii</i> (Honduras pine)
Tree subcanopy	Palm tree	<i>Sabal palmetto</i> (cabbage palmetto)
Herb (field)	Graminoid	<i>Axonopus fissifolius</i> (common carpetgrass)

Global

<u>Stratum</u>	<u>Lifeform</u>	<u>Species</u>
Tree canopy	Needle-leaved tree	<i>Pinus elliottii</i> var. <i>elliottii</i> (Honduras pine)

Characteristic Species

Fort Pulaski National Monument: *Axonopus fissifolius* (common carpetgrass), *Ilex vomitoria* (yaupon), *Pinus elliottii* var. *elliottii* (Honduras pine), *Sabal palmetto* (cabbage palmetto)

Global: *Pinus elliottii* var. *elliottii* (Honduras pine)

Other Noteworthy Species

Fort Pulaski National Monument

<u>Exotic and Invasive Species</u>	<u>I-Rank</u>	<u>Note</u>
<i>Sporobolus indicus</i> (smut grass)	-	exotic
<i>Triadica sebifera</i> (tallowtree)	High	exotic
<i>Wahlenbergia marginata</i> (southern bellflower)	-	exotic

Conservation Status Rank

Global Rank & Reasons: GNA (cultural) (8-Aug-2000). This community represents vegetation which has been planted in its current location and/or is treated with intensive management or manipulation. It is not a conservation priority and does not receive a conservation rank.

Classification

Status: Standard

Classification Confidence: 2 - Moderate

Fort Pulaski National Monument Comments: Although of little botanical importance, this pine-dominated community has apparent significance for nesting birds on Cockspur Island (Southeastern Wildlife Services 1981).

Global Comments: These are cultivated forests and are not considered natural or semi-natural vegetation. They are maintained as plantations for the harvest of forest products (lumber, pulpwood). Slash pine has been introduced into North Carolina and eastern Texas, and now reproduces naturally in these locations.

Element Distribution

Fort Pulaski National Monument Range: This association is found near the picnic area.

Global Range: This planted vegetation may be found throughout the lower south, from North Carolina to Texas, in the Coastal Plain and some adjacent ecoregions.

Nations: US

States/Provinces: AL, AR, FL, GA, KY?, LA, MS, NC, SC, TX, VA?

TNC Ecoregions: 40:C, 41:C, 43:P, 52:P, 53:C, 55:P, 56:P, 57:C, 58:P

USFS Ecoregions: 231:C, 232Br:CCC, 232Ca:CCC, 232Cb:CCC, 232Ce:CCC, 232Ci:CCC
Federal Lands: DOD (Fort Benning, Fort Bragg, Fort Gordon, Fort Stewart); NPS (Cape Hatteras, Fort Pulaski, Timucuan?); USFS (Angelina, Apalachicola, Conecuh, Kisatchie, Ocala, Osceola, Sabine)

Element Sources

Fort Pulaski National Monument Plots: FOPU.1.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: A.S. Weakley

References: Govus 1998, Schotz pers. comm., Southeastern Ecology Working Group n.d., Southeastern Wildlife Services 1981

Lowland or submontane cold-deciduous forest (I.B.2.N.a.)

***Triadica sebifera* Forest Alliance (A.257)**

Alliance Concept

Summary: Exotic-dominated upland forests often developed on old fields, dredge spoil, or other disturbed sites. Understory and herb layers often poorly developed. Range so far restricted to Texas, Louisiana, and Alabama, but possible in other parts of the southeastern United States.

Alliance Description

Range: The range of this alliance is so far restricted to Texas, Louisiana, and Alabama, but it could possibly be found in other parts of the southeastern United States, including Florida, Georgia, Mississippi, and South Carolina.

Nations: US

Subnations: AL, FL?, GA, LA, MS?, SC?, TX

TNC Ecoregions: 31:C, 40:C, 41:C, 53:C, 55:P

USFS Ecoregions: 231E:C?, 231Fa:CCP, 231Fb:CCC, 232Ba:CC?, 232Bg:CC?, 232Bj:CC?, 232Bs:CCC, 232Ce:CCC, 232Dc:CCC, 232Dd:CCC, 232E:CP, 232F:CP, 232Ga:CPP, 234An:???, 255Da:CCC, 255Db:CCP, 255Dc:CCC

Federal Lands: NPS (Fort Pulaski); USFWS (Anahuac, Brazoria)

(CEGL007037) Tallowtree Upland Forest

***Triadica sebifera* Upland Forest**

Chinese Tallowtree Upland Forest

NVC Classification

Physiognomic Class	Forest (I)
Physiognomic Subclass	Deciduous forest (I.B.)
Physiognomic Group	Cold-deciduous forest (I.B.2.)
Physiognomic Subgroup	Natural/Semi-natural cold-deciduous forest (I.B.2.N.)
Formation	Lowland or submontane cold-deciduous forest (I.B.2.N.a.)
Alliance	<i>Triadica sebifera</i> Forest Alliance (A.257)
Alliance (English name)	Chinese Tallowtree Forest Alliance
Association	<i>Triadica sebifera</i> Upland Forest
Association (English name)	Chinese Tallowtree Upland Forest
Association (Common name)	Tallowtree Upland Forest

Element Concept

Global Summary: *Triadica sebifera* (tallowtree)-dominated upland forests, often developed on old fields, dredge spoil, or other disturbed sites. Understory and herb layers often poorly developed. This association is defined broadly until further information is available to refine its classification.

Environmental Description

Fort Pulaski National Monument Environment: This invasive community was sampled once along the entrance road into the park. The site is a flat lowland with somewhat poorly drained clay loam soil. Evidence of disturbance includes the presence of exotic plants. In addition, many dead plants were observed in the understory, possibly due to allelopathic influences.

Vegetation Description

Fort Pulaski National Monument Vegetation: The moderately dense (70% cover) tree canopy, 15-20 m tall, is dominated by the invasive exotic *Triadica sebifera* (tallowtree), as is the sparse (20% cover) subcanopy (10-15 m). Additional tree species include *Juniperus virginiana* var. *silicicola* (southern redcedar) and *Celtis laevigata* (sugarberry). The moderate (50% cover) tall shrub layer (2-5 m) is dominated by *Ilex vomitoria* (yaupon) and *Sabal palmetto* (cabbage palmetto) and includes *Juniperus virginiana* var. *silicicola* (southern redcedar), *Morella cerifera* (Wax myrtle), *Rubus trivialis* (southern dewberry), and *Melia azedarach* (Chinaberrytree) (exotic). The herbaceous layer is very sparse (<5% cover).

Most Abundant Species

Fort Pulaski National Monument

Stratum

Tree (canopy & subcanopy)
Tall shrub/sapling
Tall shrub/sapling

Lifeform

Broad-leaved deciduous tree
Broad-leaved evergreen shrub
Palm shrub

Species

Triadica sebifera (tallowtree)
Ilex vomitoria (yaupon)
Sabal palmetto (cabbage palmetto)

Characteristic Species

Fort Pulaski National Monument: *Juniperus virginiana* var. *silicicola* (southern redcedar), *Morella cerifera* (wax myrtle), *Sabal palmetto* (cabbage palmetto), *Triadica sebifera* (tallowtree)

Other Noteworthy Species

Fort Pulaski National Monument

Exotic and Invasive Species

Lonicera japonica (Japanese honeysuckle)
Melia azedarach (Chinaberrytree)
Triadica sebifera (tallowtree)

I-Rank

High/Medium
Medium/Low
High

Note

exotic
exotic
exotic

Conservation Status Rank

Global Rank & Reasons: GNA (invasive) (1-Dec-1997).

Classification

Status: Standard

Classification Confidence: 1 - Strong

Element Distribution

Fort Pulaski National Monument Range: This community is found on Cockspur Island, along the entrance road from Highway 80 to the Visitor Center.

Global Range: The range of this association includes Texas, Louisiana, Alabama, and Georgia, but it could possibly be found in other parts of the southeastern United States, including Florida, Mississippi and South Carolina.

Nations: US

States/Provinces: AL, FL?, GA, LA, MS?, SC?, TX

TNC Ecoregions: 31:C, 40:C, 41:C, 53:C, 55:P

USFS Ecoregions: 231E:C?, 231Fa:CCP, 231Fb:CCC, 232Bs:CCC, 232Ce:CCC, 232Dc:CCC, 232Dd:CCC, 232E:CP, 232F:CP, 255Da:CCC, 255Db:CCP, 255Dc:CCC

Federal Lands: NPS (Fort Pulaski); USFWS (Anahuac, Brazoria)

Element Sources

Fort Pulaski National Monument Plots: FOPU.14.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: A.S. Weakley and L.M. Smith

References: Schotz pers. comm., Southeastern Ecology Working Group n.d.

During the field verification step of photo-interpretation (see more below), the SECN Remote Sensing Specialist identified an additional vegetation community that had not been included in the original Vegetation Classification developed by NatureServe. The Tallowtree Seasonally Flooded Forest Alliance (A.335) was added to the Vegetation Classification based on a small (but mappable) wetland inclusion found within the Wax myrtle-Carolina Laurel Cherry-Toothache-tree Shrubland. This area is dominated by the exotic Tallowtree (*Triadica sebifera*) and also contains Carolina Willow (*Salix caroliniana*), Wax myrtle (*Morella cerifera*), Virginia saltmarsh mallow (*Kosteletzkya virginica*), smallspike false nettle (*Boehmeria cylindrica*), peppervine (*Ampelopsis arborea*), saltmarsh morning-glory (*Ipomoea sagittata*), and various wetland grasses and sedges.

Seasonally flooded cold-deciduous forest (I.B.2.N)

***Triadica sebifera* Seasonally Flooded Forest Alliance (A.335)**

Alliance Description

Seasonally flooded forests dominated by the exotic *Triadica sebifera* (= *Sapium sebiferum*) (Euphorbiaceae). This tree was introduced into the lower Gulf Coastal Plain as an ornamental and is spreading. It can overtake native vegetation types to become the strong dominant. Other wetland trees may be present, including *Acer rubrum* var. *drummondii*, *Salix nigra*, and *Salix caroliniana*. Various wetland grasses, sedges, and forbs are present in these forests as well.

Note: The Alliance Description is the only information currently available for this alliance.

Woodland (II.)

Temperate broad-leaved evergreen woodland (II. A.2.N.a)

Quercus virginiana – *Juniperus virginiana* – (*Sabal palmetto*) Woodland Alliance (A.479)

Alliance Concept

Summary: This alliance accommodates calciphilic communities occurring on coastal sands, Amerindian shell middens, and natural shell deposits of the Outer Coastal Plain. Composition varies along a latitudinal gradient; associations have been defined for two separate geographic zones, one of the Atlantic Coast from northern Florida north to Virginia, and another in the Gulf Coast from Florida west to Texas. This vegetation occurs under more severe and unstable conditions than related forests, which are found in the *Quercus virginiana* - (*Sabal palmetto*) Forest Alliance (A.55).

Similar Alliances:

- *Quercus virginiana* - (*Sabal palmetto*) Forest Alliance (A.55)

Similar Alliance Comments: This vegetation occurs under more severe conditions than related forests, which are found in the *Quercus virginiana* - (*Sabal palmetto*) Forest Alliance (A.55).

Related Concepts:

- Live Oak: 89 (Eyre 1980) I

Alliance Description

Environment: This alliance accommodates calciphilic communities occurring on coastal sands, on Amerindian shell middens, and natural shell deposits of the Outer Coastal Plain.

Dynamics: This vegetation occurs under more severe and unstable conditions than related forests.

Alliance Distribution

Range: Associations have been defined for two separate geographic zones, one of the Atlantic Coast from northern Florida north to southeastern Virginia, and another in the Gulf Coast from Florida west to Texas.

Nations: US

Subnations: AL, FL, GA, LA, MS, NC, SC, TX?, VA

TNC Ecoregions: 31:C, 53:C, 54:P, 55:C, 56:C, 57:C, 58:C

USFS Ecoregions: 231Fb:PPP, 232Ce:CCC, 232Ch:CCC, 232Ci:CCP, 232Dc:CCC, 232Eb:CCC, 232Ec:CCC, 232Gb:CCC, 255Dc:PPP, 411Ad:PPP, 411Ae:PPP, 411Af:PPP, 411Ag:PPP

Federal Lands: NPS (Canaveral, Fort Pulaski, Timucuan); USFS (Francis Marion); USFWS (Back Bay?)

(CEGL003525) South Atlantic Coastal Shell Midden Woodland

Juniperus virginiana var. *silicicola* - *Zanthoxylum clava-herculis* - *Quercus virginiana* - (*Sabal palmetto*)/*Sageretia minutiflora* - (*Sideroxylon tenax*) Woodland
 Coastal Redcedar - Toothache-tree - Live Oak - (Cabbage Palmetto)/Small-flower Mock Buckthorn - (Tough Bumelia) Woodland

NVC Classification

Physiognomic Class	Woodland (II)
Physiognomic Subclass	Evergreen woodland (II.A.)
Physiognomic Group	Temperate broad-leaved evergreen woodland (II.A.2.)
Physiognomic Subgroup	Natural/Semi-natural temperate broad-leaved evergreen woodland (II.A.2.N.)
Formation	Temperate broad-leaved evergreen woodland (II.A.2.N.a.)
Alliance	<i>Quercus virginiana</i> - <i>Juniperus virginiana</i> - (<i>Sabal palmetto</i>) Woodland Alliance (A.479)
Alliance (English name)	Live Oak - Eastern Redcedar - (Cabbage Palmetto) Woodland Alliance
Association	<i>Juniperus virginiana</i> var. <i>silicicola</i> - <i>Zanthoxylum clava-herculis</i> - <i>Quercus virginiana</i> - (<i>Sabal palmetto</i>) / <i>Sageretia minutiflora</i> - (<i>Sideroxylon tenax</i>) Woodland
Association (English name)	Coastal Redcedar - Toothache-tree - Live Oak - (Cabbage Palmetto) / Small-flower Mock Buckthorn - (Tough Bumelia) Woodland
Association (Common name)	South Atlantic Coastal Shell Midden Woodland
Ecological System(s):	Southern Atlantic Coastal Plain Maritime Forest (CES203.537)

Element Concept

Global Summary: This calciphilic community occurs on calcareous coastal sands, shell hashes, Amerindian shell middens, and natural shell deposits, from mid-peninsular Florida (Cape Canaveral) north to South Carolina (southern Atlantic Coast), in maritime-influenced landscapes. The canopy consists of *Juniperus virginiana* var. *silicicola* (southern redcedar), *Sabal palmetto* (cabbage palmetto), *Celtis laevigata* var. *laevigata* (sugarberry), *Zanthoxylum clava-herculis* (Hercules' club), *Quercus virginiana* (live oak), *Osmanthus americanus* (devilwood), and other species. Shrubs are prominent and may include *Ilex vomitoria* (yaupon), *Yucca aloifolia* (aloe yucca), *Sageretia minutiflora* (small-flower mock buckthorn), *Forestiera segregata* (Florida swampprivet), *Frangula caroliniana* (Carolina buckthorn), *Opuntia humifusa* var. *austrina* (devil's-tongue), and others. Composition is very variable from site to site, depending on environmental conditions and accidents of establishment and persistence.

Environmental Description

Fort Pulaski National Monument Environment: This coastal shell midden woodland was sampled twice in the park. The sites are gently sloping oyster shell middens of unknown age with dry-mesic, well-drained sand soil. The unvegetated surface is made up of leaf litter (25-50% cover), wood (5% cover), bare soil (0-5% cover), and shells (40-70% cover). Evidence of disturbance includes the presence of exotic plants. The four large elliptical mounds suggest, from aerial view, to be of aboriginal origin. Regardless of origin, these oyster shell heaps support vegetation typical of shell midden sites along the Sea Islands section of the south Atlantic Coast (Govus 1998).

Global Environment: This calciphilic community occurs on Amerindian shell middens and natural shell deposits, from mid-peninsular Florida (Cape Canaveral) north to South Carolina (southern Atlantic Coast), and occurs along the ecotone between tidal marshes and upland maritime forests (C. Nordman pers. obs.).

Vegetation Description

Fort Pulaski National Monument Vegetation: The moderate to dense (70-80% cover) tree canopy, 2-15 m tall, is dominated by *Juniperus virginiana* var. *silicicola* (southern redcedar). Additional tree species may include *Forestiera segregata* (Florida swampprivet) (state-imperiled), *Sabal palmetto* (cabbage palmetto), *Ilex vomitoria* (yaupon), *Celtis laevigata* (sugarberry), and *Morus alba* (white mulberry) (exotic). The sparse (20% cover) shrub layer (0.5-5 m) includes tall shrubs *Forestiera segregata* (Florida swampprivet), *Sideroxylon tenax* (tough bully), *Ilex vomitoria* (yaupon), *Osmanthus americanus* (devilwood), and *Rhus copallinum* (flameleaf sumac), and short shrubs *Baccharis halimifolia* (eastern baccharis), *Borrchia frutescens* (bushy seaside tansy), *Sabal palmetto* (cabbage palmetto), and *Zanthoxylum clava-herculis* (Hercules' club). The herbaceous layer is very sparse (<5% cover) and includes *Solidago sempervirens* (seaside goldenrod). Other species also found include *Prunus caroliniana* (Carolina laurelcherry), *Persea borbonia* (redbay), and *Baccharis halimifolia* (eastern baccharis) (Govus 1998).

Global Vegetation: The canopy of this calciphilic community consists of *Juniperus virginiana* var. *silicicola* (southern redcedar), *Sabal palmetto* (cabbage palmetto), *Celtis laevigata* var. *laevigata* (sugarberry), *Zanthoxylum clava-herculis* (Hercules' club), *Quercus virginiana* (live oak), *Osmanthus americanus* (devilwood), and other species. Shrubs are prominent and may include *Ilex vomitoria* (yaupon), *Yucca aloifolia* (aloe yucca), *Sageretia minutiflora* (small-flower mock buckthorn), *Forestiera segregata* (Florida swampprivet), *Frangula caroliniana* (Carolina buckthorn), *Opuntia humifusa* var. *austrina* (devil's-tongue), and others. Composition is very variable from site to site, depending on environmental conditions and accidents of establishment and persistence.

Most Abundant Species

Fort Pulaski National Monument

<u>Stratum</u>	<u>Lifeform</u>	<u>Species</u>
Tree subcanopy	Needle-leaved tree	<i>Juniperus virginiana</i> var. <i>silicicola</i> (southern redcedar)
Tree subcanopy	Broad-leaved evergreen tree	<i>Forestiera segregata</i> (Florida swampprivet)
Tall shrub/sapling	Broad-leaved evergreen shrub	<i>Forestiera segregata</i> (Florida swampprivet)
Short shrub/sapling	Palm shrub	<i>Sabal palmetto</i> (cabbage palmetto)

Characteristic Species

Fort Pulaski National Monument: *Baccharis halimifolia* (eastern baccharis), *Borrchia frutescens* (bushy seaside tansy), *Forestiera segregata* (Florida swampprivet), *Ilex vomitoria* (yaupon), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Opuntia stricta* var. *stricta*, *Sabal palmetto* (cabbage palmetto), *Solidago sempervirens* (seaside goldenrod), *Zanthoxylum clava-herculis* (Hercules' club)

Other Noteworthy Species

Fort Pulaski National Monument

Exotic and Invasive Species

Ligustrum lucidum (glossy privet)

Morus alba (white mulberry)

Other Plant Species

Forestiera segregata (Florida swampprivet)

I-Rank

-

High/Medium

GRank

G4

Note

exotic

exotic

Note

GA state-imperiled

Global

Other Plant Species

Harrisia simpsonii (Simpson's applecactus)

Lantana depressa var. *floridana* (Florida shrubverbena)

Sideroxylon tenax (tough bully) G3?

GRank

G2

G2T1

Note

imperiled

critically imperiled

Conservation Status Rank

Global Rank & Reasons: G2? (16-Jan-2001). This community is naturally rare because of the relative rarity of the edaphic conditions on which it depends. Examples on isolated islands remain in excellent to good condition (they may have had *Quercus virginiana* removed for ship-building or *Juniperus virginiana* var. *silicicola* removed for pencil manufacture). Other examples on larger and more accessible barrier islands, spits, or peninsulas have in some cases been destroyed by coastal development. Some examples have been altered by exotics, such as *Triadica sebifera*.

Classification

Status: Standard

Classification Confidence: 2 - Moderate

Global Comments: *Sideroxylon tenax* is included in the name as a geographic indicator of the southern Atlantic Coast and distinguishing the type from temperate shell middens on the Gulf Coast. *Zanthoxylum clava-herculis* is an environmental indicator of the high calcium substrate.

Global Similar Associations:

- *Juniperus virginiana* var. *silicicola* - (*Quercus virginiana*, *Sabal palmetto*) Forest (CEGL007813)
- *Quercus virginiana* - (*Sabal palmetto*) Tropical Shell Midden Woodland (CEGL008411)-- occurs farther south, is codominated by tropical species, and lacks *Juniperus virginiana* var. *silicicola*.

Global Related Concepts:

- IB8b. Temperate Shell Midden Woodland (Allard 1990) B

Element Distribution

Fort Pulaski National Monument Range: This association is found on McQueens Island north of Highway 80 and near the Lazaretto Creek fishing pier, on shell deposits near the mouth of the south channel of the Savannah River on northeast McQueens Island (Govus 1998).

Global Range: This calciphilic community occurs on calcareous coastal sands, shell hashes, Amerindian shell middens, and natural shell deposits, from mid-peninsular Florida (Cape Canaveral) north to South Carolina (southern Atlantic Coast).

Nations: US

States/Provinces: FL, GA, NC, SC
TNC Ecoregions: 55:C, 56:C, 57:C
USFS Ecoregions: 232Ce:CCC, 232Gb:CCC
Federal Lands: NPS (Canaveral, Fort Pulaski, Timucuan); USFS (Francis Marion)

Element Sources

Fort Pulaski National Monument Plots: FOPU.8, FOPU.9.
Local Description Authors: M.J. Russo and C.W. Nordman
Global Description Authors: D.J. Allard, mod. A.S. Weakley and H. Summer
References: Allard 1990, Dorroh 1971, Govus 1998, Nelson 1986, Nordman pers. comm., Peet et al. unpubl. data 2002, Southeastern Ecology Working Group n.d.

***Sabal palmetto* Temperate Woodland Alliance (A.479)**

Alliance Concept

Summary: This alliance consists of upland palm savannahs dominated by open to scattered canopy of *Sabal palmetto* (cabbage palmetto). In some cases, these communities occupy narrow strips of upland adjacent to marsh. They have a dense shrubby understory and a high water table. Severe conditions, likely including periodic hurricane damage (by wind and flooding), preclude forest development. They occur in the Ashepoo-Combahee-Edisto (ACE) Basin and may be most common in the Sea Islands Section of the Coastal Plain. In southern Florida, vegetation of this alliance occurs on similar sites to *Pinus elliottii* var. *densa* (Florida slash pine) communities but with more frequent fire (2- to 3-year return time) favoring the palm species *Sabal palmetto* (cabbage palmetto) and *Serenoa repens* (saw palmetto).

Classification Comments: This alliance includes an association (CEGL003796) previously placed in III.A.1.N.d *Serenoa repens* Wooded Shrubland Alliance (A.727).

Similar Alliances:

- *Sabal palmetto* Saturated Woodland Alliance (A.488)
- *Sabal palmetto* Seasonally Flooded Tropical Woodland Alliance (A.1983)--is attributed to the Bahamas and not to the United States.

Related Concepts:

- Cabbage Palmetto: 74 (Eyre 1980) I

Alliance Description

Environment: In some cases, these communities occupy narrow strips of upland adjacent to marsh. They have a dense shrubby understory and a high water table. They occur in the ACE (Ashepoo-Combahee-Edisto) Basin and may be most common in the Sea Islands Section of the Coastal Plain.

Vegetation: This alliance consists of upland palm savannahs dominated by open to scattered canopy of *Sabal palmetto* (cabbage palmetto). In southern Florida, vegetation of this alliance occurs on similar sites to *Pinus elliottii* var. *densa* (Florida slash pine) communities but with

more frequent fire (2- to 3-year return time) favoring the palm species *Sabal palmetto* (cabbage palmetto) and *Serenoa repens* (saw palmetto).

Dynamics: Severe conditions, likely including periodic hurricane damage (by wind and flooding), preclude forest development in the habitat of this alliance. More frequent fire (2- to 3-year return time) favors the palm species *Sabal palmetto* and *Serenoa repens* instead of *Pinus elliottii* var. *densa*.

Alliance Distribution

Range: This alliance is found in Florida, Georgia, and South Carolina.

Nations: US

Subnations: FL, GA, SC

TNC Ecoregions: 53:P, 54:C, 55:C, 56:C, 57:P

USFS Ecoregions: 232Ce:CCC, 232Da:CCC, 232Dc:CCP, 232Ga:CC?, 232Gb:CCC, 411Ac:CCP, 411Af:CCC

Federal Lands: NPS (Canaveral, Fort Pulaski); USFS (Ocala)

(CEGL003526) Cabbage Palmetto Woodland

***Sabal palmetto* - (*Juniperus virginiana* var. *silicicola*) Woodland**

Cabbage Palmetto - (Coastal Redcedar) Woodland

NVC Classification

Physiognomic Class	Woodland (II)
Physiognomic Subclass	Evergreen woodland (II.A.)
Physiognomic Group	Temperate broad-leaved evergreen woodland (II.A.2.)
Physiognomic Subgroup	Natural/Semi-natural temperate broad-leaved evergreen woodland (II.A.2.N.)
Formation	Temperate broad-leaved evergreen woodland (II.A.2.N.a.)
Alliance	<i>Sabal palmetto</i> Temperate Woodland Alliance (A.481)
Alliance (English name)	Cabbage Palmetto Temperate Woodland Alliance
Association	<i>Sabal palmetto</i> - (<i>Juniperus virginiana</i> var. <i>silicicola</i>) Woodland
Association (English name)	Cabbage Palmetto - (Coastal Redcedar) Woodland
Association (Common name)	Cabbage Palmetto Woodland
Ecological System(s):	Southern Atlantic Coastal Plain Maritime Forest (CES203.537)

Element Concept

Global Summary: This community consists of pure to mixed, open stands of *Sabal palmetto* (cabbage palmetto), often with *Juniperus virginiana* var. *silicicola* (southern redcedar). Shrub and herb species can include *Yucca aloifolia* (aloe yucca), *Rhus copallinum* (flameleaf sumac), *Ilex vomitoria* (yaupon), and *Morella cerifera* (wax myrtle). This community is described from the outer South Atlantic Coastal Plain of Georgia and South Carolina, where it occurs in exposed situations on marsh hammock islands. This, or related vegetation, could occur in the adjacent Mid-Atlantic Coastal Plain, East Gulf Coastal Plain and/or Florida Peninsula. More information is needed on the range and variability of this type.

Environmental Description

Fort Pulaski National Monument Environment: This Cabbage Palmetto Woodland was sampled once in the park. The site is a gently sloping ridge with dry-mesic, well-drained clay loam soil. The unvegetated surface is dominated by leaf litter (90% cover) with some wood (4% cover) and bare soil (6% cover). Evidence of disturbance includes the presence of spoil mounds from ditching and exotic plants.

Global Environment: This community occurs in exposed situations on marsh hammock islands.

Vegetation Description

Fort Pulaski National Monument Vegetation: The moderate (60% cover) tree canopy, 5-10 m tall, is dominated by *Celtis laevigata* (sugarberry), *Ilex vomitoria* (yaupon), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Sabal palmetto* (cabbage palmetto), and *Triadica sebifera* (tallowtree) (exotic). The moderate (60% cover) shrub layer (1-5 m) includes dominant *Ilex vomitoria* (yaupon) along with *Morella cerifera* (wax myrtle) and *Rubus trivialis* (southern dewberry). No herbaceous layer was observed, except the vine *Ampelopsis arborea* (peppervine). Other plants documented previously in this community include *Smilax auriculata* (earleaf greenbrier), *Rhus copallinum* (flameleaf sumac), and *Yucca aloifolia* (aloe yucca) (Govus 1998).

Global Vegetation: Examples of this association are pure to mixed, open stands of *Sabal palmetto* (cabbage palmetto), often with *Juniperus virginiana* var. *silicicola* (southern redcedar). Shrub and herb species can include *Yucca aloifolia* (aloe yucca), *Rhus copallinum* (flameleaf sumac), *Ilex vomitoria* (yaupon), and *Morella cerifera* (wax myrtle).

Most Abundant Species

Fort Pulaski National Monument

<u>Stratum</u>	<u>Lifeform</u>	<u>Species</u>
Tree subcanopy	Needle-leaved tree	<i>Juniperus virginiana</i> var. <i>silicicola</i> (southern redcedar)
Tree subcanopy	Broad-leaved deciduous tree	<i>Celtis laevigata</i> (sugarberry), <i>Triadica sebifera</i> (tallowtree)
Tree subcanopy	Broad-leaved evergreen tree	<i>Ilex vomitoria</i> (yaupon)
Tree subcanopy	Palm tree	<i>Sabal palmetto</i> (cabbage palmetto)
Tall shrub/sapling	Broad-leaved evergreen shrub	<i>Ilex vomitoria</i> (yaupon)

Characteristic Species

Fort Pulaski National Monument: *Ampelopsis arborea* (peppervine), *Ilex vomitoria* (yaupon), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Morella cerifera* (wax myrtle), *Sabal palmetto* (cabbage palmetto)

Other Noteworthy Species

Fort Pulaski National Monument

<u>Exotic and Invasive Species</u>	<u>I-Rank</u>	<u>Note</u>
<i>Triadica sebifera</i> (tallowtree)	High	exotic

Conservation Status Rank

Global Rank & Reasons: G2? (31-Jan-2001). This community appears to be rare and restricted, but is poorly understood and circumscribed. As defined, its occurrence is limited to barrier island and barrier spit situations, where it is negatively affected by coastal development. It is also affected by natural disturbances in the form of hurricanes, which probably help maintain this community's structure and composition, as *Juniperus virginiana* var. *silicicola* and *Sabal palmetto* are favored by hurricane disturbance relative to other coastal woody species.

Classification

Status: Standard

Classification Confidence: 2 - Moderate

Global Similar Associations:

- *Juniperus virginiana* var. *silicicola* - (*Quercus virginiana*, *Sabal palmetto*) Forest (CEGL007813)

Element Distribution

Fort Pulaski National Monument Range: This community is located near the picnic area and along the north edge of Cockspur Island, adjacent the high marsh communities (Govus 1998).

Global Range: This community is described from the outer South Atlantic Coastal Plain of Georgia and South Carolina. This, or related vegetation, could occur in the adjacent Mid-Atlantic Coastal Plain, East Gulf Coastal Plain and/or Florida Peninsula.

Nations: US

States/Provinces: FL?, GA, SC

TNC Ecoregions: 53:P, 55:P, 56:C, 57:P

USFS Ecoregions: 232Ce:CCC, 232Da:CPP, 232Dc:CPP, 232Ga:CP?, 232Gb:CPP

Federal Lands: NPS (Fort Pulaski); USFS (Ocala?)

Element Sources

Fort Pulaski National Monument Plots: FOPU.3.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: Southeastern Ecology Group

References: EPA 2004, Govus 1998, Keys et al. 1995, Peet et al. unpubl. data 2002, Southeastern Ecology Working Group n.d.

Shrubland (III.)

Sclerophyllous temperate broad-leaved evergreen shrubland (III. A.2.N.c)

***Quercus virginiana* – *Ilex vomitoria* – (*Morella cerifera*) Shrubland Alliance (A.785)**

Alliance Concept

Summary: This alliance consists of upland maritime dune shrublands of southeastern Coastal Plain (southern, mid-Atlantic, and Gulf coasts). Vegetation of this alliance is characteristically 0.5-3 m tall, typically with a single, extremely dense stratum, with *Quercus virginiana* (live oak) dominant or codominant, and often also with species such as *Ilex vomitoria* (yaupon), *Morella cerifera* (wax myrtle), *Quercus geminata* (sand live oak) (in range), *Serenoa repens* (saw palmetto) (in range), *Smilax auriculata* (earleaf greenbrier), *Prunus caroliniana* (Carolina laurelcherry), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Smilax bona-nox* (saw greenbrier), *Persea borbonia* (redbay), and *Zanthoxylum clava-herculis* (Hercules' club). It occupies linear strands on frontal dune ridges and on the leeward sides and tops of rear dunes, where it is protected from saltwater flooding and extreme salt spray. It also occurs on low dunes and rises on the back sides of barrier islands, as upland rises associated with *Juncus roemerianus* (needlegrass rush) marshes. Examples of this vegetation in Mississippi contain *Baccharis halimifolia* (eastern baccharis).

Classification Comments: This alliance apparently occurs on Cat Island, Mississippi, as scrub thickets with *Ilex vomitoria* dominant and an admixture of *Quercus virginiana* and/or *Quercus geminata*. Occurrences of this (or a closely related alliance) in Louisiana may be better classified as a dwarf forest (*Quercus virginiana* / *Ilex vomitoria*). In Florida, this alliance is best developed on the Atlantic Coast.

Similar Alliances:

- *Morella cerifera* Saturated Shrubland Alliance (A.1906)--includes related vegetation north of the distribution of *Ilex vomitoria*.

Related Concepts:

- Barrier Island Relict Dune Shrubland/Scrub Oak (Wieland 1994b) ?
- Live Oak: 89 (Eyre 1980) I
- Maritime Shrub (Schafale and Weakley 1990) I
- Maritime dune scrub forest (Ambrose 1990a) ?
- Maritime dune shrub vegetation (Ambrose 1990a) ?
- Maritime interdune shrub vegetation (Ambrose 1990a) ?

Alliance Description

Vegetation: Vegetation of this alliance is characteristically 0.5-3 m tall, typically with a single, extremely dense stratum, with *Quercus virginiana* (live oak) dominant or codominant, and often also with species such as *Ilex vomitoria* (yaupon), *Morella cerifera* (wax myrtle), *Quercus geminata* (sand live oak) (in range), *Serenoa repens* (saw palmetto) (in range), *Smilax auriculata* (earleaf greenbrier), *Prunus caroliniana* (Carolina laurelcherry), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Smilax bona-nox* (saw greenbrier), *Persea borbonia* (redbay), and *Zanthoxylum clava-herculis* (Hercules' club). It occupies linear strands on frontal dune ridges and on the leeward sides and tops of rear dunes, where it is protected from saltwater flooding and extreme salt spray. It also occurs on low dunes and rises on the back sides of barrier islands, as upland rises associated with *Juncus roemerianus* (needlegrass rush) marshes. Examples of this vegetation in Mississippi contain *Baccharis halimifolia* (eastern baccharis).

Alliance Distribution

Range: This alliance consists of upland maritime dune shrublands of southeastern Coastal Plain (southern, mid-Atlantic, and Gulf coasts).

Nations: US

Subnations: AL, FL, GA, LA?, MS, NC, SC, VA

TNC Ecoregions: 53:C, 55:?, 56:C, 57:C

USFS Ecoregions: 232Cb:CCC, 232Ce:CCC, 232Ci:CCC, 232Dc:CCC, 232Dd:CCC, 232Gb:CCC

Federal Lands: DOD (Camp Lejeune?); NPS (Canaveral, Cape Hatteras, Cape Lookout, Fort Pulaski, Fort Sumter, Gulf Islands); USFWS (Back Bay?, Bon Secour, Cape Romain?, Pea Island?)

(CEGL004784) Wax myrtle - Carolina Laurel Cherry - Toothache-tree Shrubland
***Morella cerifera* - *Prunus caroliniana* - *Zanthoxylum clava-herculis* Shrubland**

NVC Classification

Physiognomic Class	Shrubland (III)
Physiognomic Subclass	Evergreen shrubland (III.A.)
Physiognomic Group	Temperate broad-leaved evergreen shrubland (III.A.2.)
Physiognomic Subgroup	Natural/Semi-natural temperate broad-leaved evergreen shrubland (III.A.2.N.)
Formation	Sclerophyllous temperate broad-leaved evergreen shrubland (III.A.2.N.c.)
Alliance	<i>Quercus virginiana</i> - <i>Ilex vomitoria</i> - (<i>Morella cerifera</i>) Shrubland Alliance (A.785)
Alliance (English name)	Live Oak - Yaupon - (Wax myrtle) Shrubland Alliance
Association	<i>Morella cerifera</i> - <i>Prunus caroliniana</i> - <i>Zanthoxylum clava-herculis</i> Shrubland
Association (English name)	Wax myrtle - Carolina Laurel Cherry - Toothache-tree Shrubland
Ecological System(s):	Central Atlantic Coastal Plain Maritime Forest (CES203.261)

Element Concept

Global Summary: Maritime shrublands dominated by *Morella cerifera* (wax myrtle), *Prunus caroliniana* (Carolina laurelcherry), and *Zanthoxylum clava-herculis* (Hercules' club). This shrub vegetation occurs on the back side of maritime dunes.

Environmental Description

Fort Pulaski National Monument Environment: This maritime shrubland was sampled once in the park. The site is a flat spoil mound with well-drained sand. The unvegetated surface is dominated by leaf litter (84% cover) with some wood (5% cover), bare soil (10% cover), and shells and debris (1% cover). Evidence of disturbance includes deer browse, many downed trees, the presence of spoil and debris mounds, and exotic plants.

Fort Sumter National Monument Environment: This maritime shrubland was sampled at one location at Fort Moultrie. The site is a flat interdunal swale that is part of an accreting barrier island. Soil is dry-mesic, well-drained sand. The unvegetated surface is dominated by leaf litter (90% cover) with some bare soil (10%). The site is vulnerable to salt spray or flooding from storm overwash under severe weather conditions.

Vegetation Description

Fort Pulaski National Monument Vegetation: The moderate (40% cover) tree canopy, 10-15 m tall, is dominated by *Celtis laevigata* (sugarberry) and includes lesser amounts of *Morella cerifera* (wax myrtle), *Prunus caroliniana* (Carolina laurelcherry), and *Triadica sebifera* (tallowtree) (exotic). The dense (80% cover) subcanopy (5-10 m) is dominated by *Prunus caroliniana* (Carolina laurelcherry) and *Morella cerifera* (wax myrtle), as is the moderately sparse (30% cover) shrub layer (1-5 m) along with *Sabal palmetto* (cabbage palmetto). The very sparse (<5% cover) herbaceous layer is comprised of *Eragrostis spectabilis* (purple lovegrass) and *Eremochloa ophiuroides* (centipede grass) (exotic). Vines include *Parthenocissus quinquefolia* (Virginia creeper).

Fort Sumter National Monument Vegetation: The moderately dense (70% cover) tall shrub layer (2-5 m tall) is dominated by *Morella cerifera* (wax myrtle) along with lesser amounts of *Zanthoxylum clava-herculis* (Hercules' club); short shrubs include *Prunus caroliniana* (Carolina laurelcherry) and *Rubus trivialis* (southern dewberry). The herbaceous layer is sparse and includes small amounts of *Andropogon virginicus* (broomsedge bluestem), *Heterotheca subaxillaris* (camphorweed), *Oenothera humifusa* (seabeach evening primrose), *Opuntia pusilla*

(cockspur pricklypear), and *Yucca filamentosa* (Adam's needle). Vines (20%) are dominated by *Smilax bona-nox* (saw greenbrier) along with *Smilax auriculata* (earleaf greenbrier) and *Vitis rotundifolia* (muscadine).

Most Abundant Species

Fort Pulaski National Monument

<u>Stratum</u>	<u>Lifeform</u>	<u>Species</u>
Tree (canopy & subcanopy)	Broad-leaved evergreen tree	<i>Morella cerifera</i> (wax myrtle), <i>Prunus caroliniana</i> (Carolina laurelcherry)
Tree canopy	Broad-leaved deciduous tree	<i>Celtis laevigata</i> (sugarberry)
Shrub/sapling (tall & short)	Broad-leaved evergreen shrub	<i>Prunus caroliniana</i> (Carolina laurelcherry)
Tall shrub/sapling	Broad-leaved evergreen shrub	<i>Ilex vomitoria</i> (yaupon)
Tall shrub/sapling	Palm shrub	<i>Sabal palmetto</i> (cabbage palmetto)

Characteristic Species

Fort Pulaski National Monument: *Morella cerifera* (wax myrtle), *Prunus caroliniana* (Carolina laurelcherry)

Fort Sumter National Monument

<u>Stratum</u>	<u>Lifeform</u>	<u>Species</u>
Tall shrub/sapling	Broad-leaved evergreen shrub	<i>Morella cerifera</i> (wax myrtle)
Herb (field)	Vine/Liana	<i>Smilax bona-nox</i> (saw greenbrier)

Other Noteworthy Species

Fort Pulaski National Monument

<u>Exotic and Invasive Species</u>	<u>I-Rank</u>	<u>Note</u>
<i>Eremochloa ophiuroides</i> (centipede grass)	-	exotic
<i>Lonicera japonica</i> (Japanese honeysuckle)	High/Medium	exotic

Conservation Status Ranks

Global Rank & Reasons: G2? (8-Oct-1997).

Classification

Status: Standard

Classification Confidence: 2 - Moderate

Global Comments: This sounds like the understory of maritime forest, possibly disturbed by hurricane killed overstory (M. Schafale pers. comm.). More information is available in the ACE data for Otter Island (South end).

Element Distribution

Fort Pulaski National Monument Range: This community is found in a spoil area on the west side of the main island.

Fort Sumter National Monument Range: This community was sampled just south of Fort Moultrie and mile east of the Station B beach access.

Nations: US

States/Provinces: GA, SC

TNC Ecoregions: 56:C

USFS Ecoregions: 232Ce:CCC

Federal Lands: NPS (Fort Pulaski, Fort Sumter)

Element Sources

Fort Pulaski National Monument Plots: FOPU.11.

Fort Sumter National Monument Plots: FOSU.2.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: K. Tassin and C. Smith

References: Schafale pers. comm., Southeastern Ecology Working Group n.d.

Saturated temperate broad-leaved evergreen shrubland (III.A.2.N.i.)

***Morella cerifera* Saturated Shrubland Alliance (A.1906)**

Alliance Concept

Summary: This alliance includes wetland dune swales and other wetland shrubland situations dominated by *Morella cerifera* (wax myrtle), including sheltered backdunes, wetland flats, and interdune swales. Associated shrubs may include *Baccharis halimifolia* (eastern baccharis), *Acer rubrum* (red maple), *Vaccinium formosum* (southern blueberry), *Rosa palustris* (swamp rose), *Ilex opaca* var. *opaca* (American holly), *Juniperus virginiana* var. *silicicola* (southern redcedar), and *Morella pensylvanica* (northern bayberry). This shrubland vegetation can form vast thickets in some cases; there may be inclusions of small wetland graminoid-dominated areas. Vines may be frequent and can include *Toxicodendron radicans* ssp. *radicans* (eastern poison ivy), *Smilax* (greenbrier) spp., *Vitis* (grape) spp., and *Parthenocissus quinquefolia* (Virginia creeper). Associated herbaceous species can include *Boehmeria cylindrica* (smallspike false nettle), *Calystegia sepium* (hedge false bindweed), *Carex albolutescens* (greenwhite sedge), *Carex hormathodes* (marsh straw sedge), *Chasmanthium laxum* (slender woodoats), *Festuca rubra* (red fescue), *Galium obtusum* (bluntleaf bedstraw), *Hydrocotyle* (hydrocotyle) spp., *Juncus dichotomus* (forked rush), *Juncus scirpoides* (needlepod rush), *Leersia virginica* (whitegrass), *Osmunda regalis* var. *spectabilis* (royal fern), *Panicum* (panicgrass) spp., *Polygonum pensylvanicum* (Pennsylvania smartweed), and *Woodwardia areolata* (netted chainfern) (in the northern part of the range); and *Andropogon glomeratus* var. *pumilus* (bushy bluestem), *Fimbristylis castanea* (marsh fimbry), *Hydrocotyle bonariensis* (largeleaf pennywort), *Juncus roemerianus* (needlegrass rush), *Muhlenbergia filipes* (gulfhairawn muhly), as well as *Spartina patens* (saltmeadow cordgrass) and *Sabatia stellaris* (rose of Plymouth) (in southern examples). This is maritime shrub vegetation which generally exists in a tension zone between more exposed or active grassland vegetation and more protected and stable maritime forests. Older stands of maritime shrub vegetation ultimately become dominated by stunted and salt-pruned *Quercus virginiana* (live oak). See also the III.A.2.N.c *Quercus virginiana* - *Ilex vomitoria* - (*Morella cerifera*) Shrubland Alliance (A.785), which generally occurs in drier and more exposed sites. Soils are deep sands of stabilized dunes, with limited if any horizon development. An overlying layer of 'muck' is reported in northern examples.

Similar Alliances:

- *Juncus dichotomus* Seasonally Flooded Herbaceous Alliance (A.1427)
- *Quercus virginiana* - *Ilex vomitoria* - (*Morella cerifera*) Shrubland Alliance (A.785)-- generally occurs in drier and more exposed sites.

Similar Alliance Comments: Older stands of maritime shrub vegetation ultimately become dominated by stunted and salt-pruned *Quercus virginiana*.

Related Concepts:

- Coastal Interdunal Swale (FNAI 1992a) ?
- Maritime Shrub (Schafale and Weakley 1990) I
- Maritime interdune shrub vegetation (Ambrose 1990a) ?
- Shrub succession community (Hill 1986) I
- Mesic shrub zone (Higgins et al. 1971) I
- Swamp thicket (Klotz 1986) I
- Thicket community (Boule 1979) I

Alliance Description

Environment: This is maritime shrub vegetation which generally exists in a tension zone between more exposed or active grassland vegetation and more protected and stable maritime forests. Older stands of maritime shrub vegetation ultimately become dominated by stunted and salt-pruned *Quercus virginiana*. Soils are deep sands of stabilized dunes, with limited if any horizon development. An overlying layer of 'muck' is reported in northern examples.

Vegetation: This alliance includes wetland dune swales and other wetland shrubland situations dominated by *Morella cerifera* (wax myrtle), including sheltered backdunes, wetland flats, and interdune swales. Associated shrubs may include *Baccharis halimifolia* (eastern baccharis), *Acer rubrum* (red maple), *Vaccinium formosum* (southern blueberry), *Rosa palustris* (swamp rose), *Ilex opaca* (American holly), *Juniperus virginiana* var. *silicicola* (southern redcedar), and *Morella pensylvanica* (northern bayberry). This shrubland vegetation can form vast thickets in some cases; there may be inclusions of small wetland graminoid-dominated areas. Vines may be frequent and can include *Toxicodendron radicans* ssp. *radicans* (eastern poison ivy), *Smilax* (greenbrier) spp., *Vitis* (grape) spp., and *Parthenocissus quinquefolia* (Virginia creeper). Associated herbaceous species can include *Boehmeria cylindrica* (smallspike false nettle), *Calystegia sepium* (hedge false bindweed), *Carex albolutescens* (greenwhite sedge), *Carex hormathodes* (marsh straw sedge), *Chasmanthium laxum* (slender woodoats), *Festuca rubra* (red fescue), *Galium obtusum* (bluntleaf bedstraw), *Hydrocotyle* (hydrocotyle) spp., *Juncus dichotomus* (forked rush), *Juncus scirpoides* (needlepod rush), *Leersia virginica* (whitegrass), *Osmunda regalis* var. *spectabilis* (royal fern), *Panicum* (panicgrass) spp., *Polygonum pensylvanicum* (Pennsylvania smartweed), and *Woodwardia areolata* (netted chainfern) (in the northern part of the range); and *Andropogon glomeratus* var. *pumilus* (bushy bluestem), *Fimbristylis castanea* (marsh fimbry), *Hydrocotyle bonariensis* (largeleaf pennywort), *Juncus roemerianus* (needlegrass rush), *Muhlenbergia filipes* (gulfhairawn muhly), as well as *Spartina patens* (saltmeadow cordgrass) and *Sabatia stellaris* (rose of Plymouth) (in southern examples).

Dynamics: Older stands of maritime shrub vegetation ultimately become dominated by stunted and salt-pruned *Quercus virginiana*.

Alliance Distribution

Range: This alliance is found in coastal areas from Louisiana and Florida north to New Jersey.

Nations: US

Subnations: DE, FL, GA, LA, MD, NC, NJ, SC, VA

TNC Ecoregions: 31:C, 54:P, 55:C, 56:C, 57:C, 58:C, 62:C

USFS Ecoregions: 232Ab:CCP, 232Ac:CCC, 232Ad:CCP, 232Bx:CCC, 232Bz:CCC, 232Cb:CCP, 232Ce:CCC, 232Ch:CCP, 232Ci:CCC, 232E:CC, 232Gb:CCC

Federal Lands: DOD (Cape Canaveral); NPS (Assateague Island, Canaveral?, Cape Hatteras, Cape Lookout, Fort Pulaski, Timucuan?); USFWS (Back Bay?, Cape Romain?, Chincoteague, Merritt Island?, Pea Island, Prime Hook)

(CEGL003839) Atlantic Coast Interdune Swale

***Morella cerifera*/Spartina patens Shrubland**

Wax myrtle/Saltmeadow Cordgrass Shrubland

NVC Classification

Physiognomic Class	Shrubland (III)
Physiognomic Subclass	Evergreen shrubland (III.A.)
Physiognomic Group	Temperate broad-leaved evergreen shrubland (III.A.2.)
Physiognomic Subgroup	Natural/Semi-natural temperate broad-leaved evergreen shrubland (III.A.2.N.)
Formation	Saturated temperate broad-leaved evergreen shrubland (III.A.2.N.i.)
Alliance	<i>Morella cerifera</i> Saturated Shrubland Alliance (A.1906)
Alliance (English name)	Wax myrtle Saturated Shrubland Alliance
Association	<i>Morella cerifera</i> / <i>Spartina patens</i> Shrubland
Association (English name)	Wax myrtle / Saltmeadow Cordgrass Shrubland
Association (Common name)	Atlantic Coast Interdune Swale
Ecological System(s):	Southern Atlantic Coastal Plain Dune and Maritime Grassland (CES203.273) Northern Atlantic Coastal Plain Dune and Swale (CES203.264) Central Atlantic Coastal Plain Maritime Forest (CES203.261)

Element Concept

Global Summary: This association occurs in wetland flats, backdunes and interdune swales along the Atlantic Coast from New Jersey to Florida. The substrate may be sand or loamy sand, sometimes with a thin layer of organic matter. It occurs beyond the reach of most storm tides but may be impacted by salt spray. The water table may often be less than half a meter below the surface. Canopy closure varies from densely closed to moderately open. This vegetation is dominated by *Morella cerifera* (wax myrtle), sometimes intermixed with *Juniperus virginiana* var. *silicicola* (southern redcedar) (from North Carolina and possibly southern Virginia southward), *Morella pensylvanica* (northern bayberry) (from northern North Carolina northward), *Rhus copallinum* (flameleaf sumac), *Baccharis halimifolia* (eastern baccharis), or *Ilex vomitoria* (yaupon). The shrub layer can vary from 2-8 m in height. Occasionally, a sparse emergent canopy of trees (such as *Juniperus virginiana* var. *silicicola* (southern redcedar) or *Sabal palmetto* (cabbage palmetto) in the southern parts of the range) may be present. The herbaceous layer is characterized by *Spartina patens* (saltmeadow cordgrass). Other ground flora associates include *Toxicodendron radicans* (eastern poison ivy), *Panicum virgatum* (switchgrass), *Andropogon virginicus* (broomsedge bluestem), *Juncus dichotomus* (forked rush), *Solidago sempervirens* (seaside goldenrod), *Smilax* (greenbrier) spp., *Parthenocissus quinquefolia* (Virginia creeper), *Vitis* (grape) spp., and *Schoenoplectus pungens* (common threesquare). Small openings may have wetland graminoids, such as *Juncus roemerianus* (needlegrass rush), *Spartina patens* (saltmeadow cordgrass), *Fimbristylis castanea* (marsh fimbry), *Andropogon glomeratus* var. *pumilus* (bushy bluestem), *Muhlenbergia filipes* (gulfhairawn muhly), etc., and forbs, such as *Hydrocotyle bonariensis* (largeleaf pennywort), *Sabatia stellaris* (rose of Plymouth), *Polygonum hydropiperoides* (swamp smartweed),

Eleocharis (spikerush) spp., etc. In other cases, few herbs are present, because of the dense, thicket-like shrub layer.

Environmental Description

USFWS Wetland System: Palustrine

Fort Pulaski National Monument Environment: This wax myrtle shrubland was sampled once in the park. The site is a gently sloping, southeast-facing lowland adjacent to a spoil area with somewhat poorly drained sand soil. The unvegetated surface is made up of leaf litter (80% cover), wood (10% cover) and bare soil (10% cover). Some sites have saturated soils (Govus 1998).

Global Environment: This association occurs on maritime barrier islands and spits, usually on more protected backdunes, barrier flats, islands within salt marshes, and interdunal swales. It is not tidally flooded and occurs beyond the reach of most storm tides, but it is impacted by salt spray. The substrate may be sand or loamy sand, sometimes with a thin layer of organic matter. The water table is often less than half a meter below the surface.

Vegetation Description

Fort Pulaski National Monument Vegetation: Shrubs 2-5 m tall form the dominant layer (80% cover). However, in this example there is also a moderate (30% cover) emergent tree layer of *Celtis laevigata* (sugarberry) and *Morella cerifera* (wax myrtle). Occasionally this association includes widely scattered *Juniperus virginiana* var. *silicicola* and *Sabal palmetto* (cabbage palmetto) (Govus 1998). The shrub layer is dominated by *Morella cerifera* (wax myrtle) along with *Ilex vomitoria* (yaupon), *Sabal palmetto* (cabbage palmetto), and *Rubus trivialis* (southern dewberry). No herbs were observed. The shrubs are so densely arranged that no herbaceous species are afforded light or space to grow (Govus 1998). Vines include *Ampelopsis arborea* (peppervine), *Parthenocissus quinquefolia* (Virginia creeper), *Smilax auriculata* (earleaf greenbrier), and *Smilax bona-nox* (saw greenbrier). In the eastern portion (within the dike system), a substantial amount of *Triadica sebifera* (tallowtree) (exotic) also occurs as widely scattered individuals (Govus 1998).

Global Vegetation: This vegetation is characterized by a moderately open to densely closed canopy of *Morella cerifera* (wax myrtle). Other canopy associates include *Baccharis halimifolia* (eastern baccharis), *Morella pensylvanica* (northern bayberry) (from northern NC northward), *Juniperus virginiana* var. *silicicola* (southern redcedar) (from NC and possibly southern VA southward), and *Rhus copallinum* (flameleaf sumac). The herbaceous layer is characterized by *Spartina patens* (saltmeadow cordgrass). Other ground flora associates include *Toxicodendron radicans* (eastern poison ivy), *Panicum virgatum* (switchgrass), *Andropogon virginicus* (broomsedge bluestem), *Juncus dichotomus* (forked rush), *Solidago sempervirens* (seaside goldenrod), *Smilax* (greenbrier) spp., *Parthenocissus quinquefolia* (Virginia creeper), *Vitis* (grape) spp., and *Schoenoplectus pungens* (common threesquare). Small openings may have wetland graminoids, such as *Juncus roemerianus* (needlegrass rush), *Spartina patens* (saltmeadow cordgrass), *Fimbristylis castanea* (marsh fimbry), *Andropogon glomeratus* var. *pumilus* (bushy bluestem), *Muhlenbergia filipes* (gulfhairawn muhly), etc., and forbs, such as *Hydrocotyle bonariensis* (largeleaf pennywort), *Sabatia stellaris* (rose of Plymouth), *Polygonum*

hydropiperoides (swamp smartweed), *Eleocharis* (spikerush) spp., etc. In other cases, few herbs are present, because of the dense, thicket-like shrub layer.

Most Abundant Species

Fort Pulaski National Monument

<u>Stratum</u>	<u>Lifeform</u>	<u>Species</u>
Tree subcanopy	Broad-leaved evergreen tree	<i>Morella cerifera</i> (wax myrtle)
Tall shrub/sapling	Broad-leaved evergreen shrub	<i>Ilex vomitoria</i> (yaupon), <i>Morella cerifera</i> (wax myrtle)

Characteristic Species

Fort Pulaski National Monument: *Ampelopsis arborea* (peppervine), *Celtis laevigata* (sugarberry), *Ilex vomitoria* (yaupon), *Morella cerifera* (wax myrtle), *Parthenocissus quinquefolia* (Virginia creeper), *Sabal palmetto* (cabbage palmetto)

Other Noteworthy Species

Fort Pulaski National Monument

<u>Exotic and Invasive Species</u>	<u>I-Rank</u>	<u>Note</u>
<i>Triadica sebifera</i> (tallowtree)	High	exotic

Conservation Status Rank

Global Rank & Reasons: G3G4 (1-Feb-2005). This association occurs in interdune flats on barrier islands and barrier spits from New Jersey south to Florida. Many occurrences have been destroyed by coastal development, as these soils are sandy enough and dry enough that they are often not considered "jurisdictional wetlands" and are therefore destroyed for development. Other occurrences have been left undeveloped but now occur as isolated areas fragmented by development. Fairly extensive occurrences are protected on Cape Hatteras National Seashore and Cape Lookout National Seashore, North Carolina.

Classification

Status: Standard

Classification Confidence: 2 - Moderate

Global Comments: This association now encompasses two previous associations, one more northerly ranging and the other more southerly ranging along the Atlantic Coast. Where the ranges of these two types overlapped, they were difficult to distinguish floristically and were thus combined.

Global Similar Associations:

- (*Morella cerifera*) - *Panicum virgatum* - *Spartina patens* Herbaceous Vegetation (CEGL004129)
- *Morella cerifera* - *Vaccinium corymbosum* Shrubland (CEGL003906)
- *Morella cerifera* / *Hydrocotyle verticillata* Shrubland (CEGL003840)

Global Related Concepts:

- Maritime Interdune Shrub Vegetation (Ambrose 1990a) ?
- Mesic shrub community (Higgins et al. 1971) B
- Salt grass sea myrtle redcedar savannah (Martin 1959b) ?
- Salt grass sea myrtle savannah (Martin 1959b) ?

- Shrub succession community (Hill 1986) B
- Thicket community (Boule 1979) =
- Upland thicket (Klotz 1986) =

Element Distribution

Fort Pulaski National Monument Range: This community is found near the dredge spoil area adjacent to the maintenance shed. It occurs along the southern edge of Cockspur Island, just above the tidal shrublands and salt panne communities of central Cockspur Island and within the dike system near the southeastern part of the island in very low-lying areas (Govus 1998).

Global Range: This type occurs along the Atlantic Coast from New Jersey south to Florida.

Nations: US

States/Provinces: DE, FL, GA, MD, NC, NJ, SC, VA

TNC Ecoregions: 56:C, 57:C, 58:C, 62:C

USFS Ecoregions: 232Ab:CCP, 232Ac:CCC, 232Bz:CCC, 232Cb:CCP, 232Ce:CCC, 232Ch:CCP, 232Ci:CCC, 232Gb:CCC

Federal Lands: DOD (Cape Canaveral); NPS (Assateague Island, Canaveral?, Cape Hatteras, Cape Lookout, Fort Pulaski, Timucuan?); USFWS (Back Bay?, Cape Romain?, Chincoteague, Merritt Island?, Pea Island, Prime Hook)

Element Sources

Fort Pulaski National Monument Plots: FOPU.15.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: A.S. Weakley, mod. J. Teague

References: Ambrose 1990a, Berdine 1998, Boule 1979, Bowman 2000, Breden et al. 2001, FNAI 1992a, Fleming 2001a, Govus 1998, Harrison 2004, Higgins et al. 1971, Hill 1986, Klotz 1986, Martin 1959b, Schafale and Weakley 1990, Southeastern Ecology Working Group n.d.

Tidal cold-deciduous shrubland (III.B.2.N.h.)

***Baccharis halimifolia* - *Iva frutescens* Tidal Shrubland Alliance (A.1023)**

Alliance Concept

Summary: This alliance includes maritime scrub communities typically dominated by *Iva frutescens* (Jesuit's bark) or *Baccharis halimifolia* (eastern baccharis) or both, growing in association with salt marshes. These communities occur primarily in estuarine margin situations, especially on the sound sides of barrier islands. Characteristically, these communities form an ecotone between salt marsh and upland vegetation or in areas within the salt marsh having slightly higher elevations and lower salinity levels than the surrounding marsh. Storm-induced disturbance causes periodic die-back of the shrubs restricting the extent of their spread. Characteristic species include *Baccharis halimifolia* (eastern baccharis), *Iva frutescens* (Jesuit's bark), *Rosa carolina* (Carolina rose), *Spartina patens* (saltmeadow cordgrass), and *Panicum virgatum* (switchgrass).

Similar Alliances:

- *Baccharis halimifolia* Saturated Shrubland Alliance (A.1015)
- *Panicum virgatum* Tidal Herbaceous Alliance (A.1476)

Related Concepts:

- *Baccharis halimifolia* (Klemas et al. 1973) ?
- *Iva frutescens*-*Baccharis halimifolia* (Good 1965) ?
- *Iva frutescens* (Klemas et al. 1973) ?
- Estuarine Intertidal: Salt Marsh (Swain and Kearsley 2001) ?
- Estuarine scrub-shrub wetland (Tiner 1985b) ?
- Salt Shrub (Schafale and Weakley 1990) I
- Salt bush - salt meadow marsh (Daiber et al. 1976) ?
- Salt marsh community (Hill 1986) I
- Salt marsh complex, marsh-upland border (Breden 1989) ?
- Salt shrub (Reschke 1990) ?
- Tidal Marsh (FNAI 1992a) I
- Salt grass - marsh-elder savannah (Martin 1959b) ?
- Salt marsh and upper border (Barry 1980) ?
- Saltbush zone (Boule 1979) ?
- Shrub succession community (Higgins et al. 1971) I
- Swamp thicket (Klotz 1986) I

Alliance Description

Environment: These communities occur primarily in estuarine margin situations, especially on the sound sides of barrier islands. Characteristically, these communities form an ecotone between salt marsh and upland vegetation or in areas within the salt marsh having slightly higher elevations and lower salinity levels than the surrounding marsh. Storm-induced disturbance causes periodic die-back of the shrubs restricting the extent of their spread.

Vegetation: This alliance includes maritime scrub communities typically dominated by *Iva frutescens* (Jesuit's bark) or *Baccharis halimifolia* (eastern baccharis) or both, growing in association with salt marshes. Characteristic species include *Baccharis halimifolia* (eastern baccharis), *Iva frutescens* (Jesuit's bark), *Rosa carolina* (Carolina rose), *Spartina patens* (saltmeadow cordgrass), and *Panicum virgatum* (switchgrass).

Dynamics: Storm-induced disturbance causes periodic die-back of the shrubs restricting the extent of their spread.

Alliance Distribution

Range: This alliance is found throughout the eastern United States from Maine to Florida and west to Texas.

Nations: US

Subnations: AL, CT, DE, FL, GA, LA, MA, MD, MS, NC, NJ, NY, RI, SC, TX, VA

TNC Ecoregions: 31:C, 53:C, 55:?, 56:C, 57:C, 58:C, 62:C

USFS Ecoregions: 212P:PP, 221Aa:CCC, 221Ab:CCC, 221Ac:CCC, 221Ad:CCC, 221Ae:CCP, 221Ak:CCC, 221Dc:CPP, 231Fb:CCC, 232Aa:CCC, 232Ab:CCC, 232Ac:CCC, 232Ad:CCC, 232Bb:CC?, 232Bc:CCP, 232Bd:CCP, 232Br:CCC, 232Bt:CCC, 232Bx:CCC, 232Bz:CCC, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC, 232Dc:CCC, 232Eb:CCC, 255Dc:CCC

Federal Lands: NPS (Assateague Island, Boston Harbor Islands, Cape Cod, Cape Hatteras, Colonial, Fire Island, Fort Pulaski, Gateway, George Washington Birthplace, Timucuan?); USFS

(Croatan); USFWS (Anahuac, Aransas, Back Bay?, Big Boggy, Bon Secour, Brazoria, Chesapeake Marshlands, Chincoteague, Matagorda Island, McFaddin, Monomoy?, Parker River, Prime Hook, San Bernard)

(CEGL003920) Coastal Salt Shrub Thicket

***Baccharis halimifolia* - *Iva frutescens* - *Morella cerifera* - (*Ilex vomitoria*) Shrubland**

Groundsel Tree - Maritime Marsh-elder - Wax Myrtle - (Yaupon) Shrubland

NVC Classification

Physiognomic Class	Shrubland (III)
Physiognomic Subclass	Deciduous shrubland (III.B.)
Physiognomic Group	Cold-deciduous shrubland (III.B.2.)
Physiognomic Subgroup	Natural/Semi-natural cold-deciduous shrubland (III.B.2.N.)
Formation	Tidal cold-deciduous shrubland (III.B.2.N.h.)
Alliance	<i>Baccharis halimifolia</i> - <i>Iva frutescens</i> Tidal Shrubland Alliance (A.1023)
Alliance (English name)	Groundsel tree - Maritime Marsh-elder Tidal Shrubland Alliance
Association	<i>Baccharis halimifolia</i> - <i>Iva frutescens</i> - <i>Morella cerifera</i> - (<i>Ilex vomitoria</i>) Shrubland
Association (English name)	Groundsel tree - Maritime Marsh-elder - Wax myrtle - (Yaupon) Shrubland
Association (Common name)	Coastal Salt Shrub Thicket
Ecological System(s):	Central Atlantic Coastal Plain Fresh and Oligohaline Tidal Marsh (CES203.376)

Element Concept

Global Summary: This shrubland, dominated by the nominal species, occurs in slightly elevated areas within salt flats and salt marshes as well as in marsh edges throughout much of the East Gulf, South Atlantic, and Mid-Atlantic coastal plains. This community is usually best developed at the upper limit of non-storm tidal inundation, on natural levees deposited by above-normal tides. The most common species are typically *Baccharis halimifolia* (eastern baccharis), *Morella cerifera* (wax myrtle), *Iva frutescens* ssp. *frutescens* (Jesuit's bark), *Yucca gloriosa* (moundlily yucca), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Lycium carolinianum* (Carolina desert-thorn), *Baccharis angustifolia* (saltwater false willow), and *Ilex vomitoria* (yaupon). Other species which may be present include *Borrichia frutescens* (bushy seaside tansy), *Fimbristylis castanea* (marsh fimbry), *Limonium carolinianum* (Carolina sealavender), and *Solidago sempervirens* (seaside goldenrod).

Environmental Description

USFWS Wetland System: Estuarine

Fort Pulaski National Monument Environment: This community occurs as a fringe shrubland between either salt panne communities and upland forests or high marsh communities and upland communities. It is especially well-developed along the southern edge of Cockspur Island where extensive salt flats grade gently into the adjacent upland communities (Govus 1998).

Vegetation Description

Fort Pulaski National Monument Vegetation: The dominant species of this association are *Baccharis halimifolia* (eastern baccharis), *Iva frutescens* (Jesuit's bark), *Morella cerifera* (wax myrtle), and *Ilex vomitoria* (yaupon), other species present include *Baccharis angustifolia* (saltwater false willow), *Limonium carolinianum* (Carolina sealavender), and *Fimbristylis castanea* (marsh fimbry) (Govus 1998).

Most Abundant Species

Fort Pulaski National Monument

Stratum

Shrub/sapling (tall & short)

Lifeform

Broad-leaved evergreen shrub

Short shrub/sapling

Broad-leaved evergreen shrub

Species

Baccharis halimifolia (eastern baccharis), *Ilex vomitoria* (yaupon)

Morella cerifera (wax myrtle)

Conservation Status Rank

Global Rank & Reasons: G4? (11-Aug-1997).

Classification

Status: Standard

Classification Confidence: 2 - Moderate

Global Comments: An example documented in South Carolina on Old Island had an emergent layer of *Juniperus virginiana* var. *silicicola*, over *Iva frutescens* and *Borrchia frutescens*. The herbaceous layer consisted of *Cynanchum angustifolium*, *Juncus roemerianus*, and *Fimbristylis castanea*. This type is not known to occur in VA (G.P. Fleming and P. Coulling pers. comm.).

Global Similar Associations:

- *Baccharis halimifolia* - *Iva frutescens* / *Panicum virgatum* Shrubland (CEGL003921)--is potentially overlapping in some areas.
- *Iva frutescens* / *Spartina patens* Shrubland (CEGL006848)

Global Related Concepts:

- Salt Shrub (Inland High Subtype) (Schafale 2000) ?

Element Distribution

Fort Pulaski National Monument Range: This community is widespread on Cockspur Island, especially along the southern edge of the island where extensive salt flats grade gently into the adjacent upland communities (Govus 1998).

Global Range: This shrubland is found throughout much of the East Gulf, South Atlantic, and Mid-Atlantic coastal plains.

Nations: US

States/Provinces: AL, FL, GA, LA, MS, NC, SC

TNC Ecoregions: 53:C, 55:?, 56:C, 57:C

USFS Ecoregions: 232Bz:CCC, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC, 232Dc:CCC, 232Eb:CCC

Federal Lands: NPS (Cape Hatteras, Fort Pulaski, Timucuan?); USFS (Croatan); USFWS (Bon Secour)

Element Sources

Fort Pulaski National Monument Plots: None.

Local Description Authors: T. Govus

Global Description Authors:

References: Coulling pers. comm., FNAI 1992a, Fleming pers. comm., Govus 1998, Lea 2002b, Schafale 2000, Schotz pers. comm., Southeastern Ecology Working Group n.d., Wolfe 1990

***Borrichia frutescens* Tidal Shrubland Alliance (A.1026)**

Alliance Concept

Summary: This alliance includes tidal areas in upper marsh edges and flats dominated by *Borrichia frutescens* (bushy seaside tansy). It occurs at spatial scales ranging from narrow bands to flats in excess of 500 ha; it often occurs either just below *Iva frutescens* (Jesuit's bark) or *Baccharis halimifolia* (eastern baccharis), or just above *Juncus roemerianus* (needlegrass rush) or *Spartina alterniflora* (smooth cordgrass). Other characteristic species include *Spartina patens* (saltmeadow cordgrass), *Iva frutescens* ssp. *frutescens* (Jesuit's bark), *Iva frutescens* ssp. *oraria* (Jesuit's bark), and *Limonium carolinianum* (Carolina sealavender). In Texas, the composition of this low-diversity, irregularly tidally flooded shrubland is variable and can include *Spartina spartinae* (gulf cordgrass), *Sporobolus virginicus* (seashore dropseed), *Monanthochloe littoralis* (shoregrass), *Sarcocornia perennis* (chickenclaws), *Limonium carolinianum* (Carolina sealavender), *Lycium carolinianum* var. *quadrifidum* (Carolina desert-thorn), *Rayjacksonia phyllocephala* (camphor daisy), and *Batis maritima* (turtleweed). Here, these communities often occur just above the lower tidal flat communities *Batis maritima* - *Sarcocornia perennis* Dwarf-shrubland (CEGL003956) or *Sarcocornia perennis* - (*Batis maritima*, *Distichlis spicata*) Dwarf-shrubland (CEGL002278).

Classification Comments: Consideration should be given to placement of this alliance in Class IV. Dwarf-shrubland. Although stature can be more than 0.5 m, it is typically 0.3-0.5 m in height.

Related Concepts:

- Glasswort-Saltwort Series (Diamond 1993) I
- Salt Shrub (Schafale and Weakley 1990) I
- Tidal Marsh (FNAI 1990) ?

Alliance Description

Environment: This alliance occurs along upper marsh edges and in tidal flats ranging in scale from narrow bands to hundreds of hectares. It is found along mainland shores and the bayside of barrier islands.

Vegetation: This alliance is dominated by *Borrichia frutescens* (bushy seaside tansy), often in monospecific stands. Other characteristic species may include *Spartina patens* (saltmeadow cordgrass), *Spartina spartinae* (gulf cordgrass), *Sporobolus virginicus* (seashore dropseed), *Monanthochloe littoralis* (shoregrass), *Iva frutescens* ssp. *frutescens* (Jesuit's bark), *Iva frutescens* ssp. *oraria* (Jesuit's bark), *Limonium carolinianum* (Carolina sealavender), *Sarcocornia perennis* (chickenclaws), *Lycium carolinianum* var. *quadrifidum* (Carolina desert-thorn), *Rayjacksonia phyllocephala* (camphor daisy), and *Batis maritima* (turtleweed).

Alliance Distribution

Range: This alliance is found throughout the southeastern United States.

Nations: US

Subnations: AL, FL, GA, LA?, MS, NC, SC, TX, VA?

TNC Ecoregions: 31:C, 53:C, 55:C, 56:C, 57:C, 58:C

USFS Ecoregions: 231Fb:CCC, 232Cb:CC?, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC, 232Dc:CCC, 232Eb:CCC, 232Gb:CCC, 255Dc:CCC

Federal Lands: NPS (Canaveral, Cape Lookout, Fort Pulaski, Timucuan?); USFS (Croatan); USFWS (Anahuac, Aransas, Back Bay?, Big Boggy, Brazoria, Laguna Atascosa, Matagorda Island, McFaddin, San Bernard, Texas Point)

(CEGL003924) Seaside Oxeye Tidal Shrub Flat

***Borrichia frutescens* / (*Spartina patens*, *Juncus roemerianus*) Shrubland**

Seaside Oxeye/(Saltmeadow Cordgrass, Black Needlerush) Shrubland

NVC Classification

Physiognomic Class	Shrubland (III)
Physiognomic Subclass	Deciduous shrubland (III.B.)
Physiognomic Group	Cold-deciduous shrubland (III.B.2.)
Physiognomic Subgroup	Natural/Semi-natural cold-deciduous shrubland (III.B.2.N.)
Formation	Tidal cold-deciduous shrubland (III.B.2.N.h.)
Alliance	<i>Borrichia frutescens</i> Tidal Shrubland Alliance (A.1026)
Alliance (English name)	Seaside Oxeye Tidal Shrubland Alliance
Association	<i>Borrichia frutescens</i> / (<i>Spartina patens</i> , <i>Juncus roemerianus</i>) Shrubland
Association (English name)	Seaside Oxeye / (Saltmeadow Cordgrass, Black Needlerush) Shrubland
Association (Common name)	Seaside Oxeye Tidal Shrub Flat
Ecological System(s):	Central Atlantic Coastal Plain Fresh and Oligohaline Tidal Marsh (CES203.376)
Ecological System(s):	Central Atlantic Coastal Plain Salt and Brackish Tidal Marsh (CES203.270)
	Atlantic Coastal Plain Embayed Region Tidal Salt and Brackish Marsh (CES203.260)
	Central and Upper Texas Coast Salt and Brackish Tidal Marsh (CES203.473)
	Gulf Coast Chenier Plain Salt and Brackish Tidal Marsh (CES203.468)

Element Concept

Global Summary: This association includes upper marsh edges and flats strongly dominated by *Borrichia frutescens* (bushy seaside tansy), ranging in size from narrow bands to flats in excess of 500 ha. It is broadly described to include vegetation ranging from the Mid- and South Atlantic Coast to the West Gulf Coast of the United States. It may occur just below *Iva frutescens* (Jesuit's bark) or *Baccharis halimifolia* (eastern baccharis) shrublands and just above marshes dominated by *Juncus roemerianus* (needlegrass rush) or *Spartina alterniflora* (smooth cordgrass). Other characteristic species may include *Spartina patens* (saltmeadow cordgrass), *Iva frutescens* (Jesuit's bark), and *Limonium carolinianum* (Carolina sealavender). In Texas, the composition of this low-diversity, irregularly tidally flooded shrubland is variable and can include *Distichlis spicata* (inland saltgrass), *Schoenoplectus robustus* (sturdy bulrush), *Sporobolus virginicus* (seashore dropseed), *Monanthochloe littoralis* (shoregrass), *Atriplex matamorenensis* (Matamoros saltbush) (in south Texas), *Sarcocornia pacifica* (Pacific swampfire), *Limonium carolinianum* (Carolina sealavender), *Lycium carolinianum* var. *quadrifidum* (Carolina desert-thorn), *Rayjacksonia phyllocephala* (camphor daisy), *Prosopis reptans* (tornillo) (in south Texas), and *Batis maritima* (turtleweed). Here, this community often occurs just above the lower tidal flat communities *Batis maritima* - *Sarcocornia pacifica* Dwarf-shrubland (CEGL003956) or *Sarcocornia pacifica* - (*Batis maritima*, *Distichlis spicata*) Dwarf-shrubland (CEGL002278).

This shrubland usually occurs as a very narrow band adjacent and below tidal shrublands. Two large examples of this association at Fort Pulaski, Georgia, are extensive enough to map and occur on tidal flats adjacent the extensive salt marsh communities of eastern Cockspur Island. Typically this community is monospecific, being made up almost exclusively of *Borrchia frutescens* (bushy seaside tansy).

Element Description

USFWS Wetland System: Estuarine

Fort Pulaski National Monument Environment: This brackish marsh shrubland was sampled at two locations in the park. The sites are flat to gently sloping, southeast- and south-facing tidal flats. Soils are moderately somewhat poorly drained clay loam and muck. The unvegetated surface is mostly made up of bare soil (89-95% cover), with some leaf litter (1-10% cover), wood (0-3% cover) and debris (1% cover). Evidence of disturbance includes deer trails. This community usually occurs as a very narrow band adjacent to and below Coastal Salt Shrub Thicket (CEGL003920) tidal shrublands on tidal flats adjacent the extensive salt marsh communities such as on eastern Cockspur Island (Govus 1998).

Global Environment: This association occurs along upper marsh edges and in tidal flats ranging in scale from narrow bands to hundreds of hectares. It is found along mainland shores and the bayside of barrier islands. In south Texas, it can also occur around the base of lomas.

Environmental Description

USFWS Wetland System: Estuarine

Fort Pulaski National Monument Environment: This brackish marsh shrubland was sampled at two locations in the park. The sites are flat to gently sloping, southeast- and south-facing tidal flats. Soils are moderately somewhat poorly drained clay loam and muck. The unvegetated surface is mostly made up of bare soil (89-95% cover), with some leaf litter (1-10% cover), wood (0-3% cover) and debris (1% cover). Evidence of disturbance includes deer trails. This community usually occurs as a very narrow band adjacent to and below Coastal Salt Shrub Thicket (CEGL003920) tidal shrublands on tidal flats adjacent the extensive salt marsh communities such as on eastern Cockspur Island (Govus 1998).

Global Environment: This association occurs along upper marsh edges and in tidal flats ranging in scale from narrow bands to hundreds of hectares. It is found along mainland shores and the bayside of barrier islands. In south Texas, it can also occur around the base of lomas.

Vegetation Description

Fort Pulaski National Monument Vegetation: This community is dominated (30-80% cover) by short shrubs (0.5-1 m). However, there may also contains a sparse (20% cover) emergent tall shrub layer (2-5 m) of *Juniperus virginiana* var. *silicicola* (southern redcedar), *Ilex vomitoria* (yaupon), *Iva frutescens* (Jesuit's bark), and *Sideroxylon tenax* (tough bully) (globally and state-vulnerable). The short shrub layer is dominated by *Borrchia frutescens* (bushy seaside tansy) and may include *Sarcocornia pacifica* (Pacific swampfire), *Pinus elliottii* (slash pine), *Sabal palmetto* (cabbage palmetto), and *Yucca aloifolia* (aloe yucca). The moderately sparse to dense (40-80% cover) herbaceous layer is dominated almost exclusively by *Distichlis spicata* (inland

saltgrass). Some occurrences of this community are monospecific, being made up almost exclusively of *Borrichia frutescens* (bushy seaside tansy) (Govus 1998).

Global Vegetation: This community is strongly dominated by *Borrichia frutescens* (bushy seaside tansy). Other characteristic species may include *Spartina patens* (saltmeadow cordgrass), *Iva frutescens* (Jesuit's bark), *Distichlis spicata* (inland saltgrass), *Schoenoplectus robustus* (sturdy bulrush), *Sporobolus virginicus* (seashore dropseed), *Monanthochloe littoralis* (shoregrass), *Sarcocornia pacifica* (Pacific swampfire), *Limonium carolinianum* (Carolina sealavender), *Lycium carolinianum* var. *quadrifidum* (Carolina desert-thorn), *Rayjacksonia phyllocephala* (camphor daisy), and *Batis maritima* (turtleweed). In Texas, *Spartina patens* (saltmeadow cordgrass) is a less important component of this community, but it often forms a fringe along the upper margin.

Most Abundant Species

Fort Pulaski National Monument

<u>Stratum</u>	<u>Lifeform</u>	<u>Species</u>
Tall shrub/sapling	Needle-leaved shrub	<i>Juniperus virginiana</i> var. <i>silicicola</i> (southern redcedar)
Short shrub/sapling	Semi-shrub	<i>Borrichia frutescens</i> (bushy seaside tansy)
Herb (field)	Semi-shrub	<i>Sarcocornia pacifica</i> (Pacific swampfire)
Herb (field)	Graminoid	<i>Distichlis spicata</i> (inland saltgrass)

Characteristic Species

Fort Pulaski National Monument: *Borrichia frutescens* (bushy seaside tansy), *Distichlis spicata* (inland saltgrass), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Sarcocornia pacifica* (Pacific swampfire)

Other Noteworthy Species

Fort Pulaski National Monument

<u>Other Plant Species</u>	<u>GRank</u>	<u>Note</u>
<i>Sideroxylon tenax</i> (tough bully)	G3?	globally vulnerable

Global

<u>Animal Species</u>	<u>GRank</u>	<u>Note</u>
<i>Grus americana</i> (whooping crane)	G1	

Conservation Status Rank

Global Rank & Reasons: G4 (10-Oct-1997). This is a common tidal flat community along the Gulf and Middle and South Atlantic coasts of the United States.

Classification

Status: Standard

Classification Confidence: 1 - Strong

Global Comments: The relationship between this community and similar *Borrichia frutescens* / *Spartina spartinae* Shrubland (CEGL004617) is poorly documented. The ranges of these two communities overlap along the middle and south Texas coast where *Borrichia frutescens* / *Spartina spartinae* Shrubland (CEGL004617) dominates less saline, higher ground than this community (CEGL003924).

NatureServe identified the perennial glasswort plants at Fort Pulaski as *Sarcocornia pacifica* using the recent key in Weakley (2007), which follows Flora of North America (FNA Editorial Committee 2003). The perennial glassworts on the U.S. East Coast have been called *Sarcocornia perennis* in the past (Kartesz 1999), and their taxonomy remains unsettled. For the time being, NatureServe will continue to follow the Kartesz (1999) nomenclature in our global descriptions and U.S. National Vegetation Classification association names.

Global Similar Associations:

- *Borrchia frutescens* / *Spartina spartinae* Shrubland (CEGL004617)

Global Related Concepts:

- Glasswort-Saltwort Series (Diamond 1993) B
- Salt Shrub (Low Subtype) (Schafale 2000) ?

Element Distribution

Fort Pulaski National Monument Range: Two large examples of this association are extensive enough to map and occur on tidal flats adjacent to the extensive salt marsh communities of eastern Cockspur Island [see 1998 vegetation map (Govus 1998)].

Global Range: This association is found in coastal areas from North Carolina to Texas.

Nations: US

States/Provinces: AL, FL, GA, LA?, MS, NC, SC, TX, VA?

TNC Ecoregions: 31:C, 53:C, 55:C, 56:C, 57:C

USFS Ecoregions: 231Fb:CCP, 232Cb:CC?, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC, 232Dc:CCC, 232Eb:CCC, 232Gb:CCC, 255Dc:CPP

Federal Lands: NPS (Canaveral, Cape Lookout, Fort Pulaski, Timucuan?); USFS (Croatan); USFWS (Anahuac, Aransas, Back Bay?, Big Boggy, Brazoria, Laguna Atascosa, Matagorda Island, McFaddin, San Bernard, Texas Point)

Element Sources

Fort Pulaski National Monument Plots: FOPU.2, FOPU.6.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: A.S. Weakley

References: Diamond 1993, FNA Editorial Committee 2003, FNAI 1990, FNAI 1992a, Govus 1998, Kartesz 1999, McAlister and McAlister 1993, McAlister and McAlister 1995, Peet et al. unpubl. data 2002, Schafale 2000, Schafale 2003b, Schafale and Weakley 1990, Schotz pers. comm., Southeastern Ecology Working Group n.d., Weakley 2007

Dwarf-Shrubland (IV.)

Tidal needle-leaved or microphyllous evergreen dwarf - shrubland (IV.A.2.N.e)

***Batis maritima* Tidal Dwarf-shrubland Alliance (A.1111)**

Alliance Concept

Summary: This alliance consists of small to large tidal flats dominated by *Batis maritima* (turtleweed), typically with saltwater-tidal irregularly flooded hydrology. Other species can include *Sarcocornia perennis* (chickenclaws), *Sporobolus virginicus* (seashore dropseed), *Borrchia frutescens* (bushy seaside tansy), *Suaeda linearis* (annual seepweed), *Lycium carolinianum* (Carolina desert-thorn), *Spartina spartinae* (gulf cordgrass), *Monanthochloe littoralis* (shoregrass), *Sesuvium portulacastrum* (shoreline seapurslane), and *Blutaparon vermiculare* (silverhead). This alliance occurs especially in salt panne situations, where tidal saltwater evaporates, augmenting salinity. This is a subtropical alliance, which ranges northwards into the warmest portions of the temperate southeastern United States (e.g., Florida and Texas). This community may occur in areas up to at least 100 hectares.

Similar Alliances:

- *Sarcocornia pacifica* - (*Distichlis spicata*, *Spartina alterniflora*) Tidal Dwarf-shrubland Alliance (A.1705)--ranges from the Gulf Coast to the northern Atlantic Coast of the United States, and though *Batis maritima* is a component in this alliance, it would only dominate small patches within occurrences.

Similar Alliance Comments: This is a subtropical alliance, dominated by *Batis maritima*, which ranges northwards into the warmest portions of the temperate southeastern United States. It is not regarded as occurring in the South Atlantic Coastal Plain. It is best developed in the United States in Florida and the south-central coast of Texas and may be distinguished from the similar alliance (A.1705) by the dominance of *Batis maritima*. These two alliances can be difficult to distinguish along the upper Texas coast where their ranges overlap.

Related Concepts:

- *Batidion maritimae* Alliance (Borhidi 1991) ?
- *Batis-Salicornia-Sesuvium* (Olmsted et al. 1981) ?
- Glasswort-Saltwort Series (Diamond 1993) I
- Saltwort Scrub -- 11.4 (Dansereau 1966) ?

Alliance Description

Environment: This alliance occurs in lower tidal flats and low areas of higher tidal flats, within the reach of regular high tides. It is regularly to irregularly flooded by shallow brackish waters. As these waters evaporate, high concentrations of salt accumulate, producing hypersaline conditions.

Vegetation: This alliance is dominated by *Batis maritima* (turtleweed) sometimes in association with *Sarcocornia perennis* (chickenclaws), *Sporobolus virginicus* (seashore dropseed), *Borrchia frutescens* (bushy seaside tansy), *Suaeda linearis* (annual seepweed), *Lycium carolinianum* (Carolina desert-thorn), *Spartina spartinae* (gulf cordgrass), *Sesuvium portulacastrum* (shoreline seapurslane), and *Blutaparon vermiculare* (silverhead). Algal mats of blue-green and sometimes green algae are characteristically present, visible even in densely vegetated pannes.

Alliance Distribution

Range: This is a subtropical alliance, which ranges northwards into the warmest portions of the temperate southeastern United States. This alliance is found in Florida, Texas, and possibly in Alabama, Louisiana, Mississippi, and the Mexican state of Tamaulipas. It is also found in the Caribbean in Puerto Rico, the Bahamas, Cuba, Jamaica, Martinique, and elsewhere in the West Indies and other tropical countries.

Nations: BS, CU, JM, MQ, MX, PR, US, XB

Subnations: AL?, FL, GA, LA?, MS?, MXTM?, TX

TNC Ecoregions: 30:C, 31:C, 53:C, 54:C, 55:C, 56:C

USFS Ecoregions: 231Fb:CCC, 232Ce:CCC, 232Da:CCC, 232Dc:CCC, 232De:CCC, 232Ed:CCC, 232Ee:CCC, 232Gb:CCC, 255Dc:CCC, 315F:CC, 411Ae:CCC, 411Af:CCC, 411Ag:CCC, M411A:CC

Federal Lands: DOD (Guantanamo Bay); NPS (Canaveral, Everglades, Fort Matanzas, Fort Pulaski, Timucuan?); USFWS (Aransas, Big Boggy, Brazoria, Laguna Atascosa, Matagorda Island, San Bernard)

(CEGL003956) Saltwort - Swampfire Dwarf-shrubland

***Batis maritima* - *Sarcocornia pacifica* Dwarf-shrubland**

NVC Classification

Physiognomic Class	Dwarf-shrubland (IV)
Physiognomic Subclass	Evergreen dwarf-shrubland (IV.A.)
Physiognomic Group	Extremely xeromorphic evergreen dwarf-shrubland (IV.A.2.)
Physiognomic Subgroup	Natural/Semi-natural extremely xeromorphic evergreen dwarf-shrubland (IV.A.2.N.)
Formation	Tidal needle-leaved or microphyllous evergreen dwarf-shrubland (IV.A.2.N.c.)
Alliance	<i>Batis maritima</i> Tidal Dwarf-shrubland Alliance (A.1111)
Alliance (English name)	Saltwort Tidal Dwarf-shrubland Alliance; Planta de Sal Dwarf-shrubland Alliance
Association	<i>Batis maritima</i> - <i>Sarcocornia pacifica</i> Dwarf-shrubland
Association (English name)	Saltwort - Swampfire Dwarf-shrubland
Ecological System(s):	South Texas Salt and Brackish Tidal Flat (CES301.461) Central and Upper Texas Coast Salt and Brackish Tidal Marsh (CES203.473) Florida Big Bend Salt and Brackish Tidal Marsh (CES203.508)

Element Concept

Global Summary: This association includes regularly to irregularly flooded tidal areas dominated (often with 95% or more cover) by *Batis maritima* (turtleweed). Other species can include *Sarcocornia pacifica* (Pacific swampfire), *Sporobolus virginicus* (seashore dropseed), *Borrhichia frutescens* (bushy seaside tansy), *Suaeda* (seepweed) sp., *Lycium carolinianum* (Carolina desert-thorn), *Spartina spartinae* (gulf cordgrass), *Monanthochloe littoralis* (shoregrass), *Sesuvium portulacastrum* (shoreline seapurslane), and *Blutaparon vermiculare* (silverhead). This community generally lacks trees, though scattered individuals of *Avicennia germinans* (black mangrove) can occur in Florida and southern Texas. It may occur as a narrow band or occupy areas up to at least 100 hectares. This is generally a subtropical community, which also ranges northwards into the warmest portions of the temperate southeastern United States.

Environmental Description

USFWS Wetland System: Estuarine

Fort Pulaski National Monument Environment: This saltwort dwarf-shrubland was sampled at one location in the park. The site is a hypersaline tidal flat with very poorly drained muck soil. The unvegetated surface is made up entirely of bare soil (100% cover), wood (2-5% cover) and bare soil (45-50% cover). These flats are only flooded during high tides.

Global Environment: This association occurs in lower tidal flats and low areas of higher tidal flats, within the reach of regular high tides. It is regularly to irregularly flooded by shallow brackish waters. As these waters evaporate, high concentrations of salt accumulate, producing hypersaline conditions.

Vegetation Description

Fort Pulaski National Monument Vegetation: The moderate (40% cover) dwarf-shrub layer (0.5-1 m tall) is dominated by *Batis maritima* (turtleweed) and additionally only includes lesser amounts of *Borrchia frutescens* (bushy seaside tansy) and *Sarcocornia pacifica* (Pacific swampfire). Other halophytes found here include *Suaeda linearis* (annual seepweed), *Sesuvium portulacastrum* (shoreline seapurslane), and *Limonium carolinianum* (Carolina sealavender) (Govus 1998).

Global Vegetation: This community is dominated by *Batis maritima* (turtleweed) sometimes in association with *Sarcocornia pacifica* (Pacific swampfire), *Sporobolus virginicus* (seashore dropseed), *Borrchia frutescens* (bushy seaside tansy), *Suaeda linearis* (annual seepweed), *Suaeda conferta* (beach seepweed) (in southern Texas), *Lycium carolinianum* (Carolina desert-thorn), *Spartina spartinae* (gulf cordgrass), *Sesuvium portulacastrum* (shoreline seapurslane), and *Blutaparum vermiculare* (silverhead). Algal mats of blue-green and sometimes green algae are characteristically present, visible even in densely vegetated pannes.

Most Abundant Species

Fort Pulaski National Monument

<u>Stratum</u>	<u>Lifeform</u>	<u>Species</u>
Herb (field)	Semi-shrub	<i>Batis maritima</i> (turtleweed)

Characteristic Species

Fort Pulaski National Monument: *Batis maritima* (turtleweed), *Borrchia frutescens* (bushy seaside tansy), *Sarcocornia pacifica* (Pacific swampfire)

Other Noteworthy Species

Global

<u>Animal Species</u>	<u>GRank</u>	<u>Note</u>
<i>Grus americana</i> (whooping crane)	G1	

Conservation Status Rank

Global Rank & Reasons: G5 (15-May-1995). This is a common tidal flat community along the Gulf Coast of Texas and Florida. It may occur as a narrow band or occupy areas up to at least 100 hectares.

Classification

Status: Standard

Classification Confidence: 2 - Moderate

Fort Pulaski National Monument Comments: NatureServe identified the perennial glasswort plants at Fort Pulaski as *Sarcocornia pacifica* using the recent key in Weakley (2007), which follows FNA Editorial Committee 2003). The perennial glassworts on the U.S. East Coast have been called *Sarcocornia perennis* in the past (Kartesz 1999), and their taxonomy remains unsettled. For the time being, NatureServe will continue to follow the Kartesz (1999) nomenclature in our global descriptions and U.S. National Vegetation Classification association names.

Global Comments: This association is currently known only from Florida, Georgia and the southern coast of Texas and may be distinguished from *Sarcocornia pacifica* - (*Batis maritima*, *Distichlis spicata*) Dwarf-shrubland (CEGL002278) by the dominance of *Batis maritima*. *Sarcocornia pacifica* - (*Batis maritima*, *Distichlis spicata*) Dwarf-shrubland (CEGL002278) ranges along the Gulf Coast of the United States, and though *Batis maritima* is a component in this association, it is rarely dominant, and then only at small scales. These two associations can be difficult to distinguish along the middle and upper Texas coast where their ranges overlap.

NatureServe identified the perennial glasswort plants at Fort Pulaski as *Sarcocornia pacifica* using the recent key in Weakley (2007), which follows FNA Editorial Committee 2003). The perennial glassworts on the U.S. East Coast have been called *Sarcocornia perennis* in the past (Kartesz 1999), and their taxonomy remains unsettled. For the time being, NatureServe will continue to follow the Kartesz (1999) nomenclature in our global descriptions and U.S. National Vegetation Classification association names.

Global Similar Associations:

- *Salicornia* (*virginica*, *bigelovii*, *maritima*) - *Spartina alterniflora* Herbaceous Vegetation (CEGL004308)
- *Sarcocornia pacifica* - (*Batis maritima*, *Distichlis spicata*) Dwarf-shrubland (CEGL002278)

Global Related Concepts:

- Glasswort-Saltwort Series (Diamond 1993) B

Element Distribution

Fort Pulaski National Monument Range: This community occurs beyond the dike system south of the fort. At least two large mappable examples of this association occur on hypersaline flats that grade into other salt panne communities along eastern and southeastern Cockspar Island (Govus 1998).

Global Range: This association is currently only known from Texas, Georgia and Florida. It may also occur in Alabama, Mississippi, and Louisiana, and the Mexican state of Tamaulipas. It does not extend to South Carolina.

Nations: MX?, US

States/Provinces: AL?, FL, GA, LA?, MS?, MXTM?, TX

TNC Ecoregions: 30:C, 31:C, 53:C, 54:C, 55:C, 56:C

USFS Ecoregions: 231Fb:CCC, 232Ce:CCC, 232Da:CCC, 232Dc:CCC, 232De:CCC, 232Ed:CCC, 232Ee:CCC, 232Gb:CCC, 255Dc:CCC, 315F:CC, 411Ae:CCC, 411Af:CCC, 411Ag:CCC, M411:C

Federal Lands: NPS (Canaveral, Everglades, Fort Matanzas, Fort Pulaski, Timucuan?); USFWS (Aransas, Big Boggy, Brazoria, Laguna Atascosa, Matagorda Island, San Bernard)

Element Sources

Fort Pulaski National Monument Plots: FOPU.17.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: A.S. Weakley and K.D. Patterson

References: Diamond 1993, FNA Editorial Committee 2003, FNAI 1992a, Govus 1998, Johnston 1952, Kartesz 1999, McAlister and McAlister 1993, McAlister and McAlister 1995, Schotz pers. comm., Southeastern Ecology Working Group n.d., Weakley 2007

***Sarcocornia pacifica* – (*Distichlis spicata*, *Spartina alterniflora*) Tidal Dwarf-shrubland Alliance (A.1705)**

Alliance Concept

Summary: Tidally flooded hypersaline flats or very shallow depressions, dominated by halophytic herbs, including *Sarcocornia perennis* (chickenclaws), *Salicornia bigelovii* (dwarf saltwort), *Salicornia virginica* (Virginia glasswort), *Distichlis spicata* (inland saltgrass), and stunted *Spartina alterniflora* (smooth cordgrass). Total vegetative cover is quite variable in pannes, from near total absence of vascular plants to a dense cover of *Salicornia bigelovii* (dwarf saltwort), *Salicornia virginica* (Virginia glasswort), *Sarcocornia perennis* (chickenclaws), or *Spartina alterniflora* (smooth cordgrass). *Limonium carolinianum* (Carolina sealavender) is another common associate. In the southern portions of its distribution, *Monanthochloe littoralis* (shoregrass) and *Batis maritima* (turtleweed) can be major components. Algal mats are characteristically present, visible even in densely vegetated pannes. Blue-green algae are an important component of this community, in some cases contributing significantly more biomass than do vascular species. The following algae were noted to occur in association with *Spartina alterniflora* (smooth cordgrass) in the littoral zone of a Massachusetts salt marsh: *Oscillatoria subuliformis*, *Oscillatoria amphibia*, *Lyngbea* spp., *Microcoleus chthonoplastes*, *Nodularia harveyana*, *Hydrocoleum lyngbyaceum*, *Symploca* spp. (Webber 1967). Diagnostic species are *Salicornia bigelovii* (dwarf saltwort), *Salicornia virginica* (Virginia glasswort), and *Sarcocornia perennis* (chickenclaws). Vegetation of this alliance tends to develop in shallow depressions within high salt marshes where drainage is poor. The depressions are flooded by high tides but as the water evaporates during low tide the salinity concentration increases forming "salt pannes." Formation of the pannes may result from ice scouring, rafting flotsam, peat compaction, or by mosquito ditch levees which create small impoundments. This community is regularly to irregularly flooded by nearby brackish water. Bare peat and/or mucky soils are prevalent (up to 85% bare soils), and standing water covers this community at high tide.

Similar Alliances:

- *Batis maritima* Tidal Dwarf-shrubland Alliance (A.1111)

Similar Alliance Comments: This is a temperate alliance, dominated by the perennial glasswort *Sarcocornia perennis*. It is best developed in the United States in the Gulf and upper to mid-Atlantic coasts and may be distinguished from the similar alliance (A.1111) by the lack of dominance of *Batis maritima*. These two alliances can be difficult to distinguish along the upper Texas coast where their ranges overlap.

Related Concepts:

- Glasswort-Saltwort Series (Diamond 1993) ?
- Salt Flat (Wieland 1994b) ?
- Salt Flat (Nelson 1986) ?
- Salt Flat (Schafale and Weakley 1990) ?

Alliance Description

Environment: Tidally flooded hypersaline flats or very shallow depressions, dominated by halophytic herbs. Vegetation of this alliance tends to develop in shallow depressions within high salt marshes where drainage is poor. The depressions are flooded by high tides but as the water evaporates during low tide the salinity concentration increases forming "salt pannes." Formation of the pannes may result from ice scouring, rafting flotsam, peat compaction, or by mosquito ditch levees which create small impoundments. This community is regularly to irregularly flooded by nearby brackish water. Bare peat and/or mucky soils are prevalent (up to 85% bare soils), and standing water covers this community at high tide.

Vegetation: Tidally flooded hypersaline flats or very shallow depressions, dominated by halophytic herbs, including *Sarcocornia perennis* (chickenclaws), *Salicornia bigelovii* (dwarf saltwort), *Salicornia virginica* (Virginia glasswort), *Distichlis spicata* (inland saltgrass), and stunted *Spartina alterniflora* (smooth cordgrass). Total vegetative cover is quite variable in pannes, from near total absence of vascular plants to a dense cover of *Salicornia bigelovii* (dwarf saltwort), *Salicornia virginica* (Virginia glasswort), *Sarcocornia perennis* (chickenclaws), or *Spartina alterniflora* (smooth cordgrass). *Limonium carolinianum* (Carolina sealavender) is another common associate. In the southern portions of its distribution, *Monanthochloe littoralis* (shoregrass) and *Batis maritima* (turtleweed) can be major components. Algal mats are characteristically present, visible even in densely vegetated pannes. Blue-green algae are an important component of this community, in some cases contributing significantly more biomass than do vascular species. The following algae were noted to occur in association with *Spartina alterniflora* (smooth cordgrass) in the littoral zone of a Massachusetts salt marsh: *Oscillatoria subuliformis*, *Oscillatoria amphibia*, *Lyngbea* spp., *Microcoleus chthonoplastes*, *Nodularia harveyana*, *Hydrocoleum lyngbyaceum*, *Symploca* spp. (Webber 1967). Diagnostic species are *Salicornia bigelovii* (dwarf saltwort), *Salicornia virginica* (Virginia glasswort), and *Sarcocornia perennis* (chickenclaws).

Alliance Distribution

Range: This alliance is found in the southern United States and adjacent Mexico, ranging from South Carolina and Florida, west to Texas and Tamaulipas, Mexico.

Nations: CA?, MX, US

Subnations: AL, FL, GA, LA, MS, MXTM, SC, TX

TNC Ecoregions: 30:P, 31:C, 41:P, 53:C, 54:P, 55:C, 56:C, 57:C, 58:C, 62:C, 63:C

USFS Ecoregions: 212C:PP, 212Db:PPP, 221Aa:CCC, 221Ab:CCC, 221Ac:CCC, 221Ad:CCC, 221Ae:CC?, 221Ak:CCC, 221B:CC, 221Dc:CPP, 231Fb:CCC, 232Aa:CCC, 232Ab:CCC, 232Ac:CCC, 232Bb:CCP, 232Bc:CCP, 232Bd:CCP, 232Bx:CC?, 232Bz:CCC, 232Cb:CC?, 232Ce:CCC, 232Ch:CCP, 232Ci:CCP, 232Db:CC?, 232Dc:CCC, 232Dd:CCC, 232De:CCC, 232Eb:CCC, 232Ed:CCC, 232Ee:CCC, 232Gb:CCC, 255Da:CCC, 255Dc:CCC, 315F:PP

Federal Lands: NPS (Acadia, Assateague Island, Canaveral, Fort Pulaski, Gulf Islands, Timucuan); USFWS (Anahuac, Aransas, Brazoria, Matagorda Island, McFaddin, San Bernard, Texas Point)

(CEGL002278) Salt Flat (Swampfire Type)

***Sarcocornia pacifica* - (*Batis maritima*, *Distichlis spicata*) Dwarf-shrubland**

Swampfire - (Saltwort, Saltgrass) Dwarf-shrubland

NVC Classification

Physiognomic Class	Dwarf-shrubland (IV)
Physiognomic Subclass	Evergreen dwarf-shrubland (IV.A.)
Physiognomic Group	Extremely xeromorphic evergreen dwarf-shrubland (IV.A.2.)
Physiognomic Subgroup	Natural/Semi-natural extremely xeromorphic evergreen dwarf-shrubland (IV.A.2.N.)
Formation	Tidal needle-leaved or microphyllous evergreen dwarf-shrubland (IV.A.2.N.c.)
Alliance	<i>Sarcocornia pacifica</i> - (<i>Distichlis spicata</i> , <i>Spartina alterniflora</i>) Tidal Dwarf-shrubland Alliance (A.1705)
Alliance (English name)	Swampfire - (Saltgrass, Saltmarsh Cordgrass) Tidal Dwarf-shrubland Alliance
Association	<i>Sarcocornia pacifica</i> - (<i>Batis maritima</i> , <i>Distichlis spicata</i>) Dwarf-shrubland
Association (English name)	Swampfire - (Saltwort, Saltgrass) Dwarf-shrubland
Association (Common name)	Salt Flat (Swampfire Type)
Ecological System(s):	Atlantic Coastal Plain Indian River Lagoon Tidal Marsh (CES203.257) Mississippi Delta Salt and Brackish Tidal Marsh (CES203.471) South Texas Salt and Brackish Tidal Flat (CES301.461) Central and Upper Texas Coast Salt and Brackish Tidal Marsh (CES203.473) Gulf Coast Chenier Plain Salt and Brackish Tidal Marsh (CES203.468)

Element Concept

Global Summary: This is one of several associations found in tidally influenced hypersaline areas along the southern Atlantic Coast and Gulf of Mexico region. In contrast to other tidal salt flats or pannes of this region, examples attributable to this type are dominated by the succulent dwarf-shrub *Sarcocornia pacifica* (Pacific swampfire) while stands dominated by *Batis* (turtleweed) or *Salicornia* (pickleweed) are accommodated by other associations [see *Batis maritima* - *Sarcocornia pacifica* Dwarf-shrubland (CEGL003956) and *Salicornia* (*virginica*, *bigelovii*, *maritima*) - *Spartina alterniflora* Herbaceous Vegetation (CEGL004308), respectively]. Examples of this association may occur in patches throughout this relatively large range. They vary somewhat locally in expression, but all tend to exhibit low vascular plant diversity, dominated by halophytic species. In addition to *Sarcocornia pacifica* (Pacific swampfire), other commonly encountered species may include *Salicornia bigelovii* (dwarf saltwort), *Distichlis spicata* (inland saltgrass), and sometimes stunted *Spartina alterniflora* (smooth cordgrass). Other typical species can include *Batis maritima* (turtleweed), *Suaeda* (seepweed) spp., *Sporobolus virginicus* (seashore dropseed), *Sesuvium portulacastrum* (shoreline seapurslane), and *Limonium carolinianum* (Carolina sealavender). Total vegetative cover is quite variable, from near total absence of vascular plants to a dense cover of the nominal species.

Algal mats are characteristically present, visible even in densely vegetated pannes. Blue-green algae may contribute significantly more biomass than vascular species.

Environmental Description

USFWS Wetland System: Estuarine

Fort Pulaski National Monument Environment: This woody glasswort dwarf-shrubland was sampled at one location in the park. The site is a tidal flat located on slightly higher ground than adjacent to cordgrass salt marsh. Soil is very poorly drained muck. The unvegetated surface is made up of equal parts leaf litter (50% cover) and bare soil (50%). This association represents the salt panne community. It is particularly well-developed on the broad gentle flats that lie along the south side of Cockspur Island. This is a hypersaline environment caused by the repeated evaporation of tidal water from these expansive shallow areas (Govus 1998).

Global Environment: Typically, this vegetation tends to develop in lower tidal flats and shallow depressions within upper tidal flats and high salt marshes where drainage is poor. These areas flood during high tides, and salt accumulates as the water evaporates during low tides, producing hypersaline conditions, forming salt pannes.

Vegetation Description

Fort Pulaski National Monument Vegetation: This sparse dwarf-shrubland is dominated (20% cover) by *Sarcocornia pacifica* (Pacific swampfire). The only other species present is *Spartina alterniflora* (smooth cordgrass). Vegetative cover varies from a total absence of vascular plants to a dense concentration of halophytic herbs, particularly *Sarcocornia pacifica* (Pacific swampfire), which is clearly dominant. *Batis maritima* (turtleweed) and *Distichlis spicata* (inland saltgrass) are also abundant and usually interspersed with dwarfed *Spartina alterniflora* (smooth cordgrass). Additional species found in this association include *Limonium carolinianum* (Carolina sealavender), *Suaeda linearis* (annual seepweed), and *Sesuvium portulacastrum* (shoreline seapurslane) (Govus 1998).

Global Vegetation: This community is dominated by the halophytic, succulent dwarf-shrub *Sarcocornia pacifica* (Pacific swampfire) and other halophytes, including *Salicornia bigelovii* (dwarf saltwort), *Distichlis spicata* (inland saltgrass), and sometimes stunted *Spartina alterniflora* (smooth cordgrass). Other typical species can include *Suaeda* (seepweed) spp., *Sporobolus virginicus* (seashore dropseed), *Sesuvium portulacastrum* (shoreline seapurslane), and *Limonium carolinianum* (Carolina sealavender). Algal mats are characteristically present, visible even in densely vegetated pannes. Blue-green algae may contribute significantly more biomass than vascular species. Texas examples, of alternately wet and dry saline soils along the Gulf Coast, may contain *Monanthochloe littoralis* (shoregrass), *Rayjacksonia phyllocephala* (camphor daisy), *Borrchia frutescens* (bushy seaside tansy), *Maytenus phyllanthoides* (Florida mayten), *Suaeda* (seepweed) spp., *Sesuvium portulacastrum* (shoreline seapurslane), and *Sporobolus virginicus* (seashore dropseed).

Most Abundant Species

Fort Pulaski National Monument

Stratum

Herb (field)

Lifeform

Semi-shrub

Species

Sarcocornia pacifica (Pacific swampfire)

Characteristic Species

Fort Pulaski National Monument: *Sarcocornia pacifica* (Pacific swampfire), *Spartina alterniflora* (smooth cordgrass)

Conservation Status Rank

Global Rank & Reasons: G4 (15-Dec-1994). This is a common tidal flat community along the Gulf and South Atlantic coasts of the United States.

Classification

Status: Standard

Classification Confidence: 2 - Moderate

Fort Pulaski National Monument Comments: NatureServe identified the perennial glasswort plants at Fort Pulaski as *Sarcocornia pacifica* using the recent key in Weakley (2007), which follows FNA Editorial Committee 2003). The perennial glassworts on the U.S. East Coast have been called *Sarcocornia perennis* in the past (Kartesz 1999), and their taxonomy remains unsettled. For the time being, NatureServe will continue to follow the Kartesz (1999) nomenclature in our global descriptions and U.S. National Vegetation Classification association names.

Global Comments: This association is dominated by *Sarcocornia pacifica* and may be distinguished from *Batis maritima* - *Sarcocornia pacifica* Dwarf-shrubland (CEGL003956) by the lack of dominance by *Batis maritima*. These two associations can be difficult to distinguish along the middle and upper Texas coast where their ranges overlap. A similar community, *Salicornia (virginica, bigelovii, maritima)* - *Spartina alterniflora* Herbaceous Vegetation (CEGL004308), occurs along the mid- and northern Atlantic Coast.

NatureServe identified the perennial glasswort plants at Fort Pulaski as *Sarcocornia pacifica* using the recent key in Weakley (2007), which follows FNA Editorial Committee 2003). The perennial glassworts on the U.S. East Coast have been called *Sarcocornia perennis* in the past (Kartesz 1999), and their taxonomy remains unsettled. For the time being, NatureServe will continue to follow the Kartesz (1999) nomenclature in our global descriptions and U.S. National Vegetation Classification association names.

Global Similar Associations:

- *Batis maritima* - *Sarcocornia pacifica* Dwarf-shrubland (CEGL003956)
- *Salicornia (virginica, bigelovii, maritima)* - *Spartina alterniflora* Herbaceous Vegetation (CEGL004308)--similar habitat with a more northerly range along the mid and northern Atlantic coasts.

Global Related Concepts:

- Glasswort-Saltwort Series (Diamond 1993) ?

Element Distribution

Fort Pulaski National Monument Range: This community is located northeast of the fort. It is widely represented on the south side of Cockspur Island and, to a lesser extent, also on McQueens Island (Govus 1998).

Global Range: This community ranges from the Atlantic Coast of South Carolina across the Gulf of Mexico region of the United States into Mexico. It ranges from South Carolina to the upper coast of Mexico.

Nations: MX, US

States/Provinces: AL:S1, FL, GA, LA, MS, MXTM, SC, TX

TNC Ecoregions: 31:C, 53:C, 55:C, 56:C, 57:?

USFS Ecoregions: 231Fb:CCC, 232Ce:CCC, 232Db:CC?, 232Dc:CCC, 232Dd:CCC, 232De:CCC, 232Eb:CCC, 232Ed:CCC, 232Ee:CCC, 232Gb:CCC, 255Da:CCC, 255Dc:CCC, 315F:PP

Federal Lands: NPS (Canaveral, Fort Pulaski, Gulf Islands, Timucuan); USFWS (Anahuac, Aransas, Brazoria, Matagorda Island, McFaddin, San Bernard, Texas Point)

Element Sources

Fort Pulaski National Monument Plots: FOPU.5.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: R.E. Evans

References: Diamond 1993, FNA Editorial Committee 2003, Govus 1998, Kartesz 1999, Peet et al. unpubl. data 2002, Schotz pers. comm., Southeastern Ecology Working Group n.d., Weakley 2007, Wieland 1994a, Wieland 1994b, Wieland 2000b

Herbaceous Vegetation (V.)

Tidal temperate or subpolar grassland (V.A.5.N.n)

***Juncus roemerianus* Tidal Herbaceous Alliance (A.1475)**

Alliance Concept

Summary: *Juncus roemerianus* (needlegrass rush)-dominated marshes with tidal hydrology. Associations in this alliance have a range of water chemistry, including saline, brackish, and (rarely) freshwater. This widely distributed salt marsh alliance occurs in association with low salt marshes or in brackish marshes, and is characterized by discrete, dense patches of vegetation usually strongly dominated by *Juncus roemerianus* (needlegrass rush) with few other associates. Associates which occur at low cover may include *Borrchia frutescens* (bushy seaside tansy), *Baccharis halimifolia* (eastern baccharis), *Spartina alterniflora* (smooth cordgrass), *Distichlis spicata* (inland saltgrass), *Schoenoplectus robustus* (sturdy bulrush), *Symphyotrichum tenuifolium* (perennial saltmarsh aster), and *Symphyotrichum subulatum* (eastern annual saltmarsh aster). This community may occur as isolated patches within low salt marsh, or may dominate vast areas at the heads of tidal creeks that drain the marsh. Its hydrology is generally irregularly tidally flooded. Soils of this association are generally poorly to very poorly drained, often with standing water atop peat accumulations (averaging 15 cm in depth) which in turn overlie gleyed sands. Degree of tidal inundation and relative elevation control the distribution of salt marsh vegetation in general; *Juncus roemerianus* (needlegrass rush) is typically found at lower elevation than the associated *Spartina patens* (saltmeadow cordgrass) marshes or mixed marshes. *Juncus roemerianus* (needlegrass rush) generally occurs on sandy substrates. The alliance may more rarely occur in freshwater, wind tidal situations. In these examples, *Juncus*

roemerianus (needlegrass rush) is the dominant, but other characteristic species may include *Thelypteris palustris* var. *pubescens* (eastern marsh fern), *Polygonum punctatum* (dotted smartweed), *Cladium mariscus* ssp. *jamaicense* (Jamaica swamp sawgrass), *Asclepias lanceolata* (fewflower milkweed), *Osmunda regalis* var. *spectabilis* (royal fern), *Eleocharis fallax* (creeping spikerush), *Boehmeria cylindrica* (smallspike false nettle), and others. This species composition is indicative of the freshwater marsh conditions, as they do not occur in more brackish and saline associations in the *Juncus roemerianus* (needlegrass rush) alliance.

Classification Comments: Few associations have been described; more may be necessary to accommodate variation in this wide-ranging alliance.

Similar Alliances:

- *Schoenoplectus americanus* Tidal Herbaceous Alliance (A.2007)
- *Spartina patens* - (*Distichlis spicata*) Tidal Herbaceous Alliance (A.1481)

Related Concepts:

- Brackish Marsh (Nelson 1986) I
- Brackish Marsh (Schafale and Weakley 1990) I
- Brackish Marsh (Wieland 1994b) I
- Brackish Marsh (Smith 1996a) I
- Brackish Marsh (Wieland 1994a) ?
- Smooth Cordgrass Series (Diamond 1993) I
- Tidal Marsh (FNAI 1992a) I

Alliance Description

Environment: *Juncus roemerianus*-dominated marshes with tidal hydrology. Associations in this alliance have a range of water chemistry, including saline, brackish, and (rarely) freshwater. This community may occur as isolated patches within low salt marsh, or may dominate vast areas at the heads of tidal creeks that drain the marsh. Its hydrology is generally irregularly tidally flooded. Soils of this association are generally poorly to very poorly drained, often with standing water atop peat accumulations (averaging 15 cm in depth) which in turn overlie gleyed sands. Degree of tidal inundation and relative elevation control the distribution of salt marsh vegetation in general; *Juncus roemerianus* is typically found at lower elevation than the associated *Spartina patens* marshes or mixed marshes. *Juncus roemerianus* generally occurs on sandy substrates. The alliance may more rarely occur in freshwater, wind tidal situations.

Vegetation: This widely distributed salt marsh alliance occurs in association with low salt marshes or in brackish marshes, and is characterized by discrete, dense patches of vegetation usually strongly dominated by *Juncus roemerianus* (needlegrass rush) with few other associates. Associates which occur at low cover may include *Borrchia frutescens* (bushy seaside tansy), *Baccharis halimifolia* (eastern baccharis), *Spartina alterniflora* (smooth cordgrass), *Distichlis spicata* (inland saltgrass), *Schoenoplectus robustus* (sturdy bulrush), *Symphyotrichum tenuifolium* (perennial saltmarsh aster), and *Symphyotrichum subulatum* (eastern annual saltmarsh aster). The alliance may more rarely occur in freshwater, wind tidal situations. In these examples, *Juncus roemerianus* (needlegrass rush) is the dominant, but other characteristic species may include *Thelypteris palustris* var. *pubescens* (eastern marsh fern), *Polygonum*

punctatum (dotted smartweed), *Cladium mariscus ssp. jamaicense* (Jamaica swamp sawgrass), *Asclepias lanceolata* (fewflower milkweed), *Osmunda regalis var. spectabilis* (royal fern), *Eleocharis fallax* (creeping spikerush), *Boehmeria cylindrica* (smallspike false nettle), and others. This species composition is indicative of the freshwater marsh conditions, as they do not occur in more brackish and saline associations in the *Juncus roemerianus* (needlegrass rush) alliance.

Alliance Distribution

Range: This alliance occurs in the southeastern Coastal Plain from Maryland south to Florida and west to Louisiana and Texas.

Nations: US

Subnations: AL, DE, FL, GA, LA?, MD, MS, NC, SC, TX, VA

TNC Ecoregions: 30:P, 31:C, 41:P, 42:P, 53:C, 54:C, 55:C, 56:C, 57:C, 58:C

USFS Ecoregions: 231Fb:CCC, 232Ad:CCC, 232Br:CCC, 232Bx:CCC, 232Bz:CCC, 232Cb:CCC, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC, 232Cj:CCC, 232Db:CCP, 232Dc:CCC, 232Dd:CCC, 232De:CCP, 232Eb:CCC, 232Ec:CC?, 232Ed:CCC, 232Gb:CCC, 255Da:CCP, 255Dc:CCC

Federal Lands: NPS (Assateague Island, Canaveral, Cape Hatteras, Cape Lookout, Fort Pulaski, Timucuan); USFS (Croatan); USFWS (Anahuac, Back Bay, Big Boggy, Bon Secour, Brazoria, Chesapeake Marshlands, McFaddin, San Bernard, Texas Point)

(CEGL004186) Needlerush High Marsh

***Juncus roemerianus* Herbaceous Vegetation**

Black Needlerush Herbaceous Vegetation

NVC Classification

Physiognomic Class	Herbaceous Vegetation (V)
Physiognomic Subclass	Perennial graminoid vegetation (V.A.)
Physiognomic Group	Temperate or subpolar grassland (V.A.5.)
Physiognomic Subgroup	Natural/Semi-natural temperate or subpolar grassland (V.A.5.N.)
Formation	Tidal temperate or subpolar grassland (V.A.5.N.n.)
Alliance	<i>Juncus roemerianus</i> Tidal Herbaceous Alliance (A.1475)
Alliance (English name)	Black Needlerush Tidal Herbaceous Alliance
Association	<i>Juncus roemerianus</i> Herbaceous Vegetation
Association (English name)	Black Needlerush Herbaceous Vegetation
Association (Common name)	Needlerush High Marsh
Ecological System(s):	Atlantic Coastal Plain Embayed Region Tidal Salt and Brackish Marsh (CES203.260) Atlantic Coastal Plain Indian River Lagoon Tidal Marsh (CES203.257) Florida Big Bend Salt and Brackish Tidal Marsh (CES203.508) Mississippi Delta Salt and Brackish Tidal Marsh (CES203.471) Northern Atlantic Coastal Plain Tidal Salt Marsh (CES203.519) Central Atlantic Coastal Plain Salt and Brackish Tidal Marsh (CES203.270) Mississippi Sound Salt and Brackish Tidal Marsh (CES203.303) Central and Upper Texas Coast Salt and Brackish Tidal Marsh (CES203.473)
Ecological System(s):	Gulf Coast Chenier Plain Salt and Brackish Tidal Marsh (CES203.468)

Element Concept

Global Summary: This broad-ranging *Juncus roemerianus* (needlegrass rush) salt marsh community is characterized by discrete, dense patches usually strongly dominated by *Juncus*

roemerianus (needlegrass rush), often with few other associates, or with low cover of *Distichlis spicata* (inland saltgrass), *Spartina alterniflora* (smooth cordgrass), *Spartina patens* (saltmeadow cordgrass), and *Limonium carolinianum* (Carolina sealavender). As currently defined, this community occurs in a variety of settings in different marsh regions including both "high" and "low" marshes. For example, large expanses of this type are found in northwest Florida at or below the mean high water line. In other regions it may be found as isolated patches within high salt marsh, or may dominate vast areas at the heads of tidal creeks. In general, the prevalence of *Juncus roemerianus* (needlegrass rush) in Florida indicates the prevalence of high marshes (above mean high water). Its hydrology is generally irregularly tidally flooded.

Environmental Description

USFWS Wetland System: Estuarine

Fort Pulaski National Monument Environment: This Needlerush High Marsh was sampled at one location in the park. The site is a tidally flooded marsh with very poorly drained muck soil. Examples of it are usually located near the low salt marsh fringe (Govus 1998). The unvegetated surface is made up of standing water (50% cover), leaf litter (10% cover) and bare soil (40% cover). Evidence of disturbance includes the presence of trails.

Global Environment: Large expanses of this type are found in northwest Florida, where approximately 60% of the salt marshes consist of monospecific stands of this type. These stands extend nearly to the water's edge and may be considered low marsh (Montague and Wiegert 1990). In other regions of Florida, including northeast Florida, Indian River (central Atlantic Coast), and south Florida, *Juncus roemerianus* occurs in smaller discrete patches of high marsh (Montague and Wiegert 1990). Presumably, this type also occurs primarily as high salt marsh along the Atlantic Coast as well (Hackney and de la Cruz 1982). In Mississippi *Juncus roemerianus* marsh was inundated less than 5.4% of the time that any portion of the marsh was flooded (Eleuterius and Eleuterius 1979). Due to longer and more frequent periods of exposure and resulting evaporation, interstitial water salinity may be higher than more frequently inundated zones. In addition, the interstitial water salinity was generally higher and more stable than flood water (Hackney and de la Cruz 1982). A range of soil conditions may be present in this broadly defined association, but soils are generally poorly to very poorly drained. In some areas, including tall-growth zones, soils were highly organic and peaty (averaging 15 cm in depth?), while shorter zones occurred on sandy soils underlain by clay (Eleuterius and Caldwell 1984). In general, *Juncus roemerianus* occurs on sandy substrates (Penfound 1952). pH at the Mississippi site was 6.2 (Hackney and de la Cruz 1982).

Vegetation Description

Fort Pulaski National Monument Vegetation: The dense (80% cover) herbaceous layer is dominated by *Juncus roemerianus* (needlegrass rush), with small amounts of *Spartina alterniflora* (smooth cordgrass). *Limonium carolinianum* (Carolina sealavender), *Distichlis spicata* (inland saltgrass), and *Symphyotrichum tenuifolium* (perennial saltmarsh aster) are also sometimes present (Govus 1998).

Global Vegetation: The very dense vegetation is typically exclusively codominated by *Juncus roemerianus* (needlegrass rush) and *Spartina alterniflora* (smooth cordgrass). Tall and short growth-form zones have been noted in several areas (Kruczynski et al. 1978, Hackney and de la

Cruz 1982, Stout 1984, Montague and Wiegert 1990). Associates that may occur at low cover include *Borrchia frutescens* (bushy seaside tansy), *Baccharis halimifolia* (eastern baccharis), *Spartina alterniflora* (smooth cordgrass), *Distichlis spicata* (inland saltgrass), *Schoenoplectus robustus* (sturdy bulrush), *Limonium carolinianum* (Carolina sealavender), *Symphyotrichum tenuifolium* (perennial saltmarsh aster), *Symphyotrichum subulatum* (eastern annual saltmarsh aster), and (in more southern occurrences) *Ipomoea sagittata* (saltmarsh morning-glory).

Most Abundant Species

Fort Pulaski National Monument

<u>Stratum</u>	<u>Lifform</u>	<u>Species</u>
Herb (field)	Graminoid	<i>Juncus roemerianus</i> (needlegrass rush)

Global

<u>Stratum</u>	<u>Lifform</u>	<u>Species</u>
Herb (field)	Graminoid	<i>Juncus roemerianus</i> (needlegrass rush)

Characteristic Species

Fort Pulaski National Monument: *Juncus roemerianus* (needlegrass rush), *Spartina alterniflora* (smooth cordgrass)

Global: *Juncus roemerianus* (needlegrass rush)

Other Noteworthy Species

Global

<u>Other Plant Species</u>	<u>GRank</u>	<u>Note</u>
<i>Ptilimnium nodosum</i> (piedmont mock bishopweed)	G2	Federally listed endangered; globally imperiled

Conservation Status Rank

Global Rank & Reasons: G5 (29-Mar-2001). This is a common coastal marsh of the Gulf and Atlantic coasts of the United States.

Classification

Status: Standard

Classification Confidence: 1 - Strong

Global Comments: This community is common on the southeastern seaboard, but large undisturbed areas are of high conservation concern. Although this community exhibits little floristic variation across its range, the associated animal species may vary to a greater extent. This community may not occur west of Texas. *Juncus roemerianus* was found to be lower in elevation than the associated *Spartina patens* type and mixed type (Cooper and Waits 1973).

Global Similar Associations:

- *Spartina alterniflora* - *Juncus roemerianus* - *Distichlis spicata* Louisianian Zone Salt Tidal Herbaceous Vegetation (CEGL004190)
- *Spartina patens* - *Distichlis spicata* - (*Juncus roemerianus*) Herbaceous Vegetation (CEGL004197)

Global Related Concepts:

- *Juncus roemerianus* Tidal Herbaceous Vegetation (Harrison 2001) =

- *Juncus roemerianus* Tidal Herbaceous Vegetation (VDNH 2003) =
- *Juncus roemerianus* Tidal Herbaceous Vegetation (Coulling 2002) =
- *Juncus roemerianus* association of the low marsh (Adams 1963) =
- *Juncus* type (Cooper and Waits 1973) =
- *Spartina* - *Distichlis* - *Juncus* associates (Penfound 1952) B
- Brackish Marsh (Wieland 1994b) B
- Brackish Marsh (Needlerush Subtype) (Schafale 2000) ?
- Irregularly flooded salt marsh (Jenkins 1974) =
- Lower high marsh (Stalter 1973a) =
- Needlerush - saltmeadow type (Nicholson and Van Deusen 1954) =
- Saline Marsh (Wieland 1994a) B
- Salt marsh (Higgins et al. 1971) B
- Salt marsh community (Hill 1986) B
- Smooth Cordgrass Series (Diamond 1993) B

Element Distribution

Fort Pulaski National Monument Range: This community is located along the rails to trails between the gate and first wooden bridge. This community is not widely developed at Fort Pulaski and does not occur in large enough patches to be mapped at a reasonable scale. Discrete examples are visible on both Cockspur and McQueens islands (Govus 1998).

Global Range: This community is widespread along the Atlantic and Gulf coasts of North America ranging from Delaware south to Florida, and west to Texas.

Nations: US

States/Provinces: AL:S2S3, DE:S2S3, FL, GA, LA?, MD:S4, MS, NC, SC, TX, VA:S5

TNC Ecoregions: 31:C, 53:C, 54:C, 55:C, 56:C, 57:C, 58:C

USFS Ecoregions: 231Fb:CCC, 232Ad:CCC, 232Br:CCC, 232Bx:CCC, 232Bz:CCC, 232Cb:CCC, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC, 232Cj:CCC, 232Db:CCP, 232Dc:CCC, 232Dd:CCC, 232De:CCP, 232Eb:CCC, 232Ec:CC?, 232Ed:CCC, 232Gb:CCC, 255Da:CCP, 255Dc:CCC

Federal Lands: NPS (Assateague Island, Canaveral, Cape Hatteras, Cape Lookout, Fort Pulaski, Timucuan); USFS (Croatan); USFWS (Anahuac, Big Boggy, Bon Secour, Brazoria, Chesapeake Marshlands, McFaddin, San Bernard, Texas Point)

Element Sources

Fort Pulaski National Monument Plots: FOPU.12.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: R.E. Evans, mod. L.A. Sneddon and M. Pyne

References: Adams 1963, Bowman 2000, Cooper and Waits 1973, Coulling 2002, Diamond 1993, Eleuterius and Caldwell 1984, Eleuterius and Eleuterius 1979, Fleming et al. 2001, Govus 1998, Hackney and de la Cruz 1981, Hackney and de la Cruz 1982, Harrison 2001, Harrison 2004, Higgins et al. 1971, Hill 1986, Jenkins 1974, Kruczynski et al. 1978, Lynch 1941, Montague and Wiegert 1990, Nelson 1986, Nicholson and Van Deusen 1954, Peet et al. unpubl. data 2002, Penfound 1952, Schafale 2000, Schafale 2003b, Schafale and Weakley 1990, Schotz

pers. comm., Smith 1996a, Southeastern Ecology Working Group n.d., Stalter 1973a, Stalter 1973b, Stout 1984, VDNH 2003, Wieland 1994a, Wieland 1994b, Wieland 2000b

***Spartina alterniflora* Tidal Herbaceous Alliance (A.1471)**

Alliance Concept

Summary: This alliance includes various tidal marshes dominated by *Spartina alterniflora* (smooth cordgrass). The hydrology is usually regularly tidally flooded. In the northern part of its range, southern Maine to Cape Hatteras, North Carolina, this alliance is generally limited to the zone between mean sea level and the mean high water level. The habitat occurs in protected inlets behind barrier beaches or in drowned river valleys. Peat depth ranges from a few feet, if the community formed over a mud flat, to 80 feet in drowned river valleys. *Spartina alterniflora* (smooth cordgrass) is limited to the low marsh zone by moderate salinity; it can withstand longer submergence than other salt marsh grasses, but still requires periodic exposure of the substrate. It also requires moderately high levels of iron (7-15 ppm). This community is commonly known as "low salt marsh," occurring as a tall grassland strongly dominated by *Spartina alterniflora* (smooth cordgrass). There is little variation in vascular plant species composition across the range. It occurs in nearly pure stands, with occasional low growing species such as *Spergularia salina* (salt sandspurry), *Salicornia* (pickleweed) spp., *Suaeda maritima* (herbaceous seepweed), and seaweeds such as *Ulva lactuca* and other algae such as *Fucus vesiculosus* and *Ascophyllum nodosum*, which grow at the bases of the *Spartina* (cordgrass) plants. Herbs of *Salicornia virginica* (Virginia glasswort) and *Salicornia bigelovii* (dwarf saltwort) can be quite common mixed in with the *Spartina* (cordgrass), often becoming more apparent later in the growing season. *Limonium carolinianum* (Carolina sealavender) is another characteristic herb, but only as scattered individuals. More detailed information is needed on the variability of the alliance in the southern parts of its range.

Similar Alliances:

- *Schoenoplectus americanus* Tidal Herbaceous Alliance (A.2007)
- *Spartina patens* - (*Distichlis spicata*) Tidal Herbaceous Alliance (A.1481)

Related Concepts:

- Brackish Marsh (Nelson 1986) I
- Estuarine Intertidal: Brackish Tidal Marsh (Swain and Kearsley 2001) ?
- Estuarine Intertidal: Salt Marsh (Swain and Kearsley 2001) ?
- Saline Marsh (Wieland 1994b) I
- Saline Marsh (Wieland 1994a) I
- Salt Marsh (Smith 1996a) ?
- Salt Marsh (Nelson 1986) ?
- Salt Marsh (Schafale and Weakley 1990) ?
- Smooth Cordgrass Marsh (Wharton 1978) ?
- Smooth Cordgrass Series (Diamond 1993) I
- Tidal Marsh (FNAI 1992a) I

Alliance Description

Environment: This alliance includes various tidal marshes dominated by *Spartina alterniflora*. The hydrology is usually regularly tidally flooded. In the northern part of its range, southern Maine to Cape Hatteras, North Carolina, this alliance is generally limited to the zone between mean sea level and the mean high water level. The habitat occurs in protected inlets behind barrier beaches or in drowned river valleys. Peat depth ranges from a few feet, if the community formed over a mud flat, to 80 feet in drowned river valleys. *Spartina alterniflora* is limited to the low marsh zone by moderate salinity; it can withstand longer submergence than other salt marsh grasses, but still requires periodic exposure of the substrate. It also requires moderately high levels of iron (7-15 ppm).

Vegetation: This community is commonly known as "low salt marsh," occurring as a tall grassland strongly dominated by *Spartina alterniflora* (smooth cordgrass). There is little variation in vascular plant species composition across the range. It occurs in nearly pure stands, with occasional low growing species such as *Spergularia salina* (salt sandspurry), *Salicornia* (pickleweed) spp., *Suaeda maritima* (herbaceous seepweed), and seaweeds such as *Ulva lactuca* and other algae such as *Fucus vesiculosus* and *Ascophyllum nodosum*, which grow at the bases of the *Spartina* (cordgrass) plants. Herbs of *Salicornia virginica* (Virginia glasswort) and *Salicornia bigelovii* (dwarf saltwort) can be quite common mixed in with the *Spartina* (cordgrass), often becoming more apparent later in the growing season. *Limonium carolinianum* (Carolina sealavender) is another characteristic herb, but only as scattered individuals.

Alliance Distribution

Range: This alliance is found in coastal states from Maine to Texas.

Nations: CA, US

Subnations: AL, CT, DE, FL, GA, LA, MA, MD, ME, MS, NB, NC, NH, NJ, NS, NY, RI, SC, TX, VA

TNC Ecoregions: 30:P, 31:C, 41:P, 53:C, 54:P, 55:P, 56:C, 57:C, 58:C, 62:C, 63:C

USFS Ecoregions: 212Cb:CCC, 212Db:CCC, 212Dc:CCC, 221Aa:CCP, 221Ab:CCC, 221Ac:CCC, 221Ad:CCC, 221Ae:CCC, 221Ak:CCC, 221Dc:CCP, 231Fb:CCC, 232Aa:CCC, 232Ab:CCC, 232Ac:CCC, 232Ad:CCC, 232Ae:CCC, 232Bb:CCP, 232Bc:CCP, 232Bd:CCP, 232Br:CCC, 232Bt:CCC, 232Bx:CCC, 232Bz:CCC, 232Cb:CCC, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC, 232Cj:CCC, 232Db:CCP, 232Dc:CCC, 232Dd:CCC, 232De:CCP, 232Eb:CCC, 232Ed:CCC, 232Ee:CCC, 232Gb:CCP, 255Da:CC?, 255Dc:CCC

Federal Lands: NPS (Acadia, Assateague Island, Boston Harbor Islands, Cape Cod, Cape Hatteras, Cape Lookout, Colonial, Fire Island, Fort Pulaski, Gateway, George Washington Birthplace, Muskeget Island, Sagamore Hill, Timucuan); USFS (Croatan); USFWS (Anahuac?, Aransas?, Back Bay?, Big Boggy, Bon Secour, Brazoria, Chesapeake Marshlands, Chincoteague, Matagorda Island, McFaddin, Monomoy, Moosehorn?, Parker River, Prime Hook, Rachel Carson, San Bernard, Texas Point)

(CEGL004191) Southern Atlantic Coast Salt Marsh
***Spartina alterniflora* Carolinian Zone Herbaceous Vegetation**
Saltmarsh Cordgrass Carolinian Zone Herbaceous Vegetation

NVC Classification

Physiognomic Class	Herbaceous Vegetation (V)
Physiognomic Subclass	Perennial graminoid vegetation (V.A.)
Physiognomic Group	Temperate or subpolar grassland (V.A.5.)
Physiognomic Subgroup	Natural/Semi-natural temperate or subpolar grassland (V.A.5.N.)
Formation	Tidal temperate or subpolar grassland (V.A.5.N.n.)
Alliance	<i>Spartina alterniflora</i> Tidal Herbaceous Alliance (A.1471)
Alliance (English name)	Saltmarsh Cordgrass Tidal Herbaceous Alliance
Association	<i>Spartina alterniflora</i> Carolinian Zone Herbaceous Vegetation
Association (English name)	Saltmarsh Cordgrass Carolinian Zone Herbaceous Vegetation
Association (Common name)	Southern Atlantic Coast Salt Marsh
Ecological System(s):	Atlantic Coastal Plain Embayed Region Tidal Salt and Brackish Marsh (CES203.260) Atlantic Coastal Plain Indian River Lagoon Tidal Marsh (CES203.257) Central Atlantic Coastal Plain Salt and Brackish Tidal Marsh (CES203.270)

Element Concept

Global Summary: This vegetation includes various tidal marshes dominated by *Spartina alterniflora* (smooth cordgrass), from Cape Hatteras, North Carolina, south to the Atlantic Coast of the Florida peninsula. The hydrology is usually regularly tidally flooded. *Spartina alterniflora* (smooth cordgrass) is limited to the low marsh zone by moderate salinity; it can withstand longer submergence than other salt marsh grasses, but still requires periodic exposure of the substrate. It also requires moderately high levels of iron (7-15 ppm). This community is commonly known as "low salt marsh," occurring as a tall grassland strongly dominated by *Spartina alterniflora* (smooth cordgrass). There is little variation in vascular plant species composition across the range. *Spartina alterniflora* (smooth cordgrass) occurs in nearly pure stands, occasionally with other low-statured species.

Environmental Description

USFWS Wetland System: Estuarine

Fort Pulaski National Monument Environment: This coastal salt marsh association was sampled at two locations but is the most abundant vegetation in the park. The sites are tidal with very poorly drained muck soil. The unvegetated surface is made up entirely of standing water (100% cover). These flats are in areas between mean high and low tides and are flooded twice a day. It also slowly grades into salt panne communities that include *Spartina alterniflora* along the extensive gentle flats along the southern edge of Cockspur Island.

Global Environment: This vegetation is found along the margins of sounds and estuaries, backs of barrier islands, and old flood tide deltas near closed inlets (Schafale and Weakley 1990). These are areas with diurnal salt tides. The hydrology is estuarine, tidally flooded, and euhaline (30-40 ppt). This range is consistent with the "Carolinian Province" of Cowardin et al. (1979).

Vegetation Description

Fort Pulaski National Monument Vegetation: The sparse to moderately dense (30-70% cover) herbaceous layer is dominated by *Spartina alterniflora* (smooth cordgrass). Other species present are *Distichlis spicata* (inland saltgrass), *Sarcocornia pacifica* (Pacific swampfire), and *Juncus roemerianus* (needlegrass rush).

Global Vegetation: This community is commonly known as "low salt marsh," occurring as tall grassland strongly dominated by *Spartina alterniflora* (smooth cordgrass). There is little variation in vascular plant species composition across the range. *Spartina alterniflora* (smooth cordgrass) occurs in nearly pure stands, occasionally with other low-statured species. *Salicornia* (pickleweed) spp. can be quite common mixed in with the *Spartina* (cordgrass), often becoming more apparent later in the growing season. *Limonium carolinianum* (Carolina sealavender) is another characteristic herb, but only as scattered individuals. More detailed information is needed on the composition of this vegetation and how it differs floristically from its northern equivalent, *Spartina alterniflora*/(*Ascophyllum nodosum*) Acadian/Virginian Zone Herbaceous Vegetation (CEGL004192).

Most Abundant Species

Fort Pulaski National Monument

Stratum

Herb (field)

Lifeform

Graminoid

Species

Spartina alterniflora (smooth cordgrass)

Global

Stratum

Herb (field)

Lifeform

Graminoid

Species

Spartina alterniflora (smooth cordgrass)

Characteristic Species

Fort Pulaski National Monument: *Sarcocornia pacifica* (Pacific swampfire), *Spartina alterniflora* (smooth cordgrass)

Global: *Spartina alterniflora* (smooth cordgrass)

Conservation Status Rank

Global Rank & Reasons: G5 (1-Dec-1997). Since it occurs over large areas, this community is in no danger of disappearing. However, its role in the functioning of ecosystems makes the presence of large areas very important. Losses of this community could have serious consequences for nutrient cycling and reproduction of estuarine and marine organisms (Schafale and Weakley 1990).

Classification

Status: Standard

Classification Confidence: 1 - Strong

Fort Pulaski National Monument Comments: NatureServe identified the perennial glasswort plants at Fort Pulaski as *Sarcocornia pacifica* using the recent key in Weakley (2007), which follows FNA Editorial Committee 2003). The perennial glassworts on the U.S. East Coast have been called *Sarcocornia perennis* in the past (Kartesz 1999), and their taxonomy remains unsettled. For the time being, NatureServe will continue to follow the Kartesz (1999) nomenclature in our global descriptions and U.S. National Vegetation Classification association names.

Global Similar Associations:

- *Spartina alterniflora*/(*Ascophyllum nodosum*) Acadian/Virginian Zone Herbaceous Vegetation (CEGL004192)

Global Related Concepts:

- Salt Marsh (Carolinian Subtype) (Schafale 2000) ?

Element Distribution

Fort Pulaski National Monument Range: This community was sampled in the salt marsh northeast of the fort and west of the gate house. This community by far occupies the greatest amount of the property present at Fort Pulaski National Monument. Over 340 acres of Cockspur Island and over 4400 acres of McQueens Island are dominated by salt marsh (Govus 1998).

Global Range: This vegetation ranges from Cape Hatteras, North Carolina, south to the Atlantic Coast of the Florida peninsula (Cowardin 1979). The northern limit corresponds with the southern limit of the Virginian province of the American Atlantic Temperate Region, a transitional area harboring animal species of both southern and northern affinities.

Nations: US

States/Provinces: FL, GA, NC, SC

TNC Ecoregions: 55:P, 56:C, 57:C

USFS Ecoregions: 232Cb:CCC, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC

Federal Lands: NPS (Cape Hatteras, Cape Lookout, Fort Pulaski, Timucuan); USFS (Croatan)

Element Sources

Fort Pulaski National Monument Plots: FOPU.7, FOPU.10.

Local Description Authors: M. J. Russo and C. W. Nordman

Global Description Authors: A. S. Weakley

References: Cowardin et al. 1979, FNA Editorial Committee 2003, FNAI 1992a, Govus 1998, Kartesz 1999, Nelson 1986, Peet et al. unpubl. data 2002, Schafale 2000, Schafale 2003b, Schafale and Weakley 1990, Southeastern Ecology Working Group n.d., Weakley 2007

***Spartina cynosuroides* Tidal Herbaceous Alliance (A.1480)**Alliance Concept

Summary: This alliance occurs as narrow, almost pure, stands of *Spartina cynosuroides* (big cordgrass) along tidal creeks and sloughs or on levees of oligohaline tidal marshes. Some *Spartina cynosuroides* (big cordgrass) communities are nearly monospecific, while others have a diverse component of other graminoids and forbs. Associated plants include *Schoenoplectus pungens* (common threesquare), *Schoenoplectus robustus* (sturdy bulrush), *Schoenoplectus tabernaemontani* (softstem bulrush), *Pontederia cordata* (pickerelweed), *Peltandra virginica* (green arrow arum), *Typha domingensis* (southern cattail), and *Typha angustifolia* (narrowleaf cattail), among others. Communities in this alliance occur mainly in the mid-Atlantic states with the northern extent of distribution being southern New England.

Classification Comments: There may be several associations, determined by salinity.

Related Concepts:

- Big Cordgrass Community Type (Odum et al. 1984) ?
- Brackish Marsh (Nelson 1986) I

- Brackish Marsh (Schafale and Weakley 1990) I
- Brackish Marsh (Wharton 1978) I
- Tidal Freshwater Marsh, Oligohaline Variant (Schafale and Weakley 1990) ?

Alliance Description

Environment: This alliance occurs as narrow, almost pure, stands of *Spartina cynosuroides* along tidal creeks and sloughs or on levees of oligohaline tidal marshes.

Vegetation: Some *Spartina cynosuroides* (big cordgrass) communities are nearly monospecific, while others have a diverse component of other graminoids and forbs. Associated plants include *Schoenoplectus pungens* (common threesquare), *Schoenoplectus robustus* (sturdy bulrush), *Schoenoplectus tabernaemontani* (softstem bulrush), *Pontederia cordata* (pickerelweed), *Peltandra virginica* (green arrow arum), *Typha domingensis* (southern cattail), and *Typha angustifolia* (narrowleaf cattail), among others.

Alliance Distribution

Range: Communities in this alliance occur mainly in the mid-Atlantic states with the northern extent of distribution being southern New England. It is found in Georgia, North Carolina, South Carolina, Connecticut, Delaware, Maryland, Massachusetts, New Jersey, New York, and Virginia.

Nations: US

Subnations: DE, GA, MA, MD, NC, NJ, SC, VA

TNC Ecoregions: 52:P, 56:C, 57:C, 58:C, 62:C

USFS Ecoregions: 221Ab:CCC, 232Ab:CCP, 232Ac:CCP, 232Ad:CCC, 232Br:CCC, 232Bt:CCC, 232Bx:CCC, 232Bz:CCC, 232Cb:CCP, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC

Federal Lands: NPS (Assateague Island, Cape Cod, Colonial, Fort Pulaski, George Washington Birthplace); USFWS (Back Bay, Chesapeake Marshlands)

(CEGL004195) Atlantic Giant Cordgrass Marsh

Spartina cynosuroides Herbaceous Vegetation

Giant Cordgrass Herbaceous Vegetation

NVC Classification

Physiognomic Class	Herbaceous Vegetation (V)
Physiognomic Subclass	Perennial graminoid vegetation (V.A.)
Physiognomic Group	Temperate or subpolar grassland (V.A.5.)
Physiognomic Subgroup	Natural/Semi-natural temperate or subpolar grassland (V.A.5.N.)
Formation	Tidal temperate or subpolar grassland (V.A.5.N.n.)
Alliance	<i>Spartina cynosuroides</i> Tidal Herbaceous Alliance (A.1480)
Alliance (English name)	Giant Cordgrass Tidal Herbaceous Alliance
Association	<i>Spartina cynosuroides</i> Herbaceous Vegetation
Association (English name)	Giant Cordgrass Herbaceous Vegetation
Association (Common name)	Atlantic Giant Cordgrass Marsh
Ecological System(s):	Northern Atlantic Coastal Plain Brackish Tidal Marsh (CES203.894) Central Atlantic Coastal Plain Fresh and Oligohaline Tidal Marsh (CES203.376) Atlantic Coastal Plain Embayed Region Tidal Freshwater Marsh (CES203.259)

Element Concept

Global Summary: This community includes narrow, almost pure stands of *Spartina cynosuroides* (big cordgrass) along tidal creeks and sloughs or on levees of oligohaline tidal marshes along the central Atlantic Coast. Some occurrences are nearly monospecific, while others have a diverse component of other graminoids and forbs. Where mixed, associated plants include *Schoenoplectus pungens* (common threesquare), *Schoenoplectus robustus* (sturdy bulrush), *Schoenoplectus americanus* (chairmaker's bulrush), *Kosteletzkya virginica* (Virginia saltmarsh mallow), *Hibiscus moscheutos* (crimson-eyed rosemallow), *Amaranthus cannabinus* (tidalmarsh amaranth), *Panicum virgatum* (switchgrass), *Peltandra virginica* (green arrow arum), and *Polygonum punctatum* (dotted smartweed). Where more saline, *Spartina alterniflora* (smooth cordgrass) and *Iva frutescens* (Jesuit's bark) can become more frequent. Where less saline, associates can include *Schoenoplectus tabernaemontani* (softstem bulrush), *Pontederia cordata* (pickerelweed), *Peltandra virginica* (green arrow arum), *Leersia oryzoides* (rice cutgrass), *Mikania scandens* (climbing hempvine), *Rumex verticillatus* (swamp dock), *Echinochloa walteri* (coast cockspur grass), *Polygonum hydropiperoides* (swamp smartweed), and *Typha angustifolia* (narrowleaf cattail) (or *Typha domingensis* (southern cattail) in the south), among others. In more disturbed areas, this association can be displaced by *Phragmites australis* (common reed).

Environmental Description

USFWS Wetland System: Estuarine

Fort Pulaski National Monument Environment: This community does not occur over a large area; typically it is located in narrow bands associated with tidal ditches and along fringe areas between tidal herbaceous vegetation and tidal shrublands (Govus 1998).

Global Environment: This association occurs along tidal creeks, guts, and levees of oligohaline to mesohaline marshes in irregularly flooded high marsh situations, but in areas that are well-drained.

Vegetation Description

Fort Pulaski National Monument Vegetation: In addition to *Spartina cynosuroides* (big cordgrass), *Schoenoplectus robustus* (sturdy bulrush) is also sometimes present (Govus 1998).

Global Vegetation: *Spartina cynosuroides* (big cordgrass) is strongly dominant in this community, often occurring in pure stands. Where stands are mixed, common associates can include *Schoenoplectus robustus* (sturdy bulrush), *Schoenoplectus pungens* (common threesquare), *Schoenoplectus americanus* (chairmaker's bulrush), *Peltandra virginica* (green arrow arum), *Kosteletzkya virginica* (Virginia saltmarsh mallow), *Amaranthus cannabinus* (tidalmarsh amaranth), *Panicum virgatum*, *Peltandra virginica* (green arrow arum), and *Polygonum punctatum* (dotted smartweed). Where more saline, *Spartina alterniflora* (smooth cordgrass) and *Iva frutescens* (Jesuit's bark) can become more frequent. Where less saline, *Schoenoplectus tabernaemontani* (softstem bulrush), *Polygonum hydropiperoides* (swamp smartweed), *Pontederia cordata* (pickerelweed), *Peltandra virginica* (green arrow arum), *Leersia oryzoides* (rice cutgrass), *Mikania scandens* (climbing hempvine), *Rumex verticillatus* (swamp dock), *Echinochloa walteri* (coast cockspur grass), and *Typha angustifolia* (narrowleaf cattail) are also common associates. In more disturbed areas, this association can be displaced by *Phragmites australis* (common reed).

Most Abundant Species

Fort Pulaski National Monument

Stratum

Herb (field)

Lifeform

Graminoid

Species

Spartina cynosuroides (big cordgrass)

Global

Stratum

Herb (field)

Lifeform

Graminoid

Species

Spartina cynosuroides (big cordgrass)

Characteristic Species

Fort Pulaski National Monument: *Spartina cynosuroides* (big cordgrass)

Global: *Spartina cynosuroides* (big cordgrass)

Conservation Status Rank

Global Rank & Reasons: G4 (13-Aug-1997).

Classification

Status: Standard

Classification Confidence: 2 - Moderate

Global Comments: *Spartina cynosuroides* - *Panicum virgatum* - *Phyla lanceolata* Herbaceous Vegetation (CEGL007741) is similar in terms of dominant and codominant species, however, associated species differ somewhat and CEGL007741 occurs in wind tidal situations in North Carolina and Virginia. North of New Jersey, *Spartina cynosuroides* is a component of brackish high marsh associations, but does not tend to dominate.

Global Similar Associations:

- *Spartina cynosuroides* - *Panicum virgatum* - *Phyla lanceolata* Herbaceous Vegetation (CEGL007741)

Global Related Concepts:

- *Spartina cynosuroides* Herbaceous Vegetation (Coulling 2002) =
- Big Cordgrass Community Type (Odum et al. 1984) ?
- Brackish Marsh (Wharton 1978) B
- Brackish tidal marsh complex (Breden 1989) ?
- Tidal Freshwater Marsh (Giant Cordgrass Subtype) (Schafale 2000) ?

Element Distribution

Fort Pulaski National Monument Range: This community does not occur over a large area; typically it is located in narrow bands associated with tidal ditches and also along fringe areas between tidal herbaceous vegetation and tidal shrublands (Govus 1998).

Global Range: This association occurs along the Atlantic Coast from New Jersey to Georgia.

Nations: US

States/Provinces: DE:S3?, GA, MA, MD:S4?, NC, NJ:S3, SC, VA

TNC Ecoregions: 56:C, 57:C, 58:C, 62:C

USFS Ecoregions: 221Ab:CCC, 232Ab:CCP, 232Ac:CCP, 232Ad:CCC, 232Br:CCC, 232Bt:CCC, 232Bx:CCC, 232Bz:CCC, 232Cb:CCP, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC

Federal Lands: NPS (Assateague Island, Cape Cod, Colonial, Fort Pulaski, George Washington Birthplace); USFWS (Back Bay, Chesapeake Marshlands)

Element Sources

Fort Pulaski National Monument Plots: None.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: S.L. Neid

References: Bowman 2000, Breden 1989, Breden et al. 2001, Clancy 1996, Coulling 2002, Fleming 2001a, Fleming and Moorhead 1998, Fleming et al. 2001, Govus 1998, Harrison 2001, Harrison 2004, Nelson 1986, Odum 1988, Odum and Smith 1981, Odum et al. 1984, Schafale 2000, Schafale 2003b, Schafale and Weakley 1990, Southeastern Ecology Working Group n.d., Wharton 1978

***Spartina patens* – (*Distichlis spicata*) Tidal Herbaceous Alliance (A.1481)**

Alliance Concept

Summary: This alliance comprises "high salt marsh" vegetation dominated or codominated by *Spartina patens* (saltmeadow cordgrass) along the Gulf and Atlantic coasts from Maine to Texas. The high salt marsh is irregularly flooded by tides and forms at slightly higher elevations than regularly flooded low marshes; they establish where peat accumulation raises the marsh surface above mean high tide. Landward the vegetation can transition to brackish marsh or upland vegetation. Vegetation of this alliance also occurs in mesohaline zones along lower reaches of tidal rivers.

Variation in codominant species occurs across the geographic range. From the Canadian maritime provinces south to Delaware (discontinuously south to Virginia), this alliance is characterized by the dominance of *Spartina patens* (saltmeadow cordgrass), *Distichlis spicata* (inland saltgrass), and *Juncus gerardii* (saltmeadow rush) and the presence of more northerly distributed marsh species such as *Puccinellia fasciculata* (saltmarsh alkaligrass), *Plantago maritima* (goose tongue), and *Triglochin maritima* (seaside arrowgrass). In brackish reaches of tidal rivers, this alliance includes *Spartina patens* (saltmeadow cordgrass)-dominated vegetation that may also be characterized by the presence of *Agrostis stolonifera* (creeping bentgrass), *Festuca rubra* (red fescue), *Symphyotrichum novi-belgii* (New York aster), *Hierochloa odorata* (vanilla grass), *Carex paleacea* (chaffy sedge), or *Spartina pectinata* (prairie cordgrass). From Delaware south to Florida, this high salt marsh coastal community is dominated by *Spartina patens* (saltmeadow cordgrass), forming meadows at slightly higher elevations in relation to the adjacent *Spartina alterniflora* (smooth cordgrass) Tidal Herbaceous Alliance (A.1471). Diagnostic species for this community are *Spartina patens* (saltmeadow cordgrass), *Distichlis spicata* (inland saltgrass), *Borrchia frutescens* (bushy seaside tansy), *Kosteletzkya virginica* (Virginia saltmarsh mallow), and *Pluchea odorata* (sweetscent). Shrub seedlings such as *Baccharis halimifolia* (eastern baccharis) and *Morella cerifera* (wax myrtle) may also be present. The associated *Juncus roemerianus* (needlegrass rush) Tidal Herbaceous Alliance (A.1475) often occurs as discrete patches which may reach substantial size.

This alliance also includes mesohaline to oligohaline marshes of the Gulf Coast of Texas and Louisiana. In these associations, *Spartina patens* (saltmeadow cordgrass) may strongly dominate, *Distichlis spicata* (inland saltgrass), *Spartina alterniflora* (smooth cordgrass), and *Spartina*

patens (saltmeadow cordgrass) may codominate, *Distichlis spicata* (inland saltgrass) may form pure stands, *Paspalum vaginatum* (seashore paspalum) may strongly dominate, or *Spartina patens* (saltmeadow cordgrass) and *Vigna luteola* (hairypod cowpea) may codominate. Other characteristic species include *Juncus roemerianus* (needlegrass rush), *Spartina spartinae* (gulf cordgrass), *Spartina cynosuroides* (big cordgrass) (within its range), *Schoenoplectus robustus* (sturdy bulrush), *Schoenoplectus americanus* (chairmaker's bulrush), *Sagittaria lancifolia* (bulltongue arrowhead), *Phragmites australis* (common reed), and *Eragrostis* (lovegrass) spp. Here, this alliance forms mosaics with *Spartina spartinae* (gulf cordgrass) and *Spartina alterniflora* (smooth cordgrass) marshes and saline herbaceous vegetation.

Western states have a different alliance for inland situations dominated by *Distichlis spicata* (inland saltgrass), the *Distichlis spicata* (inland saltgrass) Intermittently Flooded Herbaceous Alliance (A.1332).

Classification Comments: This may represent multiple zones; more research is needed.

Similar Alliances:

- *Distichlis spicata* Intermittently Flooded Herbaceous Alliance (A.1332)
- *Distichlis spicata* Tidal Herbaceous Alliance (A.1882)
- *Juncus roemerianus* Tidal Herbaceous Alliance (A.1475)
- *Schoenoplectus americanus* Tidal Herbaceous Alliance (A.2007)
- *Spartina alterniflora* Tidal Herbaceous Alliance (A.1471)

Similar Alliance Comments: From Delaware south to Florida, this high salt marsh coastal community, dominated by *Spartina patens*, forms meadows at slightly higher elevations in relation to the adjacent *Spartina alterniflora* Tidal Herbaceous Alliance (A.1471). The associated *Juncus roemerianus* Tidal Herbaceous Alliance (A.1475) occurs as discrete patches which may reach substantial size. Western states have a different alliance for inland situations, the *Distichlis spicata* Intermittently Flooded Herbaceous Alliance (A.1332). In coastal Texas and possibly elsewhere along the Gulf Coast, the relationship between this alliance and the similar *Schoenoplectus americanus* Tidal Herbaceous Alliance (A.2007) is not well understood. Dominance patterns are likely related to gradients in salinity and hydrology. *Schoenoplectus americanus* may dominate in areas with higher average water levels, lower salinities, and lower frequency of flooding than areas typically dominated by *Spartina patens* or *Distichlis spicata*.

Related Concepts:

- Brackish Marsh (Wieland 1994b) I
- Brackish Marsh (Nelson 1986) I
- Estuarine Intertidal: Brackish Tidal Marsh (Swain and Kearsley 2001) ?
- Estuarine Intertidal: Salt Marsh (Swain and Kearsley 2001) ?
- Intermediate Marsh (Smith 1996a) ?
- Marshhay Cordgrass Series (Diamond 1993) I
- Salt Marsh (Wieland 1994b) I
- Salt Marsh (Schafale and Weakley 1990) I
- Salt Marsh (Nelson 1986) I
- Salt Marsh (Smith 1996a) I

- Saltgrass-Cordgrass Series (Diamond 1993) ?

Alliance Description

Environment: The high salt marsh is irregularly flooded by tides and forms at slightly higher elevations than regularly flooded low marshes; they establish where peat accumulation raises the marsh surface above mean high tide. Landward the vegetation can transition to brackish marsh or upland vegetation. Vegetation of this alliance also occurs in mesohaline zones along lower reaches of tidal rivers. The substrate is peat of variable depths overlying sand.

Vegetation: Variation in codominant species occurs across the geographic range. From the Canadian maritime provinces south to Delaware (discontinuously south to Virginia), this alliance is characterized by the dominance of *Spartina patens* (saltmeadow cordgrass), *Distichlis spicata* (inland saltgrass), and *Juncus gerardii* (saltmeadow rush) and the presence of more northerly distributed marsh species such as *Puccinellia fasciculata* (saltmarsh alkaligrass), *Plantago maritima* (goose tongue), and *Triglochin maritima* (seaside arrowgrass). In brackish reaches of tidal rivers, this alliance includes *Spartina patens* (saltmeadow cordgrass)-dominated vegetation that may also be characterized by the presence of *Agrostis stolonifera* (creeping bentgrass), *Festuca rubra* (red fescue), *Symphyotrichum novi-belgii* (New York aster), *Hierochloa odorata* (vanilla grass), *Carex paleacea* (chaffy sedge), or *Spartina pectinata* (prairie cordgrass). From Delaware south to Florida, this high salt marsh coastal community is dominated by *Spartina patens* (saltmeadow cordgrass), forming meadows at slightly higher elevations in relation to the adjacent *Spartina alterniflora* (smooth cordgrass) Tidal Herbaceous Alliance (A.1471). *Distichlis spicata* (inland saltgrass), *Limonium carolinianum* (Carolina sealavender), *Agalinis maritima* (saltmarsh false foxglove), *Salicornia virginica* (Virginia glasswort), *Sabatia stellaris* (rose of Plymouth), *Borrichia frutescens* (bushy seaside tansy), *Lythrum lineare* (wand lythrum), *Schoenoplectus pungens* (common threesquare), *Eleocharis rostellata* (beaked spikerush), *Solidago sempervirens* (seaside goldenrod), *Fimbristylis castanea* (marsh fimbry), *Pluchea odorata* (sweetscent), *Hibiscus moscheutos* (crimson-eyed rosemallow), and *Atriplex prostrata* (triangle orache) are characteristic associates. Diagnostic species are *Spartina patens* (saltmeadow cordgrass), *Distichlis spicata* (inland saltgrass), *Borrichia frutescens* (bushy seaside tansy), *Kosteletzkya virginica* (Virginia saltmarsh mallow), and *Pluchea odorata* (sweetscent). This alliance also includes mesohaline to oligohaline marshes of the Gulf Coast of Texas and Louisiana. In these associations, *Spartina patens* (saltmeadow cordgrass) may strongly dominate, *Distichlis spicata* (inland saltgrass), *Spartina alterniflora* (smooth cordgrass), and *Spartina patens* (saltmeadow cordgrass) may codominate, *Distichlis spicata* (inland saltgrass) may form pure stands, *Paspalum vaginatum* (seashore paspalum) may strongly dominate, or *Spartina patens* (saltmeadow cordgrass) and *Vigna luteola* (hairypod cowpea) may codominate. Other characteristic species include *Juncus roemerianus* (needlegrass rush), *Spartina spartinae* (gulf cordgrass), *Spartina cynosuroides* (big cordgrass) (within its range), *Schoenoplectus robustus* (sturdy bulrush), *Schoenoplectus americanus* (chairmaker's bulrush), *Sagittaria lancifolia* (bulltongue arrowhead), *Phragmites australis* (common reed), and *Eragrostis* (lovegrass) spp. Here, this alliance forms mosaics with *Spartina spartinae* (gulf cordgrass) and *Spartina alterniflora* (smooth cordgrass) marshes and saline herbaceous vegetation.

Dynamics: Dominance patterns are likely related to gradients in salinity and hydrology and are not well understood.

Alliance Distribution

Range: This alliance is found along the Atlantic and Gulf coasts from Maine south to Florida and west to Texas.

Nations: US

Subnations: AL, CT, DE, FL, GA, LA, MA, MD, ME, MS, NC, NH, NJ, NY, RI, SC, TX, VA

TNC Ecoregions: 30:P, 31:C, 41:P, 53:C, 54:P, 55:P, 56:C, 57:C, 58:C, 62:C, 63:C

USFS Ecoregions: 212Cb:CCC, 212Db:CCC, 212Dc:CCC, 221Aa:CCC, 221Ab:CCC, 221Ac:CCC, 221Ad:CCC, 221Ae:CCC, 221Ak:CCC, 221Dc:CCC, 231Fb:CCC, 232Aa:CCC, 232Ab:CCC, 232Ac:CCC, 232Ad:CCC, 232Bb:CCP, 232Bc:CCP, 232Bd:CCP, 232Br:CCC, 232Bx:CCC, 232Bz:CCC, 232Cb:CCC, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC, 232Cj:CCC, 232Db:CCP, 232Dc:CCC, 232Dd:CCC, 232De:CCP, 232Eb:CCC, 232Ed:CCC, 232Ee:CCC, 232Gb:CCP, 255Da:CCP, 255Dc:CCC, 315F:CC

Federal Lands: NPS (Acadia, Assateague Island, Boston Harbor Islands, Cape Cod, Cape Hatteras, Cape Lookout, Chincoteague, Fire Island, Fort Pulaski, Gateway, Muskeget Island?, Sagamore Hill); USFWS (Anahuac, Back Bay?, Big Boggy, Brazoria, Chesapeake Marshlands, Matagorda Island, McFaddin, Moosehorn, Parker River?, Prime Hook, Rachel Carson, San Bernard, Texas Point)

(CEGL004197) Mid-Atlantic High Salt Marsh

***Spartina patens* - *Distichlis spicata* - (*Juncus roemerianus*) Herbaceous Vegetation**

Saltmeadow Cordgrass - Saltgrass - (Black Needlerush) Herbaceous Vegetation

NVC Classification

Physiognomic Class	Herbaceous Vegetation (V)
Physiognomic Subclass	Perennial graminoid vegetation (V.A.)
Physiognomic Group	Temperate or subpolar grassland (V.A.5.)
Physiognomic Subgroup	Natural/Semi-natural temperate or subpolar grassland (V.A.5.N.)
Formation	Tidal temperate or subpolar grassland (V.A.5.N.n.)
Alliance	<i>Spartina patens</i> - (<i>Distichlis spicata</i>) Tidal Herbaceous Alliance (A.1481)
Alliance (English name)	Saltmeadow Cordgrass - (Saltgrass) Tidal Herbaceous Alliance
Association	<i>Spartina patens</i> - <i>Distichlis spicata</i> - (<i>Juncus roemerianus</i>) Herbaceous Vegetation
Association (English name)	Saltmeadow Cordgrass - Saltgrass - (Black Needlerush) Herbaceous Vegetation
Association (Common name)	Mid-Atlantic High Salt Marsh
Ecological System(s):	Atlantic Coastal Plain Embayed Region Tidal Salt and Brackish Marsh (CES203.260) Northern Atlantic Coastal Plain Tidal Salt Marsh (CES203.519) Central Atlantic Coastal Plain Salt and Brackish Tidal Marsh (CES203.270)

Element Concept

Global Summary: This coastal community is an irregularly tidally flooded high salt marsh of the mid- and southern Atlantic Coast from Delaware south to Florida. It is dominated by *Spartina patens* (saltmeadow cordgrass), which forms meadows with a distinct "cowlicked" appearance. These meadows occur at slightly higher elevations than adjacent, regularly flooded low salt marsh occupying the zone extending from mean high tide landward approximately to the limit of high spring tides. The substrate is peat of variable depths overlying sand. *Distichlis spicata* (inland saltgrass) can be codominant. Additional associated species that generally occur in low abundance can include *Limonium carolinianum* (Carolina sealavender), *Agalinis maritima* (saltmarsh false foxglove), *Salicornia virginica* (Virginia glasswort), *Juncus roemerianus*

(needlegrass rush), *Sabatia stellaris* (rose of Plymouth), *Borrichia frutescens* (bushy seaside tansy), *Lythrum lineare* (wand lythrum), *Solidago sempervirens* (seaside goldenrod), *Pluchea odorata* (sweetscent), *Hibiscus moscheutos* ssp. *moscheutos* (crimsoneyed rosemallow), or *Atriplex prostrata* (triangle orache). Shrub seedlings of *Baccharis halimifolia* (eastern baccharis), *Iva frutescens* (Jesuit's bark), and/or *Morella cerifera* (wax myrtle) may occur sporadically. Diagnostic species are *Spartina patens* (saltmeadow cordgrass), *Distichlis spicata*, *Borrichia frutescens* (bushy seaside tansy), *Kosteletzkya virginica* (Virginia saltmarsh mallow), and *Pluchea odorata* (sweetscent). One unusual Virginia occurrence is dominated by *Plantago maritima* var. *juncoides* (goose tongue).

Environmental Description

USFWS Wetland System: Estuarine

Fort Pulaski National Monument Environment: This community is found on infrequently tidally flooded terraces (Govus 1998).

Global Environment: This high salt marsh association generally occurs behind barrier beaches, but also in the outer reaches of estuaries, occupying the zone extending from mean high tide landward approximately to the limit of high spring tides. They are often adjacent to low salt marshes dominated by *Spartina alterniflora* (tall form), which are regularly flooded by diurnal tides. *Spartina patens*-dominated high marshes form very dense peat with high organic matter content. Peat forms over sand, silt or bedrock.

Vegetation Description

Fort Pulaski National Monument Vegetation: This is one of the most diverse and scenically attractive natural communities present on the park. In addition to *Spartina patens* (saltmeadow cordgrass), *Distichlis spicata* (inland saltgrass), and *Borrichia frutescens* (bushy seaside tansy), other species found here include *Solidago sempervirens* (seaside goldenrod), *Limonium carolinianum* (Carolina sealavender), *Cynanchum angustifolium* (gulf coast swallow-wort), *Paspalum distichum* (knotgrass), *Fimbristylis* (fimbry) spp., and, along the upper edges in overwash areas, *Atriplex prostrata* (triangle orache) (Govus 1998).

Global Vegetation: This high salt marsh is dominated by *Spartina patens* (saltmeadow cordgrass) with *Distichlis spicata* (inland saltgrass) occurring as a common associate or a codominant species. Associated species that generally occur in low abundance can include *Limonium carolinianum* (Carolina sealavender), *Agalinis maritima* (saltmarsh false foxglove), *Salicornia virginica* (Virginia glasswort), *Sabatia stellaris* (rose of Plymouth), *Borrichia frutescens* (bushy seaside tansy), *Lythrum lineare* (wand lythrum), *Juncus roemerianus* (needlegrass rush), *Solidago sempervirens* (seaside goldenrod), *Pluchea odorata* (sweetscent), *Hibiscus moscheutos* ssp. *moscheutos* (crimsoneyed rosemallow), and/or *Atriplex prostrata* (triangle orache). Shrub seedlings of *Baccharis halimifolia* (eastern baccharis), *Iva frutescens* (Jesuit's bark), and/or *Morella cerifera* (wax myrtle) may occur sporadically.

Most Abundant Species

Fort Pulaski National Monument

Stratum

Herb (field)

Herb (field)

Lifeform

Forb

Graminoid

Species

Borrichia frutescens (bushy seaside tansy)

Spartina patens (saltmeadow cordgrass)

Global

Stratum

Herb (field)

Lifeform

Graminoid

Species

Distichlis spicata (inland saltgrass), *Spartina patens* (saltmeadow cordgrass)

Characteristic Species

Fort Pulaski National Monument: *Borrichia frutescens* (bushy seaside tansy), *Spartina patens* (saltmeadow cordgrass)

Global: *Distichlis spicata* (inland saltgrass), *Kosteletzkya virginica* (Virginia saltmarsh mallow), *Pluchea odorata* (sweetscent), *Spartina patens* (saltmeadow cordgrass)

Conservation Status Rank

Global Rank & Reasons: G4G5 (13-May-2002). Although widespread on the eastern seaboard, examples of this community that have not been impacted by ditching, dredging and filling, mosquito spraying, and other activities are relatively uncommon.

Classification

Status: Standard

Classification Confidence: 2 - Moderate

Global Comments: Although *Juncus roemerianus* may not be present or have minor cover in some stands, it is a differential species and was chosen as a nominal of this association to distinguish it from *Spartina patens* - *Distichlis spicata* - (*Juncus gerardii*) Herbaceous Vegetation (CEGL006006), which occurs to the north. *Spartina patens* - *Distichlis spicata* - (*Juncus roemerianus*) Herbaceous Vegetation (CEGL004197) is differentiated by the absence or relative infrequent occurrence of *Juncus gerardii*, *Plantago maritima*, and *Triglochin maritima* (in pannes), and by the importance of species of southern distribution, such as *Borrichia frutescens*, *Kosteletzkya virginica*, *Fimbristylis castanea*, and *Lythrum lineare*. If discrete patches of *Juncus roemerianus* occur in substantial size (several acres), the community is considered to be *Juncus roemerianus* Herbaceous Vegetation (CEGL004186). Currently, some unusually diverse variants exist in Virginia that may warrant recognition as a separate association.

Global Similar Associations:

- *Juncus roemerianus* Herbaceous Vegetation (CEGL004186)
- *Spartina alterniflora* - *Distichlis spicata* Tidal Herbaceous Vegetation [Provisional] (CEGL006586)
- *Spartina patens* - *Distichlis spicata* - (*Juncus gerardii*) Herbaceous Vegetation (CEGL006006)

Global Related Concepts:

- *Aster tenuifolius* - *Distichlis spicata* - *Fimbristylis castanea* - *Borrichia frutescens* - *Spartina patens* association (Adams 1963) =
- *Spartina* - *Distichlis* - *Juncus* associates (Penfound 1952) =
- *Spartina patens* - *Distichlis spicata* - *Sarcocornia perennis* - *Limonium carolinianum* Herbaceous Vegetation (Coulling 2002) F
- *Spartina patens* - *Distichlis spicata* Herbaceous Vegetation (Harrison 2001) I

- *Spartina patens* - *Distichlis spicata* Herbaceous Vegetation (Coulling 2002) F
- *Spartina patens* - *Distichlis spicata* high marsh (Clancy 1993b) =
- *Spartina patens* type (Cooper and Waits 1973) =
- Brackish Marsh (Salt Meadow Cordgrass Subtype) (Schafale 2000) ?
- High marsh (Cooper 1974) =
- Salt marsh (Higgins et al. 1971) B
- Salt marsh community (Hill 1986) B

Element Distribution

Fort Pulaski National Monument Range: This community is found in large areas, primarily along the infrequently tidally flooded terraces of north central Cockspear Island (Govus 1998).

Global Range: This association occurs along the Atlantic Coast from Delaware to Florida.

Nations: US

States/Provinces: DE:S4, FL, GA, MD, NC, NJ, SC, VA

TNC Ecoregions: 56:C, 57:C, 58:C, 62:C

USFS Ecoregions: 232Br:CCC, 232Bx:CCC, 232Bz:CCC, 232Ce:CCC, 232Ch:CCC, 232Ci:CCC

Federal Lands: NPS (Assateague Island, Cape Hatteras, Cape Lookout, Fort Pulaski); USFWS (Back Bay?, Chesapeake Marshlands, Chincoteague, Prime Hook)

Element Sources

Fort Pulaski National Monument Plots: None.

Local Description Authors: M.J. Russo and C.W. Nordman

Global Description Authors: L.A. Sneddon, mod. S.L. Neid

References: Adams 1963, Bowman 2000, Clancy 1993b, Clancy 1996, Cooper 1974, Cooper and Waits 1973, Coulling 2002, Eastern Ecology Working Group n.d., Fleming 2001a, Fleming et al. 2001, Govus 1998, Harrison 2001, Harrison 2004, Higgins et al. 1971, Hill 1986, Nelson 1986, Peet et al. unpubl. data 2002, Penfound 1952, Schafale 2000, Schafale 2003b, Schafale and Weakley 1990

Bibliography

- Abrahamson, W. G., A. F. Johnson, J. N. Layne, and P. A. Peroni. 1984. Vegetation of the Archbold Biological Station, Florida: An example of the southern Lake Wales Ridge. *Florida Scientist* 47:209-250.
- Adams, D. A. 1963. Factors influencing vascular plant zonation in North Carolina salt marshes. *Ecology* 44:445-456.
- Allard, D. J. 1990. Southeastern United States ecological community classification. Interim report, Version 1.2. The Nature Conservancy, Southeast Regional Office, Chapel Hill, NC. 96 pp.
- Ambrose, J. 1990a. Georgia's natural communities--A preliminary list. Unpublished document. Georgia Natural Heritage Inventory. 5 pp.
- Au, S. 1974. Vegetation and ecological processes on Shackleford Bank, North Carolina. USDI National Park Service, Scientific Monograph No. 6.
- Austin, D. F., and K. Coleman-Marais. 1977. Vegetation of southeastern Florida. II. Boca Raton hammock site. *Florida Scientist* 40:331-338.
- Barbour, M. G., M. Rejmanek, A. F. Johnson, and B. M. Pavlik. 1987. Beach vegetation and plant distribution patterns along the northern Gulf of Mexico. *Phytocoenologia* 15:201-234.
- Barry, J. F. 1980. Natural vegetation of South Carolina. University of South Carolina Press, Columbia. 214 pp.
- Bellis, V. J. 1992. Floristic continuity among the maritime forests of the Atlantic Coast of the United States. Pages 21-29 in: C. A. Cole and F. K. Turner, editors. Barrier island ecology of the mid-Atlantic Coast: A symposium. Technical Report NPS/SERCAHA/NRTR-93/04.
- Berdine, M. A. 1998. Maryland vegetation classification. Maryland Department of Natural Resources, Annapolis, MD.
- Borhidi, A. 1991. Phytogeography and vegetation ecology of Cuba. Akademiai Kiado. Budapest, Hungary. 858 pp. plus color plates and map by A. Borhidi and O. Muniz (1970) inside of back cover.
- Boule, M. E. 1979. The vegetation of Fisherman Island, Virginia. *Castanea* 44:98-108.
- Bourdeau, P. F., and H. J. Oosting. 1959. The maritime live oak forest in North Carolina. *Ecology* 40:148-152.

- Bowman, P. 2000. Draft classification for Delaware. Unpublished draft. Delaware Natural Heritage Program.
- Breden, T. F. 1989. A preliminary natural community classification for New Jersey. Pages 157-191 in: E. F. Karlin, editor. New Jersey's rare and endangered plants and animals. Institute for Environmental Studies, Ramapo College, Mahwah, NJ. 280 pp.
- Breden, T. F., Y. R. Alger, K. S. Walz, and A. G. Windisch. 2001. Classification of vegetation communities of New Jersey: Second iteration. Association for Biodiversity Information and New Jersey Natural Heritage Program, Office of Natural Lands Management, Division of Parks and Forestry, New Jersey Department of Environmental Protection, Trenton.
- Clancy, K. 1993b. A preliminary classification of the natural communities of Delaware. Unpublished draft, Delaware Natural Heritage Inventory, Division of Parks and Recreation, Dover. 30 pp.
- Clancy, K. 1996. Natural communities of Delaware. Unpublished review draft. Delaware Natural Heritage Program, Division of Fish and Wildlife, Delaware Division of Natural Resources and Environmental Control, Smyrna, DE. 52 pp.
- Clewell, A. F. 1971. The vegetation of the Apalachicola National Forest: An ecological perspective. Unpublished document. USDA Forest Service, Tallahassee, FL. 152 pp.
- Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K. Snow, and J. Teague. 2003. Ecological systems of the United States: A working classification of U.S. terrestrial systems. NatureServe, Arlington, VA.
- Cooper, A. S. 1974. Salt marshes. In: H. T. Odum and B. J. Copeland, editors. Coastal Ecological Systems of the United States. II. The Conservation Foundation, Washington, DC.
- Cooper, A. W., and E. D. Waits. 1973. Vegetation types in an irregularly flooded salt marsh on the North Carolina Outer Banks. *Journal of the Elisha Mitchell Scientific Society* 89:78-91.
- Coulling, P. P. 2002. A preliminary classification of tidal marsh, shrub swamp, and hardwood swamp vegetation and assorted non-tidal, chiefly non-maritime, herbaceous wetland communities of the Virginia Coastal Plain. October 2002. Virginia Department of Conservation and Recreation, Division of Natural Heritage. Natural Heritage Technical Report 02-18. 30 pp.
- Coulling, Phil. Personal communication. Vegetation Ecologist. Department of Conservation & Recreation, 217 Governor St., Richmond, VA 23219.

- Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish and Wildlife Service, Biological Service Program. FWS/OBS-79/31. Washington, DC. 103 pp.
- Daiber, F. C., L. L. Thornton, K. A. Bolster, T. G. Campbell, O. W. Crichton, G. L. Esposito, D. R. Jones, and J. M. Tyrawski. 1976. An atlas of Delaware's wetlands and estuarine resources. College of Marine Studies, University of Delaware, Newark. 528 pp.
- Dansereau, P. 1966. Studies on the vegetation of Puerto Rico. Part I. Description and integration of the plant-communities. University of Puerto Rico, Institute of Caribbean Sciences. Special Publication No. 1. Mayagüez, Puerto Rico. 287 pp.
- Diamond, D. D. 1993. Classification of the plant communities of Texas (series level). Unpublished document. Texas Natural Heritage Program, Austin. 25 pp.
- Dorroh, R. J. 1971. The vegetation of Indian shell mounds and rings of the South Carolina coast. M.S. thesis, University of South Carolina, Columbia. 68 pp.
- Duever, L. C., and S. Brinson. 1984b. Florida natural communities. Florida Game and Freshwater Fish Commission, Nongame Wildlife Program, Natural Areas Inventory, Tallahassee. 8 pp.
- EPA [Environmental Protection Agency]. 2004. Level III and IV Ecoregions of EPA Region 4. U.S. Environmental Protection Agency, National Health and Environmental Effects Research Laboratory, Western Ecology Division, Corvallis, OR. Scale 1:2,000,000.
- Eastern Ecology Working Group of NatureServe. No date. International Ecological Classification Standard: International Vegetation Classification. Terrestrial Vegetation. NatureServe, Boston, MA.
- Eleuterius, L. N. 1972. The marshes of Mississippi. *Castanea* 37:153-168.
- Eleuterius, L. N., and C. K. Eleuterius. 1979. Tide levels and salt marsh zonation. *Bulletin of Marine Science* 29:394-400.
- Eleuterius, L. N., and E. G. Otvos, Jr. 1979. Floristic and geologic aspects of Indian middens in salt marshes of Hancock County, Mississippi. *Sida* 8:102-112.
- Eleuterius, L. N., and J. D. Caldwell. 1984. Flowering phenology of tidal marsh plants in Mississippi. *Castanea* 49:172-179.
- Eyre, F. H., editor. 1980. Forest cover types of the United States and Canada. Society of American Foresters, Washington, DC. 148 pp.

- FNA Editorial Committee [Flora of North America Editorial Committee], editors. 2003. Flora of North America, north of Mexico. Volume 4. Magnoliophyta: Caryophyllidae, part 1. Oxford Univ. Press, New York. 559 pp.
- FNAI [Florida Natural Areas Inventory]. 1990. Guide to the natural communities of Florida. Florida Natural Areas Inventory and Florida Department of Natural Resources, Tallahassee. 111 pp.
- FNAI [Florida Natural Areas Inventory]. 1992a. Natural communities. Unpublished document. The Nature Conservancy, Florida Natural Areas Inventory, Tallahassee. 6 pp.
- FNAI [Florida Natural Areas Inventory]. 1992b. Natural community classification. Unpublished document. The Nature Conservancy, Florida Natural Areas Inventory, Tallahassee. 16 pp.
- Fleming, G. P. 1998. Virginia natural community framework, version January 30, 1998. Unpublished document. Virginia Department of Conservation and Recreation, Division of Natural Heritage, Richmond. 6 pp.
- Fleming, G. P. 2001a. Community types of Coastal Plain calcareous ravines in Virginia. Preliminary analysis and classification. Virginia Department of Conservation and Recreation, Division of Natural Heritage, Richmond, VA. 4 pp.
- Fleming, G. P., P. P. Coulling, D. P. Walton, K. M. McCoy, and M. R. Parrish. 2001. The natural communities of Virginia: Classification of ecological community groups. First approximation. Natural Heritage Technical Report 01-1. Virginia Department of Conservation and Recreation, Division of Natural Heritage, Richmond, VA. Unpublished report. January 2001. 76 pp.
- Fleming, G. P., and W. H. Moorhead, III. 1998. Comparative wetlands ecology study of the Great Dismal Swamp, Northwest River, and North Landing River in Virginia. Natural Heritage Technical Report 98-9. Virginia Department of Conservation and Recreation, Division of Natural Heritage, Richmond. Unpublished report submitted to the U.S. Environmental Protection Agency. 181 pp. plus appendices.
- Fleming, Gary P. Personal communication. Ecologist, Virginia Department of Conservation and Recreation, Division of Natural Heritage, Richmond, VA.
- Gaddy, L. L. 1981. Overview: Botany Bay Island, South Carolina. National Audubon Society. 20 pp.
- Gawler, S. C. 2001. Natural landscapes of Maine: Natural community profiles. Open (non-forested) types. Final review draft, July 2001. Maine Natural Areas Program. Department of Conservation. Augusta, ME.
- Godfrey, P. J. 1976. Comparative ecology of east coast barrier islands: Hydrology, soil, vegetation. Pages 5-31 in: Technical Proceedings of the 1976 Barrier Island Workshop. The Conservation Foundation. Washington, DC.

- Good, R. E. 1965. Salt marsh vegetation, Cape May, New Jersey. *Bulletin of the New Jersey Academy of Science* 10:1-11.
- Gosselink, J. G. 1984. The ecology of delta marshes of coastal Louisiana: A community profile. USDI Fish and Wildlife Service. FWS/OBS-84/09. 134 pp.
- Gould, E., and J. Ewan. 1975. Phytogeographic and ecologic relationships of the flora of Breton Island (St. Bernard Parish, Louisiana) *Tulane Studies in Zoology and Botany* 19:26-36.
- Govus, T. E. 1998. Fort Pulaski National Monument Inventory. Final report. Purchase Order # 1443PX509097564. Prepared for National Park Service, Southeast Region, Atlanta, GA. 41 pp.
- Hackney, C. T., and A. A. de la Cruz. 1981. Effects of fire on brackish marsh communities: Management implications. *Wetlands* (81-09):75-86.
- Hackney, C. T., and A. A. de la Cruz. 1982. The structure and function of brackish marshes in the north central Gulf of Mexico: A ten year case study. Pages 89-107 in: B. Gopal et al., editors. *Wetlands ecology and management*. National Institute of Ecology. International Science Publication, Jaipur, India.
- Harrison, J. W. 2001. Herbaceous tidal wetland communities of Maryland's eastern shore: Identification, assessment and monitoring. Report submitted to the U.S. EPA (Clean Water Act 1998 State Wetlands Protection Development Grant Program). Biodiversity Program, Maryland Department of Natural Resources, Wildlife and Heritage Division. 30 June 2001. [U.S. EPA Reference Wetland Natural communities of Maryland's Herbaceous Tidal Wetlands Grant #CD993724].
- Harrison, J. W. compiler. 2004. Classification of vegetation communities of Maryland: First iteration. A subset of the International Classification of Ecological Communities: Terrestrial Vegetation of the United States, NatureServe. Maryland Natural Heritage Program, Maryland Department of Natural Resources, Annapolis. 243 pp.
- Harshberger, J. W. 1914a. The vegetation of south Florida south of 27 degrees 30 minutes north, exclusive of the Florida Keys. *Transactions of the Wagner Free Institute of Science Philadelphia* 7:51-189.
- Higgins, E. A. T., R. D. Rappleye, and R. G. Brown. 1971. The flora and ecology of Assateague Island. *University of Maryland Experiment Station Bulletin A-172*. 70 pp.
- Hill, S. R. 1986. An annotated checklist of the vascular flora of Assateague Island (Maryland and Virginia). *Castanea* 5:265-305.
- Hillestad, H. O., J. R. Bozeman, A. S. Johnson, C. W. Berisford, and J. I. Richardson. 1975. The ecology of the Cumberland Island National Seashore, Camden County, Georgia. Technical Report Series No. 75-5. Georgia Marine Sciences Center, Skidway Island, GA.

- Hosier, P. E. 1975. Dunes and marsh vegetation. Pages D3-D45 in: W. M. Campbell and J. M. Dean, directors. Environmental inventory of Kiawah Island. Environmental Research Center, Inc. Columbia, SC.
- Huck, R. B. 1987. Plant communities along an edaphic continuum in a central Florida watershed. *Florida Scientist* 50:111-128.
- Jenkins, D. 1974. Natural areas of the Chesapeake Bay region: Ecological priorities. Smithsonian Institute, Ecology Program, Center for Natural Areas Ecology.
- Johnson, A. F., J. W. Muller, and K. A. Bettinger. 1990b. An assessment of Florida's remaining coastal upland natural communities: Southeast Florida. The Nature Conservancy, Florida Natural Areas Inventory, Tallahassee. 10 pp. plus appendices.
- Johnson, A. F., and J. W. Muller. 1993a. An assessment of Florida's remaining coastal upland natural communities: Final summary report. The Nature Conservancy, Florida Natural Areas Inventory, Tallahassee. 37 pp.
- Johnson, A. F., and J. W. Muller. 1993b. An assessment of Florida's remaining coastal upland natural communities: Northeast Florida. The Nature Conservancy, Florida Natural Areas Inventory, Tallahassee. 10 pp. plus appendices.
- Johnson, A. F., and M. G. Barbour. 1990. Dunes and maritime forests. Pages 429-480 in: R. L. Myers and J. J. Ewel, editors. *Ecosystems of Florida*. University of Central Florida Press, Orlando.
- Johnson, A. S., H. O. Hillestad, S. F. Shanholtzer, and G. F. Shanholtzer. 1974. An ecological survey of the coastal region of Georgia. USDI National Park Service. Science Monograph Series, No. 3. 233 pp. plus maps.
- Johnston, M. C. 1952. Vegetation of eastern Cameron County, Texas. M.S. thesis, University of Texas, Austin. 127 pp.
- Kartesz, J. T. 1999. A synonymized checklist and atlas with biological attributes for the vascular flora of the United States, Canada, and Greenland. First edition. In: J. T. Kartesz and C. A. Meacham. *Synthesis of the North American Flora*, Version 1.0. North Carolina Botanical Garden, Chapel Hill, NC.
- Keys, J. E., Jr., C. A. Carpenter, S. L. Hooks, F. G. Koenig, W. H. McNab, W. E. Russell, and M-L. Smith. 1995. Ecological units of the eastern United States - first approximation (map and booklet of map unit tables). Presentation scale 1:3,500,000, colored. USDA Forest Service, Atlanta, GA.
- Klemas, V., F. C. Daiber, D. S. Bartlett, O. W. Crichton, and A. O. Fornes. 1973. Coastal vegetation of Delaware. University of Delaware, College of Marine Studies. 27 pp.

- Klotz, L. H. 1986. The vascular flora of Wallops Island and Wallops Mainland, Virginia. *Castanea* 51:306-326.
- Kruczynski, W. L., C. B. Subrahmanyam, and S. H. Drake. 1978. Studies on the plant community of a north Florida salt marsh. *Bulletin of Marine Science* 28:707-715.
- Kurz, H. 1942. Florida dunes and scrub, vegetation and geology. Florida Department of Conservation, Geologic Survey. Geologic Survey Bulletin No. 23. Tallahassee. 154 pp.
- Kurz, H., and K. Wagner. 1957. Tidal marshes of the Gulf and Atlantic coasts of northern Florida and Charleston, South Carolina. Florida State University Studies, No. 24. Tallahassee, FL. 168 pp.
- Laessle, A. M. 1942. Plant communities of the Welaka area. University of Florida Biological Sciences Series No. 4. 143 pp.
- LeGrand, H. E., Jr., C. C. Frost, and J. O. Fussell, III. 1992. Regional inventory for critical natural areas, wetland ecosystems, and endangered species habitats of the Albemarle-Pamlico estuarine region: Phase 2. North Carolina Department of Environment, Health, and Natural Resources, Division of Parks and Recreation, Natural Heritage Program, Raleigh. 506 pp.
- Lea, C. 2002b. Vegetation classification of Assateague Island National Seashore, Addendum (version 2) - March 1, 2002. Unpublished draft. National Park Service. 31 pp.
- Lemaire, R. J. 1961. A preliminary annotated checklist of the vascular plants of the marshes and included higher lands of St. Bernard Parish, Louisiana. *Proceedings of the Louisiana Academy of Sciences* 24:56-70.
- Lewis, I. F. 1917. The vegetation of Shackleford Bank. North Carolina Geologic and Economic Survey. Economic Paper No. 46. 32 pp.
- Lynch, J. J. 1941. The place of burning in management of Gulf Coast wildlife refuges. *Journal of Wildlife Management* 5:454-457.
- Martin, W. E. 1959b. The vegetation of Island Beach State Park, New Jersey. *Ecological Monographs* 29:1-46.
- McAlister, W. H., and M. K. McAlister. 1993. A naturalist's guide: Matagorda Island. University of Texas Press, Austin. 354 pp.
- McAlister, W. H., and M. K. McAlister. 1995. A naturalist's guide: Aransas. University of Texas Press, Austin. 354 pp.
- Metzler, K. J., and J. Barrett. 1992. Connecticut community classification. Unpublished draft. Connecticut Department of Environmental Protection, Natural Resources Center, Natural Diversity Database, Hartford.

- Metzler, K. J., and J. P. Barrett. 2001. Vegetation classification for Connecticut. Draft 5/21/2001. Connecticut Department of Environmental Protection, Natural Resources Center, Natural Diversity Database, Hartford.
- Montague, C. L., and R. G. Wiegert. 1990. Salt marshes. Pages 481-516 in: R. L. Myers and J. J. Ewel, editors. *Ecosystems of Florida*. University of Central Florida Press, Orlando.
- Montz, G. N. 1981. Annotated checklist of plants of the coastal beaches, islands and barrier islands of Louisiana. U.S. Army Corps of Engineers, New Orleans District.
- NatureServe Ecology - Southeastern United States. No date. Unpublished data. NatureServe, Durham, NC.
- Nelson, J. B. 1986. The natural communities of South Carolina: Initial classification and description. South Carolina Wildlife and Marine Resources Department, Division of Wildlife and Freshwater Fisheries, Columbia, SC. 55 pp.
- Nicholson, W. R., and R. D. Van Deusen. 1954. Marshes of Maryland. Maryland Game and Inland Fish Commission. Resource Study Report No. 6. Baltimore, MD.
- Nordman, Carl W. Personal communication. Regional Ecologist. NatureServe, Southern Resource Office, Durham, NC.
- Odum, W. E. 1988. Comparative ecology of tidal freshwater and salt marshes. *Annual Review of Ecological Systems* 19:147-176.
- Odum, W. E., T. J. Smith, III, J. K. Hoover, and C. C. McIvor. 1984. The ecology of tidal freshwater marshes of the United States east coast: A community profile. USDI Fish & Wildlife Service. FWS/OBS-83/17. 176 pp.
- Odum, W. E., and T. J. Smith. 1981. Ecology of tidal, low salinity ecosystems. Pages 36-44 in: R. C. Carey, P. S. Markovits, and J. B. Kirkwood, editors. *Proceedings of the workshop on coastal ecosystems of the southeastern United States*. USDI Fish & Wildlife Service, Office of Biological Services. Report No. FWS/OBS-80/59. Washington, DC.
- Olmsted, I. C., L. L. Loope, and R. P. Russell. 1981. Vegetation of the southern coastal region of Everglades National Park between Flamingo and Joe Bay. USDI National Park Service, Everglades National Park, Southern Florida Research Center. Report T-620. Homestead, FL. 18 pp.
- Oosting, H. J., and W. D. Billings. 1942. Factors affecting vegetational zonation on coastal dunes. *Ecology* 23:131-142.
- Peet, R. K., T. R. Wentworth, M. P. Schafale, and A.S. Weakley. 2002. Unpublished data of the North Carolina Vegetation Survey. University of North Carolina, Chapel Hill.

- Penfound, W. T. 1952. Southern swamps and marshes. *Botanical Review* 7:413-446.
- Platt, W. J., and M. W. Schwartz. 1990. Temperate hardwood forests. Pages 194-229 in: R. L. Myers and J. J. Ewel, editors. *Ecosystems of Florida*. University of Central Florida Press, Orlando.
- Rayner, D. A. 1984. Inventory of natural areas in Colleton County, South Carolina. Pages 61-93 in: *Inventory of botanical natural areas in Colleton, Dorchester, Horry, and Jasper counties, South Carolina*. South Carolina Wildlife and Marine Resources Department, Columbia.
- Rayner, D. A., and W. T. Batson. 1976. Maritime closed dunes vegetation in South Carolina. *Castanea* 41:58-70.
- Reschke, C. 1990. *Ecological communities of New York State*. New York Natural Heritage Program. New York State Department of Environmental Conservation. Latham, NY. 96 pp.
- SCWMRD [South Carolina Wildlife and Marine Resources Department]. 1984. *Inventory of botanical natural areas in Colleton, Dorchester, Horry, and Jasper counties, South Carolina*. South Carolina Wildlife and Marine Resources Department, Columbia. 144 pp.
- Sandifer, P. A., J. V. Miglarese, D. R. Calder, J. J. Manzi, and L. A. Barclay, editors. 1980. *Ecological characterization of the Sea Island coastal region of South Carolina and Georgia*. Volume III. Biological features of the characterization area. USDI Fish & Wildlife Service, Office of Biological Service. FWS/OBS-79/42. Washington, DC. 620 pp.
- Schafale, M. 2000. Fourth approximation guide. Coastal Plain. January 2000 draft. North Carolina Natural Heritage Program, Raleigh.
- Schafale, M. 2003b. Fourth approximation guide. Coastal Plain communities. March 2003 draft. North Carolina Natural Heritage Program, Raleigh.
- Schafale, M. P., and A. S. Weakley. 1990. *Classification of the natural communities of North Carolina*. Third approximation. North Carolina Department of Environment, Health, and Natural Resources, Division of Parks and Recreation, Natural Heritage Program, Raleigh. 325 pp.
- Schafale, Mike P. Personal communication. Ecologist, North Carolina Department of Environment, Health, and Natural Resources, Division of Parks and Recreation, Natural Heritage Program, Raleigh.
- Schotz, Al. Personal communication. Community Ecologist. Alabama Natural Heritage Program. Huntingdon College, Massey Hall, 1500 East Fairview Avenue, Montgomery, AL 36106-2148.

- Sharitz, R. R. 1975. Forest communities of Kiawah Island. Pages F1-F39 in: W. M. Campbell, J. M. Dean, and W. D. Chamberlain, editors. Environmental inventory of Kiawah Island. Environmental Research Center, Inc., Columbia, SC.
- Smith, L. M., compiler. 1996a. Natural plant communities in Louisiana currently recognized by the Louisiana Natural Heritage Program. Unpublished document. Louisiana Department of Wildlife and Fisheries, Natural Heritage Program, Baton Rouge. 2 pp.
- Smith, Latimore M. Personal communication. Natural Heritage Program Ecologist. Louisiana Department of Wildlife and Fisheries, Natural Heritage Program, Baton Rouge.
- Sneddon, L., M. Anderson, and K. Metzler. 1996. Community alliances and elements of the Eastern Region. Unpublished report. The Nature Conservancy, Eastern Heritage Task Force, Boston, MA. 235 pp.
- Southeastern Ecology Working Group of NatureServe. No date. International Ecological Classification Standard: International Vegetation Classification. Terrestrial Vegetation. NatureServe, Durham, NC.
- Southeastern Wildlife Services. 1981. An inventory of the vegetation and vertebrate fauna of Fort Pulaski National Monument. 2 volumes. Report to the National Park Service, Fort Pulaski National Monument, Savannah, GA.
- Sperduto, D. D. 2000b. A classification of wetland natural communities in New Hampshire. New Hampshire Natural Heritage Inventory, Department of Resources and Economic Development, Division of Forests and Lands. Concord, NH. 156 pp.
- Stalter, R. 1973a. The flora of Turtle Island, Jasper Co., South Carolina. *Castanea* 38:35-37.
- Stalter, R. 1973b. Factors influencing the distribution of vegetation of the Cooper River Estuary. *Castanea* 38:18-24.
- Stout, J. P. 1984. The ecology of irregularly flooded salt marshes of the northeastern Gulf of Mexico: A community profile. USDI Fish and Wildlife Service, Minerals Management Service. Biological Report 85 (7.1). 98 pp.
- Swain, P. C., and J. B. Kearsley. 2001. Classification of natural communities of Massachusetts. September 2001 draft. Natural Heritage and Endangered Species Program, Massachusetts Division of Fisheries and Wildlife. Westborough, MA.
- Tiner, R. W., Jr. 1977. An inventory of South Carolina's coastal marshes. South Carolina Marine Resources Center. Technical Report 23. Columbia, SC. 33 pp.
- Tiner, R. W., Jr. 1985b. Wetlands of New Jersey. USDI Fish & Wildlife Service, National Wetlands Inventory, Newton Corner, MA. 117 pp.

- VDNH [Virginia Division of Natural Heritage]. 2003. The natural communities of Virginia: Hierarchical classification of community types. Unpublished document, working list of November 2003. Virginia Department of Conservation and Recreation, Division of Natural Heritage, Ecology Group, Richmond.
- Weakley, A. S. 2007. Flora of the Carolinas, Virginia, Georgia, and surrounding areas. Unpublished working draft of 11 January 2007. University of North Carolina Herbarium (NCU), North Carolina Botanical Garden, University of North Carolina, Chapel Hill. [<http://www.herbarium.unc.edu/flora.htm>]
- Webber, E. E. 1967. Blue-green algae from a Massachusetts salt marsh. *Bulletin of the Torrey Botanical Club* 94:99-106.
- Wentworth, T. R., M. P. Schafale, A. S. Weakley, R. K. Peet, P. S. White, and C. C. Frost. 1993. A preliminary classification of North Carolina barrier island forests. Pages 31-46 in: C. A. Cole and F. K. Turner, editors. *Barrier island ecology of the mid-Atlantic Coast: A symposium*. USDI National Park Service. Technical Report NPS/SERCAHA/NRTR-93/04. Atlanta, GA.
- Wharton, C. H. 1978. The natural environments of Georgia. Georgia Department of Natural Resources, Atlanta. 227 pp.
- Wieland, R. G. 1994a. Marine and estuarine habitat types and associated ecological communities of the Mississippi Coast. Mississippi Department of Wildlife, Fisheries, and Parks. Museum of Natural Science. Museum Technical Report 25. Jackson, MS. 270 pp.
- Wieland, R. G. 1994b. Mississippi Natural Heritage Program: Ecological communities. Unpublished document. Mississippi Department of Wildlife, Fisheries, and Parks, Museum of Natural Science, Natural Heritage Program, Jackson, MS. 7 pp.
- Wieland, R. G. 2000b. Ecological communities of Mississippi: Mississippi Natural Heritage Program. Unpublished document. Mississippi Department of Wildlife, Fisheries, and Parks, Museum of Natural Science, Natural Heritage Program, Jackson, MS. 8 pp.
- Wolfe, S. H., editor. 1990. An ecological characterization of the Florida Springs Coast: Pithlachascotee to Waccasassa rivers. USDI Fish & Wildlife Service, Biological Report 90(21). Slidell, LA. 323 pp.

Appendix C: Field Key to Vegetation Types Draft Key to the National Vegetation Classification (NVC) Associations that Occur or Potentially occur at Fort Pulaski and Fort Sumter National Monuments

May 2008, updated August 2012
NatureServe, Durham, NC/SECN Staff

Associations which are documented from Fort Pulaski and/or Fort Sumter National Monuments are in **bold type**. For each association, the common name is given, with the Element Code in parentheses (CEGL00####) and the Alliance Code in brackets [A.###].

KEY TO KEYS

1. Vegetation dominated by trees, forests or woodlands.**KEY A – FORESTS & WOODLANDS**
1. Vegetation not dominated by trees, trees absent or very sparse..... 2
2. Shrub dominated and vine dominated areas, with few or no trees. This includes areas that may be dominated by shrub species of very large size (technically tree size of 4cm in DBH or larger, and over 1.5m in height). Shrubs of large size may include *Morella cerifera*, *Juniperus virginiana* var. *silicicola*, *Prunus caroliniana*, *Zanthoxylon clava-herculis*, and *Ilex vomitoria*.**KEY B – SHRUBLANDS**
2. Vegetation dominated by herbaceous plants, including sparse vegetation of herbaceous plants, such as may occur between the foredunes and the beach..... **KEY C – GRAMINOID VEGETATION**

KEY A –FORESTS & WOODLANDS

1. Forest dominated by *Pinus elliottii* var. *elliottii* (Slash Pine).....**Slash Pine Plantation (CEGL007170) [A.95]**
1. Forest not dominated by *Pinus elliottii* var. *elliottii* (Slash Pine)..... 2
2. Forest dominated by *Triadica sebifera* (Tallowtree) 3
2. Forest dominated by evergreen or deciduous trees, some combination of *Quercus virginiana* (live oak), *Sabal palmetto* (Cabbage palmetto), and/or *Juniperus virginiana* var. *silicicola* (southern redcedar) 4
3. Upland forest dominated by *Triadica sebifera* (Tallowtree) (FOPU only)
Tallowtree Upland Forest (CEGL007037) [A.257]
3. Wetland forest dominated by *Triadica sebifera* (Tallowtree). *Salix caroliniana* (coastal plain willow) may be sub-dominant (FOSU ONLY) ***Triadica sebifera* Seasonally Flooded Forest Alliance [A.335]**
4. Open stands dominated by *Celtis laevigata* (sugarberry), *Ilex vomitoria* (yaupon), *Juniperus virginiana* var. *silicicola* (southern redcedar), *Sabal palmetto* (cabbage palmetto), and *Triadica sebifera* (tallowtree) (exotic) **Cabbage Palmetto Woodland (CEGL003526) [A.481]**
4. Closed forest stands dominated by *Juniperus virginiana* var. *silicicola* (southern redcedar) 5

5. Shell midden (on McQueens Island) dominated by *Juniperus virginiana* var. *silicicola* (southern redcedar). Additional tree species may include *Forestiera segregata* (Florida swampprivet), *Sabal palmetto* (cabbage palmetto), *Ilex vomitoria* (yaupon), *Celtis laevigata* (sugarberry)
 **South Atlantic Coastal Shell Midden Woodland (CEGL003525) [A.479]**
5. Hammock forest dominated by *Juniperus virginiana* var. *silicicola* and *Quercus virginiana* with *Sabal palmetto*, *Ulmus americana*, *Ilex vomitoria*, or *Pinus taeda* 6
6. Hammock forest (on Cockspur Island) mostly on old dredge spoil with dense (70-90% cover) tree canopy, 10-20 m tall, dominated by *Juniperus virginiana* var. *silicicola* (southern redcedar) along with *Sabal palmetto* (cabbage palmetto) and *Ulmus americana* (American elm) or *Ilex vomitoria* (yaupon)
 **Cedar - Live Oak - Cabbage Palmetto Marsh Hammock (CEGL007813) [A.55]**
6. Hammock forest, with species composition similar to the Cedar- Live Oak – Cabbage Marsh Hammock, but with an overstory containing *Pinus taeda* (loblolly pine).....***Quercus virginiana* – (*Sabal palmetto*) Forest Alliance [A.55]**

KEY B –SHRUBLANDS

1. Tidally influenced sparse to moderate (20 - 40% cover) dwarf-shrubland (generally < 1 m tall) dominated by *Batis maritima* (turtleweed) and/or *Sarcocornia pacifica* (Pacific swampfire) and may include lesser amounts of *Borrchia frutescens* (bushy seaside tansy or seaside oxeye). These sites include open salt flats. Other halophytes may include *Suaeda linearis* (annual seepweed), *Sesuvium portulacastrum* (shoreline seapurslane), and *Limonium carolinianum* (Carolina sealavender) 2
1. Shrublands (tidal and non-tidal), not dominated by *Batis maritima* (turtleweed) and/or *Sarcocornia pacifica* (Pacific swampfire) 3
2. Tidally influenced sparse to moderate (20 - 40% cover) dwarf-shrubland (generally < 0.5 m tall) dominated by *Batis maritima* (turtleweed) and additionally only includes lesser amounts of *Borrchia frutescens* (bushy seaside tansy or seaside oxeye) and *Sarcocornia pacifica* (Pacific swampfire)
 **Saltwort - Swampfire Dwarf-shrubland (CEGL003956) [A.1111]**
2. Tidally influenced sparse (~20% or more) dwarf-shrubland is dominated by *Sarcocornia pacifica* (Pacific swampfire). Other plants may include *Batis maritima* (turtleweed) and *Distichlis spicata* (inland saltgrass) and dwarfed *Spartina alterniflora* (smooth cordgrass).....
Salt Flat (Swampfire Type) (CEGL002278) [A.1705]
3. Shrublands which are generally not tidally influenced, dominated by *Morella cerifera* (Wax myrtle), or *Prunus caroliniana* (Carolina laurelcherry) along with lesser amounts of *Zanthoxylum clava-herculis* (Hercules' club), *Celtis laevigata* (sugarberry), and *Sabal palmetto* (cabbage palmetto). Vines include may include *Parthenocissus quinquefolia* (Virginia creeper), *Smilax auriculata* (earleaf greenbrier), and *Smilax bona-nox* (saw greenbrier) 4
3. Tidal fringe shrublands, generally at upper edge of tidal influence, dominated by *Borrchia frutescens* (bushy seaside tansy or seaside oxeye), or some combination of *Baccharis halimifolia* (Groundsel tree), *Iva frutescens* (maritime marsh-elder), and *Morella cerifera* (wax myrtle), and *Ilex vomitoria* (yaupon) .. 5

4. *Morella cerifera* (wax myrtle) dominates the shrub layers; additional shrubs may include *Ilex vomitoria* (Yaupon), and *Juniperus virginiana* var. *silicicola* (southern redcedar), *Celtis laevigata* (sugarberry), *Sabal palmetto* (cabbage palmetto), and *Rubus trivialis* (southern dewberry). There may be a moderate emergent tree layer of *Celtis laevigata* and/or *Morella cerifera*. Vines include *Ampelopsis arborea* (peppervine), *Parthenocissus quinquefolia* (Virginia creeper), *Smilax auriculata* (earleaf greenbrier), and *Smilax bona-nox* (saw greenbrier).**Atlantic Coast Interdune Swale (CEGL003839) [A.1906]**

4. Vegetation is dominated by *Morella cerifera* (wax myrtle) and *Prunus caroliniana* (Carolina laurelcherry) along with lesser amounts of *Zanthoxylum clava-herculis* (Hercules' club); and *Rubus trivialis* (southern dewberry). Some trees of *Celtis laevigata* (sugarberry) and tree sized *Prunus caroliniana* (Carolina laurelcherry) may be present. Vines include *Vitis rotundifolia* (muscadine), *Parthenocissus quinquefolia* (Virginia creeper), *Smilax auriculata* (earleaf greenbrier), and *Smilax bona-nox* (saw greenbrier)
Wax myrtle – Carolina Laurel Cherry – Toothache-tree Shrubland (CEGL004784) [A.785]

5. Vegetation is a tidal shrubland dominated by a short shrub layer (0.5-1 m tall) of Seaside Oxeye (*Borrchia frutescens*). The herbaceous layer, when present, is dominated by *Distichlis spicata* (inland saltgrass). Additional scattered emergent shrubs may include *Juniperus virginiana* var. *silicicola* (southern redcedar), *Ilex vomitoria* (yaupon), *Iva frutescens* (Jesuit's bark), and *Sideroxylon tenax* (tough bully) (globally and state-vulnerable). ...**Seaside Oxeye Tidal Shrub Flat (CEGL003924) [A.1026]**

5. Vegetation on interdunal swales and/or salt flats that are subject to irregular storm and tidal flooding. Vegetation is not dominated by *Borrchia frutescens* (bushy seaside tansy or seaside oxeye), but characterized by some combination of *Baccharis halimifolia* (Groundsel tree), *Iva frutescens* (Maritime Marsh-elder), and *Morella cerifera* (wax myrtle) and *Ilex vomitoria* (yaupon). Other species present may include *Baccharis angustifolia* (saltwater false willow), *Limonium carolinianum* (Carolina sealavender), and *Fimbristylis castanea* (marsh fimbry).....**Coastal Salt Shrub Thicket (CEGL003920) [A.1023]**

KEY C – GRAMINOID VEGETATION

1. Sparse vegetation of dunes and fore dunes and interdunal swales, generally dominated by *Uniola paniculata* (sea oats) or *Heterotheca subaxillaris* (camphorweed). Vines (5% - 40% cover) are dominated by *Smilax auriculata* (earleaf greenbrier), *Strophostyles helvula* (trailing wild bean), or *Hydrocotyle bonariensis* (beach pennywort). *Spartina patens* (saltmeadow cordgrass), *Cenchrus incertus* (coast sandspur), *Panicum amarum* (bitter panicgrass), *Ipomoea imperati* (beach morning-glory), and *Oenothera drummondii* (beach evening primrose) may also be common
***Uniola paniculata* Temperate Herbaceous Alliance [A.1119]**

1. Tidal salt marshes, including the higher areas with less tidal influence sometimes called salt flats 2

2. High salt marsh, with vegetation dominated by *Juncus roemerianus* (Black Needlerush)
..... **Needlerush High Marsh (CEGL004186) [A.1475]**

2. Salt marsh dominated by cordgrasses (*Spartina* spp.) or *Distichlis spicata* (inland saltgrass)..... 3

3. Tidal salt marsh vegetation, dominated by *Spartina alterniflora* (Salt marsh Cordgrass). Very extensive on McQueens Island **Southern Atlantic Coast Salt Marsh (CEGL004191)**

3. Vegetation dominated by grasses other than *Spartina alterniflora* (Salt marsh Cordgrass)..... 4

4. Grassland vegetation, dominated by *Spartina patens* or *Spartina cynosuroides* 5
4. Grassland vegetation, dominated by *Phragmites australis* (common reed), generally in brackish marshes **Reed Tidal Marsh (CEGL004187)**
5. Vegetation dominated or codominated *Spartina patens* (saltmeadow cordgrass), *Distichlis spicata* (inland saltgrass), and/or *Borrchia frutescens* (bushy seaside tansy or seaside oxeye) **Mid-Atlantic High Salt Marsh (CEGL004197) [A.1481]**
5. Vegetation narrow, almost pure stands of *Spartina cynosuroides* (big cordgrass) along tidal creeks, ditches and sloughs **Atlantic Giant Cordgrass Marsh (CEGL004195) [A.1480]**

Appendix D: Photo-Interpretation Guide

Photo-Interpretation Guide to the Vegetated Map Classes of Fort Pulaski National Monument

I. Forest

Map Class 1: Cedar – Live Oak – Cabbage Palmetto Marsh Hammock

Map Class 2: Slash Pine Plantation

Map Class 3: Tallowtree Upland Forest

II. Woodland

Map Class 4: South Atlantic Coastal Shell Midden Woodland

Map Class 5: Cabbage Palmetto Woodland

III. Shrubland

Map Class 6: Seaside Oxeye Tidal Shrub Flat

IV. Dwarf-Shrubland

Map Class 7: Salt Flat (Swampfire Type)

Map Class 8: Saltwort – Swampfire Dwarf-Shrubland

V. Herbaceous

Map Class 9: Needlerush High Marsh

Map Class 10: Southern Atlantic Coast Salt Marsh

Map Class 11: Mid-Atlantic High Salt Marsh

Map Class 12: Atlantic Giant Cordgrass Marsh

I. Forest

MAP CLASS 1

Cedar – Live Oak – Cabbage Palmetto Marsh Hammock:



Association:

Juniperus virginiana var. *silicicola* – (*Quercus virginiana*, *Sabal palmetto*) Forest

Characteristic species:

- *Juniperus virginiana* var. *silicicola*
- *Celtis laevigata*
- *Ilex vomitoria*
- *Quercus virginiana*
- *Sabal palmetto*
- *Morella cerifera*

Map Class Statistics

Frequency: 21 polygons

Total area: 52.9 ha

Average polygon size: 2.5 ha

Producer's accuracy: 56%

User's accuracy: 89%

Interpretation

The Cedar – Live Oak – Cabbage Palmetto Marsh Hammock has a relatively closed canopy on the imagery (e.g., not much spacing between the canopy trees) and contains several species that

appear different shades of pink/red/purple on the imagery. The Cabbage Palm has the smallest crown size of the canopy trees present and is light pink with a white speckled appearance. Coastal redcedar is a dark red/purple color and has a fairly large, usually assymetric crown. Live oak has a rather large assymetric crown and is bright red. All three of these species are usually visible within this association.



Example of photo signature for Cedar – Live Oak – Cabbage Palmetto Marsh Hammock

MAP CLASS 2

Slash Pine Plantation:



Association:

Pinus elliottii var. *elliottii* Planted Forest

Characteristic species

- *Pinus elliottii* var. *elliottii*
- *Sabal palmetto*

Map Class Statistics

Frequency: 2 polygons

Total area: 1.9 ha

Average polygon size: 0.97 ha

Producer's accuracy: 100%

User's accuracy: 100%

Interpretation

Slash pine has a relatively small, circular crown and appears dark red/magenta on the imagery.



Example of photo signature for Slash Pine Planted Forest

MAP CLASS 3

Tallowtree Upland Forest:



Association:

Triadica sebifera Upland Forest

Characteristic species

- *Triadica sebifera*
- *Ilex vomitoria*
- *Sabal palmetto*

Map Class Statistics

Frequency: 8 polygons

Total area: 4.3 ha


Average polygon size: 0.53 ha

Producer's accuracy: 36%

User's accuracy: 100%

Interpretation

Tallowtree has a distinctive pink color on the imagery and a circular crown of small to moderate size.



Example of photo signature for Tallowtree Upland Forest (inside yellow polygon).

II. Woodland

MAP CLASS 4

South Atlantic Coastal Shell Midden Woodland:



Characteristic species

- *Juniperus virginiana* var. *silicicola*
- *Forestiera segregata*
- *Sabal palmetto*
- *Ilex vomitoria*
- *Celtis laevigata*

Map Class Statistics

Frequency: 5 polygons

Total area: 3.95 ha

Average polygon size: 0.79 ha

Producer's accuracy: 99.1%

User's accuracy: 100%

Interpretation

From aerial view, there are four elliptical mounds composed of this association that are surrounded by salt marsh. These mounds are fairly easy to decipher due to their unique shape and position within the landscape. The canopy appears mainly dark red in color, due to the large presence of *Juniperus vir. var. sil.* dominant in the canopy.



Example of photo signature for South Atlantic Coastal Shell Midden Woodland.

MAP CLASS 5

Cabbage Palmetto Woodland:



Association

Sabal palmetto – (*Juniperus virginiana* var. *silicicola*) Woodland

Abundant species

- *Juniperus virginiana* var. *silicicola*
- *Celtis laevigata*
- *Triadica sebifera*
- *Ilex vomitoria*
- *Sabal palmetto*

Map Class Statistics

Frequency: 12 polygons

Total area: 13.7 ha

Average polygon size: 1.1 ha

Producer's accuracy: 35.2%

User's accuracy: 36%

Interpretation

Cabbage Palmetto Woodland has a somewhat similar photo signature to that of the Cedar – Live Oak – Cabbage Palmetto Marsh Hammock, with two main differences being that the Cabbage Palmetto Woodland has a more open canopy and generally a larger presence of *Sabal palmetto* in the canopy than the Cedar – Live Oak – Cabbage Palmetto Marsh Hammock. *Sabal palmetto* has a very distinctive appearance of a small, circular crown that is pink in color with an appearance

of white speckles. Additionally, *Juniperus virginiana* var. *silicicola* is also usually prominent in the canopy, and it has a dark purple color and large, assymetric crown.



Example of photo signature for Cabbage Palmetto Woodland.

III. Shrubland

MAP CLASS 6

Seaside Oxeye Tidal Shrub Flat



Association

Borrichia frutescens/ (*Spartina patens*, *Juncus roemerianus*) Shrubland

Characteristic species

- *Borrichia frutescens*
- *Distichlis spicata*
- *Sarcocornia pacifica*

Map Class Statistics

Frequency: 16 polygons

Total area: 22.1 ha


Average polygon size: 1.4 ha

Producer's accuracy: 100%

User's accuracy: 68%

Interpretation

This community is typically dominated by *Borrichia frutescens* and appears pink with a smooth to moderately rough texture on the imagery. At Fort Pulaski, this community is frequently found along tidal fringe areas.



Example of photo signature of Seaside Oxeye Tidal Shrub Flat (inside blue polygon).

IV. Dwarf-Shrubland

MAP CLASS 7

Salt Flat (Swampfire Type)



Association

Sarcocornia pacifica – (*Batis maritima*, *Distichlis spicata*) Dwarf-shrubland

Characteristic species

- *Sarcocornia pacifica*
- *Spartina alterniflora*

Map Class Statistics

Frequency: 16 polygons

Total area: 58.7 ha

Average polygon size: 3.7 ha

Producer's accuracy: 8.3%

User's accuracy: 33%

Interpretation

The appearance of this association is gray/pale pink color (a mixture of these two colors is present) and is very smooth in texture. Within the landscape, this association occurs on lower tidal flats where high concentrations of salt accumulate and produce hypersaline conditions.



Example of the photo signature for the Salt Flat (Swampfire Type) Dwarf-shrubland

MAP CLASS 8

Saltwort - Swampfire Dwarf-shrubland



Association

Batis maritima-*Sarcocornia pacifica* Dwarf-shrubland

Characteristic species

- *Batis maritima*
- *Sarcocornia pacifica*
- *Borrchia frutescens*

Map Class Statistics

Frequency: 4 polygons

Total area: 20.2 ha


Average polygon size: 5.1ha

Producer's accuracy: 12.5%

User's accuracy: 80%

Interpretation

The appearance of this association is gray/bright pink color (a mixture of these two colors is present) and is very smooth in texture. The photo signature for this association differs slightly from the Salt Flat (Swampfire Type) association because the pink is brighter in color. Within the landscape, this association occurs on lower tidal flats where high concentrations of salt accumulate and produce hypersaline conditions. This association often occurs alongside the Salt Flat (Swampfire Type).



Example of the photo signature for the Saltwort-Swampfire Dwarf-shrubland (inside blue polygon)

V. Herbaceous

MAP CLASS 9

Needlerush High Marsh



Association

Juncus roemerianus Herbaceous Vegetation

Abundant species

- *Juncus roemerianus*

Map Class Statistics

Frequency: 20 polygons

Total area: 91.4 ha


Average polygon size: 4.6 ha

Producer's accuracy: 0%

User's accuracy: 0%

Interpretation

Needlerush High Marsh has a generally smooth, velvety appearance on the imagery and is a dark pink to brown color. Often stands of Needlerush form circular patterns that are distinguishable on the imagery. Needlerush High Marsh is slightly less smooth in texture and darker than the Southern Atlantic Coast Salt Marsh.



Example of photo signature of Needle Rush High Marsh (inside blue polygon)

MAP CLASS 10

Southern Atlantic Coast Salt Marsh



Association

Spartina alterniflora Carolinian Zone Herbaceous Vegetation

Abundant species

- *Spartina alterniflora*

Map Class Statistics

Frequency: 57 polygons

Total area: 1,488.3 ha

Average polygon size: 26 ha

Producer's accuracy: 99.6%

User's accuracy: 93%

Interpretation

Southern Atlantic Coast Salt Marsh appears very smooth in texture and is pinkish/gray in color. This community occupies the most area within Fort Pulaski National Monument boundaries and occurs on flats located between mean high and low tides that are flooded regularly twice a day.



Example of photo signature of Southern Atlantic Coast Salt Marsh.

MAP CLASS 11

Mid-Atlantic High Salt Marsh



Association

Spartina patens – *Distichlis spicata* – (*Juncus roemerianus*) Herbaceous Vegetation

Abundant species

- *Spartina patens*
- *Distichlis spicata*
- *Borrichia frutescens*

Map Class Statistics

Frequency: 2 polygons

Total area: 2.0 ha

Average polygon size: 1.0 ha

Producer's accuracy: 100%

User's accuracy: 60%

Interpretation

Mid-Atlantic High Salt Marsh usually appears bright pink on the imagery and is smooth to somewhat rough in texture.



Example of photo signature of Mid-Atlantic High Salt Marsh (inside blue polygon).

MAP CLASS 12

Atlantic Giant Cordgrass Marsh



Association

Spartina cynosuroides Tidal Herbaceous Vegetation

Abundant species

- *Spartina cynosuroides*

Map Class Statistics

Frequency: 14 polygons

Total area: 75.1 ha


Average polygon size: 5.36 ha

Producer's accuracy: 0%

User's accuracy: 0%

Interpretation

The Giant Atlantic Cordgrass Marsh appears dark pink to bright red on the imagery and has a mottled appearance. It is typically located in narrow bands associated with tidal ditches or along fringe areas between tidal herbaceous vegetation and tidal shrublands.



Example of photo signature of Atlantic Giant Cordgrass Marsh (inside blue polygon)

Appendix E: Field Reconnaissance Form

FORT PULASKI NATIONAL MONUMENT FIELD RECONNAISSANCE SHEET

I. LOCATION DATA

1. Point #: _____ 2. Date: _____ 3. Observers: _____
4. Photo (s) #: _____
5. Coordinates: X: _____ Y: _____
6. GPS Error: _____ (m) 7. Number of satellites: _____ 8. Elevation: _____

II. ENVIRONMENTAL/ SITE DESCRIPTION

1. Environmental Comments (*site-specific information (if applicable) such as hydrology, topographic position, unvegetated surfaces, recent disturbances, etc.*)

III. VEGETATION DESCRIPTION

Leaf phenology (of dominant stratum)

- Trees and Shrubs
☐ Evergreen
☐ Cold-deciduous
☐ Drought-deciduous
☐ Mixed evergreen - cold-deciduous
☐ Mixed evergreen - drought-deciduous

- Herbs
☐ Annual
☐ Perennial

Leaf Type (of dominant stratum)

- ☐ Broad-leaved
☐ Needle-leaved
☐ Mixed broad-leaved/Needle-leaved
☐ Microphyllous
☐ Graminoid
☐ Forb
☐ Pteridophyte

Physiognomic class

- ☐ Forest
☐ Woodland
☐ Shrubland
☐ Dwarf Shrubland
☐ Herbaceous
☐ Nonvascular
☐ Sparsely Vegetated

IV. VEGETATION COMPOSITION (list **dominant species only** and indicate which stratum it occurs in by checking the appropriate stratum next to the species name)

[illegible]

V. VEGETATION ASSOCIATION

1. Primary and secondary (if there is a close call or multiple associations within the same polygon) vegetation classification(s) within radius of observation point, based on field key:

Appendix F: Accuracy Assessment Form

I. LOCATION DATA:

1. AA Point Code: _____ 2. Date: _____

3. Observers: _____

4. Photo (s) #: _____

5. Coordinates: X: _____ Y: _____

6. Estimated Accuracy (m): _____ 7. Number of Satellites: _____

II. ENVIRONMENTAL/ SITE DESCRIPTION:

1. Environmental Comments (site specific information such as hydrology, topographic position, unvegetated surfaces, recent disturbances, etc.)

III. VEGETATION DESCRIPTION:

Leaf phenology (of dominant stratum)	Leaf Type (of dominant stratum)	Physiognomic class
Trees and Shrubs:	___ Broad-leaved	___ Forest
___ Evergreen	___ Needle-leaved	___ Woodland
___ Cold-deciduous	___ Mixed broad-	___ Shrubland
___ Drought-deciduous	leaved/Needle leaved	___ Dwarf Shrubland
___ Mixed evergreen - cold-	___ Microphyllous	___ Herbaceous
deciduous	___ Graminoid	___ Nonvascular
___ Mixed evergreen - drought-	___ Forb	___ Sparsely Vegetated
deciduous	___ Pteridophyte	
Herbs: Annual Perennial		

IV. VEGETATION COMPOSITION: (list dominant species only and indicate which stratum it occurs in by checking the appropriate stratum next to the species name)

[illegible]

V. VEGETATION ASSOCIATION:

1. Primary and secondary (if there is a close call or multiple associations within the same polygon) vegetation classification(s) within radius of observation point, based on field key:

2. Are there other vegetation associations present within the radius observation area in addition to what was listed above?

3. Representativeness: Does the assigned association represent the vegetation well? If not, why? Please rank the fit as "Good" "Fair" or "Poor."

V. SOIL CHARACTERISTICS:

Soil Notes:

VI. CLASSIFICATION COMMENTS *(use this space to provide additional comments about problems or ease in applying the vegetation key at this site, rationale for choice of association if there was doubt, etc.)*

Appendix G: Accuracy Assessment Contingency Table

Explanation of the Contingency Matrices

The accuracy assessment (AA) contingency matrix for the Fort Pulaski National Monument (FOPU) vegetation map layer is designed in a row and column format so as to compare the results of the vegetation types represented on the map layer to the vegetation types as verified on the ground. Map class codes are used to identify vegetation types for ease of comparison during the analysis. (A crosswalk between map classes and vegetation types is provided.)

The sample contingency table provides an initial summary of the AA data by displaying counts of observations, with sample data values (vegetation map classes) as rows and reference data values (vegetation types as identified on the ground) as columns. The values in the shaded cells along the diagonal represent counts for correctly classified observations, where the reference data (column) matches the mapped vegetation type (row). User's accuracy was calculated by dividing the number of samples that agreed with their corresponding map class by the total number of samples in that class. Producer's accuracy was calculated by dividing the number of samples that agreed with their corresponding map class by the total number of samples whose field call belonged to that category.

Measures of accuracy for the AA are defined on the population contingency table. Similar to the sample contingency table, rows in the population contingency table are defined by the sample data values, and the columns are defined by the reference data values. Unlike the sample contingency table, the values in each cell are the proportion of the target area in the corresponding true and mapped vegetation classes, rather than the raw counts of observations. The population and sample contingency tables are reported together as a summary of between-class error relationships and class accuracy statistics (Lea and Curtis 2010).

Field Determination																		
Map Units		A	B	C	D	E	F	G	H	I	J	K	L	M	N	Totals	User's Accuracy	90% Confidence Interval
Photo-interpreter's Determination	A	0	3	1	0	0	0	0	0	0	0	0	1	0	0	5	0%	- + 0% 10%
	B	2	4	5	0	0	0	0	0	0	0	0	0	0	0	11	36%	8% 65%
	C	0	2	16	0	0	0	0	0	0	0	0	0	0	0	18	89%	74% 100%
	D	0	0	0	3	0	2	0	0	0	0	0	0	0	0	5	60%	14% 100%
	E	0	0	0	0	0	0	0	0	0	0	5	0	0	0	5	0%	0% 10%
	F	0	0	0	0	0	3	5	0	0	0	1	0	0	0	9	33%	2% 65%
	G	0	0	0	0	0	1	4	0	0	0	0	0	0	0	5	80%	41% 100%
	H	0	0	0	0	0	0	0	15	0	0	5	0	0	2	22	68%	50% 87%
	I	0	0	0	0	0	0	0	0	3	0	0	0	0	0	3	100%	83% 100%
	J	0	0	0	0	0	0	0	0	0	5	0	0	0	0	5	100%	90% 100%
	K	0	0	0	0	1	1	0	0	0	0	26	0	0	0	28	93%	83% 100%
	L	0	0	0	0	0	0	0	0	0	0	0	4	0	0	4	100%	88% 100%
	M	0	4	4	0	0	0	0	0	0	0	0	1	0	0	9	0%	0% 100%
	N	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%	0% 0%
Totals		2	9	22	3	0	6	9	15	3	5	37	6	0				
Producer's Accuracy		0%	44%	73%	100%	0%	50%	44%	100%	100%	100%	70%	100%	0%			83 Total Correct Points	
90% Confidence Interval		-	0%	12%	55%	83%	0%	8%	12%	100%	83%	90%	57%	100%	0%			129 Total Points
Interval		+	25%	100%	91%	117%	0%	100%	77%	100%	100%	100%	84%	100%	100%			

Key to Map Class Codes

A= Atlantic Coast Interdune Swale

B= Cabbage Palmetto Woodland

C= Cedar-Live oak - Cabbage Palmetto Marsh Hammock

D= Mid-Atlantic High Salt Marsh

E= Needlerush High Marsh

F= Salt Flat (Swampfire Type)

G= Saltwort-Swampfire Dwarf-Shrubland

H= Seaside Oxeye Tidal Shrub Flat

I= Slash Pine Plantation

J= South Atlantic Coastal Shell Midden Woodland

K= Southern Atlantic Coast Salt Marsh

L= Tallowtree Upland Forest

M= Wax myrtle - Carolina Laurel Cherry - Toothache-Tree Shrubland

N= Giant Atlantic Cordgrass Marsh

Reference Data (Observations Classified as Vegetation Type on Ground)																			
(Observations Selected Within the Map Class)																Class Area (Ha)	Estimate (Users' Accuracy)	Limit, 90% Conf	Limit, 90% Conf. Int.
	A	B	C	D	E	F	G	H	I	J	K	L	M	N	Row Total				
A	0	0.0014	0.0005	0	0	0	0	0	0	0	0.005	0	0	0	0.0069	4.1	0%	0%	10%
B	0.0019	0.0038	0.0048	0	0	0	0	0	0	0	0	0	0	0	0.0105	18.2	36%	8%	65%
C	0	0.0015	0.0120	0	0	0	0	0	0	0	0	0	0	0	0.0135	23.2	89%	74%	100%
D	0	0	0	0.0008	0	0.0006	0	0	0	0	0	0	0	0	0.0014	2.4	60%	14%	100%
E	0	0	0	0	0	0	0	0	0	0	0.0022	0	0	0	0.0022	3.8	0%	0%	10%
F	0	0	0	0	0	0.0031	0.0052	0	0	0	0.0010	0	0	0	0.0093	16.2	33%	2%	65%
G	0	0	0	0	0	0.0002	0.0007	0	0	0	0	0	0	0	0.0009	1.6	80%	41%	100%
H	0	0	0	0	0	0	0	0.0007	0	0	0.0002	0	0	0.0001	0.0010	1.90	68%	50%	87%
I	0	0	0	0	0	0	0	0	0.0011	0	0	0	0	0	0.0011	1.93	100%	83%	100%
J	0	0	0	0	0	0	0	0	0	0.0020	0	0	0	0	0.0020	3.43	100%	90%	100%
K	0	0	0	0	0.0338	0.0338	0	0	0	0	0.8778	0	0	0	0.9454	1633.4	93%	83%	100%
L	0	0	0	0	0	0	0	0	0	0	0	0.0008	0	0	0.0008	1.44	100%	88%	100%
M	0	0.0041	0.0041	0	0	0	0	0	0	0	0	0.0010	0	0	0.0092	16.1	0%	0%	6%
N	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0000	0.0	0%	0%	0%
Column Total	0.0019	0.0108	0.0214	0.0008	0.0338	0.0377	0.0059	0.0007	0.0011	0.0020	0.8862	0.0018	0	0.0001	1.0	1727.70	Point Estimate, Overall Accuracy (with 90% Confidence Interval): 90.3% (82.7% - 97.9%) Point Estimate, Overall Accuracy Kappa Statistic (with 90% Confidence Interval): 59% (51.3% - 66.6%)		
Estimated True Map Class Area (Ha)	3.31	18.9	37.0	1.44	58.3	65.0	10.3	1.30	1.93	3.43	1522.8	4.10	0	0.17					
Point Estimate, Mean Producers' Accuracy	0%	35.2%	56.0%	100%	0%	8.3%	12.5%	100%	100%	99.1%	99.6%	35.6%	0%	0%					
Lower Limit, 90% Conf. Int.	0%	34.5%	55.8%	99.9%	0%	4.6%	12.2%	100%	100%	98.1%	98.7%	35.6%	0%	0%					
Upper Limit, 90% Conf. Int.	0%	35.9%	56.3%	100%	0%	12.0%	12.7%	100%	100%	100%	100%	35.6%	0%	0%					

Key to Map Class Codes

A= Atlantic Coast Interdune Swale

B= Cabbage Palmetto Woodland

C= Cedar-Live Oak Cabbage Palmetto Marsh Hammock

D= Mid-Atlantic High Salt Marsh

E= Needlerush High Marsh

F= Salt Flat (Swampfire Type)

G= Saltwort-Swampfire Dwarf Shrubland

H= Seaside Oxeye Tidal Shrub Flat

I=Slash Pine Plantation

J=South Atlantic Coastal Shell Midden Woodland

K=Southern Atlantic Coast Salt Marsh

L=Tallowtree Upland Forest

M=Wax-myrtle - Carolina Laurel Cherry - Toothache Tree Shrubland

N=Giant Atlantic Cordgrass Marsh

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS 348/122523, October 2013

National Park Service
U.S. Department of the Interior



Natural Resource Stewardship and Science

1201 Oakridge Drive, Suite 150
Fort Collins, CO 80525

www.nature.nps.gov

EXPERIENCE YOUR AMERICA™