


Study Plan for Vertebrate and Vascular Plant Inventories

Appalachian Highlands Network and Cumberland/Piedmont Network, National Park Service

**Prepared by: Becky Nichols, Mike Jenkins, Janet Rock,
Keith Langdon, and Teresa Leibfreid**



Appalachian Highlands Network

Big South Fork National River and Recreation Area (BISO)
Blue Ridge Parkway (BLRI)
Great Smoky Mountains National Park (GRSM)
Obed Wild and Scenic River (OBRI)

Cumberland/Piedmont Network

Abraham Lincoln Birthplace National Historic Site (ABLI)
Carl Sandburg Home National Historic Site (CARL)
Chickamauga and Chattanooga National Military Park (CHCH)
Cowpens National Battlefield (COWP)
Cumberland Gap National Historical Park (CUGA)
Fort Donelson National Battlefield (FODO)
Guilford Courthouse National Military Park (GUCO)
Kings Mountain National Military Park (KIMO)
Little River Canyon National Preserve (LIRI)
Mammoth Cave National Park (MACA)
Ninety Six National Historic Site (NISI)
Russell Cave National Monument (RUCA)
Shiloh National Military Park (SHIL)
Stones River National Battlefield (STRI)

Table of Contents

I. Introduction	2
-Descriptions of Each Park	3
II. Project Description.....	9
A. Sources of Existing Information	9
-NRBib.....	9
-Maps and GIS Layers	10
B. Park-Specific Objectives	10
-Special Habitats and Special Concern Species	14
C. Overall Sampling Strategy - Terrestrial	18
-Sample Sites	18
Small Parks	20
Large Parks	20
River Canyon Parks	20
Parkways	21
-Stratification	21
-General Sampling Design.....	23
D. Overall Sampling Strategy – Aquatic	25
E. Sampling Strategy for Each Taxon.....	25
-Plants	25
-Amphibians and Reptiles.....	27
-Birds	28
-Mammals	28
-Fish	29
F. Data Analysis.....	29
-Estimation of Inventory Completeness.....	29
-Species Distribution Maps.....	30
-Regional Analysis.....	30
III. Logistical Support and Compliance	31
IV. Budget	32
V. Completion Schedule.....	34
VI. Voucher Specimens	37
VII. Products and Deliverables	37
VIII. Literature Cited	39
IX. Appendices	41
A. NRBib References	41
B. Status of Data Deliverables	70
C. Species Lists with Common Names	77
D. Southeast Regional Director’s memo.....	79
E. I&M Coordinator Roles and Responsibilities.....	81

I. INTRODUCTION

This study plan is for inventories of vertebrate and vascular plant species located in parks of the Southeast Region of the National Park Service. Two networks are included in this plan: Appalachian Highlands Network and Cumberland/Piedmont Network. These combined networks encompass 18 parks which are broadly similar ecologically, have a history of working together, and are relatively close to one another, which allows for efficiencies in sampling; therefore, it is reasonable to include them both into one comprehensive plan. In this study plan, we outline the specific steps for completing inventories as described in the 1999 NPS Inventory and Monitoring publication, *Guidelines for Biological Monitoring* (National Park Service 1999).

The main goal of this project is to provide park managers with comprehensive, scientifically-based information about species in their parks. This type of information is lacking in most park units containing significant natural resources, and will be invaluable for making sound management decisions, as well as provide a framework for future monitoring strategies. These biological inventories will be designed to meet three basic objectives; to:

- 1) document at least 90% of the species estimated to occur in each park, along with their associated habitats
- 2) describe the distribution and relative abundance of species of special concern, and
- 3) provide baseline information to develop a general monitoring strategy

As discussed in this plan, these inventories will be partially conducted using a structured statistically valid study design. Although this method is less efficient at generating a species list than traditional, or non-structured, sampling, it allows for inferences about species distribution and relative abundance to be made. However, since structured designs require intensive site-specific sampling, comparatively more resources are used to gain more information about fewer species. Non-structured sampling maximizes the number of species inventoried but provides little information to managers regarding distribution, species interactions, etc. Since this study is focused on inventorying 90% of all species, the structured design, along with non-structured, will together likely meet study objectives.

Parks within the combined networks range in size from 47 ha (Abraham Lincoln Birthplace National Historic Site) to 211,097 ha (Great Smoky Mountains National Park), occur in seven states, and cover a total of 342,648 ha. Resources among the parks are extremely diverse, culturally as well as naturally. Eleven parks are cultural and include Revolutionary War sites, Civil War sites, historic home sites, and monuments. Although these sites typically are smaller than other parks in the networks and were created to protect cultural resources, they often preserve unique natural areas as well. The remaining seven parks were created for their natural and/or recreational resources.

A team assembled from GRSM (Keith Langdon, Mike Jenkins, Becky Nichols, Janet Rock) and MACA (Teresa Leibfreid) visited each of the 18 parks (excluding GRSM) from March-June, 2000. Following are brief descriptions of each park, in alphabetical order (a table of species scientific names/common names is presented in Appendix C):

Abraham Lincoln Birthplace National Historic Site (ABLI) (Kentucky)

Abraham Lincoln, our country's 16th president, was born in 1809 in a small cabin on Sinking Spring Farm in Kentucky. The cabin now is preserved inside a memorial building which was erected in 1909. The 47 ha site, which includes the memorial and the farm, was established as a national park in 1916 and designated as Abraham Lincoln Birthplace National Historic Site in 1959. Lincoln and his family lived in this cabin for two years when they moved a short distance away to Knob Creek, a 91 ha tract of land currently being acquired by the Park.

ABLI contains mixed deciduous woods, open fields, and a ~2 ha area of old-growth forest. The old growth is represented by large specimens of black cherry, black oak, and white ash, and the understory contains a mix of native shrubs (Carolina buckthorn) and aggressive non-natives (Oriental bittersweet). The Park's visitor center complex is landscaped with non-native plants including Amur honeysuckle and much of the woods near the birthplace monument currently are being invaded by Japanese honeysuckle.

ABLI is acquiring a 91 ha tract of land 16 km north of the current Park area, in the Knob Creek drainage. This site is in a different physiographic province from ABLI and adds considerably to the natural resources diversity. Knob Creek contains rich slopes of red buckeye and chinquapin oak, with a lush understory nearly free of non-native plant species. The Knob Creek tract also contains ridgetop woods, hay fields, a floodplain, Knob Creek, and south-facing rock outcrops.

Big South Fork National River and Recreation Area (BISO) (Kentucky, Tennessee)

Mining, agriculture, and logging practices of the early 20th century stripped much of the Cumberland Plateau of its marketable trees and accessible coal. Big South Fork National River and Recreation Area was created in 1974 as an effort to curtail the effects that these practices were having on the landscape, to provide economic recreational opportunities for the region, and to maintain the Big South Fork of the Cumberland River as a free-flowing river. The Big South Fork and its tributaries pass through 144 km of scenic gorges and valleys containing a wide range of natural and historic features.

This 50,586 ha site now is lush with second-growth forest concealing old homesteads, mining entrances, and logging roads. Much of the Park consists of a massive gorge carved into the Cumberland Plateau by the Big South Fork. Upstream topography is characterized by dendritic drainages that form narrow v-shaped gorges. Downstream, sheer bluffs dominate the gorge rim, towering over mixed-hardwood talus slopes and the river floodplain below, which is typified by river birch and American sycamore. The area above the gorge is relatively flat and is dominated by oak species, hickory species, and red maple with Virginia pine common on dry ridges and cliff edges. Mountain laurel is common in the understories of these dry forests. Mesic ravines contain large components of American beech, sugar maple, and yellow birch. Eastern hemlock and *Rhododendron* are common along narrow gorges and small streams.

Blue Ridge Parkway (BLRI) (North Carolina, Virginia)

Designed as a scenic highway, the 750 km Blue Ridge Parkway encompasses 35,894 ha and ranges in elevation from 198 m to 1,844 m as it winds along the crest of the southern Appalachian Mountains. The Parkway connects Shenandoah and Great Smoky Mountains National Parks and passes through George Washington, Jefferson, Pisgah, and Nantahala National Forests. Authorized in the 1930's and taking over 50 years to complete, the Parkway provides tremendous scenic vistas while also protecting diverse high elevation sites along a latitudinal transect.

Lower elevations along the Parkway are dominated by oak and oak-pine forests with rich cove forests common in topographically sheltered areas. As elevation increases, northern hardwood species such as American beech, yellow buckeye and yellow birch become common. At the highest elevations, spruce-fir forest dominates, although much of the Fraser fir overstory has been killed by the balsam woolly adelgid. Interspersed among the forests are small unique habitats such as grassy balds, heath balds, mountain bogs, and beech gaps.

Carl Sandburg Home National Historic Site (CARL) (North Carolina)

In 1968, Carl Sandburg Home National Historic Site became the first park to honor a poet. Carl Sandburg was a biographer, a folksinger, a lecturer, a Pulitzer Prize winning author, and was known as the Poet of the People, often writing about social justice. The Sandburgs moved to this farm, named Connemara, in 1945 and remained there for 22 years.

This 107 ha site contains relatively steep and rugged terrain, as well as low areas which were cleared for agricultural purposes. Second-growth deciduous hardwoods occupy ~80 ha of the upland areas in the Park. Specifically, this vegetational community is comprised primarily of oak, hickory, tulip poplar, maple, black gum, and white pine. Small streams originate on Big Glassy and Little Glassy Mountains and flow through the site; however, they are dammed at several locations to form small lakes and ponds. Wooded slopes contain granitic outcrops with an array of plant species specific to this unique habitat.

Chickamauga and Chattanooga National Military Park (CHCH) (Georgia, Tennessee)

During the fall of 1863, Union and Confederate armies fought for control of the city of Chattanooga which was a key rail center and gateway to the heart of the Confederacy. In September, a two-day battle was fought near Chickamauga Creek which resulted in the retreat of Union forces. This was considered the last major Confederate victory in the Western theatre but General Grant and his Union forces ultimately gained control of Chattanooga two months later. The Park consists of over 3,318 ha straddling the Georgia/Tennessee border. It was established in 1890 and is the first and largest of the national military parks. Only the three largest units of CHCH (i.e., Chickamauga Battlefield, Lookout Mountain, and the Sherman Reservation) will be covered by this inventory plan.

Chickamauga Battlefield, the largest unit of the Park, contains limestone cedar glades, creek bluffs, sagponds, beaver and quarry ponds, and open fields which are either cut twice a year for hay or mowed regularly. The oak-hickory forests surrounding the glades are comprised of post and northern red oak, loblolly pine, and pignut hickory. Lookout Mountain, a 1,120 ha section of the Park is comprised of a maintained lawn, sandstone cliffs, and rich limestone slopes. The slopes are dominated by white ash, northern red oak, and hickory species. Cave

Spring is located near the northwest boundary of this unit and the woods along Lookout Creek contain floodplain species such as hackberry. The Sherman Reservation, a 20 ha tract, is predominantly a disturbed second-growth woodland.

Cowpens National Battlefield (COWP) (South Carolina)

Early in 1781, a pivotal battle in the Revolutionary War occurred at this frontier pasturing ground. At the battle of Cowpens, the Continental Army was victorious over a larger force of British regulars, and it was a link in a chain of British defeats in the South that led ultimately to final defeat at Yorktown.

Topographically, Cowpens is relatively flat with a few small streams that form deeply cut channels. The 341 ha battlefield is a mixture of open fields and small woodlands. The open fields consist of mown fescue (an exotic grass) and dense thickets of multiflora rose (an aggressive exotic). Common trees across the Park include white oak, hickory species, southern red oak, and Virginia pine, with alder and black willow in riparian areas. A management plan currently is being developed to restore the battlefield to its historic vegetation structure and composition, which included open forests dominated by larger open-grown trees.

Cumberland Gap National Historical Park (CUGA) (Kentucky, Tennessee, Virginia)

Cumberland Gap, located where the borders of Kentucky, Tennessee, and Virginia meet, is a natural pass which has been used as a transportation corridor since prehistoric times. Migratory large game created a path through the gap long before people arrived. The Gap later became a Native American route to hunting grounds in Kentucky; however, it is best known for its importance in opening the interior of our nation to settlement by early colonists. Cumberland Gap National Historical Park was authorized by Congress in 1940 and encompasses 8,274 ha.

This diverse Park, which encompasses much of the east-west oriented Cumberland Mountain, contains 23 caves, streams, an historic settlement, beaver dams, bogs, rock outcrops, and a limestone cliff face. Much of the Park is covered by second-growth forests of mixed deciduous and oak-hickory species. Common species on mesic sites include tulip poplar, northern red oak, and American beech. On drier sites, chestnut oak, white oak, and hickories are common. An unusual bog complex on the east end of the Park is dominated by white pine and eastern hemlock.

Fort Donelson National Battlefield (FODO) (Tennessee)

During the Civil War, this Confederate fort guarded the strategic Cumberland River. Early in 1862 the Union Army, under the command of General Grant, captured Fort Donelson along with approximately 13,000 Confederate soldiers. This was the first major victory of the Civil War for the Union Army and set the stage for the Union's invasion of the deep South.

The battlefield site is 226 ha and is bordered by the impounded Cumberland River to the north, or what currently is known as Lake Barkley. FODO is a topographically diverse Park

ranging from dry ridges to bottomlands, and a mosaic of forest types occur depending upon topography. In general, hills and ridges are divided by deep ravines that contain a diverse vegetation community and include tulip poplar, sugar maple, and American beech. Dry slopes are dominated mainly by oak and oak-hickory forests. The vegetation is considered transitional between mixed-deciduous forests to the drier oak-hickory forests.

Great Smoky Mountains National Park (GRSM) (North Carolina, Tennessee)

Established by Congress in 1934, Great Smoky Mountains National Park is internationally renowned as a center of biological diversity within North America. Complex ecological gradients combine to create a diverse mosaic of biotic communities in this 211,097 ha Park. The biological importance of GRSM led to its designation as an International Biosphere Reserve in 1976, and as a World Heritage Site in 1983.

Elevations in the park range from 1,200 m to over 2,670 m. At lower elevations, tulip poplar dominates large areas that historically were cleared and farmed. In sheltered rich coves (typically with northerly aspects), yellow buckeye, sugar maple, white basswood, and tulip poplar dominate the overstory. In coves with steeper v-shaped drainages, silver bell and hemlock dominate the canopy and *Rhododendron* often forms a thick, often impenetrable understory layer. Drier slopes (south and west facing) are dominated by chestnut oak with an understory of mountain laurel. Dry ridges typically have a large component of pine (pitch, Virginia, and Table Mountain) mixed with dry site oaks (chestnut, scarlet, and black). At higher elevations, the northern hardwood forest is prevalent, which is composed of sugar maple, yellow buckeye, yellow birch, and American beech. At the highest elevations in the Park, red spruce forests (over 1,900 m) and red spruce-Fraser fir forests (over 2,400 m) dominate. Scattered throughout the Park are small unique communities such as grassy balds, heath balds, beech gaps, and small wetlands.

Guilford Courthouse National Military Park (GUCO) (North Carolina)

In 1781, Nathanael Greene, commanding General of the Continental Army's Southern Department, was defeated at Guilford Courthouse by the British General Lord Cornwallis. Although Greene's army suffered defeat, his losses were slight, whereas Cornwallis suffered overwhelming losses, a situation that greatly hastened the end of the Revolutionary War. This 89 ha site preserves the battlefields and commemorates the battle with 28 monuments.

At the time of the battle, this area was predominantly under cultivation. Today, GUCO contains mowed fields and flat upland woods of mixed pine and hardwoods, particularly oak species. A small stream made up of two tributaries flows north out of the Park.

Kings Mountain National Military Park (KIMO) (South Carolina)

Kings Mountain is a rocky spur of the Blue Ridge Mountains that rises 46 m above the surrounding area. In 1780, British Major Patrick Ferguson and his loyalist militia were severely defeated by a small band of patriot forces, turning the tide on England's attempt to conquer the South. Congress established this 1,597 ha site to become a National Military Park in 1931.

Topographically, KIMO is characterized by a series of ridges that generally run from southwest to northeast. Elevations in the Park range from 197 m to 324 m above sea level and dendritic drainages create numerous ravines. Tulip poplar, sweet gum, black walnut, and American sycamore dominate floodplain forests in the Park. On mesic and dry-mesic slopes, overstories contain white oak, red maple, and tulip poplar, and understories contain large components of flowering dogwood. Typical dry-site species include chestnut oak, scarlet oak, and shortleaf pine, with post oak and blackjack oak also common. Fire suppression and other changes in land use have drastically increased the density of trees at KIMO compared to that at the time of the battle. A management plan is underway to restore the battlefield to its historic vegetation structure.

Little River Canyon National Preserve (LIRI) (Alabama)

This 5,543 ha preserve protects the nation's longest mountaintop river, which flows for almost its entire length down the middle of Lookout Mountain in northeast Alabama. The free-flowing Little River is one of the cleanest, wildest waterways in the South and its canyons are some of the deepest (183 m) in the Southeast. This is the newest park unit in either network, being authorized in 1992.

Little River Canyon can be divided into four communities: oak-hickory forests, canyon shoulders, sandstone rock outcrops, and riparian areas. Oak-hickory forests occupy deep soils above the canyon shoulders. Downslope, shortleaf and loblolly pine are common, grading into Virginia pine on the glade-like canyon shoulders. Sandstone rock outcrops are common along the canyon shoulder and mainly harbor stunted Virginia or scrub pine. The shrub layer consists of sparkleberry, fringe tree, Georgia holly, and black gum. Riparian areas usually are narrow except in broader channels where oxbows exist, with woods of mainly red maple, beech, umbrella magnolia, sycamore, and river birch.

Mammoth Cave National Park (MACA) (Kentucky)

This is the longest recorded cave system in the world with more than 538 km explored and mapped. Geologists estimate that there could be as many as 960 km of yet undiscovered passageways. The cave ecosystem is considered one of the world's most diverse, but the Park also contains tremendous above-ground diversity. Rivers, bluffs, sinkholes, cave entrances, and ridgetops all are habitats in which are found many distinct plant and animal communities. The Green River, the main watercourse through the Park, is known as one of the most diverse rivers in North America, containing 82 fish species and providing habitat for federally endangered freshwater mussels. Mammoth Cave was authorized as a National Park in 1941, designated as a World Heritage Site in 1981, and as an International Biosphere Reserve in 1990. The Park is 21,380 ha in size.

MACA features mostly second-growth forests and small areas of old growth. American beech trees dominate mesic hollows, joined by tulip poplar and sugar maple on lower and middle slopes. White and black oaks, along with three species of hickory, occupy upland mesic sites and slopes. Old fields cover approximately 45% of the Park. These sites are largely dominated by eastern red cedar and/or Virginia pine mixed with deciduous trees along their outer margins.

Ninety Six National Historic Site (NISI) (South Carolina)

This Revolutionary War site preserves the frontier village of Ninety Six and the earthworks associated with its role as a British outpost. It was the scene of repeated confrontations between loyalists and patriots, and one of the longest sieges of the war conducted by the Continental Army. Through the mid-18th century, Ninety Six was the economic and political center of the region. It supposedly was named by traders for the distance between Charleston and the Cherokee town of Keowee.

The site is 400 ha, comprised primarily of mixed woods, riparian areas along Spring Branch, and some grassy areas, particularly around the earthworks. Bottomland forest along Ninety Six Creek contains white oak, water oak, willow oak, shagbark hickory, sweetgum, and sycamore. Oak-hickory forest dominates the upland slopes. Star Fort Lake is an 11 ha impoundment within the Site. Other water resources include 0.4 ha Little Pond and several springs in the Spring Branch area. Other communities include open grassy fields, a slash pine plantation, and abandoned early-seral fields.

Obed Wild and Scenic River (OBRI) (Tennessee)

The Obed River, Clear Creek, Daddy's Creek, and Emory River together comprise the 72 km of wild and scenic river within OBRI. These water courses have cut through the sandstone of the Cumberland Plateau to form a rugged landscape of steep gorges with cliffs rising as much as 160 m above the stream channel. This Park, which received its Wild and Scenic River designation in 1976, has an area of 2,046 ha.

OBRI and BISO share similar geological features, historic land use, and some vegetational characteristics as well. Like BISO, oak, hickory, and pine species combine with red maple to dominate overstories in the dry flat areas above cliff-lines. Mountain laurel is common in the understories of these dry forests. American beech, sugar maple, and yellow birch are typical in mesic ravines and eastern hemlock and *Rhododendron* are common in narrow gorges. River birch and American sycamore are common overstory components in the major stream bottoms.

Russell Cave National Monument (RUCA) (Alabama)

Russell Cave has one of the longest and most complete archeological records in the eastern United States. Artifacts indicate intermittent human habitation for almost 9,000 years. Varying styles of spears and arrow points show that different bands of Indians used the cave as a permanent home, as winter quarters, or, in the case of nomadic tribes, as a stopover. Russell Cave National Monument was established in 1961 when 125 ha were donated to the NPS by the National Geographic Society.

This site is 125 ha in size and contains the cave, a stream, sinkholes, and sandstone outcrops. The woods near the cave entrance are comprised of blue ash, yellow buckeye, white basswood, and tulip poplar. The stream floodplain consists of boxelder and tulip poplar. Wooded slopes contain a diverse canopy that includes black cherry, white oak, yellow buckeye, American beech, black locust, and hickory species.

Shiloh National Military Park (SHIL) (Tennessee)

The first major battle of the Civil War's Western theatre occurred at Shiloh in 1862. It was a two-day battle resulting in nearly 24,000 casualties and a decisive victory for Union forces, who later went on to seize control of the Confederate railway system in Corinth, Mississippi. The Park was established in 1894 to preserve the battle scene and is approximately 1,607 ha.

SHIL contains not only the fields where the major portion of the battle was fought, but also alluvial floodplains, mixed bottomland and swamp forests, steep river bluffs, and rock outcrops. Water resources in the Park include Bloody Pond, natural springs, the Tennessee River, and Owl Creek. Uplands are predominantly mixed oak forests and old-fields. Hardwoods dominate the river bluff slopes and ravines.

Stones River National Battlefield (STRI) (Tennessee)

One of the bloodiest battles of the Civil War occurred at Stones River during late 1862-early 1863 where 81,000 men fought for control of middle Tennessee. Although the battle was considered to be tactically indecisive, General Rosecrans claimed victory for the Union forces after General Bragg and his Confederate troops withdrew. This 287 ha site also encompasses Stones River National Cemetery which contains over 6,000 Union graves.

STRI is characterized by upland hardwood forests and successional communities dominated by eastern red cedar. Unique habitats include rock outcrops, hay fields, numerous limestone cedar glades containing endemic plant species, and a small tract of oak-hickory woods. West Fork Stones River, and its tributary Lytle Creek, have associated limestone bluffs, floodplains, and springs.

II. PROJECT DESCRIPTION

A. Sources of Existing Information

NRBib

The Natural Resources Bibliography (NRBib) contains a wide variety of literature, including published papers, proceedings from meetings, government documents, research reports, hand-written notes, species lists, information compiled by local volunteers, etc.; however, not all of this information is useful in determining the current status of inventories due to the fact that it is often unverifiable. Searches using NRBib produced a large amount of inventory information, especially considering not all sources have yet been entered. Some parks, such as MACA and BLRI, have extensive amounts of information, although it often is taxon-specific (e.g., numerous bird citations, but none for fish). Other parks, such as NISI and GUCO, don't have any lists.

In addition to searching NRBib, we gathered as much inventory information as possible during our visits to each park. Discussions with park managers and visual field inspections using available species lists helped us determine the level of completeness for each inventory. Through this process, we determined that the natural resource inventories of the 18 network parks are in various stages of completion - a few parks having all inventories completed, while others have virtually no information (Table 1). Additionally, some species lists were 20 years

old or older and were considered to be out-of-date, and while most completed inventories include fully accessioned specimens, some do not, such as OBRI, which has a set of vouchers that have not been mounted or labeled and have not been placed in a herbarium.

GRSM is one park with completed systematic inventories of vascular plants and the vertebrate groups; however, the status of some of its special concern species are not well known. Here, GRSM will be included only in discussions of unique habitats and inventories of special concern species. The most current, pertinent, and verifiable literature for each taxonomic group for each park is listed in Appendix A.

Maps and GIS Layers

Digital data layers exist for most parks in these two networks (Table 2) which will be useful during the stratification phase of sample site determination. One serious limitation is that only five parks have the highest resolution 10 m x 10 m Digital Elevation Models (DEM's), as opposed to 30 m x 30 m DEM's. Also, only four parks have a completed vegetation map, and only one has a completed geology map.

B. Park-Specific Objectives

The overall objectives of the phase 1 NPS biological inventories are to document at least 90% of vertebrate and vascular plant species and to produce detailed distribution maps for species of special management concern. Each park has specific issues and objectives brought forward during a workshop at MACA in July, 1999. Park managers were asked to identify key issues, unique habitats, and special concern species. Some of the major issues to emerge (Table 3) included a lack of updated verified species checklists, a lack of information about unique habitats, and threats such as air pollution, visitor impacts, oil and gas extraction, adjacent land development, exotic species, etc. Of the 16 basic issue categories (mammals, birds, fish, reptiles/amphibians, mollusks, insects, plants, T&E species, vegetation maps, lichens, wetlands, soils, geology, exotics, caves and karst, fire management), seven were selected as high priority. They are: plants, birds, T&E species, reptiles and amphibians, mussels, vegetation maps, and exotic plants. A breakdown of park-specific objectives is presented in Table 3.

Table 1. Level of inventory completeness (by park) for the six phase 1 taxonomic groups. "Complete" indicates inventories which are at or above 90% of species documented (based on documentation and field verification).

Park	Plants	Amphibians	Reptiles	Birds	Mammals / Bats	Fish
ABLI	incomplete ^a	none	none	incomplete ^b	none / none	none

BISO	incomplete ^b	incomplete ^{bc}	incomplete ^{bc}	complete	none / incomplete ^{bc}	incomplete ^b
BLRI	incomplete ^b	none	none	complete	incomplete / incomplete ^{bc}	incomplete ^b
CARL	incomplete ^a	none	none	incomplete ^b	none / none	incomplete ^b
CHCH	incomplete ^{ab}	none	none	none	none / none	incomplete ^b
COWP	incomplete ^a	in progress	in progress	incomplete ^b	in progress / none	incomplete ^b
CUGA	incomplete ^{ab}	incomplete ^{bc}	incomplete ^{bc}	incomplete ^c	incomplete ^{bc} /incomplete ^b	incomplete ^b
FODO	in progress	none	none	none	none / none	none
GRSM	complete	complete	complete	complete	complete / complete	complete
GUCO	none	incomplete ^{bc}	none	incomplete ^b	none / none	none
KIMO	complete	in progress	in progress	none	incomplete / none	incomplete ^{bc}
LIRI	complete	incomplete ^b	incomplete ^b	none	none / none	complete
MACA	in progress	incomplete ^{bc}	incomplete ^{bc}	incomplete ^b	none / incomplete ^c	complete
NISI	in progress	none	none	incomplete ^{ab}	none / none	none
OBRI	complete	incomplete ^{bc}	incomplete ^{bc}	incomplete ^b	incomplete / incomplete	incomplete ^b
RUCA	complete	none	none	none	none / none	none
SHIL	complete	incomplete ^b	incomplete ^b	incomplete ^{ab}	incomplete ^{bc} /in progress	complete
STRI	complete	none	none	incomplete ^c	none / none	incomplete ^b

^a visual inspection

^b discussions with park managers

^c checklist 20 years old or older

Table 2. Available GIS layers and maps.

Park	Landsat TM	Geology	NWI	Vegetation	Species of Concern	DRG	DOQQ	DLG HYP	DLG HYD	DLG BDY	DLG RDTR	DLG PITR	DLG RR	DEM	Other
ABLI		P	X			X	X	P	X	X	X	X	X	X-10m	1999 hydrological study
BISO	X	scan only	X		P	X	X	X	X	X	X	X	X	X-10m	Cliff line and streamhead habitat, archaeology
BLRI			X		P	X	X	P	X	X	X	X	X	X-10m	Soils in progress
CARL			P			X	X	P	X	X	X	X	X	X-30m	Viewsheds
CHCH			X	X	P	X	X		X	X	X	X	X	X-30m	Cedar glade study
COWP			P		P	X	X	X	X	X	X	X	X	X-30m	Exotics
CUGA	X	P	X		P	X	X	X	X	X	X	X	X	X-30m	
FODO			X			X	X	X	X	X	X	X	X	X-30m	
GRSM	X	P	P	P	P	X	X	P	X	X	X	X	X	X-30m P-10m	Spot99, soils in progress, fire/disturbance history
GUCO			X			X	X		X	X	X	X	X	X-30m	
KIMO			X	X	P	X	X	X	X	X	X	X	X	X-30m	
LIRI			P		P	X	X	X	X	X	X	X	X	X-30m	
MACA	X	X	X	P	P	X	X	P	X	X	X	X	X	X-30m	1998 habitat map-30m DEMs, Ikonos on order
NISI			X			X	X	X	X	X	X	X	X	X-30m	
OBRI	X		X		P	X	X	X	X	X	X	X	X	X-10m	
RUCA			P	X	P	X	X	X	X	X	X	X	X	X-30m	
SHIL			X	X	P	X	X		X	X	X	X	X	X-30m	
STRI			X		P	X	X		X	X	X	X	X	X-10m	Cedar glade study

P = Partial / In Progress

X = Complete either at park, I&M Program, or available through USGS, USFWS, etc.

BDY = Boundary

DEM = Digital Elevation Model

DLG = Digital Line Graph

DOQQ = Digital OrthoQuarterQuads

DRG = Digital Raster Graphic

HYD = Hydrology

HYP = Hypsography

Landsat TM = Thematic Mapper

NWI = National Wetland Inventory

PITR = Pipes & Transmission Lines

RDTR = Roads & Trails

RR = Railroad

Table 3. Outcome of the July, 1999 scoping workshop at MACA attended by Appalachian Cluster park representatives.

Park	Key Issues	Objectives
ABLI	• lack of baseline data • development on adjacent lands • new land acquisition	• inventory / map habitats for plants and vertebrates • determine distributions of special concern species • complete inventories at Knob Creek
BISO	• development on adjacent lands • exotics • visitor use impacts • acid mine drainage • oil and gas extraction	• map unique habitats • inventory special concern species
BLRI	• exotics • visitor use impacts • parkway maintenance and development • lack of data on rare species	• compile existing data • map habitats • complete inventories, particularly at high elevations
CARL	• lack of baseline data • exotics • ANCS+ backlog	• inventory / map habitats of vertebrates • control exotics
CHCH	• development on adjacent lands • exotics • protection of species of concern	• inventory plants and vertebrates
COWP	• lack of baseline data • exotics • air quality	• update inventories • complete all inventories
CUGA	• exotics • air quality (acid deposition) • acid mine drainage • Cumberland Gap restoration	• update inventories • complete all inventories • complete vegetation mapping • inventory bogs
FODO	• lack of baseline data • exotics • protection of species of concern • lack of wetland / habitat data	• complete inventories • map habitats
GRSM	• exotics • air quality • habitat loss / fragmentation	• inventory special concern species • inventory exotics • conduct salamander research
GUCO	• lack of baseline data • development on adjacent lands	• verify checklists • complete inventories • map habitats
KIMO	• lack of baseline data • exotics	• complete inventories • map habitats (particularly wetlands)

		• map invasive exotics • QA/QC on plant list
LIRI	• lack of baseline data • status of species of concern	• complete vegetation mapping • inventory vertebrates • inventory special concern species
MACA	• exotics • air / water quality • status of species of concern • loss of open habitat for barren species • river impoundment from dam	• map habitats • complete vegetation mapping • map special concern species • map exotics • prescribed fire
NISI	• lack of baseline data • protection of rare species	• inventory plants and vertebrates • map special concern species
OBRI	• development on adjacent lands • water quality • threat of dam	• map unique habitats • inventory special concern species
RUCA	• lack of baseline data • status of species of concern	• inventory vertebrates • QA/QC on plant list
SHIL	• exotics • air quality • external threats from paper mill, agriculture	• inventory amphibians, reptiles, birds, insects, bats • status of species of concern
STRI	• development on adjacent lands • exotics • air / water quality	• inventory vertebrates

Special Habitats and Special Concern Species

Network park managers considered special habitats and special concern species to be key issues. “Special concern species” was selected as one of the highest ranked categories (Table 3) during the 1999 scoping workshop. Several parks (e.g., BLRI, BISO, GRSM, MACA) have some information on federally listed Threatened and Endangered (T&E) species while others have virtually none (e.g., KIMO, NISI). Parks that were at least aware of the presence of listed species often had no information on distribution within their parks.

Parks without general inventories or with incomplete inventories will be given higher funding priority than those that do. Because this is a phase 1 inventory program, aimed at discovering 90% of the species present, inventories of special concern species will be completed following the comprehensive surveys. Park managers expressed a need for maps of special habitats and the locations of special concern species. Location information will be compiled for park managers (not to be distributed) after general inventories are complete.

All parks contain special habitats that will be incorporated into the sampling design (see “General Sampling Design” section). Often these habitats contain special concern species,

including federally listed species. It is important to identify these habitats during the initial inventory phase in order to not “miss” rarer species in observing the 90% rule. GIS layers describing the distribution of special habitats do not exist. Table 4 provides a summary of special habitats known to occur within each Network park.

Most park managers are aware of the presence of federally listed T&E species but usually are not familiar with the distribution of these species within their boundaries. Parks having the same listed species (e.g., Indiana bat, gray bat) would benefit from a regional expert (in this case, a small mammal specialist) so as to promote an evenness of effort and economy. Table 4 summarizes T&E species at each park.

As a federal agency, we are mandated by the Endangered Species Act and NPS Management Policies to identify and protect federal, state, and local T&E species on park lands. This also includes species which are rare or unique, declining, sensitive, or candidates for listing. Protecting these species will involve determining their distribution, critical habitat, and other life history information, such as seasonality. Active management may be required in order to maintain or increase the distribution and abundance of these species and their habitats.

Table 4. Summary of documented special habitats and federally listed species by park.

Park	Special Habitats	Threatened and Endangered Species
ABLI	• Old-growth forests • Karst (sinking springs, caves)	Unknown
BISO	• Old-growth forests • River gorge • Cliff lines • Streamheads, seeps, wet depressions • Rock outcrops	• Chaffseed (<i>Schwalbea americana</i>) (a flowering plant) • Cumberland sandwort (<i>Minuartia cumberlandensis</i>) • Cumberland rosemary (<i>Conradina vertillata</i>) • Virginia spiraea (<i>Spiraea virginiana</i>) • Cumberland elktoe (<i>Alasmidaonta atropurpurea</i>) (a mollusk) • Cumberlandian combshell (<i>Epioblasma brevidens</i>) (a mollusk) • Littlewing pearlymussel (<i>Pegias fabula</i>) (a mollusk) • Oyster mussel (<i>Epioblasma capsaeformis</i>) (a mollusk) • Tan riffleshell (<i>Epioblasma florentina walkeri</i>) (a mollusk) • Cumberland bean (<i>Villosa trabalis</i>) (a mollusk) • Duskytail darter (<i>Etheostoma percnurum</i>) (a fish) • Red-cockaded woodpecker (<i>Picoides borealis</i>)
BLRI	• Rock outcrops/cliffs • High elevation communities • Wetlands (including bogs)	• Eastern cougar (<i>Felis concolor</i>) • Carolina northern flying squirrel (<i>Glaucomys sabrinus coloratus</i>) • Virginia big-eared bat (<i>Corynorhinus townsendii</i>)

		• Bog turtle (<i>Clemmys muhlenbergii</i>) • Blue Ridge goldenrod (<i>Solidago spithamea</i>) • Braun's rock cress (<i>Arabis perstellata</i>) • Mountain bluet (<i>Houstonia purpurea</i> var. <i>montana</i>) • Spreading avens (<i>Geum radiatum</i>) (a flowering plant) • Small whorled pogonia (<i>Isotria medeoloides</i>) • Swamp-pink (<i>Helonias bulbata</i>) • Heller's blazing star (<i>Liatris helleri</i>) • Virginia spiraea (<i>Spiraea virginiana</i>) • Rock gnome lichen (<i>Gymnoderma lineare</i>)
CARL	• Granitic domes • Seeps and ponds	Unknown
CHCH	• Glades • Cliffs • Caves	• Large-flowered skullcap (<i>Scutellaria montana</i>) • Gray bat (<i>Myotis grisescens</i>)
COWP	• Wetlands • Springs	• Dwarf-flowered heartleaf (<i>Hexastylis naniflora</i>)
CUGA	• Mountain meadows • Bogs • Caves	• Indiana bat (<i>Myotis sodalis</i>) • Blackside dace (<i>Phoxinus cumberlandensis</i>) (a fish)
FODO	• Old-growth forest • Floodplains • Wetlands • Springs	Unknown
GRSM	• Old-growth forest • Grassy balds • Caves • High elevation communities • Geo-chemically different sites • Rock outcrops	• Spreading avens (<i>Geum radiatum</i>) (a flowering plant) • Rock gnome lichen (<i>Gymnoderma lineare</i>) • Virginia spiraea (<i>Spiraea virginiana</i>) • Carolina northern flying squirrel (<i>Glaucomys sabrinus coloratus</i>) • Indiana bat (<i>Myotis sodalis</i>) • Gray bat (<i>Myotis grisescens</i>) • Red-cockaded woodpecker (<i>Picoides borealis</i>) • Spotfin chub (<i>Cyprinella monacha</i>) (a fish) • Smoky madtom (<i>Noturus baileyi</i>) (a fish) • Yellowfin madtom (<i>Noturus flavipinnis</i>) (a fish) • Spruce-fir moss spider (<i>Microhexura montivaga</i>)
GUCO	• Moist areas	Unknown
KIMO	• Floodplains • Wetlands • Rock outcrops • Power line	Unknown

	• Old mine shaft • Small waterfalls	
LIRI	• Canyon floor • Glades • River • Bogs	• Green pitcher plant (<i>Sarracenia oreophila</i>) • Harperella (<i>Ptilmnum nodosum</i>) (a flowering plant) • Kral's water-plantain (<i>Saggitaria secundifolia</i>) (a flowering plant) • Blue shiner (<i>Cyprinella caerulea</i>) (a fish)
MACA	• Karst/cave • Glades • Bogs • River islands • Sinkholes • Hemlock hollows • Barrens remnants • Small lake • Upland swamps • Sandstone cliff lines	• Eggert's sunflower (<i>Helianthus eggertii</i>) • Indiana bat (<i>Myotis sodalis</i>) • Gray bat (<i>Myotis grisescens</i>) • Fanshell (<i>Cyprogenia stegaria</i>) (a mollusk) • Northern riffleshell (<i>Epioblasma torulosa rangiana</i>) (a mollusk) • Cracking pearlymussel (<i>Hemistena lata</i>) (a mollusk) • Ring pink (<i>Obovaria retusa</i>) (a mollusk) • Clubshell (<i>Pleurobema clava</i>) (a mollusk) • Rough pigtoe (<i>Pleurobema plenum</i>) (a mollusk) • Mammoth Cave shrimp (<i>Palaemonias ganteri</i>) • Bald eagle (<i>Haliaeetus leucocephalus</i>)
NISI	• Rock outcrops • Wetlands (bogs, swamps) • Lake	Unknown
OBRI	• Old-growth forest • Cliff lines • River gorge • Seeps, wet depressions • Rock outcrops	• Cumberland rosemary (<i>Conradina verticillata</i>) • Purple bean (<i>Villosa perpurpurea</i>) (a mollusk) • Alabama lampshell (<i>Lampsilis virescens</i>) (a mollusk) • Spotfin chub (<i>Cyprinella monacha</i>) (a fish)
RUCA	• Cave	
SHIL	• Old-growth forests • Floodplains • Wetlands (swamps, springs)	• Indiana bat (<i>Myotis sodalis</i>) • Gray bat (<i>Myotis grisescens</i>)
STRI	• Cedar glades, barrens • Karst • Bluffs • River • Ponds	• Tennessee coneflower (<i>Echinacea tennesseensis</i>)

C. Overall Sampling Strategy - Terrestrial

For the purpose of this plan, terrestrial and aquatic systems each will have separate sampling designs. The 17 network parks have been divided into four sampling groups: 1) small parks (< 1,000 ha), 2) large parks (1,000-21,000 ha), 3) river canyon parks, and 4) parkways (Table 5).

Within all parks, the emphasis is to sample the dominant community and land types in addition to special habitats. Once sample sites are established, a plot design will be created that allows sampling of vascular plants, mammals, birds, amphibians, and reptiles at the same sites. When possible, we have incorporated the recommendations outlined in the report “Guidance for the Design of Sampling Schemes for Inventorying and Monitoring Biological Resources in National Parks.” This report was the product of a panel of sampling design experts convened in Fort Collins, Colorado in February 2000.

The sampling strategy for all parks will consist of three steps: 1) the establishment of a coarse network of grid points across the entire park, 2) the location of additional sample sites using a stratified random design (Thompson 1992), and 3) intensive searches of small areas of special habitats. The use of a coarse grid to establish sample sites ensures that they are distributed evenly across an entire area; however, one of the inherent weaknesses of a grid design is that it tends to over-sample common habitats and consistently misses rare and spatially-scattered habitats (Fortin et al. 1989, Legendre and Fortin 1989). Since rare habitats often are characterized by high biological diversity, sampling these areas is critical. In this strategy we recommend using a widely-spaced grid with the understanding that it will not provide sufficient representation of all habitats, but over-sampling of common habitats will be avoided. Once grid points are selected and overlaid onto the map of selected strata (discussion on recommended strata follows), the number of sample points by strata will be tallied. Additional sample sites will be randomly located within each under-represented strata.

Sample Sites

During the planning stage, it is not possible to know the exact number of sample sites we will require to achieve the goals of this inventory. The number of sample sites needed within a park will depend upon the variability of habitat, species distributions, and sampling effort. Therefore, the ultimate number of sample sites will be determined through an iterative process. Data from the first sampling year for a given taxon will be examined to determine how many additional sample sites are needed. Relationships between species accumulation and area, and species accumulation and effort will be examined to determine when sufficient sampling has occurred within a park. If data suggest that additional sampling is needed within a strata for a given taxon, additional sample sites will be randomly located and inventoried.

In the initial grid-based design, we have selected grid-point intervals that are unlikely to over-sample particular strata or divert excessive resources from the stratified samples, but provide a representative sample of common habitats in each park. Overall, the grid-point density per park is based on size. The following increments will be used: <1,000 ha; 1,000-5,000 ha; 5,000-8,000 ha; 8,000-20,000 ha; 20,000-50,000 ha; and >50,000 ha. If grid-point sampling requires an

Table 5. Initial grid sampling intensity by park group.

Park group	Area (ha)	Sample points	Grid density	Grid size (km)
Small Parks				
ABLI ¹	47	15	1 sample point/3 ha	0.18 x 0.18
GUCO	89	15	1 sample point/6 ha	0.24 x 0.24
CARL	107	15	1 sample point/7 ha	0.27 x 0.27
RUCA	125	15	1 sample point/8 ha	0.28 x 0.28
FODO	226	15	1 sample point/15 ha	0.38 x 0.38
STRI	287	15	1 sample point/19 ha	0.43 x 0.43
COWP	341	15	1 sample point/23 ha	0.48 x 0.48
NISI	400	15	1 sample point/27 ha	0.52 x 0.52
Large Parks				
KIMO	1,597	20	1 sample point/80 ha	0.90 x 0.90
SHIL	1,607	20	1 sample point/80 ha	0.90 x 0.90
CHCH				
• Chickamauga Battlefield	2,179	27	1 sample point/81 ha	0.90 x 0.90
• Lookout Mountain	1,108	14	1 sample point/79 ha	0.90 x 0.90
• Sherman Reservation	24	3	1 sample point/8 ha	0.28 x 0.28
CUGA	8,274	39	1 sample point/212 ha	1.45 x 1.45
MACA	21,380	44	1 sample point/486 ha	2.20 x 2.20
River Canyon Parks²				
OBRI	2,046	25	1 sample point/82 ha	0.90 x 0.90
LIRI	5,543	35	1 sample point/158 ha	1.25 x 1.25
BISO	50,586	53	1 sample point/954 ha	3.10 x 3.10
Parkways				
BLRI (road corridor) ³	22,239	47	1 sample point/473 ha	2.18 x 2.18
• Humpback Rocks	1,214	15	1 sample point/81 ha	0.90 x 0.90
• Peaks of Otter	1,659	20	1 sample point/83 ha	0.90 x 0.90
• Devil's Backbone	453	15	1 sample point/30 ha	0.55 x 0.55
• Smart View	282	15	1 sample point/19 ha	0.43 x 0.43
• Rock Castle Gorge	1,452	18	1 sample point/81 ha	0.90 x 0.90
• Fisher Peak	708	15	1 sample point/47 ha	0.70 x 0.70
• Cumberland Knob	324	15	1 sample point/22 ha	0.46 x 0.46
• Mahogany Rock	584	15	1 sample point/39 ha	0.62 x 0.62
• Doughton Park	2,320	29	1 sample point/80 ha	0.90 x 0.90
• E. B. Jeffress Park	418	15	1 sample point/28 ha	0.53 x 0.53
• Moses Cone Mem. Park	1,421	18	1 sample point/79 ha	0.90 x 0.90
• Julian Price Mem. Park	1,726	21	1 sample point/82 ha	0.90 x 0.90
• Linville Falls	547	15	1 sample point/36 ha	0.60 x 0.60
• Craggy Gardens	238	15	1 sample point/16 ha	0.40 x 0.40

¹ ABLI is in the process of acquiring a 91 ha tract at Knob Creek. If successful, sample size must be adjusted.

² River corridor included in area estimate.

³ Sampling points located along road corridor every 10 miles (16.1 km). Units within BLRI are separated.

excessive amount of resources in a given park, the grid-point interval may be increased to ensure that sufficient resources remain for stratified and specialized habitat sampling.

Small Parks

The unusual topographic, geologic, or vegetation characteristics found at these sites were recognized by historic figures such as military commanders, aboriginal leaders, and a poet. Many of these figures now are commemorated through the creation of these parks, which also preserve the unique natural areas within them. They include battlefields, archeological, and home site parks, and range in size from 47 to 400 ha. Within these small parks, we propose establishing 15 grid-point sample sites per park (Table 5). Grid densities will range from 1 sample point/3 ha at ABLI to 1 sample point/20 ha at NISI. Grid-point intervals will vary from 0.18 km (180 m) at ABLI to 0.45 km (450 m) at NISI. Since this grid interval produces a large number of sample sites relative to overall park size, we anticipate that relatively few additional sample points will be needed to sufficiently represent major strata. All parks within this group contain small areas of special habitats (Table 4) that will require intensive sampling.

Large Parks

This category includes the larger battlefields, historic parks, and natural resource parks, which are all large enough to contain a wide range of community types and topography. For parks in this group that are less than 3,000 ha (i.e., KIMO, SHIL), we will use a 0.90 km (750 m) grid-point interval (Table 5). A 1.45 km x 1.45 km grid will be used to select sample points at CUGA, and a 2.2 km x 2.2 km grid will be used at MACA. With these intervals, sample site densities range from 1 sample point/81 ha at KIMO and SHIL to 1 sample point/484 ha at MACA. Using an iterative approach, stratified sampling will ensure that delineated strata are adequately sampled. As with the smaller parks, specialized habitat will be differentiated and sampled separately. Some areas that are extremely hazardous to sample, such as cliff faces, will not be included in the grid or stratified sampling, but will be sampled as access permits. For example, due to its extremely steep slopes, the Lookout Mountain unit of CHCH will be sampled as trail access safely permits.

River Canyon Parks

This category includes parks containing riparian areas within steep canyons. Typically, the areas above the canyons are fairly level and accessible; however, steep canyon walls make accessing riparian areas difficult and present considerable hazards for sampling. For these reasons, sampling in the river canyon parks will be done in three parts. In the areas above the canyon, initially we will use a coarse grid to select sample points. Grid-point intervals will vary from 0.90 km at OBRI (1 sample point/81 ha) to 3.1 km at BISO (1 sample point/961 ha) (Table 5). Additional sample points above the canyon will be selected through stratified random sampling. In the riparian areas, we propose randomly selecting sample points equivalent to the number of points per hectare sampled above the canyon. Additional sampling points will be assigned above

the canyon and in the riparian areas if species accumulation curves suggest that more sample points are needed. Due to their hazardous nature, the canyon walls will be sampled as safe access allows. As with other parks, special habitats such as rock shelters, riffles, and river islands will be sampled separately to ensure that their species diversity is captured.

Parkways

In the combined networks, there is one parkway (BLRI). BLRI offers a unique challenge since it covers a geographic gradient (North Carolina to Virginia) but often is only a few hundred meters wide. This gradient results in considerable community and species diversity. BLRI also contains 14 sub-units, such as Doughton Park and Peaks of Otter, ranging in size from 238 ha to 2,320 ha. Consequently, BLRI will require greater sampling intensity than a non-linear park of the same size. Most of the Parkway will be sampled as a transect with sampling points established on each side of the road every 10 mile (16.1 km). Road mileposts will be used to locate transect points which will result in approximately 46 sample points along the road corridor. Additional points may be added based upon stratification. Sub-units will be sampled with a grid-point interval equal to that used to sample individual parks of similar size. If needed, we will select additional sample points through stratified random sampling. In addition, special habitats such as wetlands, rock outcrops, and riparian areas will be sampled more intensely using additional plots. A random subsample of sites will be selected from any special habitat type that exists in large numbers.

Stratification

Due to their small size and less varied topography and biotic communities, we do not anticipate that small parks will require stratified sampling. If it becomes evident that additional sampling is needed in small parks, they may be stratified and sampled based upon available data layers. In large parks, stratifying the landscapes into biologically significant units will be an important step in completing this inventory. If the sole purpose of this project was to inventory vascular plants and vertebrates in each park, stratification by vegetation cover type would provide the most efficient and logical means of stratifying complex landscapes. However, since sample sites established in this study may eventually serve as locations for long-term monitoring, the use of vegetation cover types is problematic. Because vegetation constantly is changing as a result of disturbance and succession, it may not be an appropriate stratification layer for long-term monitoring programs that require strata to remain fixed. This issue is of particular importance in some historic parks (e.g., COWP, KIMO) that are developing plans to restore historic vegetation structure and species composition. Therefore, we will use topography and edaphic characteristics to stratify samples when possible (Table 6). When other data layers are unavailable or inappropriate, we will use broad vegetation classes that are more stable than narrower classification units which often encompass transitional community types. For example, SHIL generally does not have sufficient topographic relief to allow stratification based upon aspect, slope, or slope position; therefore, we will use the Park's existing vegetation map to develop strata. We recommend reducing and combining the 18 vegetation classes into 5 broad classes: 1) evergreen forest, 2) dry deciduous forest, 3) wet deciduous forest, 4) dry mixed forest, and 5) wet mixed forest.

Many stratification schemes will use DEM's to derive slope, slope position, and aspect. Currently, many parks in the combined networks only have 30 m x 30 m DEM's; however, as 10 m x 10 m DEM's become available we recommend their use. In the fairly rugged Ozark Mountains of southern Missouri, Hammer et al. (1995) found that a 30 m resolution DEM correctly classified field-measured slopes into 5% slope classes only 25% of the time. The same DEM placed field-measured slopes into correct or adjacent slope classes 69% of the time. Therefore, if 30 m x 30 m DEM's must be used, we recommend that the broadest possible slope classes be used.

Since many of the parks in the combined networks have variable topography that strongly influences species distribution, elevation, slope, slope position, and aspect will be used to stratify. Where available, soil and geologic spatial data also may be used to extract better-defined strata. For example, MACA will be stratified based upon an existing habitat map that combines slope and aspect with geologic features (Olson and Franz 1998). Elevation will be used to stratify BLRI which ranges in elevation from 198 to 1,829 m. Other parks in the combined network (e.g., KIMO, CUGA, CHCH) do not have strong enough gradients to delineate strata based purely upon elevation; therefore, GIS software will be used to combine aspect, slope, and slope position into landform units such as dry slopes, mesic slopes, floodplains, etc. Table 6 offers specific spatial data layers we recommend for stratifying each park. It is impossible to predict the number of stratified sample plots this inventory will require before sampling strata have been selected.

Table 6. Recommended sampling strata and spatial data layers available to create strata.

Park group	Recommended strata	Available data layers
Large Parks		
KIMO	Aspect, slope, slope position	30 m x 30 m DEM
SHIL	Vegetation map-broad classes	Vegetation map
CHCH		
• Chickamauga Battlefield	Aspect, slope, slope position	30 m x 30 m DEM
• Lookout Mountain	Aspect, slope, slope position	30 m x 30 m DEM
• Sherman Reservation	Aspect, slope, slope position	30 m x 30 m DEM
CUGA	Aspect, slope, slope position	30 m x 30 m DEM 10 m x 10 m DEM (partial)
MACA	Slope, aspect, geology	Habitat classification map 30 m x 30 m DEM
River Canyon Parks		
OBRI	Aspect, slope, slope position	10 m x 10 m DEM
LIRI	Aspect, slope, slope position	30 m x 30 m DEM
BISO	Aspect, slope, slope position	10 m x 10 m DEM
Parkways		
BLRI (road corridor)	Elevation	10 m x 10 m DEM
• Humpback Rocks	Aspect, slope, slope position	10 m x 10 m DEM
• Peaks of Otter	Aspect, slope, slope position	10 m x 10 m DEM
• Devil's Backbone	Aspect, slope, slope position	10 m x 10 m DEM
• Smart View	Aspect, slope, slope position	10 m x 10 m DEM
• Rock Castle Gorge	Aspect, slope, slope position	10 m x 10 m DEM
• Fisher Peak	Aspect, slope, slope position	10 m x 10 m DEM
• Cumberland Knob	Aspect, slope, slope position	10 m x 10 m DEM
• Mahogany Rock	Aspect, slope, slope position	10 m x 10 m DEM
• Doughton Park	Aspect, slope, slope position	10 m x 10 m DEM
• E. B. Jeffress Park	Aspect, slope, slope position	10 m x 10 m DEM
• Moses Cone Mem. Park	Aspect, slope, slope position	10 m x 10 m DEM
• Julian Price Mem. Park	Aspect, slope, slope position	10 m x 10 m DEM
• Linville Falls	Aspect, slope, slope position	10 m x 10 m DEM
• Craggy Gardens	Aspect, slope, slope position	10 m x 10 m DEM

General Sampling Design

In the proposed sampling design, inventories of vascular plants and vertebrates will be centered on the sample points selected with the use of: 1) grids, and 2) stratified random sampling. Once established, each of these points will be classified within The Nature Conservancy's National Vegetation Classification System (Anderson et al. 1998, Grossman et al. 1998, Weakley et al. 2000). Topographic features (aspect, slope, slope position) and recent disturbances also will be noted. This type of information is critical to mapping species distributions and determining species co-occurrences. Sampling will be conducted within a 1 ha circular plot (radius = 56.4 m) extending concentrically from each sample point (Figure 1). If the 1 ha plot extends into multiple vegetation community types, each major type will be classified within the 1 ha plot. If

the 1 ha plot is divided among two or more vegetation types, all major types should be inventoried.

Whenever possible, all target species will be inventoried within the 1 ha plot [a similar 1 ha plot design has been used for the All Taxa Biodiversity Inventory (ATBI) at GRSM and has provided sufficient area for the establishment of vegetation plots, small mammal trap lines/grids, and cover board arrays]. Some types of sampling (e.g., bird point-count surveys) may require sampling outside of the 1 ha plots, but we recommend that, whenever possible, at least one point be within each 1 ha plot. Due to time and financial constraints, some inventory techniques (e.g., pitfall and funnel traps) used to sample some taxa (e.g., reptiles and amphibians) may be impractical for use on all 1 ha plots. In addition, some taxa may not require sampling on all plots to reach the 90% inventory goal. Therefore, if sampling on all plots is excessive, inefficient, or impractical for a given type of sampling or taxa, researchers may select a subset of plots for standardized sampling. To allow all species inventoried to be linked to a geographic location, researchers will record the UTM's of each search site using a GPS unit.




Figure 1. General plot design and potential inventory methods for vascular plants and terrestrial vertebrates.

Small, special habitats, such as seeps and outcrops, will be located from existing park records and from observations made while conducting structured (i.e., grid-point, stratified) sampling.

GIS layers for these habitats do not exist. These areas are difficult to sample with a structured design; consequently, they will be sampled through intensive searches consisting of walking transects and sweeps or by installing smaller specialized plots. Species-specific searches may be appropriate in some areas, and if additional special habitat sampling is desired, usually in larger parks, a multistage design may be required (Thompson 1992).

D. Overall Sampling Strategy – Aquatic

The sampling strategy for aquatic resources will differ from the terrestrial design just described, even in a case where a stream flows through a plot established for terrestrial sampling. The primary aquatic resources in these network parks are lower order streams, and in some cases, only small portions of streams flow through the park property, in which case it is feasible that the entire length could be sampled. In cases where this is not possible, the streams will be stratified by order. Firstly, the total length of all stream segments of a particular order will be determined (e.g., 500 m of 1st order streams, 1,200 m of 2nd order streams, etc.). Sample sites then will be located at n equal intervals along the entire length of streams of each category, depending upon resources available. Special habitats have a possibility of being missed in this type of design; therefore, these areas will be sampled separately.

E. Sampling Strategy for Each Taxon

Determining the timing of projects and budget allocations for this project required having information on typical sampling methods and seasons for each taxonomic group. We held a scoping meeting on 17-18 July, 2000 at GRSM with subject matter experts (primarily USGS-BRD scientists) to ascertain this data (Table 7).

Plants

Plants in the 18 parks will be surveyed through structured sampling within the 1 ha plot established at each sample point and through intensive searches of unique habitats. Within each 1 ha plot, a 20 m x 50 m (0.1 ha) subplot will be established based upon the general design used by The Nature Conservancy in vegetation mapping and vascular plant inventory projects (TNC and ESRI 1994, Grossman et al. 1998). Within each plot, all vascular plant species will be identified and recorded according to the height strata in which they occur. Because the number of strata will vary by forest type, the researcher will identify and record the height of all vegetation strata observed on each plot. Total coverage of each species will be estimated visually (vertical projection onto the ground), and recorded within 10 cover classes [Table 8 (Peet et al. 1998)]. The same 10 cover classes also will be used to estimate the cover of each species by strata. While estimating species cover does increase the time needed to complete a plot, the added data will allow for better estimations of species distribution and diversity. The larger 1 ha plot will be searched for species not identified in the 0.1 ha subplot. Plots will be visited three times during the April to September growing season to ensure that the full suite of vascular plants are identified.

Within rare habitats (e.g., seeps, rock outcrops, small glades) that are too small and irregular for the standard plot design, the 1 ha plot will not be used. A smaller subplot will be sampled based upon 10 m x 10 m sub-sample units or modules. For example, in a small glade, a 20 m x 20 m plot (four 10 m x 10 m modules) may be used to best capture diversity. For a long, thin rock outcrop, a series of three 10 m x 10 m modules (10 m x 30 m subplot) might best be used for sampling. In some cases, even a very small plot may be impractical for sampling. In these instances, the entire habitat will be inventoried through walking sweeps and inspection of microhabitats.

Table 7. Typical sampling protocols for plants and vertebrates obtained during the network scoping meeting, July, 2000.

	Plants	Amphibians	Reptiles	Birds	Mammals	Fish
Sampling methods	- nested plots @ grids - walking transects - blocked survey	- night driving - trapping - cover boards - time / area constrained searches - drift fence arrays - litter bags - call surveys - pitfall / funnel traps - egg mass counts	- passive trapping - roadkills - cover boards - time / area constrained searches - drift fence arrays - pitfall / funnel traps - live traps	point count survey and mapping	- trap lines - scent posts - pitfall traps - Anabat - scat / track surveys - litter raking (shrews)	- backpack electrofisher (small, large streams) - seines (small, large streams, lakes) - boat-mounted electrofisher (large streams, lakes)
Sampling season	- spring - summer - fall	- spring - summer - year-round (species specific)	summer	- spring (May-June) - winter (Dec-Jan)	cooler months	- late summer - early fall
Voucher types	- specimens, with duplicates in park and other repository - non-destructive samples for rare species - photos	- specimens - non-destructive tissue samples - photos	- specimens - photos	- photos - audio	- specimens / skulls - photos - tracks	specimens

Table 8. Cover-abundance scale classes used in plot sampling of vascular plant species (after Peet et al. 1998).

Cover range	Scale value	Class midpoint
Missing but nearby	--	--
Solitary or few individuals	1	0.3
0-1%	2	0.5
1-2%	3	1.5
2-5%	4	3.5
5-10%	5	7.5
10-25%	6	17.5
25-50%	7	37.5
50-75%	8	62.5
75-95%	9	85.0
95-100%	10	97.5

Amphibians and Reptiles

For the past three years, USGS-BRD scientists have been working on a study of how to inventory the diverse amphibian fauna in GRSM. When applicable, their methodologies may be utilized in other parks for the purpose of these inventories. A variety of collecting methods will be employed to sample herpetofauna, both within and outside of the grid plots. Typically, complete species inventories of amphibians and reptiles involve searching for and collecting specimens in all possible microhabitats, both during the day and night to coincide with diurnal activity patterns. Because timing is critical to the success of an amphibian and reptile inventory, seasonal patterns of activity will be considered as well. Within the plots, the above described bird inventory design will be set up in a way so that it can be used for herpetofauna as well. Transect lines used for bird counts will be flagged and used as fixed width (5-10 m) natural cover transects.

Both amphibians and reptiles can be sampled efficiently with time and/or area constraint surveys, drift fence arrays, and cover boards. Pitfall trap arrays, used in conjunction with funnel traps, will sample snakes, lizards, and fossorial species rarely encountered visually. Road-cruising surveys will be used to sample night-active species such as snakes and toads. Additionally, roadkill specimens can provide valuable data on species composition. Specific aquatic methods will include litter bags in, or near, streams for salamanders, call surveys for frogs, egg counts for some salamanders and frogs, and live traps for some turtles.

Birds

The point count is a common method used for assessing bird species in an area. This is a count of all birds detected visually or aurally by a single observer at equidistant points in a line transect during a specified period (e.g., 10 minutes). We recommend that a line transect and three point counts be conducted at each plot, with points being a minimum of 250 m apart. Observers should do a 10 minute variable radius count on an outer point in the plot, then record all birds on a variable width transect as they walk to the center point of the plot, conduct another 10 minute variable radius count, and continue along the transect to the third point, where they conduct the third 10 minute count. Counts typically are made under acceptable weather conditions during the breeding season (May-June), although other seasons frequently are sampled as well. Bird species will be identified and recorded, along with their sex and age if desired, and their distance from the observer will be estimated as closely as possible. Distance data may then be grouped into intervals during analysis. Adding the distance factor to a simple point count allows for estimation of detection rates, an important component of an accurate survey (see Fancy and Sauer 2000). The time of first detection will be listed in two categories: within the first 3 minutes, or during the last 2 minutes (given a 5 minute count period). All of this information will be mapped on a bulls-eye data sheet, which depicts a circle around a given point and is marked with distances and directions. Each point will be geographically referenced, and will be at least 250 m apart for off-road counts, and approximately 1 km apart for roadside counts to avoid encountering the same birds.

Mammals

There is no assurance that a species count based on any particular sampling method fairly represents the true number of species in an area; therefore, a variety of techniques will be employed for determining mammal species richness. Trapping may provide a good estimate for small herbivorous or granivorous mammals, but it will provide no data or only limited data on larger mammals and on certain small mammal taxa that are difficult to capture without specialized trapping techniques.

Sherman live-traps often are used for sampling small nocturnal rodents; however, depending on the behavior of certain species, this method may not capture everything. For example, shrews rarely are captured in live-traps; therefore, other methods must be employed, such as raking litter and turning cover. For larger mammals, scat and track surveys, attractants and scent posts can be used in conjunction with infra-red triggered photography, and the resulting photograph can be used as a voucher.

For conducting bat inventories, mist nets are the most common collecting device, often placed at entrances to roosts and over water sources used by bats. Each species of bat has its own specific vocalization, and the latest technology in bat research is a device for recording these calls. Referred to as Anabat, this audio recording device, along with accompanying software, often can accurately determine bat species based on recorded calls.

Fish

When inventorying fish, it is important to recognize and thoroughly sample all habitat types. Streams in these network parks vary in size but most are lower order and will not require extensive equipment. Seines, backpack electrofishers, or both will be employed for smaller streams, depending on local geomorphology and in-stream cover. Larger streams will require the same techniques as small streams, but a boat-mounted electrofisher may be needed for deep water. Lakes, which are found only in three network parks (NISI, CARL, and MACA), will be sampled by seine and a boat-mounted electrofisher.

Early fall and late summer are the ideal seasons for sampling lotic systems. Series of specimens representing all size classes will be collected, except in the case of rare species with localized populations, when only the minimum necessary will be collected.

F. Data Analysis

A variety of techniques will be used to analyze, summarize, and report data collected in this inventory. GIS maps will be incorporated into final reports to spatially represent distributions of species within parks. Data also will be analyzed across the combined networks to determine how species are regionally distributed.

Field data sheets will be reviewed for errors after data are collected and will be photocopied and stored at multiple locations until data entry is complete. Data will be entered in the I&M Database Template of Microsoft Access, or a compatible database, and data screening programs may be used to detect potential data entry errors. Copies of all data files will be stored with individual researchers and at a centralized location within the combined networks.

Estimation of Inventory Completeness

Given that parks in these two networks usually are anomalous areas of the southern landscape, it is impossible to accurately predict the number of species in any of these groups. Therefore, throughout the course of the inventory, data analysis will be conducted as an iterative process to determine when 90% of the species in each taxon have been inventoried. The number of species inventoried will be examined as a function of sampling effort, as defined for each taxonomic group. For vascular plants, species richness will be plotted against total area or number of plots sampled. For other groups, such as small mammals, sampling effort may be quantified as the number of trap nights or the number of sampling days. Data will be examined for each taxonomic group following the sampling of grid-point plots to help determine how many stratified sample plots should be selected to reach the 90% inventory goal.

Species richness vs. sampling effort relationships will be examined by strata within each park. Examination by strata will allow researchers to pinpoint which strata need additional sampling and which have received sufficient attention. In addition to plotting species richness against sampling effort, we also will use the Lincoln-Peterson Index (Southwood 1978) as a supporting analysis of inventory completeness. This analysis technique is based on mark-recapture methods (Thompson 1992) and allows the estimation of species richness based upon two species lists

generated independently at two different times. Related programs to estimate species richness are available from the USGS Patuxent Wildlife Research Center at <http://www.mbr-pwrc.usgs.gov/software.html>. These programs were written by J. E. Hines based on methods described by Burnham and Overton (1979).

Many of the parks in the combined networks have had partial inventories completed for vascular plants and vertebrates. Since these inventories usually were not based on a systematic design, assessing their completeness is difficult. To determine completeness, a subset of grid-point and stratified sample plots (5-10 plots per strata) will be inventoried for each taxonomic group. If unlisted species do not comprise more than 10% of all species encountered, then the inventory will be considered 90% complete. If unlisted species comprise greater than 10% of total species, a more extensive inventory will be conducted.

Species Distribution Maps

The proposed study design will allow the collection of systematic information about the distribution of species. Species occurrences for all vertebrates at each park will be mapped based upon presence/absence data. For vascular plants, cover class midpoints (Table 8) will be averaged by species at each plot and used to represent species importance. These values will be used to compare individual species importance among community types and create quantitative distribution maps. GIS can be used to plot maps of predicted species abundance or predicted probability of species occurrence. The predictor variables from regression or discriminant analysis (respectively) then can be used as GIS layers to estimate these probability values at many grid points on the map. Inventory data will be used with site data (slope, aspect, slope position, etc.) and vegetation community classification data to model the distribution of species of special interest (e.g., exotics, threatened species) based upon encounter probabilities. For all special concern species, maps with geo-referenced coordinates for all inventoried populations also will be supplied.

Regional Analysis

In addition to examining species distribution within parks, this study will examine the distribution of individual species across the combined networks. Because many units of the National Park Service serve as relatively small islands of biological diversity that often are under pressure from surrounding land use, it is critical to understand which species may be endemic to one or a few units. This sort of information will not only help protect a species at a given NPS unit, but also will improve management of biological diversity at the regional level. For each species, a list of parks at which it occurs will be compiled.

III. LOGISTICAL SUPPORT AND COMPLIANCE

During site visits to each of the 17 parks, meetings were held with natural resource personnel and usually the superintendents. A series of questions were asked regarding the following topics: recent disturbances, new land acquisitions, special events that researchers may interfere with (e.g., battle re-enactments), restrictions on plot markings, available housing and workspace,

safety hazards, space/facilities for storing collections, and other topics. A variety of responses were received, but overall, local park personnel are very enthusiastic about this project and will make every effort to accommodate visiting scientists and provide the needed facilities.

With regard to housing and workspace, some parks have adequate facilities on-site that scientists are welcome to use. Parks that can provide both housing and workspace include ABLI, BISO, CUGA, CHCH, COWP, FODO, GRSM, GUCO, KIMO, MACA, and NISI. Those that can provide workspace only are BLRI, CARL, OBRI, and STRI. A few parks have nothing available (LIRI, RUCA, SHIL), in which case researchers may need to stay in local hotels, campgrounds, bed & breakfasts, etc.

SAFETY is of the utmost concern for scientists visiting these parks. It is assumed that they will be experienced and familiar with field conditions; however, we still felt that this was an important question to ask at each park. Some of the potential safety issues which were brought up during our park visits include the following: venomous insects and snakes, ticks (may carry Lyme disease), hazard trees, abandoned mine shafts, cave openings, difficult terrain, hunting seasons, busy roads, and crime in adjacent communities. Park-specific safety information will be made available to scientists working in those areas.

All investigators must apply for and be approved for a collecting permit in those parks where they will conduct research. These permits will instruct the principal investigator as to the terms that are allowed by the NPS and the specific park for which work is proposed. Federal law requires that actions which could affect natural or cultural resources on federal lands, including research, inventories, and monitoring, be preceded by proper assessment and documentation of potential impacts. The type of inventories discussed in this plan typically do not cause concern regarding the National Historic Preservation Act; however, it is at each park's discretion to restrict activities if it is felt that a Section 106 compliance issue has arisen. With regard to the National Environmental Policy Act (NEPA), these inventories are categorically excluded except when the Endangered Species Act is an issue. Researchers who anticipate handling endangered species must obtain a permit from the U. S. Fish and Wildlife Service and be familiar with park, state, and federal regulations regarding the handling, disposition, and recovery plans for listed species. Additionally, it is incumbent upon all scientists involved in these inventories to be aware of T&E species that they may encounter.

IV. BUDGET (omitted)

V. COMPLETION SCHEDULE

The timeline for completion of biological inventories in these two networks (Table 10) is based on anticipated allocation of funds, the existing levels of inventory completion, the level of funding necessary to carry out inventories for specific taxa, and the urgency for some taxonomic groups (e.g., amphibians).

Table 10. Schedule of events per year for plot establishment and conducting inventories for each taxonomic group.

	FY 00	FY 01	FY 02	FY 03	FY 04	FY 05
Plot setup	• planning • air photos • held BRD workshop	• complete air photos for ~13 parks • plot set-up (grid)	• plot set-up (stratified) • specialized habitat selection	• additional stratified plots • habitat type report		
Plants	• evaluated checklists for completeness • gathered literature • held BRD workshop	• begin structured sampling of plots in 8 parks: 7 with partial lists (ABLI, BISO, BLRI, CARL, CHCH, COWP, CUGA), and 1 with no list (GUCO)	• sample stratified plots and specialized habitats	• sample stratified plots • data analysis • final report		
Amphibians	• gathered literature • held BRD workshop	• obligate funds	• begin field work in 6 largest parks (BISO, BLRI, CHCH, CUGA, LIRI, MACA), excluding GRSM	• continue work in original 6 parks • begin sampling in remaining 11 parks	• complete field work in all parks • data analysis • final report	
Reptiles	• gathered literature • held BRD workshop	• obligate funds	• begin opportunistic collections in each park by local park staff • record reptile occurrences found in amphibian sampling	• begin comprehensive field sampling	• finish field work through summer • data analysis • final report	
Birds	• gathered		• begin breeding bird	• continue bird work	• data analysis	

	literature • held BRD workshop		field work		• final report	
Mammals	• gathered literature • held BRD workshop			• begin mammal work	• data analysis • final report	
Bats	• gathered literature • held BRD workshop			• begin Anabat surveys, supplemented with mist net captures. • winter cave surveys	• complete bat work • data analysis • final report	
Fish	• gathered literature • held BRD workshop			• begin fish sampling	• complete field work • data analysis • final report	
Special concern species		• incidental observations as part of concurrent sampling (in all groups) within standardized design	• incidental observation as part of concurrent sampling (in all groups) within standardized design	• incidental observation as part of concurrent sampling (in all groups) within standardized design	• incidental observation as part of concurrent sampling (in all groups) within standardized design	• full comprehensive specialized surveys on species of high management concern

VI. VOUCHER SPECIMENS

Research permits are required for all collections, even if the specimens are not intended for deposit. Permit request forms and permits are available from each park. The permit process serves as a record that identifies collector(s), purpose of the collection, and clearly states any restrictions. Federally threatened and endangered species (T&E) will not be killed.

Collections will be attempted for all species (except for T&E or if a specimen already exists). High quality voucher photographs will be used for medium and large mammals and birds unless an animal or bird is found dead (e.g., roadkill). Voucher photographs will be clearly labeled with the park name, species name, specific location, date, and photographer. Copies will be made and permanently archived in two places.

If collection is planned, curation must be written into the project (attached to the permit request). Many network parks do not have the space or proper facilities to serve as a repository for voucher specimens; therefore, arrangements will be park-specific depending upon needs and preferences. We plan to have cooperators catalog species information into NPSpecies or a compatible database, which hopefully will allow NPS staff to easily obtain the records and transfer to ANCS+. Collected specimens then will be deposited in park collections or university natural history museums.

VII. PRODUCTS AND DELIVERABLES

Data management will continue with the documentation of existing lists, datasets, GIS layers, and bibliographic records for both networks (the status of data deliverables is presented in Appendix B). A cooperative agreement has been established with Western Kentucky University (WKU) through Spring 2001 to continue data mining and development of NPSpecies and NRBib (as of 1 December 2000, data from 17 parks have been entered into NPSpecies, and bibliographic information has been entered for all 18 parks). Voucher documentation, a process which is just beginning, will also continue as part of this cooperative agreement. This work is being conducted at MACA (30 miles from WKU). Also, the Cumberland/Piedmont Network is establishing two positions in FY01, both at MACA. These positions are: I&M Coordinator and I&M Data Manager. Data management and implementation of the inventory study plan (development of scopes-of-work, cooperative agreements for the inventory field work, etc.) for both networks will be coordinated through these positions. The Appalachian Highlands Network also will hire an I&M Coordinator and Data Manager in FY01 (see Appendices D and E). These positions will be funded out of the Network *Monitoring* budget.

Standardized products will be written into scopes-of-work for all inventory projects, and copies of all products will be delivered to parks in the networks, to the regional I&M Coordinator, and to the servicewide I&M Program Manager. Annual reports summarizing inventory progress will be prepared by the two network I&M Coordinators. Assistance with completion of the databases and GIS layers will be provided by the Network Data Managers where needed, such as with park-level ANCS+ entry, cataloging datasets, and incorporation of GIS layers into the NPS

Theme Manager. The inventory deliverables will include the following:

1. Species data: NPSpecies format
2. Field data: I&M Database Template (MSAccess) format (under development) with any necessary modifications to capture special needs
3. GIS layers: ArcView format, organized into NPS Theme Manager directory structure
4. Metadata: FGDC format
5. Voucher data: ANCS+ format and NPSpecies, voucher section format
6. References: NPBib format linked to NPSpecies
7. Datasets: Dataset Catalog, NPBib, or whatever is recommended to catalog datasets
8. Annual/Final reports: MS Word

The networks' two new I&M coordinators will oversee the overall coordination of this project and form a board of directors together with the superintendents of network parks. A technical committee, consisting of park natural resource managers and outside scientists, plus the five members of the study plan development team (i.e., K. Langdon, T. Leibfreid, M. Jenkins, B. Nichols, J. Rock), will serve in an advisory capacity and provide reviews of policies, plans, progress reports, etc. An annual meeting of the board of directors and the technical committee would be advantageous.

When selecting scientists to conduct these inventories, an agreed upon selection priority should be followed. We suggest that potential cooperators include: USGS-BRD scientists, not-for profit organizations, CESU's, major universities, and perhaps NPS staff. Parks may have input on the selection process, particularly if they have a pre-existing working relationship with a local university and would like it to continue, in which case priority would be considered for that institution. For each taxonomic group, a coordinating institution may be established to ensure standardization and quality of deliverables across all parks. We strongly recommend that these inventories be conducted by well qualified scientists and researchers to obtain the highest quality product.

VIII. LITERATURE CITED

- Anderson, M., P. Bourgeron, M. T. Bryer, R. Crawford, L. Engelking, D. Faber-Langendoen, M. Gallyoun, D. H. Grossman, S. Landaal, K. Metzler, K. D. Patterson, M. Pyne, M. Reid, L. Sneddon, and A. S. Weakley. 1998. Terrestrial vegetation of the United States, volume 2. The national vegetation classification system: list of types. The Nature Conservancy, Arlington, VA.
- Burnham, K. P. and W. S. Overton. 1979. Robust estimation of population size when capture probabilities vary among animals. *Ecology* 60(5): 927-936.
- Fancy, S. G. and J. R. Sauer. 2000. Recommended methods for inventorying and monitoring landbirds in national parks. National Park Service, Fort Collins. 13 p.
- Fortin, M. J., P. Drapeau, and P. Legendre. 1989. Spatial autocorrelation and sampling design in plant ecology. *Vegetatio* 83: 209-222.
- Grossman, D. H., D. Faber-Langendoen, A. S. Weakley, M. Anderson, P. Bourgeron, R. Crawford, K. Goodin, S. Landaal, K. Metzler, K. Patterson, M. Pyne, M. Reid, and L. Sneddon. 1998. Terrestrial vegetation of the United States, volume 1. The national vegetation classification system: list of types. The Nature Conservancy, Arlington, VA.
- Hammer, R. D., F. J. Young, N. C. Wollenhaupt, T. L. Barney, and T. W. Haithcoate. 1995. Slope class maps from soil survey and digital elevation models. *Soil Society of America Journal* 59: 509-519.
- Legendre, P. and M. J. Fortin. 1989. Spatial pattern and ecological analysis. *Vegetatio* 80: 107-138.
- National Park Service. 1999. Guidelines for biological inventories. Inventory and Monitoring Program, National Park Service. www1.nature.nps.gov/sfancy.
- Olson, R. and M. Franz. 1998. A vegetation habitat classification for Mammoth Cave National Park. *In: Proceedings of Mammoth Cave National Park's Seventh Science Conference*, pp. 19-25. Mammoth Cave, KY.
- Peet, R. K., T. R. Wentworth, and P. S. White. 1998. A flexible multipurpose method for recording vegetation composition and structure. *Castanea* 63: 262-274.
- Southwood, T. R. E. 1978. Ecological methods with particular reference to insect populations, second edition. Chapman and Hall, London. 524 p.
- Thompson, S. K. 1992. Sampling. John Wiley and Sons, Inc., New York. 343 p.

- TNC [The Nature Conservancy] and ESRI [Environmental Systems Research Institute]. 1994. Field methods of vegetation mapping: NBS/NPS Vegetation Mapping Program. The Nature Conservancy, Arlington, VA.
- Weakley, A. S., K. D. Patterson, M. Pyne, J. Teague, R. Evans, and M. J. Russo. 2000. International classification of ecological communities: terrestrial vegetation of the southeastern United States. (working draft) Association for Biodiversity Information / The Nature Conservancy, Durham, NC.

IX. APPENDICES

Appendix A: Checklist of references obtained from NRBib and park files for each park in the combined networks.

ABLI

Plants:

Johnson, K. 2000. Invasive plant management in smaller parks of the Appalachian Cluster.

McKinney, L. E. 1993. The flora and vegetation of Abraham Lincoln Birthplace National Historic Site. Kentucky State Nature Preserves Commission. Cooperative Agreement no. CA-5530-2-9003. 25 p.

Herpetofauna:

no information

Birds:

Sturgeon, M. and S. Kistler. 1994. Bird list of Abraham Lincoln Birthplace National Historic Site. Hodgenville, KY.

Mammals:

no information

Fish:

no information

BISO

Plants:

Allawos, J. G. 1994. The vascular flora of north White Oak Creek gorge, Scott and Fentress Counties, Tennessee. University of Tennessee, Knoxville. 167 p.

DeSelm, H. R. Flora and vegetation of the barrens of the Cumberland Plateau of Tennessee. *In:* Snyder, D. H. (ed.), Proceedings of the contributed paper sessions of the fourth annual symposium on the natural history of the lower Tennessee and Cumberland River valleys. The Center for Field Biology, Austin Peay State University, Clarksville. 38 p.

Medley, M. E. and D. F. Harker, Jr. 1979. The flora and vegetation of the river bars of the Big South Fork of the Cumberland River in McCreary County, Kentucky. *In:* Annual meeting of the Kentucky Academy of Sciences at Northern Kentucky University; Highland Heights. 25 p.

Patrick, T. S. 1979. Upper Cumberland River flora project. Final report. 170+ p.

Wofford, B. E., T. S. Patrick, L. R. Phillippe, and D. H. Webb. 1979. The vascular flora of Savage Gulf, Tennessee. *SIDA* 8(2): 135-151.

Birds:

Ganier, A. F. 1937. Summer birds of Pickett Forest. *Migrant* 8: 24-27.

Nicholson, C. P. 1982. The birds of Pickett County, Tennessee. *Migrant* 53(2): 25-26.

Stedman, S. J. 1995. Report on breeding bird surveys and censuses conducted in Big South Fork National River and Recreation Area - May-June, 1995. Tennessee Technological University. 511 p.

Stedman, S. J. 1995. Spring neotropical migrant survey in the Big South Fork National River and Recreation Area, Kentucky and Tennessee. *Kentucky Warbler* 71(2): 27-32.

Herpetofauna:

Echtarnacht, A. C. 1979. Herpetofauna of potential natural areas of the eastern upper Cumberland River drainage in Tennessee. Final report. 129 p.

Stephens, D. E. 1979. A survey of the amphibians and reptiles of McCreary County, Kentucky. Eastern Kentucky University, Richmond. 176 p.

Mammals:

Army Corps of Engineers. 1976. Final environmental impact statement: Establishment, administration, and maintenance of the Big South Fork National River and Recreation Area, Tennessee and Kentucky. U.S. Dept. of Army, Nashville District, COE.

Fish:

Branson, B. A. 1975. The fishes of the Daniel Boone National Forest. 75+ p.

Brazinski, M. J. 1979. Fish populations of the New River, Tennessee: A system receiving acid coal mine drainage. Tennessee Technological University, Cookeville. 68 p.

Comiskey, C. E. 1970. The fishes of the Big South Fork of the Cumberland River system. University of Tennessee, Knoxville. 92 p.

Comiskey, C. E. and D. A. Etnier. 1972. Fishes of the Big South Fork of the Cumberland River. *Journal of the Tennessee Academy of Science* 47(4): 140-146.

Etnier, D. A. 1976. A check list of the fishes of Tennessee. 6 p.

Kirsch, P. H. 1893. Notes on a collection of fishes from the southern tributaries of the Cumberland River in Kentucky and Tennessee. *Bulletin of the United States Fish Commission* 11: 258-268.

O'Bara, C. J., R. D. Estes, F. J. Bulow, J. D. Rector, and L. E. Rider. Range extension and distribution of rare fishes of the Big South Fork of the Cumberland River, Tennessee and Kentucky. 7 p.

O'Bara, C. J., W. L. Pennington, and W. P. Bonner. 1982. A survey of water quality, benthic macroinvertebrates and fish for sixteen streams within the Big South Fork National River and Recreation Area. United States Army Corps of Engineers, Nashville, TN. Contract # DACW62-81-C-0162. 175+ p.

O'Bara, C. J., J. D. Rector, and F. J. Bulow. 1984. Distribution of uncommon fishes of the Big South Fork of the Cumberland River, Tennessee and Kentucky. *Journal of the Tennessee Academy of Science* 59: 69-71.

General:

Anonymous. 1995. Historic and current records of federally and state protected species occurring in the Big South Fork NRR as of September 1, 1995. 13 p.

Big South Fork National River and Recreation Area Biological Conservation Database. 316 records.

Shoup, C. S. and J. H. Peyton. 1940. Collections from the drainage of the Big South Fork of the Cumberland River in Tennessee. *Journal of the Tennessee Academy of Science* 15: 106-116.

BLRI

Plants:

Anonymous. Checklist of 140 wildflowers of Balsam Gap sub-district, Blue Ridge Parkway. 65+ p.

Anonymous. Checklist of vascular plants found in western Carolina counties adjacent to the Blue Ridge Parkway, Balsam Gap sub-district. 39 p.

Anonymous. A descriptive list of some plants found on the Blue Ridge Parkway from Oteen north to Singe Cat Ridge overlook, North Carolina. 25 p.

Bancroft, W. L. 1968. Checklist of vascular plants of the Blue Ridge Parkway, mile post 305-355. 16 p.

Bliss, D. C. 1970. Blue Ridge Parkway flora: preliminary checklist. Randolph-Macon Woman's College, Lynchburg, VA. 56 p.

Bliss, D. C. 1970. Reference lists of native plants of some selected forest communities along the Blue Ridge Parkway. Randolph-Macon Woman's College, Lynchburg, VA. 18 p.

Brennan, T. M. 1987. Assay of plant species in a Blue Ridge Parkway vista, MP 36.5, 1985. *In: Summary of resource management activities and scientific studies relating to the Blue Ridge Parkway, 1983-1986*, pp. 19-26.

- Cason, A. L. 1994. A floristic survey of milepost 104 to milepost 144, Blue Ridge Parkway. National Park Service, Blue Ridge Parkway, report # 94-2. 30 p.
- Coleman, D. A. and J. D. Pittillo. 1987. Important plant habitats of the Blue Ridge Parkway, Roanoke to Rock Fish Gap, Virginia. 110 p.
- Dalmas, T. H. 1972. The flora of the headwaters of Cornelius Creek, Botetourt County, Virginia. 22 p.
- Edwards, R. L. 1996. Plant diversity survey of Rock Castle Gorge backcountry area, Blue Ridge Parkway, Virginia. Virginia Polytechnic Institute and State University, Blacksburg. Final report, contract # CA-5140-2-9003. 56 p.
- Kelsey, H. P. 1946. Flora of Grandfather Mountain. National Parks Magazine: Apr-Jun.
- Kendrick, R. A. 1978. Wildflowers of the Rocky Knob sub-district. 30 p.
- Kendrick, R. A. 1987. An annotated checklist of the wildflowers of the Rocky Knob district (revised 1984). *In: Summary of resource management activities and scientific studies relating to the Blue Ridge Parkway, 1983-1986*, pp. 27-64.
- Kendrick, R. A. 1990. An annotated checklist of selected trees and vines along Hardwood Cove self-guiding nature trail in the Rocky Knob district. *In: Resource management activities and scientific studies relating to the Blue Ridge Parkway 1987-1989*, pp. 39-45.
- Kendrick, R. A. 1990. Checklist of the trees of the Rocky Knob district. *In: Resource management activities and scientific studies relating to the Blue Ridge Parkway 1987-1989*, pp. 34-38.
- Kendrick, R. A. 1990. Showy shrubs and trees of the Rocky Knob district. *In: Resource management activities and scientific studies relating to the Blue Ridge Parkway 1987-1989*, pp. 48-52.
- Krochmal, A. 1976. A survey of plants along the Blue Ridge Parkway from Asheville to Mt. Pisgah and Julian Price Park: a preliminary report. 40+ p.
- Livengood, J. M. 1972. Vascular flora of the Sim's Pond area. M. S. Thesis, Appalachian State University, Boone, NC. 27 p.
- National Park Service. Exotic plant species control. *In: Resource management activities and scientific studies relating to the Blue Ridge Parkway 1987-1989*, pp. 76-79.
- National Park Service. 1979. Rakes Mill Pond [wildflower list]. 2 p.
- National Park Service. 1997. Wildflower checklist of the Plateau district, Blue Ridge Parkway. 10 p.
- National Park Service. 1970. Wildflowers of Fancy Gap sub-district. 6 p.

- National Park Service. Wildflowers of the Rocky Knob sub-district, Blue Ridge Parkway, mile posts 165 to 217. 4 p.
- Olson, C. S. 1994. Checklist of blooming plants within five wildflower display areas in the Asheville District, Blue Ridge Parkway. National Park Service, Blue Ridge Parkway, report # 94-4. 5 p.
- Pittillo, J. D. and T. E. Govus. 1978. Important plant habitats of the Blue Ridge Parkway from the Great Smoky Mountain National Park to Roanoke, Virginia. 480 p.
- Ramsey, G. W. 1994. Plant diversity survey of the Peaks of Otter backcountry area, Blue Ridge Parkway, Virginia. Lynchburg College, Lynchburg, VA. Final report, contract # CA 5140-1-9002, SUB A #1. 71 p.
- Smathers, G. A. Native orchids of the Blue Ridge Parkway in North Carolina [checklist]. 2 p. MAY CONTAIN SENSITIVE INFORMATION
- Smathers, G. A. 1958. The flora of the Blue Ridge Parkway in North Carolina. 8 p.
- Stewart, N. 1957. Small woods plants and shrubs. National Park Service, Blue Ridge Parkway. 305 p.
- Teague, B. 1985. Vascular plants - Blue Ridge Parkway. 30 p.
- Teague, B. 1987. Preliminary checklist of the vascular flora of the Blue Ridge Parkway, 1986. *In*: Summary of resource management activities and scientific studies relating to the Blue Ridge Parkway, 1983-1986, pp. 79-109.
- Thacker, W. 1967. Checklist of plants, Crabtree Meadows, Blue Ridge Parkway. 4 p.
- Wayt, J. 1987. A preliminary checklist of the vascular flora of the Tanawha Trail in the Cone-Price district. *In*: Summary of resource management activities and scientific studies relating to the Blue Ridge Parkway, 1983-1986, pp. 10-18.
- Wiser, S. K. 1994. High-elevation cliffs and outcrops of the Southern Appalachians: vascular plants and biogeography. *Castanea* 59(2): 85-116.
- Birds:
- Anonymous. 1990. Birds most likely seen: Doughton Park, Blue Ridge Parkway, Alleghany County, NC. 2 p.
- Bachmann, J. 1986. Breeding bird atlas project - Rocky Knob district. 12 p.
- Bancroft, W. L. 196?. Blue Ridge Parkway birds, mile post 305-355 [check list]. 5 p.
- Kendrick, R. A. 1978. Birds of the Rocky Knob sub-district, Blue Ridge Parkway, and their seasonal occurrence. 8 p.

Papahronis, F. B. 1990. Checklist of nesting birds observed in Rocky Knob picnic area. *In: Resource management activities and scientific studies relating to the Blue Ridge Parkway 1987-1989*, pp. 94-96.

Roanoke Valley Bird Club. 1968. Preliminary checklist of birds, Virginia Peaks of Otter area, Blue Ridge Parkway, National Park Service. 14 p.

Herpetofauna:

Bancroft, W. L. 196?. Blue Ridge Parkway reptiles and amphibians, mile post 305-355 [check list]. 2 p.

Howard, J. 1987. Annotated checklist of amphibians and reptiles of the Rocky Knob district. *In: Summary of resource management activities and scientific studies relating to the Blue Ridge Parkway, 1983-1986*, pp. 111-119.

Olsen, T. C. 1967. List of reptiles and amphibians, Blue Ridge Parkway. 52 p.

Petranka, J. W., F. Scott, and C. K. Smith. 1995. Protocol manual for long-term monitoring of amphibians in GRSM-BLRI-CUGA-MACA. Cooperative Agreement # CA-5140-1-9001, sub-agreement # CA-5140-3-9002.

Witt, W. L. 1972. Amphibians and reptiles of Blue Ridge Parkway from Roanoke to Shenandoah National Park, a checklist. 113 p.

Mammals:

Bancroft, W. L. 196?. Blue Ridge Parkway mammals, mile post 305-355 [check list]. 2 p.

Chamberlain, J. L. 1969. Mammals of the Blue Ridge Parkway. Utica College, Utica NY. 57 p.

Fish:

Compilation of individual surveys maintained by stream name - Blue Ridge Parkway. (B. Teague, pers. comm.).

General:

Cherry, R. Threatened, endangered or rare species, Bluffs district, Blue Ridge Parkway. 85+ p.

National Park Service. 1979. Endangered and threatened plants and animals along the Blue Ridge Parkway in the counties of Bedford, Botetourt and Rockbridge, Virginia. 5 p.

Randall, J. F. 1990. Biological survey of the area below Trout Lake Dam. *In: Resource management activities and scientific studies relating to the Blue Ridge Parkway 1987-1989*, pp. 62-65.

Smith, A. B. 1990. Field survey of rare and unusual plants and animals, Fisher Peak, Virginia and North Carolina. Final report, contract # PX514081056. 14 p.

Smith, C. and J. Day. 1979. Natural resource inventory of Parkway land between milepost 187 and 187.4. 92 p.

Smith, S. M. 1990. Stream survey of Otter Creek, Amherst County, VA. Virginia Department of Game and Inland Fisheries. 6 p.

Wright, R. A. S. and B. Teague. 1992. Flora checklist of Purgatory Mountain Overlook wildflower display area, Blue Ridge Parkway. National Park Service, Blue Ridge Parkway, report # 92-6. 11 p.

CARL

Plants:

Blaha, M. and A. Ulinski. 1994. Plant identification study, Carl Sandburg Home National Historic Site, Flat Rock, N.C. 1993-1994. 1510 p.

Jones, C. W. 1992. Flowering trees checklist, Carl Sandburg Home NHS. 142 p.

Jones, C. W. 1992. Wildflower checklist, Carl Sandburg Home NHS. 142 p.

Remaley, T. 1998. Survey of invasive exotic plants, Carl Sandburg National Historic Site.

Birds:

Anonymous. 1991. Master list of birds sighted at CARL as of 3/20/91. 2 p.

Audubon Society. 1994. Birds of Carl Sandburg Home National Historic Site, Flat Rock, NC, National Park Service. [brochure-updated 1999 by Ed Caldwell]

Herpetofauna:

no information

Mammals:

no information

Fish:

Master's Thesis on file. Carl Sandburg National Historic Site. (Warren Weber, pers. comm.)

General:

Anonymous. Wildlife sighted at Carl Sandburg Home National Historic Site 'Connemara'. 2p.

LeGrand, H. 1992. Potential species. North Carolina Natural Heritage Program. (S. Benjamin, pers. comm.)

CHCH

Plants:

- Dickson, C. C. and C. F. Nixon. 1989. Taxa of special concern, Sunset Rock area, Chickamauga and Chattanooga National Military Park, with attached flora. 35+ p.
- Gallyoun, M., G. Meyer, A. Andreu, and W. Slocumb. 1995. Mapping vegetation communities with the Nature Conservancy's vegetation classification system on five small national parks in the southeastern U.S.A. Chapel Hill, NC. 100+ p.
- Johnson, K. 1985. Protected plants - Chickamauga Battlefield. 1 p.
- Rogers, C. L., M. J. Ratnaswamy, and R. J. Warren. 1993. Vegetation communities of Chickamauga Battlefield National Military Park, Georgia. University of Georgia, Athens. Technical report NPS/SERCHCH/NRTR-93/11. 75 p.
- Rosen, A., I. Owen, and S. P. Bratton. 1982. Exotic plant survey of the Lookout Mountain unit at Chickamauga and Chattanooga National Military Park. National Park Service, Atlanta, GA. 14 p.
- Sutter, R., S. Benjamin, S. Rollins, G. Livingstone, and N. Rudd. 1994. Baseline monitoring of calcareous glades at Chickamauga-Chattanooga National Military Park. The Nature Conservancy, Southern Heritage Task Force, Chapel Hill, NC. 100+ p.
- Van Horn, G. S. 1980. Additions to the cedar glade flora of northwest Georgia. *Castanea* 45: 134-137.
- Van Horn, G. S. 1981. A checklist of the vascular plants of Chickamauga and Chattanooga National Military Park. *Journal of the Tennessee Academy of Science* 56(3): 92-99.

Birds:

- Anonymous. 1963. The birds of Chickamauga Park. 11 p.
- Anonymous. 1994. Birds of Chickamauga and Chattanooga National Military Park. 7 p.
- Comstock, R. L. 1963. A list of birds compiled by Rock L. Comstock while stationed at Chickamauga Battlefield 1958-1963. 11 p.

Herpetofauna:

no information

Mammals:

no information

Fish:

no information

General:

Hobbs, H. H., III. 1993. Assessment of the ecological resources of the caves of Russell Cave National Monument, Jackson county, Alabama and of selected caves at the Lookout Mountain unit of Chickamauga-Chattanooga National Military Park, Dade County, Georgia.

COWPPlants:

Bratton, S. P. and T. Butler. 1982. The distribution of exotic plant species at Cowpens National Battlefield. 18 p.

King, C. W. 1997. Botanical survey of Cowpens National Battlefield. University of Southern Mississippi. 18 p.

Newberry, G. 2000. Rare plant distribution and proposed collection and inventory of herbaceous plants. University of South Carolina-Spartanburg.

Patton, E. G. 1996. Ten notable trees on the interpretive walking trail of Cowpens National Battlefield. 6 p.

Birds:

Piedmont Audubon Society. 1997. Cowpens Bird Checklist.

Herpetofauna:

Thomas, R. B. 1999. Proposed herpetofaunal inventory of the Cowpens National Battlefield. University of South Carolina-Spartanburg. Partial funding received through Small Park NRPP, FY2000.

Mammals:

Ferris, D. K. 2000. Proposed mammal inventory of the Cowpens National Battlefield. University of South Carolina-Spartanburg. Partial funding received through Small Park NRPP, FY2000.

Fish:

no information

CUGAPlants:

Bratton, S. Exotic plant study.

Ettman, J. K. 1976. An annotated checklist of the Orchidaceae of Bell County, Kentucky. Annals of Kentucky Natural History 3.

Hinkle, C. R. 1975. A preliminary study of the flora and vegetation of Cumberland Gap

National Historical Park. University of Tennessee, Knoxville. 236 p.

Pounds, L., T. S. Patrick, and C. R. Hinkle. 1989. Rare plant assessment and checklist for Cumberland Gap National Historical Park. National Park Service, contract # PX-546042127. 79 p.

Birds:

Barbour, R. W., W. H. Davis, and R. A. Kuehne. 1979. The vertebrate fauna of Cumberland Gap National Historical Park. Final report, contract # C X 5000 71 232. 82 p.

Herpetofauna:

Burger, W. L. 1958. Herpetological specimens collected in the Cumberland Gap National Historical Park and vicinity. College of William and Mary, Williamsburg, VA. 9 p.

Nicholson, C. S. 1978. Distribution patterns of Amphibia in Cumberland Gap National Historical Park. University of Arizona, Tucson. 162 p.

Petranka, J. W., F. Scott, and C. K. Smith. 1995. Protocol manual for long-term monitoring of amphibians in GRSM-BLRI-CUGA-MACA. Cooperative Agreement # CA-5140-1-9001, sub-agreement # CA-5140-3-9002.

Mammals:

Barbour, R. W., W. H. Davis, and R. A. Kuehne. 1979. The vertebrate fauna of Cumberland Gap National Historical Park. Final report, contract # C X 5000 71 232. 82 p.

Currie, R. Bat lists from Cumberland Gap National Historical Park (ongoing). Kentucky Department of Fish and Game. (R. Collier, pers. comm.).

Fish:

Stephens, D. E. 1996. The 1996 survey of the fish fauna in Little Yellow Creek (Bell County, Kentucky). 15+ p.

Woodall, B., E. Woodall, and J. E. Copeland. 1984. A preliminary study of the water quality and ichthyofauna of selected streams in Cumberland Gap National Historical Park. 113 p.

General:

Barbour, R. W., W. H. Davis, and R. A. Kuehne. 1979. The vertebrate fauna of Cumberland Gap National Historical Park. Final report, contract # C X 5000 71 232. 82 p.

FODO

Plants:

Chester, E. W. 1986. The vascular flora of Fort Donelson National Military Park, Stewart County, Tennessee. University of Tennessee, Knoxville. Research/Resources

Management Report SER-80. 86 p.

Birds:

U. S. Fish and Wildlife Service. 1999. Cross Creeks National Wildlife Refuge bird list. Dover, TN. [brochure].

Herpetofauna:

no information

Mammals:

no information

Fish:

no information

GRSM

Plants:

Amoroso, J. L. 1997. Natural Heritage Program list of the rare plant species of North Carolina. North Carolina Division of Parks and Recreation, 1997 edition. 78 p.

Anonymous. 1989. Checklist of flowering plants of Great Smoky Mountains National Park. 33 p.

Anonymous. 1988. Ozone sensitive plant list for Great Smoky Mountains National Park - 1988. 1p.

DeFoe, D., K. R. Langdon, and J. Rock. 1989. Flowering plants of the Great Smoky Mountains National Park -- revised 1995. National Park Service. 48 p.

Drew, W. B. 1937. Notes on asters in east Tennessee. *Castanea* 2: 75-82.

Hoffman, H. L. 1952. Checklist of vascular plants of the Great Smoky Mountains. 44 p.

Hoffman, H. L. 1966. Notes on vascular plant families in the Great Smoky Mountains. *Castanea* 31(4): 301-310.

Hoffman, H. L. 1966. Supplement to checklist, vascular plants, Great Smoky Mountains. *Castanea* 31(4): 307-310.

Hoffman, H. L., R. E. Shanks, and A. J. Sharp. 1958. Flowering plants of the Great Smoky Mountains, a checklist of families and genera. University of Tennessee, Knoxville. 14 p.

Jennison, H. M. 1929. A preliminary checklist of the spring wildflowers and ferns of Tennessee. *Journal of the Tennessee Academy of Science* 4(2 suppl.): 1-32.

Mark, A. F. 1959. The flora of the grass balds and fields of the southern Appalachian

Mountains. *Castanea* 24: 1-21.

Pyne, M. and A. Shea. 1996. Tennessee Natural Heritage Program rare vascular plant list. Tennessee Division of Natural Heritage, Nashville. 1996 edition. 21 p.

Sharp, A. J. and A. Baker. First and interesting reports of flowering plants in Tennessee. *Castanea*: 178-185.

Stevenson, G. B. 1967. Spring wildflowers of the Great Smoky Mountains. Buckhorn Press, Gatlinburg, TN. 16 p.

White, P. S. 1982. Checklist of flowering plants of Great Smoky Mountains National Park. 32 p.

White, P. S. 1982. The flora of Great Smoky Mountains National Park; an annotated checklist of the vascular plants and a review of previous floristic work. National Park Service, Research\Resources Management Report SER-55. 217 p.

White, P. S. 1982. New and noteworthy plants from Great Smoky Mountains National Park, North Carolina and Tennessee. *Castanea* 47: 78-83.

White, P. S. 1986. State and nationally listed vascular plants of Great Smoky Mountains National Park. 23 p.

White, P. S. and S. P. Bratton. 1979. Cataloochee access road vegetation description. 88 p.

White, P. S. and B. E. Wofford. 1984. Rare native Tennessee vascular plants in the flora of Great Smoky Mountains National Park. *Journal of the Tennessee Academy of Science* 59(3): 61-64.

Birds:

Aldrich, J. W. and P. Goodrum. 1946. Breeding bird population of a cove hardwood forest in the Great Smoky Mountains: special report. U. S. Fish and Wildlife Service.

Alsop, F. J., III. 1995. Birds of the Great Smoky Mountains [checklist]. Great Smoky Mountains National Park, Natural History Association. 8 p.

Anonymous. Bird observations in Great Smoky Mountains National Park: 1938-1948. 17 p.

Ganier, A. F. 1926. Summer birds of the Great Smoky Mountains. *Journal of the Tennessee Academy of Science* 1(2): 34-40.

Stevenson, G. B. 1985. Birds of Great Smoky Mountains National Park. 32 p.

Stupka, A. 1963. Notes on the birds of Great Smoky Mountains National Park. University of Tennessee Press, Knoxville. 242 p.

Herpetofauna:

Anonymous. 1993. Amphibians and reptiles of the Great Smoky Mountains [checklist]. Great Smoky Mountains National Park, Natural History Association. 2 p.

Anonymous. 1991. Checklist of amphibians and reptiles of Great Smoky Mountains National Park. 3 p.

Depoe, C. E., J. B. J. Funderburg, and T. L. Quay. 1961. The reptiles and amphibians of North Carolina. *Journal of the Elisha Mitchell Scientific Society* 27(2): 125-136.

King, W. 1944. Additions to the list of amphibians and reptiles of Great Smoky Mountains National Park. *Copeia* 4: 2 p.

Necker, W. L. 1934. Contribution to the herpetology of the Smoky Mountains of Tennessee. *Bulletin of the Chicago Academy of Sciences* 6(1): 1-4.

Petranka, J. W., F. Scott, and C. K. Smith. 1995. Protocol manual for long-term monitoring of amphibians in GRSM-BLRI-CUGA-MACA. Cooperative Agreement # CA-5140-1-9001, sub-agreement # CA-5140-3-9002.

Schenck, M. J. 1938. An annotated index to salamanders in Willis King's field notes, Great Smoky Mountains National Park, 1935 - 1938. 30+ p.

Mammals:

Anonymous. 1989. Bat survey summary reports, Great Smoky Mountains National Park. 5 p.

Anonymous. 1964. Checklist of mammals of Great Smoky Mountains National Park. 3 p.

Anonymous. 1942. Mammals known to occur in the Great Smokies National Park. 1 p.

Komarek, E. V. and R. Komarek. 1935. Checklist of mammals of the Smokies. Chicago Academy of Sciences. 2 p.

Komarek, E. V. and R. Komarek. 1938. Mammals of the Great Smoky Mountains. *Bulletin of the Chicago Academy of Sciences* 5(6): 137-164.

Linzey, D. W. 1995. Mammals of Great Smoky Mountains National Park. McDonald & Woodward Publishing, Blacksburg VA. 140 p.

Linzey, D. W. 1995. Mammals of the Great Smoky Mountains National Park. *Journal of the Elisha Mitchell Scientific Society* 111(1): 1-81.

Linzey, D. W. and A. V. Linzey. 1968. Mammals of the Great Smoky Mountains National Park. *Journal of the Elisha Mitchell Society* 84(3): 383-414.

Fish:

Douglas, C. B. 1980. Fishes of Great Smoky Mountains National Park. 12 p.

Hazard, F. O. and W. King. 1932. Checklist of fishes. Wilmington College. 3 p.

Lennon, R. E. 1961. An annotated list of the fishes of Great Smoky Mountains National Park. Fish Control Laboratory, La Crosse, Wisconsin. 20 p.

Lennon, R. E. 1960. Fishes of Great Smoky Mountains National Park. U. S. Fish and Wildlife Service. 7 p.

Simbeck, D. J. 1990. Distribution of the fishes of the Great Smoky Mountains National Park. Southern Bindery, Knoxville, TN.

General:

Anonymous. 1998. Aquatic species of Great Smoky Mountains National Park. 14 p.

LeGrand, H. E., Jr. and S. P. Hall. 1997. Natural Heritage Program list of the rare animal species of North Carolina. North Carolina Division of Parks and Recreation, 1997 edition, 81 p.

GUCO

Plants:

Locket, W. E. A resource managers guide to exotic plant control at Guilford Courthouse N.M.P. (SERO)

Novac. 1976. Checklist of trees and shrubs.

Remaley, T. 1996. Survey of invasive exotic plants.

Seifert. 1938. Trees of Guilford Courthouse National Military Park.

Birds:

no information

Herpetofauna:

no information

Mammals:

no information

Fish:

no information

General:

Shahady, T. and F. Zirkle. 1983. A survey of the vertebrate animals inhabiting the Guilford Courthouse National Military Park. Guilford College. Unpublished report (SERO).

KIMO

Plants:

Barnett-Lawrence, M. 1993. Checklist of the vascular vegetation of Kings Mountain, Kennesaw Mountain, Chickamauga and Chattanooga, Shiloh, and Russell Cave. *In: Mapping vegetation communities with the Nature Conservancy's vegetation classification system on five small national parks in the southeastern U.S.A.* Chapel Hill, NC. 100+ p.

Gallyoun, M., G. Meyer, A. Andreu, and W. Slocumb. 1995. Mapping vegetation communities with the Nature Conservancy's vegetation classification system on five small national parks in the southeastern U.S.A. Chapel Hill, NC. 100+ p.

Johnson, K. and T. Remaley. 1996. Small parks exotic plant project report. Appalachian Cluster.

Kennemore, D. E. 1995. Floristics of the Kings Mountain National Military Park and the Kings Mountain State Park. University of South Carolina, Columbia. 103 p.

Birds:

Student Conservation Associate. 1999. Database of birds for Kings Mountain NMP.

Taylor, K. B. 1998. Species lists for Kings Mountain National Military Park.

Herpetofauna:

Fields, S. Probable herpetofauna of York County and surrounding counties. Museum of York County.

Taylor, K. B. 1998. Species lists for Kings Mountain National Military Park.

Thomas, R. B. 1999. Herpetofaunal inventory of the Kings Mountain National Military Park. University of South Carolina-Spartanburg. (ongoing) Funded through Small Park NRPP, FY2000.

Mammals:

Student Conservation Associate. 1999. Database of mammal species for Kings Mountain National Military Park.

Taylor, K. B. 1998. Species lists for Kings Mountain National Military Park.

Fish:

no information

LIRI

Plants:

Dickson, C. C. 1992. A vascular and non-vascular flora of Little River on Lookout Mountain in northeast Alabama. M. S. Thesis, Jacksonville State University, Jacksonville, AL. 180 p.

Whetstone, R. D., S. J. Threlkeld, C. C. Newsome, D. D. Spaulding, T. L. Ballard, J. W. Higginbotham, and J. M. Ballard. 1997. Vascular flora of Little River Canyon National Preserve, Cherokee and Dekalb Counties, Alabama. National Park Service, Technical Report. 173 p.

Birds:

no information

Herpetofauna:

Dunaway, M. J. 1995. A herpetofaunal survey of Little River Canyon, Alabama. M. S. Thesis, Jacksonville State University, Jacksonville, AL.

Mammals:

no information

Fish:

Ballard, T. L. and J. M. Pierson. 1996. The fishes of the Little River drainage in Alabama [DRAFT]. 20+ p.

Dobson, T. L. 1994. An ichthyofaunal survey of the Little River drainage in Alabama with notes on *Cyprinella caerulea*. Jacksonville State University, Jacksonville, AL.

General:

Alabama Natural Heritage Program. 1996. Vertebrates potentially occurring in Little River Canyon National Preserve, Cherokee and DeKalb Counties, Alabama. 10 p.

Belue, M. 2000. Spreadsheet of federally/state listed occurrence records for Little River Canyon National Preserve. (ongoing)

National Park Service. 1991. Special resource study: Little River Canyon area, Cherokee, Dekalb and Etowah Counties, Alabama. Division of Planning, Design, and Compliance, Southeast Region.

MACA

Plants:

Anonymous. 1949. Accession list of the plants in the museum herbarium, Mammoth Cave National Park, Mammoth Cave, Kentucky. 22 p.

Anonymous. 1954. Accession list of the plants in the museum herbarium, Mammoth Cave National Park, Mammoth Cave, Kentucky. 21 p.

- Anonymous. 1990. Wildflowers and ferns identified in Big Woods May 29, 1990. Mammoth Cave National Park, Kentucky. 6 p.
- Anonymous. 1992. Checklist of common flowers, Mammoth Cave National Park. 3 p.
- Badger, K. S., B. Jones, and J. Taylor. 1996. Vegetation inventory and classification of Mammoth Cave National Park. Ball State University.
- Bennett, J. P. Plants in Mammoth Cave National Park that may be susceptible to ozone damage. 2 p.
- Binneweis, F. W. 1938. Trees, vines, and shrubs found in Mammoth Cave National Park. 1 p.
- Braun, E. L. 1941. Notes on Kentucky plants. *Castanea* 6: 28-30.
- Call, R. E. 1897. Note on the flora of Mammoth Cave, Kentucky. *Journal of the Cincinnati Society of Natural History* 19(2): 79-80.
- Cissell, J. F. Common poisonous plants found in the Mammoth Cave area. 3 p.
- Cooke, C. E. Flower check list, Mammoth Cave National Park. 3 p.
- Cooke, C. E. Tree check list - Mammoth Cave National Park. 2 p.
- Davies, P. A. 1951. Ferns and fern allies that have been found or may possibly be found in the Mammoth Cave National Park, Kentucky. University of Louisville, Louisville, KY. 7 p.
- Davies, P. A. 1955. A preliminary list of the vascular plants of Mammoth Cave National Park. *Castanea* 20: 107-127.
- Deskins, R. L. 1984. Additions: trees and woody shrubs-Mammoth Cave National Park. 1 p.
- Ellsworth, I. J. Shrubs of the Mammoth Cave National Park area. Mammoth Cave, KY: Mammoth Cave National Park. 1 p.
- Ellsworth, I. J. Trees found in Mammoth Cave National Park. Mammoth Cave, KY: Mammoth Cave National Park. 1 p.
- Faller, A. 1975. The plant ecology of Mammoth Cave National Park, Kentucky. Ph.D. Dissertation, Indiana State University, Terre Haute, IN. 285 p.
- Hambly, F. N. 1966. Mammoth Cave National Park vegetative cover map [hand-colored]. Mammoth Cave, KY: Mammoth Cave National Park. 1 map; 34' x 21'; 1:31680.

- Hussey, J. 1876. Report on the botany of Barren and Edmonson counties, Kentucky. Kentucky Geological Survey, series II, pp. 27-58.
- Leeds, A. A. 1934. Ferns found in park area. The Academy of Natural Sciences, Philadelphia, PA. 3 p.
- Leonard, E. C. 1947. Plants of Mammoth Cave National Park. 1 p.
- Lix, H. W. and C. V. Norton. Ferns of Mammoth Cave National Park. Mammoth Cave, KY: Mammoth Cave National Park. 1 p.
- Maugans, J. Mammoth Cave National Park wildflower checklist additions. 1 p.
- McInteer, B. B. 1944. Revegetation of the abandoned fields of Mammoth Cave National Park. American Midland Naturalist 31: 87-94.
- McKinney, L. E., M. Evans, and K. A. Nicely. 1990. The flora of Mammoth Cave National Park. Proceedings of Mammoth Cave National Park's First Annual Science Conference: Karst hydrology; Mammoth Cave National Park, Mammoth Cave, KY, pp. 17-26.
- McKinney, L. E., M. Evans, and K. A. Nicely. 1991. The flora of Mammoth Cave National Park. 50 p.
- McKinney, L. E., M. Evans, and K. A. Nicely. 1990. Floristics and vegetational patterns of Mammoth Cave.
- Nelson, T. J. 1936. Wildflowers observed in bloom at Mammoth Cave National Park. Mammoth Cave, KY: Mammoth Cave National Park. 3 p.
- Olson, R., M. Franz, and G. Ghitter. 2000. A vegetation map of Mammoth Cave National Park using satellite remote sensing data. *In*: Proceedings of the 8th Mammoth Cave Science Conference. National Park Service (in press).
- Ray, W. E. A classification of the shrubs of Mammoth Cave National Park area. 2 p.
- Ray, W. E. 1933. A classification of the trees of Mammoth Cave National Park area. 4 p.
- Rissanen, W. 1953. Partial list of plants found in flower at Mammoth Cave National Park, Kentucky from June 29th to July 5th, 1953. Mammoth Cave, KY: Mammoth Cave National Park. 1 p.
- Seymour, R. 1996. A contribution to the recorded flora of Mammoth Cave National Park. Proceedings of the Fifth Annual Mammoth Cave National Park Science Conference; Mammoth Cave National Park. Bowling Green, KY, pp. 1-10.
- Seymour, W. R., Jr. A partial listing of the vascular flora in the area around Sloans-Crossing Pond in Mammoth Cave National Park. Mammoth Cave, KY: Mammoth Cave National Park. 7 p.

Seymour, W. R., Jr. 1994. A study and census of grasses, ferns, and herbaceous flowering plants in Mammoth Cave National Park.

Seymour, W. R., Jr. 1996. A vascular plant checklist for Mammoth Cave National Park. 1996.

Short, T. 1978. Common wildflowers of southern Kentucky. U. S. Army Corps of Engineers, Louisville, KY. 29 p.

Throne, A. L. 1956. Plant families represented in the herbarium of Mammoth Cave National Park. 5 p.

White, P. S. 1979. Nationally endangered and threatened plant species reported from Mammoth Cave National Park. 53 p.

Birds:

Anonymous. 1934. Bird census taken December 27, 1934 in (proposed) Mammoth Cave, National Park, Edmonson County, Kentucky. 272 p.

Anonymous. 1935. A complete list of birds and mammals acquired since June 1, 1934 in (proposed) Mammoth Cave National Park. 148 p.

Anonymous. 1992. Checklist of birds, Mammoth Cave National Park. National Park Service. 2 p.

Barbour, R. W., C. T. Peterson, D. Rust, H. E. Shadowen, and A. L. Whitt, Jr. 1973. Kentucky birds: a finding guide. University Press of Kentucky, Lexington, KY. 306 p. ISBN: 0813112818.

Burell, J. S. and A. L. Stamm. 1984. Kentucky's 1984 eagle census. Kentucky Warbler 60(3): 46-48.

Hibbard, C. W. 193?. Birds observed in the proposed Mammoth Cave National Park with notes. 7 p.

Palmer-Ball, Jr., B. L. 1996. The Kentucky breeding bird atlas. University Press of Kentucky, Lexington.

Shadowen, H. E. 1988. Midwinter bird counts at Mammoth Cave National Park. Kentucky Warbler 64(1): 20-21.

Wilson, G. 1941. A preliminary check-list of the birds of the Mammoth Cave National Park. Kentucky Warbler 17(2): 9.

Wilson, G. 1942. Summer birds in Mammoth Cave National Park. Kentucky Warbler 18(4): 2.

Wilson, G. 1950. Bird changes in Mammoth Cave National Park, 1938-1949. Kentucky

Warbler 16(2): 17-24.

Wilson, G. 1953. Birds of the Mammoth Cave National Park. Eastern National Park and Monument Association, Bowling Green, KY. 27 p.

Wilson, G. 1958. Bird changes in Mammoth Cave National Park, 1950-57. Kentucky Warbler 34(1): 3-6.

Wilson, G. 1958. Big spring lists. Kentucky Warbler 34(3): 39-42.

Wilson, G. 1958. Mid-winter bird count 1957-'58. Kentucky Warbler 34(1): 8-16.

Wilson, G. 1959. Big spring lists. Kentucky Warbler 35(3): 49-52.

Wilson, G. 1959. Notes on Mammoth Cave count. Kentucky Warbler 35(1): 5.

Wilson, G. 1960. Big spring lists. Kentucky Warbler 36(4): 57-61.

Wilson, G. 1960. Mid-winter bird count, 1959-1960. Kentucky Warbler 36(1): 9-16.

Wilson, G. 1961. 1960-'61 mid-winter bird count. Kentucky Warbler 37(1): 9-17.

Wilson, G. 1961. Additions to 'Birds of the Mammoth Cave National Park'. Kentucky Warbler 37(1): 7-9.

Wilson, G. 1961. Big spring lists, 1961. Kentucky Warbler 37(3): 51-54.

Wilson, G. 1962. Birds of south-central Kentucky. Kentucky Warbler 38: 3-24.

Wilson, G. 1968. Birds and their habitats in Mammoth Cave National Park. Eastern National Park and Monument Association, Louisville, KY. 39 p.

Wilson, R. C. 1985. Vertebrate remains in Kentucky caves. *In*: Dougherty, P. H. (ed.), Caves and karst of Kentucky. Publication 12, Series XI. Kentucky Geologic Survey in cooperation with the National Speleological Society, Lexington, KY, pp. 168-175.

Herpetofauna:

Anonymous. 1961. Amphibians and reptiles seen and/or collected in Mammoth Cave National Park, 24-27 March 1961. 71 p.

Anonymous. 1991. Checklist of reptiles and amphibians, Mammoth Cave National Park. Mammoth Cave National Park: National Park Service. 2 p.

Brandon, R. 1961. Amphibians and reptiles not reported from but perhaps occurring in Mammoth Cave National Park. 1 p.

Hale, M. F. 1982. Amphibians of Mammoth Cave National Park. M. S. Thesis, University of Louisville.

Hibbard, C. W. 1936. The amphibians and reptiles of Mammoth Cave National Park proposed. Transactions of the Kansas Academy of Sciences 39: 277-281.

Scott, A. F., J. W. Petranks, and C. K. Smith. 1994. Implementation of a plan for long-term monitoring of amphibians in selected parks of the southeast region. Cooperative Agreement # CA-5140-1-9001, sub-agreement # CA-5140-3-9002.

Williams, J. A. 1988. A comparative checklist of the reptiles and amphibians of Mammoth Cave National Park, Kentucky. Mammoth Cave, KY: Mammoth Cave National Park. 17 p.

Mammals:

Anonymous. Mammals of the Mammoth Cave National Park area. 1 p.

Anonymous. 1935. A complete list of birds and mammals acquired since June 1, 1934 in (proposed) Mammoth Cave National Park. 148 p.

Anonymous. 1966. Bats from Kentucky in USNM. 96 p.

Anonymous. 1987. Bat species found in Mammoth Cave National Park, Kentucky. 1 p.

Anonymous. 1992. Checklist of mammals, Mammoth Cave National Park. Mammoth Cave National Park: National Park Service. 2 p.

Davis, W. H. 1963. Operation Chiroptera. Bat Banding News 4(4): 30-31.

Dilley, W. E. 1961. List of park mammals. 3 p.

Hall, J. S. 1961. Species of bats known to occur in the Mammoth Cave National Park area. 3 p.

Harvey, M. J., J. R. MacGregor, and R. R. Currie. 1991. Distribution and status of Chiroptera in Kentucky and Tennessee. Journal of the Tennessee Academy of Science 66(4): 191-193.

Hibbard, C. W. 193?. Mammals observed or represented by skins from proposed Mammoth Cave National Park, Kentucky. 2 p.

Hibbard, C. W. 1935. New mammal records for Mammoth Cave National Park, Kentucky. Journal of Mammalogy 16(4): 329.

Hibbard, C. W. 1935. Special report on the bat situation at Mammoth Cave National Park with recommendations. 23 p.

MacGregor, J. Bats in the Warren County area. Kentucky Department of Fish and Wildlife Resources. 4 p.

Rebar, C. 1993. Small mammal trapping project at Onyx Meadows. Murray State University.

Thomas, S. 1997. Allegheny woodrat monitoring in Kentucky. (ongoing) Nongame Wildlife Program, Kentucky Department of Fish and Wildlife Resources.

Van Gelder, R. G. 1982. Mammals of the national parks. Johns Hopkins University Press, Baltimore, MD. 310 p. ISBN: 0801826896.

Fish:

Burr, B. M. and M. L. Warren, Jr. 1986. A distributional atlas of Kentucky fishes. Kentucky Nature Preserves Commission, Frankfort, KY. 398 p.

Charles, J. R. 1962. Biological and limnological study of streams. Kentucky Department of Fish and Wildlife Resources. State of Kentucky Project # F-18-R-3.

Cicerello, R. R. and R. R. Hannan. 1991. Survey and review of the fishes of Mammoth Cave National Park, Kentucky. Kentucky State Nature Preserves Commission, Philadelphia, PA. 42 p.

Kuehne, R. A. 1966. Depauperate fish faunas of sinking creeks near Mammoth Cave, Kentucky. *Copeia* 21(2): 306-311.

Laflin, B. and M. W. Vincent. 1983. Aquatic survey - Green and Nolin Rivers.

Laflin, B. D. 1988. Ichthyofauna studies of Mammoth Cave National Park during 1970 through 1987. 21 p.

Laflin, B. D. 1985. Ichthyofauna studies of Mammoth Cave National Park, during 1984. 19 p.

Laflin, B. D. 1987. Ichthyofauna studies of Mammoth Cave National Park, during 1986. 17 p.

Laflin, B. D. 1984. Preliminary ichthyofauna studies of Mammoth Cave National Park. 29 p.

Laflin, B. D. and M. W. Vincent. Ichthyofauna studies of Mammoth Cave National Park, 1970 through 1987.

Lewis, J. J. 1981. Aquatic communities in the historic section of Mammoth Cave. 10 p.

Pearson, W. and R. Jones. 1998. Faunal inventory of subterranean streams and development of a cave aquatic biomonitoring program. University of Louisville.

Retzer, M. E., B. M. Burr, and M. L. Warren, Jr. 1983. Fishes of the lower Green River drainage, Kentucky. Kentucky Nature Preserves Commission, Frankfort, KY. 48 p.

General:

- Anonymous. 1987. Animal species in Hart, Barren, and Edmonson Counties, Kentucky, listed by Kentucky's Department of Fish and Wildlife Resources as endangered or threatened. 1 p.
- Anonymous. 1988. Mammoth Cave National Park resource base inventory. 64 p.
- Anonymous. 1993. Endangered, threatened and special concern plants and animals, Mammoth Cave area, International Biosphere Reserve. Kentucky Nature Conservation's biological conservation database. 173 p.
- Bailey, V. 1933. Cave life of Kentucky, mainly in the Mammoth Cave region. The University Press, Notre Dame, IN. 256 p.
- Banks, N. R. E. 1897. Some notes on the fauna of Mammoth Cave, Kentucky. *American Naturalist* 31: 380-382.
- Barr, T. C. 1964. The fauna of Mammoth Cave: a study in community composition and evolution. *Proceedings of the National Speleological Society: Abstracts of papers presented at the Cleveland meeting, December 27, 1963; Cleveland, OH*, pp. 78-79.
- Barr, T. C. 1985. Cave life of Kentucky. *In: Dougherty, P. H. (ed.), Caves and karst of Kentucky. Publication 12, Series XI. Kentucky Geologic Survey in cooperation with the National Speleological Society, Lexington, KY*, pp. 146-167.
- Berry, K. 1969. Wildlife checklist of Mammoth Cave National Park. 12 p.
- Black, K. E. 1979. Presence of listed or proposed endangered or threatened species within the Green and Barren Rivers navigational study area. U. S. Fish and Wildlife Service, Atlanta, GA. 33 p.
- Conn, D. B. 1981. Dwellers of the caves: Kentucky's hidden wildlife. *Kentucky Happy Hunting Ground* 37(6): 24-27.
- Hester, F. E. 1988. Servicewide list of threatened and endangered species. National Park Service, Washington, D. C. 319 p.
- Hickling, W. C. 1981. A list of endangered and threatened species within the study area of the 201 wastewater facility plan of the Mammoth Cave area in Hart, Barren, Edmonson and Warren Counties, Kentucky. 123 p.
- House, S. R. 1995. Resources inventory of less-extensive caves: Mammoth Cave National Park, Kentucky. *Proceedings of Mammoth Cave National Park's Fourth Science Conference; Mammoth Cave National Park. Mammoth Cave, KY*, pp. 207-213.
- Hoyt, R. D. 1995. Biological specimen cataloging project: Mammoth Cave National Park, Kentucky. Western Kentucky University. 57 p.

- Ives, J. D. 1930. Cave animals of Tennessee. *Journal of the Tennessee Academy of Science* 5(3): 112-124.
- Komarek, E. V. and R. V. Komarek. 1934. A preliminary report on the faunal relations in the eastern national parks. 29 p.
- LaVoie, K. H., R. C. Wilson, and E. A. Lisowski. 1987. Mammoth Cave National Park cave fauna survey. University of Michigan, Flint.
- Lisowski, E. A., J. Oberlies, and R. Wilson. 1979. Biological survey of the Cascade Hall area, Mammoth Cave. Mammoth Cave National Park, Mammoth Cave, KY. 211 p.
- McSween, J. L. 1990. Organisms in the karst water of Mammoth Cave. 13 p.
- Packard, A. S., Jr. 1871. The Mammoth Cave and its inhabitants. *American Naturalist* 5: 739-761.
- Poulson, T. L. Biodiversity in the Mammoth Cave region. *In*: Sakowsky, I. (ed.), Symposium on biodiversity in karst; Nashville, TN. 7 p.
- Poulson, T. L. 1993. Cave animals of Mammoth Cave National Park (revised June 1993). 35 p.
- Poulson, T. L. and D. C. Culver. 1968. Diversity in aquatic and terrestrial cave communities of Flint Ridge, Mammoth Cave National Park. *Amer. Zool.* 7: 809.
- Smith, S. Q. 1976. Endangered and threatened wildlife and plants. 112 p.
- Springer, T. 1992. Listing of endangered species in Edmonson and Hart Counties effective October 1992. 157 p.
- Strange, R. N. 1982. Biological survey for east boundary road improvement. Mammoth Cave National Park, Mammoth Cave, KY. 14 p.
- Sutton, M. 1995. Cave fauna of the Mammoth Cave region: a pictorial field guide. 26 p.
- Sutton, M. R., R. S. House, and J. D. Borden. 1994. Resources inventory of less-extensive caves within Mammoth Cave National Park. Proceedings of Mammoth Cave National Park's Third Science Conference. Mammoth Cave National Park, Mammoth Cave, KY, pp. 117-124.
- Watkins, G. R. and J. S. Watkins. Great Onyx Cave survey, job # 1109. 22 p.

NISI

Plants:

Proposed plant inventory. Funded through Small Park NRPP. FY2000.

Birds:

National Park Service. 1994. Birds observed at Ninety-Six National Historic Site.

Wilson, R. S. 1996. Birds observed at Star Fort National Historic Site. Patterson and Carolina Bird Club.

Herpetofauna:

no information

Mammals:

no information

Fish:

no information

OBRIPlants:

DeSelm, H. R. Flora and vegetation of the barrens of the Cumberland Plateau of Tennessee. *In*: Snyder, D. H., (ed.), Proceedings of the contributed paper sessions of the fourth annual symposium on the natural history of lower Tennessee and Cumberland River valleys. Austin Peay State University, Clarksville, TN. 38 p.

Patrick, T. S., B. E. Wofford, and D. H. Webb. 1983. State records and other recent noteworthy collections of Tennessee plants IV. *Castanea* 48: 109-116.

Schmalzer, P. A. 1982. Vegetation of the Obed Wild and Scenic River, Tennessee and a comparison of reciprocal averaging, ordination, and binary discriminant analysis. University of Tennessee, Knoxville. 235 p.

Schmalzer, P. A. 1988. Vegetation of the Obed River gorge system, Cumberland Plateau, Tennessee. *Castanea* 53(1): 1-32.

Schmalzer, P. A. 1989. Vegetation and flora of the Obed River gorge system, Cumberland Plateau, Tennessee. *Journal of the Tennessee Academy of Science* 64(3): 161-168.

Schmalzer, P. A. and H. R. DeSelm. 1982. Vegetation, endangered and threatened plants, critical plant habitats and vascular flora of the Obed Wild and Scenic River. Final Report. 168 p.

Schmalzer, P. A., T. S. Patrick, and H. R. DeSelm. 1985. Vascular flora of the Obed Wild and Scenic River, Tennessee. *Castanea* 50: 71-88.

Wofford, B. E., T. S. Patrick, L. R. Phillippe, and D. H. Webb. 1979. The vascular flora of Savage Gulf, Tennessee. *SIDA* 8(2): 135-151.

Birds:

Stedman, B. 1998. Check-list of Putnam County, TN. (ongoing).

Taylor, C. A., R. S. McKittrick, and M. R. Pelton. Terrestrial vertebrate inventory of the Obed Wild and Scenic River, Tennessee. National Park Service, contract # CX500091150. 75 p.

Herpetofauna:

Gentry, G. 1941. Herpetology collections from counties in the vicinity of the Obed River drainage in Tennessee. *Journal of the Tennessee Academy of Science* 27(1): 153-158.

Taylor, C. A., R. S. McKittrick, and M. R. Pelton. Terrestrial vertebrate inventory of the Obed Wild and Scenic River, Tennessee. National Park Service, contract # CX500091150. 75 p.

Mammals:

Taylor, C. A., R. S. McKittrick, and M. R. Pelton. Terrestrial vertebrate inventory of the Obed Wild and Scenic River, Tennessee. National Park Service, contract # CX500091150. 75 p.

Fish:

Taylor, C. A., R. S. McKittrick, and M. R. Pelton. Terrestrial vertebrate inventory of the Obed Wild and Scenic River, Tennessee. National Park Service, contract # CX500091150. 75 p.

Tennessee Valley Authority, Division of Forestry, Fisheries, and Wildlife Development. 1968. Tennessee valley streams: their fish, bottom fauna, and aquatic habitat - Emory River. Tennessee Valley Authority, Norris, TN. 20+ p.

RUCA

Plants:

Barnett-Lawrence, M. 1993. Checklist of the vascular vegetation of Kings Mountain, Kennesaw Mountain, Chickamauga and Chattanooga, Shiloh, and Russell Cave. *In*: Mapping vegetation communities with the Nature Conservancy's vegetation classification system on five small national parks in the southeastern U.S.A. Chapel Hill. 100+ p.

DiPetro, T. L. 1995. Vascular plant inventory and occurrence records for rare plants at Russel Cave National Monument.

Gallyoun, M., G. Meyer, A. Andreu, and W. Slocumb. 1995. Mapping vegetation communities with the Nature Conservancy's vegetation classification system on five small national parks in the southeastern U.S.A. Chapel Hill, NC. 100+ p.

Birds:

Keeler, A. V. 1994. Bird checklist {7/11/94—7/17/94}, Russell Cave National Monument-Alabama. 1 p.

Herpetofauna:

no information

Mammals:

no information

Fish:

no information

General:

Hobbs, H. H. 1994. Assessment of the ecological resources of the caves of Russell Cave National Monument, Jackson County, Alabama and of selected caves at the Lookout Mountain unit of Chickamauga-Chattanooga National Military Park, Dade County, Georgia and Hamilton County, Tennessee. Final report. 184 p.

SHILPlants:

Butler, T. and P. S. White. 1981. Exotic woody plants of Shiloh National Military Park, Tennessee: a population survey of aggressive species. National Park Service, Research/Resources Management Report # 51. 23 p.

Gallyoun, M., G. Meyer, A. Andreu,, and W. Slocumb. 1995. Mapping vegetation communities with the Nature Conservancy's vegetation classification system on five small national parks in the southeastern U.S.A. Chapel Hill. 100+ p.

Jones, R. L. 1983. Woody flora of Shiloh National Military Park. Castanea 48: 289-299.

Jones, R. L. and P. S. White. 1981. The vascular flora of Shiloh National Military Park, Hardin County, Tennessee. National Park Service, Research/Resources Management Report # 50. 49 p.

National Park Service. 1984. Flora list for Shiloh National Military Park. 13 p.

Birds:

Coffey, B., Jr. and J. W. Howell. 1954. A list of birds. National Park Service. 10 p.

Tennessee Ornithological Society. 1998. Birds of Shiloh National Military Park.

Herpetofauna:

Pritts, G. R. 1999. An investigation and assessment of amphibian and reptile fauna of Shiloh National Military Park, Hardin County, Tennessee.

Mammals:

Kennedy, M. L. 1984. Mammals of Shiloh National Military Park. 300+ p.

Fish:

Higgins, L. K. 1998. Ichthyological investigations in Shiloh National Military Park, Hardin County, Tennessee.

General:

The Nature Conservancy. Species list for Shiloh National Military Park. 18 p.

STRIPlants:

Hogan, T., R. Sutter, and N. Rudd. 1996. Vascular plant inventory, baseline and photo point monitoring, and rare species monitoring of the calcareous glades of Stones River National Battlefield.

Hogan, T. L. and M. Webber. 1999. Vascular flora of Stones River National Battlefield, including notes on natural communities and rare species.

Birds:

Walker, K. 197?. A study of the relative abundance of summer birds observed at Stones River National Battlefield. 10 p.

Herpetofauna:

no information

Mammals:

no information

Fish:

Mullen, D. 1995. Lytle Creek ichthyofauna - Fortress Rosecrans qualitative survey/electrofishing. Middle Tennessee State University. (ongoing)

Appendix B: Status of data deliverables for the Appalachian Highlands and Cumberland/Piedmont Networks.

Data documents were gathered during the 17 park visits from March – June 2000. This information includes published reports, birding brochures, and park-level databases. The entry of this information continues at Mammoth Cave National Park (MACA), through a cooperative agreement with Western Kentucky University. Oversight and workload associated with data deliverables will be continued through the following: the existing agreement (Oct - March 2001), the MACA data manager, and the new network data managers which will be hired during FY01. An overview of work is listed in Table 1, a summary of progress with NPSpecies is listed in Table 2, and a CD containing preliminary GIS files and NPSpecies data (for our progress-to-date) was sent to the I&M Program Manager on 7 October, 2000.

Some data management protocols established by our networks:

1) References:

All new references are entered into NRBib and NPSpecies, using the park 4 letter acronym followed by: XXX1, XXX2, XXX3, etc., in order to differentiate what we have entered. For multi-park references we used: MULT-XXX1, MULT-XXX2, etc.

For example: Barbara Stedman's "*Checklist of breeding birds located at Obed Wild and Scenic River*", was entered into NRBib and NPSpecies with the code: OBRI-XXX2.

For multi-park references: Kristine Johnson's "*Invasive plant management in smaller parks of the Appalachian Cluster*", was entered into NRBib and NPSpecies with the code: MULT-XXX1.

2) Habitat:

For park-level habitat information (where a species was found in the park), data was entered into the "Park Status-Details" field. Several lists have broad descriptions like: ABLI's plants: "open woods", "disturbed area", "old field".

For example: *Park Status: present in park Details: open woods*

3) Seasonal abundance:

Almost all bird lists have abundance by season. We used the "Abundance-Details" field to capture this information.

For example: ABLI's list had species name, spr, sum, fall, win, habitat.

Abundance: Common Details: spr/sum/fall (means common in spr/sum/fall)

Abundance: Abundant Details: spr/sum; common fall (means abundant in spr/sum and common in fall)

4) Residency:

Several lists had season associated with residency. For example, “winter resident”, “summer resident”, “permanent”. We used the “Residency-Details” field to capture season.

For example: ABLI’s Northern harrier:

Residency: Resident

Details: Winter

5) Data source:

All of our detailed information has come from an attached reference. Since it is unclear how references will eventually be numbered, we are using the following format in the “Data Source” field: *author, date*.

6) Entered by:

We added our initials to any existing records, in case there is some attempt to track the WASO entered records. The format is: 3 lower case letters followed by the park acronym in upper case. For example: *trl-MACA*. Please advise on whether to overwrite existing initials.

7) BONAP:

In August we received an updated BONAP file called Bonap-County. After all valid plants has been linked to a reference in the initial database, the remainder are deleted (plants with no reference leftover from initial BONAP file). We then import from the updated BONAP file, for “New Records Only”, and mark those as unconfirmed. This allows us to merge what we have worked on with the new referenced BONAP records.

8) NPS Theme Manager:

The new NPS Theme Manager is nearing distribution and will provide a tool for networks to incorporate new I&M GIS layers. We plan to make use of this format, instead of the Arcview Project file and DRAFT Mapping Products mentioned in our pre-proposal deliverables section.

9) Data Set Catalog:

The new NPS-Dataset Catalog is nearing distribution. It will provide a tool for networks to track electronic data (databases, spreadsheets, GIS layers, etc) and will provide metadata. We plan to make use of this tool and will evaluate its effectiveness for compliance with the FGDC standard (as compared to other software available, such as SMMS).

10) Vouchers:

Data entry into the voucher section of NPSpecies has not begun. We have noted known park and offsite collections and plan to begin work soon. We would like more guidance on what WASO is planning to cover for offsite collections. Does verification involve visual inspection or are written lists of specimens good enough? Also, for park collections, is there a master ANCS database that can be used to download all species records for all parks? This would save a lot of time. For Kentucky plant collections, we have a contact in the Kentucky Native Plant Society who has developed a database of herbarium collections across the state, and will supply us a copy next month.

11) Sensitive Species Flag:

As of yet, no records have been flagged as sensitive. We would like some direction on how this field is used, when they should be flagged, and by whom. Does this flag only pertain to the reference, species, or both? Will it be used to filter records on the internet?

12) Threatened and Endangered Status:

Some species such as ABLI's *Empidonax traillii*, the willow flycatcher, are showing as having federal status. However, it is the southwestern willow flycatcher that is listed, not this one. How do we change this, or do we just report the need to correct it?

Table 1. Status of data/GIS products (as of 28 September, 2000) for inventory study plan.

Park	NRBib (inventory related records)	Cataloging datasets / metadata	NPSpecies	GIS layers	Aerial photos
ABLI	3 Records (add exotic plant study)	Waiting on new dataset catalog software, some metadata done	Link plants and vouchers (206), add birds (126)	Habitat from plant report (BW), hydrology study, poster layout on CD	On order
BISO	23 Records (under review)	Waiting on new dataset catalog software	Link plants (584 in xls) and vouchers	Poster layout on CD	On order
BLRI	57 Records (under review)	Waiting on new dataset catalog software	Updated lists for all groups received 9/12	Poster layout on CD	On order
CARL	9 Records (add exotics: Remaley '98 and Blaha '99, plants)	Waiting on new dataset catalog software	Link plants and vouchers (354), add birds (43)	Poster layout on CD	On order
CHCH	12 Records (add Hobbs, GaNHP plants)	Waiting on new dataset catalog software, some metadata done	Link plants and vouchers (585), add birds	Veg-map, species of concern, habitat / glade, poster layout on CD	On order
COWP	7 Records (add current proposals for vertebrate / plant inventories)	Waiting on new dataset catalog software	Link plants (343), add birds (200)	Poster layout on CD	On order
CUGA	13 Records (add Petranka, amphibians; Currie, bats)	Waiting on new dataset catalog software	Link plants (783)	Poster layout on CD	On order
FODO	2 Records (add Chester, new plant book; 2001 plant study; bird list)	Waiting on new dataset catalog software	Link plants (645), add vouchers, add birds (251)	Poster layout on CD	On order
GRSM	48 Records (under review)	Waiting on new dataset catalog software	All groups need links: (2000+ plants, 250 birds, 63 amphibians, 60 reptiles, 77 mammals, 89 fish)	Poster layout on CD	

GUCO	5 Records (added '38, '76 trees and shrubs, '96 exotics, '83 vertebrates)	Waiting on new dataset catalog software	Link plants	Poster layout on CD	On order
KIMO	11 Records (add SCA database, two herpetofaunal studies)	Waiting on new dataset catalog software, some metadata done	Link plants (586) and vouchers	Veg-map, species of concern, poster layout on CD	On order
LIRI	8 Records (add spreadsheet)	Waiting on new dataset catalog software	Link plants and vouchers	Poster layout on CD	On order
MACA	128 Records (add Bird Atlas)	Yes, 120 records in old format. Waiting on new dataset catalog software. Some metadata done	Link plants (1164) and vouchers	Habitat map, Veg-map, species of concern, poster layout on CD	On order
NISI	3 Records (add 2001 plant proposal; two bird references; Ogelthorpe Oak study?)	Waiting on new dataset catalog software	Add birds (118)	Poster layout on CD	On order
OBRI	15 Records (add birds; plants; exotics)	Waiting on new dataset catalog software	Link plants (755) and vouchers	Poster layout on CD	On order
RUCA	5 Records (add Dipetro, plants)	Waiting on new dataset catalog software	Link plants and vouchers (460), add birds (28)	Veg-map, species of concern, poster layout on CD	On order
SHIL	11 Records (update Higgins '98, fish; add Higgins '88, mussels)	Waiting on new dataset catalog software. Some metadata done	Link plants (717), add birds (204), link vouchers	Veg-map, species of concern, poster layout on CD	On order
STRI	4 Records (add Hogan/Webber, plants '99; Mullen, fish)	Waiting on new dataset catalog software	Link plants and vouchers (195)	Poster layout on CD	On order

Table 2. NPSpecies progress as of 28 September, 2000

Park								
ABLI	206 linked to references with habitat. Updated with BONAP	126 added with habitat/seasonal abundance	No list	No list	No list	No list	Need plant search WKU, EKU through IK	3
BISO	Need to link plants (584 in xls)	262 to check	33 to check	42 to check	61 to check	94 to check	BISO Herbarium	23
BLRI	Need to link plants (1,096 in xls)	315 to enter	35 to enter	33 to enter	64 to enter	54 to enter	At several institutions	40
CARL	354 linked to references. Updated with BONAP	Added 64 with habitat/seasonal abundance	No list	5 to check	10 to check	7 to check	CARL Herbarium	14
CHCH	Need to link plants (585 in xls)	Need to add bird list on the way 9/13/00	32 to check	39 to check	45 to check	4 to check	Plant search UTChat	29
COWP	343 linked to references. Updated with BONAP	75 added with habitat/seasonal abundance	No list	No list	No list	No list	Plants at Univ. So. Miss. (?)	10
CUGA	Need to link plants (783 in xls)	94 to check	26 to check	18 to check	38 to check	34 to check	Plants at UT, not accessioned	28
FODO	645 linked to references. Updated with BONAP	251 birds added (Cross Creek). Christmas count received 9/13/00	No list	No list	No list	No list	Austin Peay Herbarium, FY00 contract for plant collection	22
GRSM	2,221 plants, no references	250 birds, no references	63, no references	60, no ref's	77, no references	89, no ref's	GRSM curation	41
GUCO	No list	48 to check	16 to check	5 to check	10 to check	No list	None known	12

KIMO	Need to link plants (586)	191 to check	No list	35 to check	39 to check	No list	KIMO Herbarium	13
LIRI	Need to link plants	238 to check	42 to check	61 to check	50 to check	74 to check	JSU Herbarium	7
MACA	Need to link plants (1164)	160 to check	45 to check	42 to check	47 to check	88 to check	MACA Herbarium and several offsite	25
NISI	No list	118 added	No list	No list	No list	No list	FY00 contract for plant collection	7
OBRI	Linked plants (755)	92 updated with habitat/seasonal abundance	36 to check	37 to check	47 to check	55 to check	BISO Herbarium, UT Fish, Tenn. Tech Fish	24
RUCA	Need to link plants (460)	14 to check, 28 to enter	8 to check	14 to check	26 to check	10 to check	RUCA Herbarium	11
SHIL	Linked plants (717)	204 added with seasonal abundance	26 to check	23 to check	56 to check	49 to check	SHIL Herbarium, SHIL Fish	17
STRI	Linked plants (195)	71 to check	No list	7 to check	10 to check	No list	STRI Herbarium, UT Herbarium	20

Appendix C: Latin names of common species presented in the text.

alder – *Alnus serrulata*
 ash, blue – *Fraxinus quadrangulata*
 ash, white – *Fraxinus americana*
 basswood, American – *Tilia americana* var. *americana*
 basswood, white – *Tilia americana* var. *heterophylla*
 beech, American – *Fagus grandifolia*
 birch, river – *Betula nigra*
 birch, yellow – *Betula alleghaniensis*
 bittersweet, Oriental – *Celastrus orbiculata*
 black gum – *Nyssa sylvatica*
 boxelder – *Acer negundo*
 buckeye, red – *Aesculus pavia*
 buckeye, yellow – *Aesculus flava*
 buckthorn, Carolina– *Rhamnus caroliniana*
 cedar, eastern red – *Juniperus virginiana*
 cherry, black – *Prunus serotina*
 dogwood, flowering – *Cornus florida*
 fescue, tall – *Festuca pratensis*
 fir, Fraser – *Abies fraseri*
 fringe tree – *Chionanthus virginicus*
 hackberry – *Celtis occidentalis*
 hemlock, eastern – *Tsuga canadensis*
 hickory – *Carya* sp.
 hickory, pignut – *Carya glabra*
 holly, Georgia – *Ilex longipes*
 honeysuckle, Amur – *Lonicera maackii*
 honeysuckle, Japanese – *Lonicera japonica*
 locust, black – *Robinia pseudoacacia*
 magnolia, umbrella – *Magnolia tripetala*

maple, red – *Acer rubrum*
maple, sugar – *Acer saccharum*
mountain laurel – *Kalmia latifolia*
oak, black – *Quercus velutina*
oak, blackjack – *Quercus marilandica*
oak, chestnut – *Quercus prinus*
oak, chinquapin – *Quercus muehlenbergii*
oak, northern red – *Quercus rubra*
oak, post – *Quercus stellata*
oak, scarlet – *Quercus coccinea*
oak, southern red – *Quercus falcata*
oak, white – *Quercus alba*
pine, loblolly – *Pinus taeda*
pine, pitch – *Pinus rigida*
pine, shortleaf – *Pinus echinata*
pine, slash – *Pinus elliottii*
pine, Table Mountain – *Pinus pungens*
pine, Virginia or scrub – *Pinus virginiana*
pine, white – *Pinus strobus*
rhododendron – *Rhododendron* sp.
rose, multiflora – *Rosa multiflora*
silverbell – *Halesia tetraptera*
sparkleberry – *Vaccinium arboreum*
spruce, red – *Picea rubens*
sweet gum – *Liquidambar styraciflua*
sycamore, American – *Platanus occidentalis*
tulip poplar – *Liriodendron tulipifera*
walnut, black – *Juglans nigra*
willow, black – *Salix nigra*

Appendix D: Southeast Regional Director's memo.

United States Department of the Interior

NATIONAL PARK SERVICE
Southeast Regional Office
Atlanta Federal Center
1924 Building
100 Alabama St., S. W.
Atlanta, Georgia 30303

IN REPLY REFER TO:
N16(SER-D)

December 8, 2000

Memorandum

To: Superintendents, Southeast Region

From: Regional Director, Southeast Region

Subject: Southeast Region Monitoring Program: Vision and Implementation

The attached document outlines the major Roles and Responsibilities for the Southeast Regional Inventory and Monitoring (I&M) Coordinator. Also included as reference material is the WASO memorandum released October 13, 2000, which served to implement the National I&M Vital Signs vision program.

Further, a list of I&M regional networks provided by WASO on the initiation of a Nation-wide Vital Signs Monitoring program of NPS is included. The term "Vital Signs" has now been approved for use by the Service in the context of the Vital Signs long term monitoring program of the Natural Resource Challenge.

A couple of clarifications with respect to the October 13, 2000, memorandum follow. The staffing indicated (eight to nine people per network) may not be possible in all networks with monitoring funds alone. It is expected that additional resources and/or funds will need to be made available by parks in each network in order to fully implement various I&M activities.

Administratively, the Southeast Region will manage the Monitoring program via line supervision of Network Coordinators through the Regional I&M Coordinator in the Division of Science and Natural Resource Management. Other Regional Offices have similar line authority supervision for Network Coordinators (see attached Roles and Responsibilities of the Regional I&M Coordinator for further detail).

This new thrust will provide funding for the National Park Service to monitor vital signs on our five Southeast Region networks of parks on a specific schedule (see below and the attachment to the WASO memorandum).

Southeast Region Schedule

The Cumberland/Piedmont Network will be the first network funded for Vital Signs Monitoring and is scheduled to receive \$415,541 when the budget is finalized this Fiscal Year. The Appalachian Highlands Network will receive partial funding of \$150,000 also in FY 2001. Both of these networks will hire two key personnel to assist them with the necessary preparation work of information and data mining, and scoping meetings.

In FY 02 the Cumberland/Piedmont Network will receive full funding of \$565,541 and the Appalachian Highlands Network will receive additional partial funding totaling \$525,657. The Gulf Coast Network will receive partial funding of \$150,000 for starting up their Monitoring program.

The Appalachian Highlands Network will receive full monitoring funds in FY 03 and thereafter will annually receive \$675,657. The Gulf Network will get additional partial funding in FY 03 of \$702,881. Both the South Florida/Caribbean Network and Southeast Coast Network are scheduled to receive \$150,000 for start-up that same year. These three networks, Gulf Coast, South Florida/ Caribbean and Southeast Coast, are all scheduled to receive full funding in FY 04 and annually thereafter of \$852,881, \$1,398,877, \$1,163,014, respectively.

Continuation of Vital Signs Monitoring via the Natural Resource Challenge is subject to future Congressional support. Satisfactory performance in these early years will have a strong bearing on completing the above funding as scheduled. I trust each Superintendent and staff will give full support and cooperation to Vital Signs Monitoring, not only because of Service goals and the promise of new funds, but because of a renewed desire in all of us to be the best stewards of our natural resources, as we are of our cultural and historic resources.

/s/ Patricia A. Hooks (Acting)
(signed copy on file)

Attachments

Appendix E: I&M Coordinator Roles and Responsibilities (attachment to Appendix D).

Southeast Region Roles and Responsibilities of the Regional Inventory and Monitoring Coordinator

Background

For the first time, there are now seven Regional Inventory and Monitoring (I&M) Coordinators dedicated to the Servicewide NPS I&M Program. This group has a great deal of education and experience directly applicable to Inventory & Monitoring.

Congress and the Executive Branch have placed a great deal of importance on this program as expressed via:

NATIONAL PARKS OMNIBUS MANAGEMENT ACT OF 1998

“The Secretary shall undertake a program of inventory and monitoring of National Park System resources to establish baseline information and to provide information on the long-term trends in the condition of National Park System resources. The monitoring program shall be developed in cooperation with other Federal monitoring and information collection efforts to ensure a cost-effective approach.”

“The trend in the condition of resources of the National Park System shall be a significant factor in the annual performance evaluation of each superintendent of a unit of the National Park System.”

FY00 Congressional Appropriations Language:

*“The Committee applauds the Service for recognizing that the preservation of the diverse natural elements and the great scenic beauty of America's national parks and other units **should be as high a priority in the Service as providing visitor services.** A major part of protecting those resources is knowing what they are, where they are, how they interact with their environment and what condition they are in. **This involves a serious commitment from the leadership of the National Park Service to insist that the superintendents carry out a systematic, consistent, professional inventory and monitoring program, along with other scientific activities, that is regularly updated to ensure that the Service makes sound resource decisions based on sound scientific data**”.*

FY2000 Appropriations Bill

*“While the Service downsizing several years ago shifted greater responsibility and authority to the parks, this decentralization did not obviate the need for the Service to operate as one agency, rather than 378 independent parks, 7 autonomous regions, and a Washington office. While the Committee is not advocating a centralization of all Service decision-making, **a concerted effort***

must be made at all levels of Park Service management, including park superintendents and regional directors, to exercise greater responsibility in implementing programs with an eye towards servicewide goals, and not individual whims.”

Each Coordinator and their phone numbers are listed below.

Regional I&M Coordinators

Dr. Steve Acker – Pacific West (206-220-4267)
 Dr. Phyllis Adams – Midwest (402-221-7294)
 Mike Britten – Intermountain (303-987-6705)
 Dr. Ellen Gray – National Capital (202-342-1443, ext. 223)
 Elizabeth Johnson – Northeast (401-874-7060)
 Sara Wesser – Alaska (907-257-2557)
 Larry West – Southeast (404-562-3113, ext. 526)

Because of the success to date of the Natural Resource Challenge (NRC) funding initiative; park, network, regional, and Servicewide I&M funding and programs and projects are rapidly increasing. In FY 2001 funding for I&M in the Southeast Region will increase to over \$ 1,600,000 and that figure will continue to increase each year until it peaks in 2004 at approximately \$4.75 million dollars per year when all 5 networks are fully funded. The NRC is forcing rapid I&M developments due to the pressing need to demonstrate progress and products quickly to ensure continued congressional support and success.

Some regional networks have experienced difficulty developing their organization and commitment to I&M projects and programs. Much of this is attributable to the lack of professional staff time and the lapse in hiring Regional I&M Coordinators.

The I&M workload at the network, regional and Servicewide levels is considerable. The major roles of the SE Regional Coordinator are outlined below:

- ❑ **Regional I&M Coordinator Roles and Functions (major)**
- ❑ Transfer information and guidance from Servicewide I&M Program and other WASO Divisions to parks and networks in their Region
- ❑ Manage regional I&M budget and relay funds to approved network plan activities in their region
- ❑ Ensure that I&M funds are spent appropriately by parks and networks in their region (e.g. the Coordinators review projects and programs, similar to the FirePro program)
- ❑ Assist the I&M Steering Committee and Servicewide I&M Program to develop NPS I&M
- ❑ Assist in developing technical guidance (e.g. contribute to Vital Signs Monitoring Handbook)

- ❑ Assist in providing outstanding examples of I&M approaches from their region (for use by other networks, regions and the Servicewide I&M Program)
- ❑ Assist in developing effective processes for implementing projects and programs (e.g. study plan development, review and approval)
- ❑ Develop regional I&M Programs
- ❑ Serve on Vital Signs Network “Boards of Directors,” called for in the draft Vital Signs Vision and Implementation Plan
- ❑ Promote I&M at the Regional Director and Superintendent Level
- ❑ Supervise Network I&M Coordinators (described below)

The roles and functions of this new position need to be incorporated into National Park Service and the Southeast Region. Each Region has been given the flexibility to manage this program with WASO FTE's and funding, with the caveat of reaching GPRA/PMDS goals.

The Regional Director has, therefore, decided that line supervision of Network Coordinators will be accomplished through the Regional Division of Science and Natural Resource Management by the I&M Coordinator. This management of the SE I&M program follows the successful precedent already established in other Regions. Line Supervision that follows the funding lines in starting a new program that touches so many different parks in the region and that has specific PMDS goals for the region has distinct advantages. Some of these are:

- ❑ Opening lines of communication, fostering professionalism and outside contacts
- ❑ Assuring information transfer across parks, networks and to outside scientists and the public.
- ❑ Facilitating budget and expenditure actions
- ❑ Continuity of information for data management and reporting to WASO and Congress
- ❑ Assuring planning and coordination for scoping and other meetings, for annual plans, Cooperative Agreements, personnel actions and budgets takes place in a timely manner
- ❑ Providing benefits to networks from the accomplishments, work, progress and ideas in other networks
- ❑ Assuring responsiveness to track and meet National Park Service goals
- ❑ Continuity of techniques for comparable results and accomplishments
- ❑ Scientific credibility of planned projects and results obtained
- ❑ Financial credibility of the program for public and Congressional oversight
- ❑ Creation of an atmosphere of mutually beneficial cooperation among the parks of the networks, between the networks and with the Regional Office.

Network I&M Coordinators will be recruited by each Network with selections determined by a panel consisting of the Regional I&M Coordinator, appropriate staff members of the network which may include members from the Board of Directors in concert with the needs identified in network plans and in consultation with WASO.

Annual performance and evaluation of the overall network accomplishments will be the responsibility of the Network I&M Coordinator and the Board of Directors for the network. The Regional I&M Coordinator will be responsible for tracking performance accomplishments of all networks for the region.