



Inventory of the Mosses, Liverworts, and Lichens of Kenai Fjords National Park, Alaska

2013 Progress Report

Natural Resource Data Series Report NPS/SWAN/NRDS—2014/635



ON THE COVER

Left: Cliff survey in Coleman Bay, Kenai Fjords NP. Photo: James Walton.

Right Top: *Diphyscium foliosum*, Coleman Bay. Photo: Martin Hutten.

Right Center: *Takakia lepidozoides*, near Aligo Point. Photo: Martin Hutten.

Right Bottom: *Siphula ceratites*, near Aligo Point. Photo: Martin Hutten.

Inventory of the Mosses, Liverworts, and Lichens of Kenai Fjords National Park, Alaska

2013 Progress Report

Natural Resource Data Series Report NPS/SWAN/NRDS—2014/635

James K. Walton¹, Martin Hutten², Stacey K. Torigoe³

¹National Park Service
Southwest Alaska Network Inventory and Monitoring Program
240 West 5th Avenue
Anchorage, Alaska 99501 USA

²National Park Service
Lassen Volcanic National Park
Mineral, California 96063 USA

³National Park Service
Haleakalā National Park
Makawao, Hawaii 96768

March 2014

U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science
Fort Collins, Colorado

The National Park Service, Natural Resource Stewardship and Science office in Fort Collins, Colorado, publishes a range of reports that address natural resource topics. These reports are of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Data Series is intended for the timely release of basic data sets and data summaries. Care has been taken to assure accuracy of raw data values, but a thorough analysis and interpretation of the data has not been completed. Consequently, the initial analyses of data in this report are provisional and subject to change.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner.

This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data. Data in this report were collected and analyzed using methods based on established, peer-reviewed protocols and were analyzed and interpreted within the guidelines of the protocols.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available from the Southwest Alaska Network (<http://science.nature.nps.gov/im/units/swan/>) and the Natural Resource Publications Management Web site (<http://www.nature.nps.gov/publications/nrpm/>) on the Internet. To receive this report in a format optimized for screen readers, please email irma@nps.gov.

Please cite this publication as:

Walton, J. K., M. Hutten, and S. K. Torigoe. 2014. Inventory of the mosses, liverworts, and lichens of Kenai Fjords National Park, Alaska: 2013 Progress Report. Natural Resource Data Series Report NPS/SWAN/NRDS—2014/635. National Park Service, Fort Collins, Colorado.

Contents

	Page
Figures	iv
Tables	iv
Appendices	iv
Abstract	v
Acknowledgements	v
Introduction	1
Methods and Results	3
Off-frame FIA epiphytic macrolichen surveys	3
Intuitively-controlled bryophyte and lichen surveys	3
Notable Discoveries and Future Work	6
Notable discoveries	6
Future work	6
Literature Cited	7

Figures

	Page
Figure 1. Bryophyte and lichen survey site locations during the 2012-2013 field seasons, Kenai Fjords NP.	2
Figure 2 (top right). Documenting epiphytic macrolichen occurrence on a Sitka spruce tree at a off-frame FIA site near Nuka Bay. Photo: Michael Shephard/NPS.	3
Figure 3 (bottom right). Low elevation Sitka spruce - mountain hemlock forests, such as this fog-drenched stand in Square Bay near McArthur Pass, were the target vegetation type for our FIA surveys. Photo: James Walton/NPS	3
Figure 4 (top). Certain rare to uncommon species, like the moss <i>Takakia lepidozioides</i> , were only found in sloping peatlands such as this one near Fire Cove. Photo: James Walton/NPS.	4
Figure 5 (center right). Bryophyte and lichen diversity was found to be high at the shoreline - forest ecotone, which required careful and patient study of substrates, such as this exposed bedrock just above the hightide line. Photo: James Walton/NPS.	4
Figure 6 (bottom right). At the end of each field day bryophyte and lichen voucher collections were organized and dried on the M/V Serac. Photo: James Walton/NPS.	4
Figure 7. <i>Kurzia trichoclados</i> , a small and pleasantly sweet aromatic liverwort that can be "sniffed out" in moist places such as shady boulder crevices or over <i>Sphagnum</i> moss. Discoveries at Crater Bay, Three Hole Bay, and Bear Cove are apparently the first for Alaska. Photo: James Walton/NPS.	5

Tables

	Page
Table 1. Ecological Systems (Boggs et al. 2008) expected to increase known bryophyte and lichen diversity in Kenai Fjords NP.	6

Appendices

	Page
Appendix A: Survey Sites.	9
Appendix B: Species List.	10
Appendix C: Potential Species List.	19
Appendix D. Voucher Collections	25

Abstract

In the summer of 2012, the Southwest Alaska Network initiated a limited bryophyte and lichen inventory in Kenai Fjords National Park that was part systematic, part opportunistic. Our focus was on low elevation Sitka spruce - mountain hemlock forest that characterizes much of the Park's coastline. We dedicated about 15 person days toward the survey of 22 sites, most of which were forested but also included select sloping peatlands, coastal cliffs, active colluvial slopes, and an alpine site. Across all sites, 297 species were documented, including 126 species of moss, 79 liverworts, and 92 lichens. Virtually all these species are new to the park as this work represents

the first systematic inventory of bryophytes and lichens conducted in the Park. While our initial focus was on a forest type that is common in Alaska, we documented a high number of unusual species, including at least three species new to Alaska and several that are considered regionally rare. While currently limited in its scope, this survey work provides critical floristic baseline information to the Park. Additionally, since the area represents a floristic region that is poorly known, further work is likely to make a significant scientific contribution to our knowledge of the Alaskan flora.

Acknowledgements

NPS employees Michael Shephard and Deb Kurtz for help with fieldwork; Christina Kriedemann for maps and guidance on land status; Amy Miller and Michael Shepherd for informative discussions on site selection as

well as helpful review of this manuscript; and Jamie Thomson, Melissa Knight, Luke Robert and Mark Kansteiner for providing safe passage to survey sites from the M/V Serac. We thank Lassen Volcanic NP for detailing their ecologist.

**A bryophyte dominated sloping peatland near Granite Passage.
Photo: James Walton/NPS.**



Introduction

The nonvascular plant flora (mosses, liverworts, lichens) of Kenai Fjords National Park (KEFJ) is virtually unknown, but thought to be rich and diverse. The Park encompasses a wide range of habitat types, as well as climate gradients known to support spectacular bryophyte (mosses and liverworts) and lichen floras elsewhere in coastal Alaska (Worley 1972, Spribille et al. 2010). KEFJ contains a unique mixture of Pacific Northwest temperate rain forest and Arctic habitat types that, due to the steep terrain, are located within close proximity of each other. Nonvascular plants are a prominent and ecologically important component in many of these habitat types. Several studies in the southern coastal Alaska region have documented an extraordinary diversity in nonvascular plant species (e.g., lichens: Spribille et al. 2010 - Klondike Goldrush NHP, Talbot et al. 1992 - Tuxedni NWR; bryophytes: Worley 1972 - SE Alaska, Schofield et al. 2002 - Tuxedni NWR), and have included many taxa new to Alaska, new to North America, and several new to science.

Bryophytes and lichens represent a significant portion of the biodiversity and comprise a considerable amount of the biomass and annual production at high latitudes (Vitt et al. 1988). They play important ecological roles in nutrient cycling (Longton 1997), and many species are known to be important indicators of change in air quality, precipitation chemistry, and climate (Nash & Gries 1995; Gough & Crock 1997). Baseline estimates of bryophyte and lichen community composition can therefore be used as an index of ecological condition across a range of sites, and can also be used for monitoring changes in condition through time.

Given the remote nature of KEFJ parkland, explorations of its bryophyte and lichen flora have largely been focused near Exit Glacier, the only part of the Park accessible by road. Species documented near the glacier have been mostly small in number and primarily represent common lichen taxa within the region (e.g., Coghill 1984, Lowthian 1987, Schindler 1990, Bryden 2001, Bryden 2002). More recently, Miller et al.

2006 made a small opportunistic survey of bryophytes and lichens on nunataks within the Park. No KEFJ bryophyte and lichen species are currently listed in NPSpecies (IRMA 2014).

The purpose of this document is to present preliminary results of 2012-2013 bryophyte and lichen surveys that were conducted primarily in the low elevation Sitka spruce - mountain hemlock forest zone that characterizes much of the KEFJ coastline. The objective of the surveys was to provide a preliminary estimate of nonvascular species richness in forested sites. Additionally, these data will be used to inform and refine preparations for a future Park-wide inventory of these two plant groups. Future inventories will help enable KEFJ to protect populations of rare and sensitive species, will improve the Park's herbarium collection, and will contribute significant baseline data not only for the Park but also for Alaska as a whole.

Study Area


KEFJ is situated on the southeastern side of the Kenai Peninsula in southcentral Alaska. The Park is a narrow geographic area, roughly 50 km wide by 120 km long and encompassing 271, 133 ha. It is oriented in a northeasterly-southwesterly direction, located between 59.4° and 60.3° N, and between 149.5° and 151.0° W, and is bounded on the west and north by the Kenai Mountains and Harding Icefield, and to the east and south by the Gulf of Alaska (**Figure 1**).

The surficial landscape of the Park reflects a combination of glacial erosion, tectonic subduction (the coastline dropped approximately two meters in one day during the 1964 Alaska earthquake), and isostatic rebound as a result of recent glacial recession since the Little Ice Age maximum ca. 1850 A.D. Predominant surficial rock types in the Park include metasedimentary flysch, sandstone, siltstone, granodiorite, and conglomerate (Bradley & Donley 1995).

Climate near KEFJ Headquarters in Seward can be generalized as humid subarctic con-

Bryophyte and lichen communities in Kenai Fjords NP play many essential ecosystem roles within the landscape.

Figure 1. Bryophyte and lichen survey site locations during the 2012-2013 field seasons, Kenai Fjords NP.



continental to suboceanic with no dry season. Precipitation peaks in September with a monthly average of 175 cm, while snowfall peaks in December with a monthly average of 214 cm. Summers are cool with an average high/low in July of 16.8°C/9.9°C, while winters are mild at -1.1°C/-6.7°C in January (WRCC 2013).

Vegetation in KEFJ includes high elevation herbaceous meadows, dwarf shrub heaths, low to tall shrubs, and sparse to dense mid- and low elevation conifer forests. Willow and alder are common in more disturbed areas such as floodplains or avalanche tracks. Many small pockets of sloping peatland exist on the outer Harris and McCarty Fjord peninsulas. Vegetation communities in KEFJ are similar to those in neighboring coastal communities of western Prince Williams Sound and in many areas established following the Little Ice Age maximum.

Methods and Results

Figure 2 (top right). Documenting epiphytic macrolichen occurrence on a Sitka spruce tree at a off-frame FIA site near Nuka Bay. Photo: Michael Shephard/NPS.

Lichens are fungi that live in intimate association with algae and/or cyanobacteria. Epiphytic lichens, those growing on trees and shrubs, are excellent indicators of changes in air quality and climate due to their unique physiology and reliance on airborne nutrients. Epiphytic lichen communities can provide a wealth of information about forest health, function, and local climatic conditions; thus, they are widely used as biological indicators for natural resource assessments.

Bryophyte and lichen species listed in this report were encountered during the establishment of off-frame Forest Inventory and Analysis (FIA) epiphytic macrolichen monitoring plots in low elevation Sitka spruce - mountain hemlock forests. We also conducted intuitively-controlled surveys for interesting bryophyte and lichen taxa in or near these FIA sites.

Off-frame FIA epiphytic macrolichen surveys

During August 2012, the first author established eleven lichen monitoring plots in mature, low elevation Sitka spruce-mountain hemlock stands (**Appendix A**) using an 'off-frame' survey method developed by the FIA Program of the U.S. Forest Service (USFS 2013). We chose this forest type to monitor because it is one of the oldest and most commonly encountered along the KEFJ coastline. Representative, spatially dispersed survey sites were initially chosen by evaluating available GIS information, including satellite imagery, topographic maps, and potential vascular vegetation maps. Actual plot locations were selected after on-the-ground reconnaissance, to ensure plot-level uniformity of our targeted vegetation type. At each 34.7m-radius circular plot, a voucher collection was made and an abundance code was recorded for each macrolichen species on woody material > 0.5 m in height and within arm's reach. Freshly fallen lichens in the litter were also collected. Surveys lasted a minimum of 30 minutes and a maximum of two hours. Surveys were terminated when no additional new species were encountered within ten minutes time.

Data summaries from our FIA lichen fieldwork will be presented in a forthcoming NPS report, so we will only note that these FIA off-frame surveys will provide a range of park-specific information, as well as contribute information to the national FIA program, which houses one of the most extensive epiphytic macrolichen community data sets in North America. These data are invaluable for using lichens as forest health indicators by exploring relationships between lichen communities and large-scale environmental factors. Our survey data can



be used for biodiversity monitoring, while multivariate gradient models can address air quality and climate questions when combined with other regional datasets.

Intuitively-controlled bryophyte and lichen surveys

Many bryophytes and lichens are micro-habitat specific and are likely to be missed using survey methods based on more general macrohabitats, such as the aforementioned FIA method. Intuitively-controlled

Figure 3 (bottom right). Low elevation Sitka spruce - mountain hemlock forests, such as this fog-drenched stand in Square Bay near McArthur Pass, were the target vegetation type for our FIA surveys. Photo: James Walton/NPS




Figure 4 (top). Certain rare to uncommon species, like the moss *Takakia lepidozioides*, were only found in sloping peatlands such as this one near Fire Cove. Photo: James Walton/NPS.



Bryophytes (mosses, liverworts) are small autotrophic cryptogams (spore-producing plants) that often flourish in high latitude ecosystems. Bryophytes, like lichens, are also important indicators of air quality, precipitation chemistry, and climate. They provide important ecosystem functions as well, including high productivity and biomass accumulation, nutrient cycling, soil thermoregulation, vascular plant facilitation, and mycorrhizal relationships.

("intuition-based") surveys are an effective and efficient means for skilled taxonomists to maximize species capture in a relatively short amount of time. The method involves surveying habitats with known patterns of species occurrence and adapting survey methods as needed to include unanticipated substrates and microhabitats. Collection of common taxa are typically not treated equally at all survey sites, but rare taxa are treated equally at all sites.

Figure 5 (center right). Bryophyte and lichen diversity was found to be high at the shoreline - forest ecotone, which required careful and patient study of substrates, such as this exposed bedrock just above the hightide line. Photo: James Walton/NPS.




Figure 6 (bottom right). At the end of each field day bryophyte and lichen voucher collections were organized and dried on the M/V Serac. Photo: James Walton/NPS.

Between August 27-31, 2012 the Southwest Alaska Inventory and Monitoring Program (SWAN) brought in the second author through a work detail agreement with Lassen Volcanic National Park. Given the second author's expertise with the western North America bryophytes, his primary objective was to develop a voucher collection of bryophytes, while the first author focussed on the FIA lichen work. The bryophyte work to date consequently is constrained by the FIA lichen surveys locations at elevations below 100 m, in or near the Sitka spruce - mountain hemlock forest zone. Time permitting, the first author also conducted intuitively-based lichen and bryophyte surveys after completing daily FIA survey work.



Additional though less intensively surveyed habitats included sloping peatlands, coastal cliffs, and active colluvial slopes. Also included in this report are a handful of specimens opportunistically encountered by the first author during a brief stopover at the McArthur Weather Station, which is located on an alpine ridge in the southern part of the Park.

In total we surveyed 22 site areas (**Figure 1, Appendix A**) over roughly 15 person-days. Upon returning from the field specimens were air dried. Specimen voucher collection information, including collector, date, site and substrate information were entered into an MS Access database in about ten person days.

Identifications were made by the first and second authors. Laboratory identification and processing of voucher collections took approximately 3-5 times as long as time spent in the field (a common "rule-of-thumb" for bryological and lichenological work). The amount of lab work required was not only due to the high diversity encountered, but also a relatively high proportion of species requiring specialized literature and advanced laboratory examination.

Figure 7. *Kurzia trichoclados*, a small and pleasantly sweet aromatic liverwort that can be "sniffed out" in moist places such as shady boulder crevices or over *Sphagnum* moss. Discoveries at Crater Bay, Three Hole Bay, and Bear Cove are apparently the first for Alaska. Photo: James Walton/NPS.



Notable Discoveries and Future Work

Notable discoveries

While several species encountered during this project begin to fill regional information gaps, the following are some of the more notable that we have identified so far:

» *Polytrichastrum papillatum* – The discovery of this moss in Coleman Bay represents the second site now known for North America. The first site discovered was in 1967 at Thomson Pass along the Richardson Highway, near Valdez, Alaska. The species is otherwise only known from the Himalayan region in Asia.

» *Kurzia trichoclados* – Discovery of this small liverwort at Crater Bay, Three Hole Bay, and Bear Cove are apparently the first for Alaska. The closest known records are from British Columbia.

» *cf. Marsupella adusta* – The identification of this small liverwort is tentative pending further comparison work with other similar taxa, but if correct would also be a new state record for Alaska. The closest known records are from British Columbia.

» *Racodium rupestre* – Our single record of this dark filamentous rock-dwelling lichen near Northwest Lagoon is apparently the first for Alaska. The closest known records are from British Columbia.

» *Takakia lepidozoides* – As a member of one of the most morphologically unusual moss genera in the world, our collections appear to be the western-most in North America for this particular species. The next nearest locality is in Cordova, Alaska, with a scattering of sites then known from SE Alaska to British Columbia.

» *Myuroclada maximowiczii* – This moss was thought to be Asian until discovered in North America near King Salmon, Alaska (Steere & Schofield 1956). A handful of additional North American sites are now also known in the high arctic near Cape Thompson and the Brooks Range in Alaska.

» *Hypogymnia duplicata* – A western North America endemic, this epiphytic lichen was previously known only in coastal areas from SE Alaska into Oregon.

Additionally, we encountered at least three lichen species listed as rare by the Alaska Natural Heritage Program (AKNHP 2014): *Cavernularia lophyrea* (G2G3/S2S3), *Stereocaulon spathuliferum* (G4G5/S2), and *Umbilicaria angulata* (G2/S3). No rare bryophyte list for Alaska currently exists.

Future work


Complete the identification of difficult and interesting specimens. – There remain several taxa new to KEFJ which were excluded from **Appendix B** because more careful study is needed before applying a species name with certainty. These include several crustose lichen taxa, and bryophyte taxa which lack certain morphological components that aid in positively identifying the species quickly. We are continuing to review these collections (approx. 100 vouchers) and will be sending several to taxonomic specialists for identification.

Review historic collections from Kenai Fjords NP. – A small set of historic bryophyte and lichen vouchers exist in the KEFJ herbarium. These should be examined and annotated. Voucher specimens made by Miller et al. (2006) should also be reviewed.

Develop list of target sites for future comprehensive inventory fieldwork. – Regional bryophyte and lichen checklists should be thoroughly reviewed to ascertain what habitats and substrates yield taxa not documented in KEFJ. Landcover documents such as Boggs et al. (2008) will be important to consult in order to fine-tune what geographic areas, ecological systems (**Table 1**), and macrohabitats are available in the Park.



Polytrichastrum papillatum



Takakia lepidozoides



Myuroclada maximowiczii

Table 1. Ecological Systems (Boggs et al. 2008) expected to increase known bryophyte and lichen diversity in Kenai Fjords NP.

» Sea Cliff
» Tree Line Forest
» Sparse Vegetation Ridge / Cliff
» Floodplain
» Sloping / Depressional Peatland
» Tall Shrub
» Alpine Active Colluvial Slope
» Alpine Dwarf Shrub
» Alpine Floodplain
» Alpine / High Alpine Herbaceous

Literature Cited

- Alaska Natural Heritage Program. 2014. Rare Lichen List. <http://aknhp.uaa.alaska.edu/botany/rare-plants-species-lists/rare-lichen-list/> (Accessed 1/27/2014).
- Bradley, D. and T. Donley. 1995. Geologic map of Kenai Fjords National Park and vicinity. [DRAFT MAP] U.S. Geological Survey. http://www.ux1.eiu.edu/~cfrbj/parks/kefj/KFNP_geol_map_.pdf (Accessed 12/30/2013).
- Boggs, K., S. C. Klein, L. Flagstad, T. Boucher, J. Grunblatt and B. Koltun. 2008. Landcover classes, ecological systems and plant associations: Kenai Fjords National Park. Natural Resource Technical Report NPS/KEFJ/NRTR-2008/136. National Park Service, Fort Collins, Colorado.
- Bryden, W. 2001. Exit Glacier Vegetation Inventory Final Report In Conjunction With Exit Glacier Carrying Capacity. [Kenai Fjords Unpublished Report].
- Bryden, W. 2002. Lichen Inventory for Exit Glacier Study Area, Kenai Fjords National Park. In Martin, E.L. 2002. Kenai Fjords National Park Exit Glacier Visitor Experience and Resource Protection Plan.
- Coghill, K. 1984. Vegetation analysis of Exit Glacier Valley. [Kenai Fjords Unpublished Report].
- Esslinger, T. L. 2012. A cumulative checklist for the lichen-forming, lichenicolous and allied fungi of the continental United States and Canada. North Dakota State University: <http://www.ndsu.edu/pubweb/~esslinge/chcklst/chcklst7.htm> (First Posted 1 December 1997, Most Recent Version (#18) 13 December 2012), Fargo, North Dakota.
- Flora of North America Editorial Committee, eds. 2007. Flora of North America North of Mexico. New York and Oxford. Volume 27.
- Flora of North America Editorial Committee, eds. *In prep.* Flora of North America North of Mexico. New York and Oxford. Volume 28 and 29.
- Gough, L. P. and J. G. Crock. 1997. Distinguishing between natural geologic and anthropogenic trace element sources, Denali National Park and Preserve. U. S. Geological Survey professional paper, U. S. G. S., Washington D. C.
- Integrated Resource Management Application. 2014. NPSpecies. <https://irma.nps.gov/NPSpecies/Search/SpeciesList/KEFJ> (Accessed 2 Jan 2014).
- Longton, R. E. 1997. The role of bryophytes and lichens in polar ecosystems. In, Woodin, S. J. and M. Marquiss (eds.), Ecology of Arctic Environments. Blackwell Science Ltd.
- Lowthian, D. 1987. Lichen collection from Exit Glacier. [Kenai Fjords collection].
- Miller A, Carlson M, Lipkin R, Spencer P. 2006. Vascular plant inventory and baseline monitoring of nunatak communities. Lake Clark National Park and Preserve and Kenai Fjords National Park. Southwest Alaska Network, National Park Service. Anchorage, AK. 43 pp.
- Nash, T. and C. Gries. 1995. The use of lichens in atmospheric deposition studies with an emphasis on the Arctic. The Science of the Total Environment. 160-161: 729-736.
- Schindler, H. 1990. Zweiter Beitrag zur Flechtenflora von Alaska (Pribilof-Insel St. Paul; Kenai-Halbinsel Katmai Nat. Park und Denali Nat. Park). Herzogia 8: 335-356.
- Schofield, W., S. S. Talbot and S. L. Talbot. 2002. Bryophytes from Tuxedni Wilderness Area, Alaska. Journal of the Hattori Botanical Laboratory 92: 91-123.
- Spribille, T., S. Pérez-Ortega, T. Tønsberg, and D. Schirokauer. 2010. Lichens and lichenicolous fungi of the Klondike Gold Rush National Historic Park, Alaska, in a global biodiversity context. The Bryologist 113(3):439-515.
- Steere, W. C. and W. B. Schofield. 1956. *Myuroclada*, a genus new to North America. The Bryologist 59(1): 1-5.
- Talbot, S., S. L. Talbot and J. W. Thomson. 1992. Lichens of Tuxedni Wilderness Area, Alaska. The Bryologist 95(1):20-30.

U.S. Forest Service. 2013. Forest Inventory and Analysis National Program. <http://www.fia.fs.fed.us/> (Accessed 12/30/2013).

Vitt, D. H., J. E. Marsh and R. B. Bovey. 1988. Mosses, Lichens & Ferns of Northwest North America. Lone Pine Publishing, Vancouver, British Columbia, Canada. 296 pp.

Western Region Climate Center. 2013. Period of Record Monthly Climate Summary, Seward, Alaska (508371): 9/1/1949 - 9/30/2012. <http://www.wrcc.dri.edu/cgi-bin/cliMAIN.pl?ak8371> (Accessed 12/30/2013).

Worley, I. A. 1972. The Bryo-geography of Southeastern Alaska. Ph.D Dissertation, The University of British Columbia, Vancouver, B. C. 729pp.

Ptilidium ciliare, a common liverwort usually found growing among mosses and other liverworts on the forest floor or over rocks and cliffs. Photo: James Walton.



Appendix A: Survey Sites.

2012-2013 survey site areas. FIA = epiphytic macrolichen survey site, with some intuitively-controlled bryophyte and lichen surveys (time permitting). INT = intuitively-controlled bryophyte and lichen surveys.

Site Area	Survey Type	Survey Date	Latitude (NAD83)	Longitude (NAD83)	Elevation (m)
McArthur WX	INT	06 May 2012	59.4726	-150.3337	398
North Arm Nuka Bay	FIA	07 August 2012	59.56793	-150.5123	44
Pilot Harbor	INT	07 August 2012	59.60275	-150.49588	4
East Arm (westside)	FIA	08 August 2012	59.50071	-150.44939	81
	INT	16 August 2013			
James Lagoon	FIA	08 August 2012	59.58743	-150.42032	5
Moonlight Bay	INT	08 August 2012	59.50245	-150.357	38
Square Bay (northside)	FIA	09 August 2012	59.48741	-150.34161	20
Chance Cove	FIA	09 August 2012	59.48638	-150.31815	24
	INT	14 August 2013			
Square Bay (southside)	FIA	10 August 2012	59.47458	-150.38519	56
	INT	15 August 2013			
Crater Bay (southside)	INT	27 August 2012	59.691	-149.785	16
	INT	17 August 2013			
Crater Bay (northside)	INT	27 August 2012	59.7073	-149.81334	91
Crater Bay	FIA	28 August 2012	59.7052	-149.78075	23
Northwest Lagoon Beach	INT	28 August 2012	59.74992	-149.85205	30
Verdant Cove	FIA	29 August 2012	59.70901	-149.7493	61
Fire Cove	FIA	29 August 2012	59.65334	-149.77316	84
	INT	18 August 2013			
Granite Passage Peatland	INT	29 August 2012	59.6448	-149.75208	31
Harris Peninsula Lakes	INT	29 August 2012	59.73121	-149.75914	89
Three Hole Bay	FIA	30 August 2012	59.74353	-149.62755	45
Coleman Bay	INT	30 August 2012	59.88228	-149.61687	76
Coleman Bay Cliffs	INT	30 August 2012	59.8762	-149.6378	3
Bear Cove	FIA	31 August 2012	59.78849	-149.61649	75
	INT	19 August 2013			
Bear Cove Notch	INT	31 August 2012	59.79963	-149.60494	69




Bartramia pomiformis



Buckiella undulata



Bryum argenteum var.
argenteum



Bartramiopsis lescurii

Appendix B: Species List.

All species listed below are documented by voucher specimens currently housed at the NPS Alaska Regional Office, Anchorage, Alaska. Nomenclature for mosses and liverworts largely follows the Flora of North America (FNA 2007, FNA, *In prep.*), and Esslinger (2012) was used for most lichens. Basic substrate information follows each species name and geographical distribution is noted. Superscript numbers represent inclusion in the following nearby checklists: 1 = Schofield et al. 2002 (Tuxedni bryophytes), 2 = Worley 1972 (SE Alaska bryophytes), 3 = Talbot et al. 1992 (Tuxedni lichens).

It is important to note that many of the determinations below are provisional and subject to change upon further investigation. In addition to the species reported below, several collections have been identified only to genus and more records will be forthcoming. All photography in **Appendix B** is from the personal archives of James Walton or Martin Hutten.

Mosses (126 taxa)

Amblystegium serpens (Hedwig) Schimper
in B.S.G. – Moist organic soil. Cosmopolitan. ^{1,2}

Amphidium lapponicum (Hedwig) Schimper – Cliff crevices and bases. Circumboreal. ^{1,2}

Andreaea blyttii Schimper in P. Bruch and W.P. Schimper – Exposed rock, subalpine. Circummontane. ^{1,2}

Andreaea nivalis Hooker – Cliff base. Circummontane. ^{1,2}

Andreaea rupestris Hedwig – Full to partially shaded rocks. Circumboreal. ^{1,2}

Antitrichia curtipendula (Hedwig) Bridel – Epiphytic on trees and shrubs. Temperate-Interrupted. ^{1,2}

Arctoa fulvella (Dicks.) Bruch & Schimper in B.S.G. – Partially shaded lakeside boulder. Circummontane. ^{1,2}

Aulacomnium palustre (Hedwig) Schwaegr. – Seepage along cliff face. Cosmopolitan. ^{1,2}

Aulacomnium turgidum (Wahlenb.) Schwaegr. – Moist organic soil. Arctic-Alpine. ^{1,2}

Bartramia ithyphylla Bridel – Moist humic soil. Arctic-Alpine. ^{1,2}

Bartramia pomiformis Hedwig – Crevices of boulders and cliffs. Circumboreal. ^{1,2}

Bartramiopsis lescurii (James) Kindberg – Shaded mineral soil and boulders. Boreal-AmphiPacifc. ^{1,2}

Blindia acuta (Hedwig) Bruch & Schimper – Moist cliff face. Boreal-Interrupted. ^{1,2}

Brachythecium albicans (Hedwig) Schimper in B.S.G. – Rock near shoreline. Circumboreal. ^{1,2}

Brachythecium frigidum (C. Müll.) Besch. Western North America. ²

Bryum argenteum Hedwig var. *argenteum* – Soil on cliff shelf. Cosmopolitan. ^{1,2}

Buckiella undulata (Hedwig) Ireland – Forest floor, base of trees, boulder ledges. Temperate-Interrupted (Europe-W. NA). ^{1,2}

Bucklandiella affinis (F. Weber & D. Mohr) Bednarek-Ochyra & Ochyra in R. Ochyra et al. – Rock. Circumtemperate. ^{1,2}

Bucklandiella heterosticha (Hedwig) Bednarek-Ochyra & Ochyra in R. Ochyra et al. – Rock. Circumboreal. ^{1,2}


Calliergon cordifolium (Hedwig) Kindberg – Periodically flooded mineral soil. Circumboreal. ^{1,2}

Calliergonella cuspidata (Hedwig) Loeske – Moist hollow. Circumboreal. ²


Campylopus atrovirens De Notaris – Moist soil, rocks, cliff bases. Temperate-Wide. ²

Climacium dendroides (Hedwig) Web. & Mohr – Moist humic soil. Circumboreal. ^{1,2}

Codriophorus acicularis (Hedwig) P. Beauvois – Moist to wet rocks, soil, cliff



Dicranella palustris



Dicranodontium denudatum



Diphyscium foliosum



Hyplocomium splendens



Hypnum plicatum

bases. Circumboreal. ^{1,2}

Codriophorus fascicularis (Hedwig) Bednarek-Ochyra & Ochyra in R. Ochyra et al. – Rock. Circumboreal. ^{1,2}

Conostomum tetragonum (Hedwig) Lindberg – Alpine sod. Arctic-Alpine. ^{1,2}

Cratoneuron filicinum (Hedwig) Spruce – Moist cliff base. Circumboreal. ^{1,2}

Dichodontium pellucidum (Hedwig) Schimper – Moist mineral soil. Boreal-Interrupted. ^{1,2}

Dicranella cerviculata (Hedwig) Schimper – Cliff bases, mineral soil, rock. Circumpolar.

Dicranella heteromalla (Hedwig) Schimper – Mineral soil. Circumboreal. ^{1,2}

Dicranella palustris (Dicks.) Crundw. ex Warb. – Periodically flooded rock. Boreal-Interrupted. ^{1,2}

Dicranella schreberiana (Hedwig) Hilf. ex Crum & Anderson – Moist organic soil. Circumboreal. ^{1,2}

Dicranodontium denudatum (Bridel) E. Britton in N. L. Britton et al. – Cliff crevices, boulder ledges, and organic soil. Circumtemperate-Interrupted. ²

Dicranum fuscescens Turner var. *fuscescens* – Organic soil. Circumboreal. ^{1,2}

Dicranum majus Turner var. *majus* – Cliff bases, soil. Boreal-Interrupted. ^{1,2}

Dicranum scoparium Hedwig – Soil. Circumboreal. ^{1,2}

Dicranum tauricum Sapjegin – Heavily decayed CWD. Temperate-Western North America/Europe Disjunct. ^{1,2}

Diphyscium foliosum (Hedwig) D. Mohr – Mineral soil in cliff crevice. Circumboreal.

Distichium capillaceum (Hedwig) Bruch & Schimper – Humic soil on cliff shelf. Cosmopolitan. ^{1,2}

Drepanocladus aduncus (Hedwig) Warnst. – Periodically flooded mineral soil. Cosmopolitan. ^{1,2}

Fissidens osmundioides Hedwig – Seepy rock crevice near shoreline. Circumboreal. ²

Funaria hygrometrica Hedwig var. *hygrometrica* – Periodically flooded mineral soil. Cosmopolitan. ^{1,2}

Grimmia ramondii (Lamarck & de Candolle) Margadant – Mesic to seepy cliff faces and rocks. Boreal-Interrupted. ²

Grimmia torquata Drummond – Rock, cliff crevices. Boreal-Interrupted. ^{1,2}

Herzogiella striatella (Bridel) Z. Iwatsuki – Organic soil. Boreal-Interrupted. ^{1,2}

Heterocladium macounii Best – Shaded cliff bases, boulders, and rootwads. North America endemic. ²

Hygrohypnum ochraceum (Turn. ex Wils.) Loeske – Wet to seepy rocks. Boreal-Interrupted. ^{1,2}

Hylocomiastrum umbratum (Hedwig) Fleisch. – Boulder. Circumboreal. ²

Hylocomium splendens (Hedwig) Schimper in B.S.G. – Boulder top and side. Circumboreal. ^{1,2}

Hypnum circinale Hooker – Conifer bases and boles, cliff face. Temperate-Western North America Endemic. ²

Hypnum dieckii Ren. & Card. in Röhl – Moist cliff face and boulder. AmphiPacific. ^{1,2}

Hypnum lindbergii Mitt. – Moderately decayed CWD. Circumboreal. ^{1,2}

Hypnum plicatum (Lindberg) Jaeg. – Cliff base and boulder top. Boreal-Interrupted. ^{1,2}

Hypnum subimponens Lesquereux – Moist rock. AmphiPacific. ^{1,2}

Isopterygiopsis pulchella (Hedwig) Iwatsuki – Organic soil. Circumboreal. ^{1,2}

Isoetecium stoloniferum Bridel – Rocks and boulders. Temperate-Western North America Endemic. ^{1,2}

Iwatsukiella leucotricha (Mitt.) Buck & Crum – Conifer and hardwood twigs and branches. Boreal-AmphiPacific. ^{1,2}



Iwatsukiella leucotricha



Kinbergia oregana



Paraleucobryum enerve



Pohlia crudoides

Kinbergia oregana (Sull.) Ochyra – Over litterfall and rock near shoreline. Temperate-Western North America Endemic. ²

Kindbergia praelonga (Hedwig) Ochyra – Moist forest floor. Circumpolar to Temperate-Interrupted. ²

Leptodictyum riparium (Hedwig) Warnstorf – Submerged along lake margin. Cosmopolitan.

Mnium thomsonii Schimper – Organic soil. Circumboreal. ^{1,2}

Myuroclada maximowiczii (G. G. Borschow) Steere & Schofield – Moist organic soil. Circumboreal.

Niphotrichum ericoides (Bridel) Bednarek-Ochyra & Ochyra in R. Ochyra et al. – Organic soil. Circumboreal to circumtemperate. ²

Oedipodium griffithianum (Dickson) Schwägrichen – Moist organic soil. Circummontane. ^{1,2}

Oligotrichum hercynicum (Hedwig) Lamarck & A. P. DeCandolle – Mineral soil. Circummontane. ^{1,2}

Oligotrichum parallelum (Mitten) Kindberg – Mineral to organic soil. Boreal-AmphiPacific. ^{1,2}

Oncophorus virens (Hedwig) Bridel – Mineral soil of upturned rootwad. Circummontane. ²

Oncophorus wahlenbergii Bridel – Mineral soil. Circumboreal. ^{1,2}

Paraleucobryum enerve (Thedenius) Loeske – Humic soil over bedrock. Circumboreal. ²

Philonotis fontana (Hedwig) Bridel var. *fontana* – Wet cliff bases, rock, and soil. Circumboreal. ^{1,2}

Plagiomnium medium (Bruch & Schimper in B.S.G.) T. Kop. – Humus. Circumboreal. ^{1,2}

Plagiothecium denticulatum (Hedwig) Schimper in B.S.G. – On moss over mineral soil. Circumboreal. ^{1,2}

Plagiothecium laetum Schimper in B.S.G. – Soil, cliff base. Boreal-Interrupted. ^{1,2}

Plagiothecium piliferum (Sw. ex Hartm.) Schimper in B.S.G. – Cliff bases, Alnus branch. Temperate-Interrupted. ²

Pleuroziopsis ruthenica (Weinmann) E. Britton – Moist to wet forest floor. Boreal-AmphiPacific. ²

Pleurozium schreberi (Bridel) Mitt. – Organic soil over boulder. Circumboreal. ^{1,2}

Pogonatum contortum (Bridel) Lesquereux – Mineral soil. Temperate-Interrupted. ²

Pogonatum dentatum (Bridel) Bridel – Mineral soil. Circumboreal. ^{1,2}

Pogonatum urnigerum (Hedwig) P. Beauvois – Mineral soil. Circumpolar. ^{1,2}

Pohlia cruda (Hedwig) Lindberg – Cliff base. Circumboreal. ^{1,2}

Pohlia crudoides (Sullivant & Lesquereux) Brotherus in A. Engler & K. Prantl – Cliff crevice, boulder. North America endemic.

Pohlia nutans (Hedwig) Lindberg – Soil. Circumboreal. ^{1,2}

Pohlia prolifera (Kindberg ex Breidl.) Lindberg ex Arnell – Cliff face. Circumboreal. ^{1,2}

Pohlia wahlenbergii (Web. & Mohr) Andrews – Moist mineral soil. Cosmopolitan. ^{1,2}

Polytrichastrum alpinum (Hedwig) G. L. Smith var. *alpinum* – Mineral and organic soil. Circumboreal. ^{1,2}

Polytrichastrum formosum (Hedwig) G.L. Smith var. *formosum* – Wet mineral soil. Circumboreal. ^{1,2}

Polytrichastrum papillatum G.L. Smith – Mineral soil. N. America (Alaska) - Asia (Kashmir, e. Nepal).

Polytrichum juniperinum Hedwig – Mineral soil. Cosmopolitan. ^{1,2}

Polytrichum piliferum Hedwig – Mineral soil. Cosmopolitan. ^{1,2}

Pseudoleskea baileyi Best & Grout – Base of shrubs, rock, and CWD. Montane-Western North America Endemic. ^{1,2}

Pseudoleskea stenophylla Ren. & Card.



Pseudotaxiphyllum elegans




Ptilium crista-castrensis



Racomitrium lanuginosum



Rhytidiadelphus loreus



Sphagnum girgensohnii

in Röll – On Alnus. Temperate-North America Endemic Disjunct. ^{1,2}

Pseudotaxiphyllum elegans (Bridel) Z.
Iwatsuki – Rock outcrop, mineral soil.
Circumtemperate. ^{1,2}

Ptilium crista-castrensis (Hedwig) De Not.
– Over boulder. Circumboreal. ^{1,2}

Ptychostomum pseudotriquetrum (Hedwig)
J. R. Spence – Cliff base. Cosmopolitan.
²

Ptychostomum weigeliai (Sprengel) J. R.
Spence – Organic soil. Circumboreal. ²

Racomitrium lanuginosum (Hedwig) Bridel
– well lit boulder, alpine sod. Circum-
boreal. ²

Rhabdoweisia crispata (Dickson ex With-
ering) Lindberg – Organic soil. Cosmo-
politan. ²

Rhizomnium glabrescens (Kindberg) T.
Kop. – Organic soil. Temperate-West-
ern North America Endemic. ^{1,2}

Rhizomnium magnifolium (Horik.) T. Kop.
– Seepy organic soil. Circumboreal. ^{1,2}

Rhizomnium nudum (Britt. & Williams) T.
Kop. – Moist soil. Montane-AmphiPa-
cific. ^{1,2}

Rhizomnium pseudopunctatum (Bruch &
Schimper) T. Kop. – Mineral soil. Cir-
cumboreal. ²

Rhytidiadelphus loreus (Hedwig) Warnst.
– Base of trees and shrubs, forest floor.
Boreal-Interrupted. ^{1,2}

Rhytidiadelphus squarrosus (Hedwig)
Warnstorf – Over soil, in depression.
Boreal-Interrupted. ^{1,2}

Rhytidiadelphus triquetrus (Hedwig)
Warnst. – Moist organic soil. Circum-
boreal. ^{1,2}

Rosulabryum capillare (Hedwig) J. R. Spen-
ce – Seepy cliff face. Cosmopolitan. ¹

Sanionia uncinata (Hedwig) Loeske - For-
est floor, over boulder. Circumboreal.
^{1,2}

Schistidium maritimum (Turner ex Robt.
Scott) Bruch & Schimper – Boulders
above high tide line. Boreal-Interrupt-

ed. ^{1,2}

Scouleria aquatica Hooker – Wet soil.
Boreal-Temperate AmphiPacific. ^{1,2}

Sphagnum alaskense R.E. Andrus & Jans-
sens – Sloping peatland. Temperate-
Western North America Endemic. ²

Sphagnum compactum Lamarck & de
Candolle – Moss-covered boulder and
sloping peatland. Circumboreal. ^{1,2}

Sphagnum girgensohnii Russow – Forest
floor. Circumboreal. ^{1,2}

Sphagnum lindbergii Schimper – Sphag-
num mire. Circumboreal. ^{1,2}

Sphagnum quinquefarium (Lindberg)
Warnstorf – Sphagnum-mire. Circum-
boreal. ²

Sphagnum russowii Warnstorf – Sphagnum
mire. Boreal-Interrupted. ^{1,2}

Sphagnum squarrosum Crome – Depres-
sion in Sphagnum mire. Circumboreal.
^{1,2}

Sphagnum tenellum (Bridel) Bory – Sphag-
num mire. Circumboreal.

Straminergon stramineum (Dickson ex
Bridel) Hedenäs – Mire. Circumboreal.
^{1,2}

Takakia lepidozoioides S. Hattori & Inoue –
Organic soil over sheltered, vertically
inclined rock. Interrupted-Circumbo-
real. ²

Tayloria serrata (Hedwig) Bruch & Schim-
per in B.S.G. – Organic soils. Circum-
boreal. ²

Tetraphis geniculata Milde – Heavily
decayed CWD and organic soil. Inter-
rupted-Circumboreal. ²

Tortella tortuosa (Hedwig) Limpricht var.
tortuosa – Cliff overhang, mineral soil.
Circumboreal. ^{1,2}

Ulota megalospora Venturi in J. Röll – Epi-
phytic on conifer twigs. North America
Endemic. ²

Ulota obtusiuscula Macoun & Kindberg
– Epiphytic over conifers and hard-
woods. Temperate-Western North
America Endemic. ^{1,2}



Anastrepta orcadensis



Barbilophozia lycopodioides



Calycularia laxa



Diplophyllum albicans

Ulota phyllantha Bridel – Epiphytic on conifer, rock. Boreal-Interrupted. ^{1,2}

Warnstorfia fluitans (Hedwig) Loeske – Partially submerged rock at lake margin. Circumboreal. ^{1,2}

Warnstorfia sarmentosa (Wahlenberg) Hedenäs – Seepy organic soil and cliff shelf. Cosmopolitan. ¹

Liverworts (79 taxa)

Anastrepta orcadensis (Hook.) Schiffn. – Mesic to moist organic soils. Circumboreal-Interrupted. ²

Anastrophyllum assimile (Mitt.) Steph. – Mesic to moist organic soils. Boreal-Interrupted. ²

Anastrophyllum minutum (Schreb.) R. M. Schust. var. *minutum* – Cliffs, conifer base, soil. Circumboreal. ²

Anastrophyllum sphenoloboides Schust. – Mineral soil. Circumboreal.

Anthelia julacea (L.) Dumort. – Soil, cliff bases, boulders, base of conifers. Boreal-Interrupted. ^{1,2}

Anthelia juratzkana (Limpr.) Trevis. – Mineral soil. Circummontane. ^{1,2}

Barbilophozia floerkei (F. Weber & D. Mohr) Loeske – Organic soil. Circummontane. ^{1,2}

Barbilophozia kunzeana (Huebener) Gams – Moist organic soil. Circumboreal. ²

Barbilophozia lycopodioides (Wallr.) Loeske – Organic soils. Circumboreal. ^{1,2}

Bazzania tricenata (Wahlenb.) Lindb. – Base of conifers, soil, boulders. Boreal-Interrupted. ²

Blepharostoma trichophyllum (L.) Dumort. subsp. *trichophyllum* – Bark. Circumboreal. ^{1,2}

Calycularia laxa Lindb. & Arnell – Wet cliff bases and soils. Boreal-Amphi-Pacific.

Calypogeia azurea Stotler & Crotz – Moist soils. Probably circumboreal. ²

Calypogeia fissa (L.) Raddi subsp. *neogaea* R. M. Schuster – Damp soil. Circumboreal.

Calypogeia integristipula Steph. – Moist soils. Temperate-Interrupted. ²

Calypogeia muelleriana (Schiffn.) Müll. Frib. – Organic soil. Circumboreal. ^{1,2}

Calypogeia neesiana (C. Massal. & Caresia) Müll. Frib. – Moist soil and rock, bark. Circumboreal. ²

Calypogeia suecica (Arnell & J. Perss.) Müll. Frib. – Moist soil. Circumboreal.

Cephalozia bicuspidata (L.) Dumort. – Organic material. Circumboreal. ^{1,2}

Cephalozia bicuspidata (L.) Dum. subsp. *ambigua* (Maas.) Schust. – Organic material. Circumboreal. ²

Cephalozia cf. *connivens* (Dicks.) Lindb. – Moist soil. Circumboreal. ²

Cephalozia lunulifolia (Dumort.) Dumort. – Base of tree, organic soil in underhang of rock. Circumboreal. ^{1,2}

Cephaloziella divaricata (Sm.) Schiffn. var. *divaricata* – Mineral soil of upturned rootwad. Cosmopolitan. ²

Conocephalum salebrosum Szwedkowski, Buczkowska & Odrykoski – Moist forest floor. Circumboreal. ^{1,2}

Diplophyllum albicans (L.) Dum. – Wet cliff face, boulder crevice. Boreal-Interrupted. ^{1,2}

Diplophyllum obtusifolium (Hook.) Dumort. – Mineral soil of upturned rootwad. Circumboreal. ²

Diplophyllum plicatum Lindb. – Cliff face. Boreal-AmphiPacific. ^{1,2}

Diplophyllum taxifolium (Wahlenb.) Dumort. – Among moss over litterfall. Circumboreal. ^{1,2}

Douinia ovata (Dicks.) Buch. – Epiphytic on conifers and hardwoods. Boreal-Interrupted. ²

Frullania nisquallensis Sull. – Epiphytic on conifer and hardwood, over boulder. Temperate-AmphiPacific. ²

Gymnocolea inflata (Huds.) Dumort. –



Frullania nisquallensis



Herbertus aduncus



Lepidozia filamentosa



Marsupella emarginata

Sphagnum mire, wet organic soils.
Circumboreal. ²

Gymnomitrium concinnatum (Lightf.)
Corda – Soil and soil over rock. Montane-Interrupted. ^{1,2}

Gymnomitrium obtusum Lindb. – Mineral soil, underhang of cliff crevice. Temperate-Wide. ^{1,2}

Harpanthus flotovianus (Nees) Nees – Wet organic soil. Circumboreal. ²

Herbertus aduncus (Dickson) S. Gray
var. *aduncus* – Cliff face, epiphytic on conifer bole. Western North America Endemic. ²

Herbertus dicranus (Taylor) Trevisan
– Boulders, cliff faces, organic soil. Circumboreal-Interrupted. ²

Jungermannia exsertifolia Stephani – Moist mineral soil. Arctic-Alpine. ^{1,2}

Kurzia trichoclados (K. Muell.) Grolle
– Boulders, over Sphagnum, organic soils. Circumboreal.

Lepidozia filamentosa (Lehm. & Lindenb.)
Lindenb. – Organic soil on cliffs or over litterfall. Amphipacific. ²

Lepidozia reptans (L.) Dumort. – Moist stump and heavily decayed CWD. Probably circumboreal. ²

Lophocolea bidentata (L.) Dumort. – Over moist litterfall. Temperate-Wide. ²

Lophozia guttulata (Lindb. & H. Arnell)
Evans – Moist log in lake margin, upturned rootwad. Circumboreal. ²

Lophozia sudetica (Nees ex Huebener)
Grolle – Moist soils and boulder. Circumboreal.

Lophozia ventricosa (Dicks.) Dumort. – Organic material. Circumboreal. ^{1,2}

Lophozia ventricosa (Dicks.) Dumort. var. *longiflora* (Nees) Macoun – Organic material. Circumboreal.

Lophozia wenzelii (Nees) Steph. var. *wenzelii* – Organic soil. Arctic-Alpine. ^{1,2}

cf. *Marsupella adusta* (Nees) Spruce – Wet seepage. Circumboreal.

Marsupella emarginata (Ehrh.) Dumort. – Moist soil. Circumboreal. ^{1,2}

Marsupella sphacelata (Giesecke ex Lindenb.) Dumort. – Soil. Montane-Interrupted. ^{1,2}

Metzgeria conjugata Lindb. – Shaded wet cliff base. Cosmopolitan. ²

Moerckia blyttii (Moerch) Brockm. – Organic soil. Montane-Interrupted. ^{1,2}

Mylia taylorii (Hook.) S. Gray – Stumps, cliff bases, humic soil, conifer boles. Circumboreal. ²

Nardia scalaris Gray – Wet soil, boulders, cliffs. Circumboreal. ^{1,2}

Odontoschisma elongatum (Lindb.) Evans – Humic soil, boulder. Circumboreal. ²

Pellia endiviifolia (Dicks.) Dumort. – Moist soil. Circumboreal. ^{1,2}

Pellia epiphylla (L.) Corda – Organic soil. Circumtemperate. ²

Pellia neesiana (Gottsche) Limpr. – Mineral soil. Circumboreal. ^{1,2}

Plagiochila porelloides (Torr. ex Nees) Lindenb. – Organic soil, over litterfall, base of confers. Circumboreal. ^{1,2}

Pleurocladula albescens (Hook.) Grolle
– Over coniferous litterfall. Arctic-Alpine. ^{1,2}

Ptilidium californicum (Aust.) Underw. – Over downed wood, base of *Vaccinium*. Western North America Endemic. ²

Ptilidium ciliare (L.) Hampe – Over ground and rocks. Circumboreal. ^{1,2}


Radula bolanderi Gott. – On snag. Western North America Endemic. ²

Radula cf. *obtusiloba* Steph. – Wet cliff base. Boreal-Amphipacific. ²

Riccardia latifrons (Lindb.) Lindb. – Heavily decayed CWD. Circumboreal. ²

Riccardia multifida (L.) A. Gray – Moist litterfall. Circumboreal. ²

Scapania bolanderi Austin – Conifer stumps and boles. Temperate-Amphipacific. ²



Scapania bolanderi



Tetralophozia setiformis



Chaenotheca furfuracea



Cladonia arbuscula subsp.
beringiana

Scapania curta (Mart.) Dumort. – Bed-rock. Circumboreal. ²

Scapania cuspiduligera (Nees) Müll. Frib. – Thin soil. Circumboreal.

Scapania paludosa (Müll. Frib.) Müll. Frib. – Over wet soil and litterfall. Circumpolar. ²

Scapanis subalpina (Nees) Dumort. – Wet rock and organic material. Circumboreal. ^{1,2}

Scapania uliginosa (Sw. ex Lindenb.) Dumort. – Soil and streamside rock. Circumboreal. ^{1,2}

Scapania umbrosa (Schr.) Dumort. – Dead, downed wood. Boreal-Interrupted. ^{1,2}

Scapania undulata (L.) Dumort. var. *undulata* – Boulder and mineral soil over boulder. Circumboreal. ^{1,2}

Schistochilopsis incisa (Schr.) Konstant. var. *incisa* – Organic soils. Circumboreal. ^{1,2}

Schistochilopsis incisa var. *opacifolia* (Culmann) Bakalin – Organic soils. Probably circumpolar.

Sphenobolopsis pearsonii (Spruce) Schust. – Mineral soil. Temperate-Interrupted.

Tetralophozia setiformis (Ehrh.) Schljakov – Alpine sod. Circumboreal. ^{1,2}

Tritomaria exsecta (Schr.) Loeske – Wet organic soil. Temperate-Wide.

Tritomaria quinquedentata (Huds.) H. Buch – Underhang of boulder. Boreal-Interrupted. ^{1,2}

Lichens (92 taxa)

Alectoria sarmentosa (Ach.) Ach. subsp. *sarmentosa* – Conifer bark, wood. Incompletely circumpolar.

Baeomyces rufus (Hudson) Rebert. – Sandy soil. Circumpolar.

Bryoria bicolor (Ehrh.) Brodo & D. Hawksw. – Conifer bark. Incompletely circumpolar.

Bryoria cervinula Motyka ex Brodo & D.

Hawksw. – Conifer branches. Western North America Endemic.

Bryoria trichodes subsp. *americana* (Motyka) Brodo & D. Hawksw. – Conifer bark and wood. North America Endemic.

Cavernularia hultenii Degel. – Conifer bark and wood. Incompletely circumpolar. ³

Cavernularia lophyrea (Ach.) Degel. – Conifer bark and wood. Western North America.

Cetraria aculeata (Schreb.) Fr. – Sandy soil. Circumpolar.

Cetraria ericetorum subsp. *reticulata* (Räsänen) Kärnefelt – Soil over ground and rock. Western North America.

Cetraria islandica subsp. *crispiformis* (Räsänen) Kärnefelt – Mineral and organic soil. Incompletely circumpolar. ³

Chaenotheca furfuracea (L.) Tibell – Hollow in roots. Incompletely circumpolar.

Cladonia amaurocraea (Flörke) Schaerer – Over moss, soil. Circumpolar. ³

Cladonia arbuscula subsp. *beringiana* Ahti – Moss over rock. W. NA - E. Eurasia

Cladonia bellidiflora (Ach.) Schaerer – Conifer bark, soil. Incompletely circumpolar. ³

Cladonia carneola (Fr.) Fr. – Conifer bark, soil. Circumpolar.

Cladonia chlorophaea group (Flörke ex Sommerf.) Sprengel – Conifer bark and wood. Circumpolar. ³

Cladonia coccifera (L.) Willd. – Mineral soil, moss over rock. Circumpolar. ³

Cladonia crispata (Ach.) Flotow var. *crispata* – Sandy soil. Circumpolar.

Cladonia gracilis subsp. *vulnerata* Ahti – Forest floor; Forest floor. W. North America - E. Eurasia.

Cladonia maxima (Asahina) Ahti – Exposed moss over boulder. Circumpolar. ³

Cladonia ochrochlora Flörke – Conifer



Cladonia uncialis



Hypogymnia duplicata



Lichenomphalia umbellifera



Lobaria linata



Nephroma isidiosum

bark and wood. Circumpolar.

Cladonia phyllophora Hoffm. – Sandy soil. Circumpolar.³

Cladonia pyxidata (L.) Hoffm. – Mineral and organic soils. Circumpolar.

Cladonia rangiferina (L.) F. H. Wigg. – Mineral and organic soils. Circumpolar.³

Cladonia squamosa Hoffm. – Conifer bark, branches, soil. Circumpolar.

Cladonia stygia (Fr.) Ruoss – On moss over exposed boulder. Circumpolar.

Cladonia uncialis (L.) F. H. Wigg. – Soil, soil over rock. Circumpolar.³

Dibaeis baeomyces (L. f.) Rambold & Hertel – Organic soils. Circumpolar.

Ephebe sp. Fr. – Exposed rock in and near streamlets. Species determination pending.

Fuscopannaria cf. *confusa* (P. M. Jørg.) P. M. Jørg. – Conifer branch. Probably circumpolar.

Graphis scripta (L.) Ach. – Alnus branches. Circumtemperate.

Hydropunctaria maura group (Wahleb.) Keller, Gueidan & Thüs – Rock above high tide line. Cosmopolitan.

Hypogymnia apinnata Goward & McCune – Conifer bark and wood. Western North America Endemic.

Hypogymnia duplicata (Ach.) Rasm. – Conifer branches. Western North America Endemic.

Hypogymnia physodes (L.) Nyl. – Conifer twig. Circumpolar.³

Hypogymnia tubulosa (Schaerer) Hav. – Conifer branch. Incompletely circumpolar.

Icmadophila ericetorum (L.) Zahlbr. – Rotten conifer stump. Circumpolar.

Leptogium lichenoides (L.) Zahlbr. – Rock. Circumpolar.

Leptogium saturninum (Dickson) Nyl. – Conifer branch. Circumpolar.³

Lichenomphalia umbellifera (L. : Fr.) Red-head, Lutzoni, Moncalvo & Vilgalys – Rotten conifer wood. Circumpolar.

Lobaria linata (Ach.) Rabenh. – Conifer bark and wood. Incompletely circumpolar.³

Lobaria oregana (Tuck.) Müll. – Conifer bark and wood. Western North America.

Lobaria pulmonaria (L.) Hoffm. – Conifer bark and wood. Incompletely circumpolar.³

Lobaria scrobiculata (Scop.) DC. – Conifer branch. Circumpolar.

Loxosporopsis corallifera Brodo, Henssen & Imshaug – Conifer bark. Western North America.

Massalongia carnosa (Dickson) Körber – Rock. Circumpolar.

Mycoblastus sanguinarius (L.) Norman – Conifer litterfall. Circumpolar.³

Nephroma bellum (Sprengel) Tuck. – Conifer bark and wood. Circumpolar.³

Nephroma helveticum subsp. *sipeanum* (Gyelnik) Goward & Ahti – Conifer and hardwood bark. Incompletely circumpolar.

Nephroma isidiosum (Nyl.) Gyelnik – Salix bark. Incompletely circumpolar.

Nephroma parile (Ach.) Ach. – Conifer branch. Circumpolar.³

Nephroma resupinatum (L.) Ach. – Rotten wood, Salix bark. Circumpolar.

Parmelia cf. *hygrophila* Goward & Ahti – Conifer wood. Western North America.

Parmelia pseudosulcata Gyelnik – Conifer bark. Western North America.

Parmelia saxatilis (L.) Ach. – Conifer bark and wood. Circumpolar.³

Parmelia squarrosa Hale – Conifer bark. Incompletely circumpolar.

Parmeliopsis hyperopta (Ach.) Arnold – Conifer bark, wood. Circumpolar.³


Peltigera britannica (Gyelnik) Holt.-Hartw. & Tønsberg – Forest floor, over moss.



Pilophorus clavatus



Platismatia lacunosa



Pseudocyphellaria crocata



Siphula ceratites

W. NA - E. Eurasia.

Peltigera collina (Ach.) Schrader – Conifer bark, rock. Probably incompletely circumpolar.³

Peltigera membranacea (Ach.) Nyl. – Moss over soil. Incompletely circumpolar.³

Peltigera neopolydactyla (Gyelnik) Gyelnik – Forest floor, over moss. Probably incompletely circumpolar.³

Peltigera praetextata (Flörke ex Sommerf.) Zopf – Over moss, rock. Circumpolar.

Phylliscum demangeonii (Moug. & Mont.) Nyl. – Rock near shoreline. Incompletely circumpolar.

Physcia caesia (Hoffm.) Fürnr. – Rock near shoreline. Circumpolar.³

Pilophorus acicularis (Ach.) Th. Fr. – Rock. W. NA - E. Eurasia.

Pilophorus clavatus Th. Fr. – Rock. W. NA - E. Eurasia.

Pilophorus nigricaulis Satô – Rock. W. NA - E. Eurasia.

Placopsis gelida (L.) Lindsay – Rock. Probably circumpolar.³

Platismatia glauca (L.) Culb. & C. Culb. – Conifer bark and wood. N. America - W. Eurasia.³

Platismatia herrei (Imshaug) Culb. & C. Culb. – Conifer bark and wood. Western North America.

Platismatia lacunosa (Ach.) W. L. Culb. & C. F. Culb. – Conifer bark and wood. Western North America.³

Platismatia norvegica (Lynge) Culb. & C. Culb. – Conifer bark, wood. N. America - E. Eurasia.

Polychidium muscicola (Ach.) Gray – Lake-side boulder above high water mark. Possibly circumpolar.

Protopannaria pezizoides (Weber) P. M. Jørg. & S. Ekman – Moss over soil. Circumpolar.³

Pseudocyphellaria anomala Brodo & Ahti – Conifer bark and wood. Western North America.

Pseudocyphellaria crocata (L.) Vainio – Salix bark. Incompletely circumpolar.

Psoroma hypnorum (Vahl) Gray – Over moss, rotten wood. Circumpolar.³

Racodium rupestre Pers. – Shaded underhang of boulder. W. NA - W. Eurasia.

Ramalina roesleri (Hochst. ex Schaerer) Hue – Conifer twig. Probably incompletely circumpolar.³

Siphula ceratites (Wahlenb.) Fr. – Moist soil. Circumpolar.

Sphaerophorus tuckermanii Räsänen – Conifer bark and wood. Apparently circumpolar.³

Stereocaulon paschale (L.) Hoffm. – Soil. Circumpolar.³

Stereocaulon spathuliferum Vainio – Rock. Incompletely circumpolar.

Stereocaulon tomentosum Fr. – Sandy soil. Circumpolar.

Stereocaulon vesuvianum Pers. – Rock. Circumpolar.

Tuckermannopsis chlorophylla (Willd.) Hale – Conifer bark and wood. Incompletely circumpolar.³

Umbilicaria angulata Tuck. – Rock. W. NA - E. Eurasia.

Umbilicaria deusta (L.) Baumg. – Rock. Circumpolar.

Umbilicaria hyperborea (Ach.) Hoffm. var. *hyperborea* – Rock. Circumpolar.

Umbilicaria torrefacta (Lightf.) Schrader – Rock. Circumpolar.

Usnea filipendula Stirton – Conifer bark. Probably circumpolar.

Usnea longissima Ach. – Conifer branches near shoreline. Incompletely circumpolar.

Appendix C: Potential Species List.

The following taxa are known to be near KEFJ but were not reported by us during the course of our present work (see **Appendix B** and **Appendix D**). This list is largely built from Schofield et al. (2002), Worley (1972), and Talbot et al. (1992), and can be used as a guide for targeting additional species that may occur within Park boundaries. Note that this list is not all-inclusive of what could be found in KEFJ. The recent lichen checklist developed for Klondike Goldrush N.H.P. by Spribille et al. (2010), reporting over 750 taxa, can also be used as an additional guide for searching out many additional lichens.

Mosses

<i>Abietinella abietina</i>	<i>Codriophorus mollis</i>
<i>Aloina brevirostris</i>	<i>Codriophorus ryszardii</i>
<i>Amphidium mougeotii</i>	<i>Codriophorus varius</i>
<i>Andreaea rothii</i>	<i>Coscinodon yukonensis</i>
<i>Anoetangium aestivum</i>	<i>Cynodontium jenneri</i>
<i>Anomobryum filiforme</i>	<i>Cynodontium polycarpon</i>
<i>Antitrichia californica</i>	<i>Cynodontium schisti</i>
<i>Aongstroemia longipes</i>	<i>Cynodontium strumiferum</i>
<i>Arctoa hyperborea</i>	<i>Cynodontium tenellum</i>
<i>Atrichum selwynii</i>	<i>Dichelyma falcatum</i>
<i>Bartramia subulata</i>	<i>Dichodontium olympicum</i>
<i>Brachydontium olympicum</i>	<i>Dicranella crispa</i>
<i>Brachythecium erythrorrhizon</i>	<i>Dicranella rufescens</i>
<i>Brachythecium plumosum</i>	<i>Dicranella subulata</i>
<i>Brachythecium rivulare</i>	<i>Dicranella varia</i>
<i>Brachythecium salebrosum</i>	<i>Dicranodontium asperulum</i>
<i>Brachythecium turgidum</i>	<i>Dicranodontium uncinatum</i>
<i>Brachythecium velutinum</i>	<i>Dicranoweisia crispula</i>
<i>Brotherella canadensis</i>	<i>Dicranum acutifolium</i>
<i>Bryhnia hultenii</i>	<i>Dicranum bonjeanii</i>
<i>Bryoerythrophyllum inaequalifolium</i>	<i>Dicranum elongatum</i>
<i>Bryoerythrophyllum recurvirostre</i>	<i>Dicranum fragilifolium</i>
<i>Bucklandiella brevipes</i>	<i>Dicranum howellii</i>
<i>Bucklandiella lawtoniae</i>	<i>Dicranum majus</i>
<i>Bucklandiella macounii</i>	<i>Dicranum muehlenbeckii</i>
<i>Bucklandiella occidentalis</i>	<i>Dicranum pallidisetum</i>
<i>Bucklandiella sudetica</i>	<i>Dicranum polysetum</i>
<i>Buxbaumia viridis</i>	<i>Dicranum spadiceum</i>
<i>Calliergon giganteum</i>	<i>Dicranum undulatum</i>
<i>Calliergon megalophyllum</i>	<i>Didymodon fallax</i>
<i>Campylium stellatum</i>	<i>Didymodon ferrugineus</i>
<i>Campylophyllum halleri</i>	<i>Didymodon giganteus</i>
<i>Campylophyllum hispidulum</i>	<i>Didymodon nigrescens</i>
<i>Campylopus gracilis</i>	<i>Didymodon rigidulus</i>
<i>Campylopus schimperi</i>	<i>Didymodon vinealis</i>
<i>Catoscopium nigratum</i>	<i>Distichium inclinatum</i>
<i>Ceratodon purpureus</i>	<i>Ditrichum ambiguum</i>
<i>Cinclidium subrotundum</i>	<i>Drepanocladus polygamus</i>
<i>Cirriphyllum cirrosum</i>	<i>Echinophyllum sachalinense</i>
<i>Claopodium bolanderi</i>	<i>Encalypta affinis</i>
<i>Claopodium crispifolium</i>	<i>Encalypta ciliata</i>
<i>Claopodium pellucinerve</i>	<i>Encalypta procera</i>
<i>Claopodium whippleanum</i>	<i>Encalypta rhaetocarpa</i>

Appendix C continued

<i>Encalypta vulgaris</i>	<i>Isopterygiopsis muelleriana</i>
<i>Entodon concinnus</i>	<i>Isothecium myosuroides</i>
<i>Eurhynchiastrum pulchellum</i>	<i>Kiaeria blyttii</i>
<i>Fissidens adianthoides</i>	<i>Kiaeria falcata</i>
<i>Fissidens bryoides</i>	<i>Kiaeria glacialis</i>
<i>Fissidens grandifrons</i>	<i>Kiaeria starkei</i>
<i>Fontinalis antipyretica</i>	<i>Leptobryum pyriforme</i>
<i>Fontinalis hypnoides</i>	<i>Lescuraea saxicola</i>
<i>Gemmabryum caespitium</i>	<i>Leskeella nervosa</i>
<i>Gemmabryum dichotomum</i>	<i>Leucolepis acanthoneura</i>
<i>Grimmia anodon</i>	<i>Loeskypnum badium</i>
<i>Grimmia anomola</i>	<i>Loeskypnum wickesiae</i>
<i>Grimmia attenuata</i>	<i>Meesia longiseta</i>
<i>Grimmia donniana</i>	<i>Meesia triquetra</i>
<i>Grimmia leibergii</i>	<i>Meesia uliginosa</i>
<i>Grimmia longirostris</i>	<i>Meiotrichum lyallii</i>
<i>Grimmia mollis</i>	<i>Mnium blyttii</i>
<i>Grimmia muehlenbeckii</i>	<i>Mnium lycopodioides</i>
<i>Grimmia ovalis</i>	<i>Mnium spinosum</i>
<i>Grimmia sessitana</i>	<i>Mnium spinulosum</i>
<i>Grimmia trichophylla</i>	<i>Myurella julacea</i>
<i>Gymnostomum aeruginosum</i>	<i>Myurella sibirica</i>
<i>Hamatocaulis vernicosus</i>	<i>Neckera douglasii</i>
<i>Haplodontium macrocarpum</i>	<i>Neckera menziesii</i>
<i>Hedwigia ciliata</i>	<i>Neckera pennata</i>
<i>Helodium blandowii</i>	<i>Niphotrichum canescens</i>
<i>Herzogiella adscendens</i>	<i>Niphotrichum elongatum</i>
<i>Heterocladium dimorphum</i>	<i>Niphotrichum muticum</i>
<i>Heterocladium procurrens</i>	<i>Oligotrichum aligerum</i>
<i>Homalothecium fulgescens</i>	<i>Orthothecium chryseum</i>
<i>Homalothecium nevadense</i>	<i>Orthothecium strictum</i>
<i>Homalothecium nuttallii</i>	<i>Orthotrichum anomalum</i>
<i>Hookeria acutifolia</i>	<i>Orthotrichum consimile</i>
<i>Hookeria lucens</i>	<i>Orthotrichum lyellii</i>
<i>Hygroamblystegium varium</i>	<i>Orthotrichum rupestre</i>
<i>Hygrohypnum alpestre</i>	<i>Orthotrichum speciosum</i>
<i>Hygrohypnum alpinum</i>	<i>Orthotrichum striatum</i>
<i>Hygrohypnum bestii</i>	<i>Orthotrichum cupulatum</i>
<i>Hygrohypnum luridum</i>	<i>Orthotrichum obtusifolium</i>
<i>Hygrohypnum molle</i>	<i>Orthotrichum pallens</i>
<i>Hygrohypnum polare</i>	<i>Orthotrichum pulchellum</i>
<i>Hymenostylium recurvirostrum</i> var. <i>recurvirostrum</i>	<i>Paludella squarrosa</i>
<i>Hymenostylium recurvirostrum</i> var. <i>insigne</i>	<i>Palustriella commutata</i>
<i>Hypnum bambergeri</i>	<i>Palustriella falcata</i>
<i>Hypnum callichroum</i>	<i>Philonotis capillaris</i>
<i>Hypnum cupressiforme</i>	<i>Philonotis fontana</i>
<i>Hypnum cupressiforme</i> var. <i>subjulaceum</i>	<i>Philonotis marchica</i>
<i>Hypnum hamulosum</i>	<i>Philonotis seriata</i>
<i>Hypnum pallens</i>	<i>Plagiobryum zierii</i>
<i>Hypnum pratense</i>	<i>Plagiomnium affine</i>
<i>Hypnum revolutum</i>	<i>Plagiomnium cuspidatum</i>
<i>Hypopterygium flavolimbatum</i>	<i>Plagiomnium ellipticum</i>
<i>Imbriobryum miniatum</i>	<i>Plagiomnium insigne</i>

Appendix C continued

<i>Plagiomnium rostratum</i>	<i>Scleropodium obtusifolium</i>
<i>Plagiopus oederianus</i>	<i>Scorpidium revolvens</i>
<i>Plagiothecium cavifolium</i>	<i>Scorpidium scorpioides</i>
<i>Platydictya jungermannioides</i>	<i>Seligeria campylopoda</i>
<i>Pohlia andalusica</i>	<i>Seligeria donniana</i>
<i>Pohlia annotina</i>	<i>Seligeria pusilla</i>
<i>Pohlia bulbifera</i>	<i>Sphagnum andersonianum</i>
<i>Pohlia columbica</i>	<i>Sphagnum angustifolium</i>
<i>Pohlia drummondii</i>	<i>Sphagnum annulatum</i>
<i>Pohlia elongata</i>	<i>Sphagnum aongstroemii</i>
<i>Pohlia filum</i>	<i>Sphagnum austinii</i>
<i>Polytrichastrum longisetum</i>	<i>Sphagnum balticum</i>
<i>Polytrichastrum sexangulare</i>	<i>Sphagnum bartlettianum</i>
<i>Polytrichum commune</i>	<i>Sphagnum brevifolium</i>
<i>Polytrichum strictum</i>	<i>Sphagnum capillifolium</i>
<i>Pseudobryum cinclidioides</i>	<i>Sphagnum centrale</i>
<i>Pseudoleskea atricha</i>	<i>Sphagnum fimbriatum</i>
<i>Pseudoleskea incurvata</i> var. <i>incurvata</i>	<i>Sphagnum fuscum</i>
<i>Pseudoleskea julacea</i>	<i>Sphagnum henryense</i>
<i>Pseudoleskea patens</i>	<i>Sphagnum inexpectatum</i>
<i>Pseudoleskea radicata</i> var. <i>radicata</i>	<i>Sphagnum inundatum</i>
<i>Pterigynandrum filiforme</i>	<i>Sphagnum lenense</i>
<i>Ptychostomum arcticum</i>	<i>Sphagnum magellanicum</i>
<i>Ptychostomum creberrimum</i>	<i>Sphagnum mendocinum</i>
<i>Ptychostomum inclinatum</i>	<i>Sphagnum obtusum</i>
<i>Ptychostomum knowltonii</i>	<i>Sphagnum pacificum</i>
<i>Ptychostomum pallens</i>	<i>Sphagnum papillosum</i>
<i>Ptychostomum pallescens</i>	<i>Sphagnum platyphyllum</i>
<i>Ptychostomum turbinatum</i>	<i>Sphagnum pseudopapillosum</i>
<i>Pylaisia polyantha</i>	<i>Sphagnum riparium</i>
<i>Pylaisia selwynii</i>	<i>Sphagnum rubellum</i>
<i>Rhizomnium andrewsianum</i>	<i>Sphagnum rubiginosum</i>
<i>Rhytidiadelphus subpinnatus</i>	<i>Sphagnum sitchense</i>
<i>Rhytidiopsis robusta</i>	<i>Sphagnum sjorsianum</i>
<i>Rhytidium rugosum</i>	<i>Sphagnum subfulvum</i>
<i>Rosulabryum flaccidum</i>	<i>Sphagnum subnitens</i>
<i>Sanionia orthothecoides</i>	<i>Sphagnum subsecundum</i>
<i>Schistidium agassizii</i>	<i>Sphagnum teres</i>
<i>Schistidium apocarpum</i>	<i>Sphagnum tongassense</i>
<i>Schistidium boreale</i>	<i>Sphagnum warnstorffii</i>
<i>Schistidium dupretii</i>	<i>Sphagnum wilfii</i>
<i>Schistidium frigidum</i>	<i>Splachnum sphaericum</i>
<i>Schistidium papillosum</i>	<i>Splachnum vasculosum</i>
<i>Schistidium platyphyllum</i>	<i>Syntrichia norvegica</i>
<i>Schistidium rivulare</i>	<i>Syntrichia ruralis</i>
<i>Schistidium strictum</i>	<i>Tayloria lingulata</i>
<i>Schistostega pennata</i>	<i>Tetraphis pellucida</i> var. <i>pellucida</i>
<i>Sciuro-hypnum latifolium</i>	<i>Tetraplodon mnioides</i>
<i>Sciuro-hypnum oedipodium</i>	<i>Tetraplodon urceolatus</i>
<i>Sciuro-hypnum populeum</i>	<i>Tetradontium brownianum</i>
<i>Sciuro-hypnum reflexum</i>	<i>Tetradontium repandum</i>
<i>Sciuro-hypnum starkei</i>	<i>Thamnobryum neckeroides</i>
<i>Scleropodium cespitans</i>	<i>Thuidium delicatulum</i> var. <i>radicans</i>

Appendix C continued

Timmia austriaca
Timmia norvegica
Tomentypnum nitens
Tortella fragilis
Tortella inclinata
Tortella tortuosa
Tortula deciduidentata
Tortula hoppeana
Tortula mucronifolia
Trachycystis flagellaris
Trematodon ambiguus
Trichostomum tenuirostre
Ulota barclayi
Ulota curvifolia
Ulota drummondii
Ulota reptans
Warnstorfia exannulata
Warnstorfia pseudostraminea
Warnstorfia sarmentosa
Warnstorfia trichophylla
Warnstorfia tundrae
Weissia controversa
Zygodon reinwardtii
Zygodon viridissimus

Liverworts

Anastrophyllum donnianum
Anastrophyllum saxicola
Aneura pinguis
Apometzgeria pubescens
Apotreubia nana
Asterella gracilis
Asterella lindenbergiana
Athalamia hyalina
Barbilophozia attenuata
Barbilophozia barbata
Barbilophozia hatcheri
Barbilophozia quadriloba
Bazzania ambigua
Bazzania pearsonii
Bazzania trilobata
Blasia pusilla
Calypogeia sphagnicola
Cephalozia catenulata
Cephalozia leucantha
Cephalozia loitlesbergeri
Cephalozia pleniceps
Cephaloziella divaricata
Cephaloziella rubella
Chiloscyphus pallescens
Chiloscyphus polyanthos

Cladopodiella fluitans
Diplophyllum imbricatum
Frullania californica
Frullania chilcootensis
Frullania franciscana
Geocalyx graveolens
Gymnomitrium alpinum
Gymnomitrium brevissimum
Gymnomitrium corallioides
Gymnomitrium pacificum
Gyrothyra underwoodiana
Haplomitrium hookeri
Harpanthus scutatus
Hygrobrella laxifolia
Jungermannia atrovirens
Jungermannia confertissima
Jungermannia hyalina
Jungermannia hyalinum
Jungermannia leiantha
Jungermannia obovata
Jungermannia pumila
Jungermannia rubra
Jungermannia sphaerocarpa
Kurzia setacea
Leiocolea bantriensis
Leiocolea gillmanii
Leiocolea heterocolpos
Lepidozia sandvicensis
Lophocolea cuspidata
Lophocolea heterophylla
Lophozia alpestris
Lophozia badensis
Lophozia bicrenata
Lophozia excisa
Lophozia obtusa
Marchantia polymorpha
Marsupella apiculata
Marsupella sparsifolia
Metzgeria leptoneura
Metzgeria temperata
Moerckia hibernica
Nardia compressa
Nardia geoscyphus
Plagiochila arctica
Plagiochila satoi
Plagiochila schofieldiana
Plagiochila semidecurrens
Pleurozia purpurea
Porella cordaeana
Porella navicularis
Porella platyphylla
Porella roellii
Preissia quadrata

Appendix C continued

Radula complanata
Reboulia hemisphaerica
Riccardia chamedryfolia
Riccardia palmata
Rivulariella gemmipara
Sauteria alpina
Scapania americana
Scapania gymnostomophila
Scapania irrigua
Scapania mucronata
Scapania paludicola
Scapania scandica
Scapania undulata
Schofieldia monticola
Tritomaria polita

Lichens

Alectoria ochroleuca
Amandinea coniops
Amandinea punctata
Arctocetraria andrejevii
Arthrorhaphis alpina
Bacidia subincompta
Baeomyces carneus
Biatorea vernalis
Bryocaulon divergens
Buellia erubescens
Caloplaca cerina
Caloplaca ferruginea
Calvitimela aglaea
Candelariella aurella
Catolechia wahlenbergii
Cetraria islandica subsp. *islandica*
Cetraria kamczatica
Cetraria laevigata
Cladonia cenotea
Cladonia cervicornis subsp. *cervicornis*
Cladonia coniocraea
Cladonia cornuta
Cladonia decorticata
Cladonia ecmocyna
Cladonia fimbriata
Cladonia furcata
Cladonia gracilis subsp. *gracilis*
Cladonia gracilis subsp. *turbinata*
Cladonia grayi
Cladonia humilis
Cladonia metacorallifera
Cladonia mitis
Cladonia pleurota
Cladonia scabriuscula

Cladonia stellaris
Cladonia stricta
Cladonia subfurcata
Cladonia sulphurina
Cladonia umbricola
Cladonia verruculosa
Collema furfuraceum
Collema nigrescens
Collema subflaccidum
Collema undulatum
Dactylina arctica
Dactylina beringica
Dermatocarpon miniatum
Dermatocarpon reticulatum
Diploschistes scruposus
Farnoldia hypocrita
Flavocetraria cucullata
Flavocetraria nivalis
Gowardia nigricans
Hafellia disciformis
Helocarpon crassipes
Hypogymnia enteromorpha
Hypogymnia inactiva
Hypogymnia vittata
Lecanora albella var. *albella*
Lecanora albellula
Lecanora cinereofusca
Lecanora circumborealis
Lecanora epanora
Lecanora hybocarpa
Lecanora impudens
Lecanora intricata
Lecanora pacifica
Lecanora polytropa
Lecanora pulicaris
Lecanora rupicola
Lecanora symmicta
Lecanora varia
Lecidea albohyalina
Lecidea auriculata
Lecidea lapicida
Lecidea tessellata
Lecidella euphorea
Lepraria spp.
Lobaria hallii
Melanelia commixta
Melanelia hepatizon
Melanelia stygia
Melanohalea elegantula
Melanohalea exasperatula
Melanohalea olivacea
Melanohalea olivaceoides
Melanohalea septentrionalis

Appendix C continued

<i>Micarea assimilata</i>	<i>Rhizocarpon inarense</i>
<i>Micarea denigrata</i>	<i>Rhizocarpon polycarpum</i>
<i>Mycoblastus alpinus</i>	<i>Rhizoplaca chrysoleuca</i>
<i>Nephroma arcticum</i>	<i>Rinodina archaea</i>
<i>Ochrolechia arborea</i>	<i>Rinodina septentrionalis</i>
<i>Ochrolechia frigida</i>	<i>Sagiolechia rhexoblephara</i>
<i>Ochrolechia laevigata</i>	<i>Scoliciosporum chlorococcum</i>
<i>Ochrolechia oregonensis</i>	<i>Solorina crocea</i>
<i>Ochrolechia szatalaensis</i>	<i>Sphaerophorus fragilis</i>
<i>Ochrolechia xanthostoma</i>	<i>Sphaerophorus globosus</i>
<i>Ophioparma lapponica</i>	<i>Staurothele drummondii</i>
<i>Parmelia omphalodes</i>	<i>Stereocaulon alpinum</i>
<i>Parmelia sulcata</i>	<i>Stereocaulon botryosum</i>
<i>Peltigera aphthosa</i>	<i>Stereocaulon condensatum</i>
<i>Peltigera canina</i>	<i>Stereocaulon grande</i>
<i>Peltigera didactyla</i>	<i>Stereocaulon intermedium</i>
<i>Peltigera horizontalis</i>	<i>Stereocaulon rivulorum</i>
<i>Peltigera malacea</i>	<i>Stereocaulon saxatile</i>
<i>Peltigera neckeri</i>	<i>Stigmidium schaereri</i>
<i>Peltigera rufescens</i>	<i>Thamnolia subuliformis</i>
<i>Peltigera scabrosa</i>	<i>Thamnolia vermicularis</i>
<i>Peltigera venosa</i>	<i>Trapeliopsis flexuosa</i>
<i>Pertusaria alpina</i>	<i>Trapeliopsis granulosa</i>
<i>Pertusaria carneopallida</i>	<i>Tuckermannopsis sepincola</i>
<i>Pertusaria dactylina</i>	<i>Umbilicaria cinereorufescens</i>
<i>Pertusaria leioplaca</i>	<i>Umbilicaria cylindrica</i>
<i>Pertusaria macounii</i>	<i>Umbilicaria hyperborea</i>
<i>Pertusaria oculata</i>	<i>Umbilicaria phaea</i>
<i>Pertusaria ophthalmiza</i>	<i>Umbilicaria vellea</i>
<i>Pertusaria panyrga</i>	<i>Usnea florida</i>
<i>Pertusaria sommerfeltii</i>	<i>Verrucaria aethiobola</i>
<i>Pertusaria subdactylina</i>	<i>Vulpicida pinastri</i>
<i>Phaeophyscia hirtella</i>	<i>Xanthoria alaskana</i>
<i>Phaeophyscia orbicularis</i>	<i>Xanthoria candelaria</i>
<i>Phaeophyscia sciastra</i>	<i>Xanthoria elegans</i>
<i>Physcia adscendens</i>	<i>Xanthoria polycarpa</i>
<i>Physcia aipolia</i>	<i>Xanthoria soorediata</i>
<i>Physcia dubia</i>	<i>Xylographa parallela</i>
<i>Physcia tenella</i>	
<i>Physconia detera</i>	
<i>Physconia muscigena</i>	
<i>Physconia perisidiosa</i>	
<i>Porpidia macrocarpa</i>	
<i>Pseudephebe minuscula</i>	
<i>Pseudephebe pubescens</i>	
<i>Ramalina farinacea</i>	
<i>Ramboldia cinnabarina</i>	
<i>Ramboldia elabens</i>	
<i>Rhizocarpon eupetraeoides</i>	
<i>Rhizocarpon eupetraeum</i>	
<i>Rhizocarpon geminatum</i>	
<i>Rhizocarpon geographicum</i>	
<i>Rhizocarpon grande</i>	

Appendix D. Voucher Collections

The following table is a current list of vouchers collections that have NPS accession and catalog numbers assigned to them under permit KEFJ-2012-SCI-0016. Survey site details can be found in Appendix A. Author abbreviations are used to denote collector: JKW = James K. Walton, MH = Martin Hutten, SKT = Stacey K. Torigoe.

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
MOSSES				
<i>Amblystegium serpens</i>	Bear Cove Notch	MH 16382	KEFJ-00278	KEFJ 19068
<i>Amphidium lapponicum</i>	Coleman Bay Cliffs	MH 16345	KEFJ-00278	KEFJ 19181
<i>Amphidium lapponicum</i>	Coleman Bay Cliffs	MH 16350	KEFJ-00278	KEFJ 19065
<i>Amphidium lapponicum</i>	Coleman Bay Cliffs	MH 16363	KEFJ-00278	KEFJ 19451
<i>Amphidium lapponicum</i>	Crater Bay (northside)	MH 16024	KEFJ-00278	KEFJ 18851
<i>Andreaea blyttii</i>	McArthur WX	JKW 17824	KEFJ-00278	KEFJ 19258
<i>Andreaea nivalis</i>	Harris Peninsula Lakes	MH 16239	KEFJ-00278	KEFJ 19004
<i>Andreaea rupestris</i>	Northwest Lagoon Beach	MH 16184	KEFJ-00278	KEFJ 19369
<i>Antitrichia curtipendula</i>	Crater Bay (northside)	MH 16048	KEFJ-00278	KEFJ 18828
<i>Arctoa fulvella</i>	Harris Peninsula Lakes	MH 16242	KEFJ-00278	KEFJ 19006
<i>Aulacomnium palustre</i>	Granite Passage Peatland	MH 16185	KEFJ-00278	KEFJ 19173
<i>Aulacomnium turgidum</i>	Granite Passage Peatland	MH 16186	KEFJ-00278	KEFJ 19174
<i>Bartramia ithyphylla</i>	Coleman Bay Cliffs	MH 16351	KEFJ-00278	KEFJ 19066
<i>Bartramia pomiformis</i>	Chance Cove	JKW 17977	KEFJ-00278	KEFJ 19336
<i>Bartramia pomiformis</i>	Crater Bay (northside)	MH 16025	KEFJ-00278	KEFJ 18850
<i>Bartramia pomiformis</i>	Crater Bay (southside)	MH 16064	KEFJ-00278	KEFJ 19164
<i>Bartramiopsis lescurii</i>	Coleman Bay	MH 16325	KEFJ-00278	KEFJ 19054
<i>Bartramiopsis lescurii</i>	Crater Bay (northside)	MH 16000	KEFJ-00278	KEFJ 18819
<i>Bartramiopsis lescurii</i>	Verdant Cove	JKW 18188	KEFJ-00278	KEFJ 19216
<i>Blindia acuta</i>	Crater Bay (northside)	MH 16011	KEFJ-00278	KEFJ 18860
<i>Brachythecium albicans</i>	Northwest Lagoon Beach	MH 16162	KEFJ-00278	KEFJ 19356
<i>Brachythecium frigidum</i>	Northwest Lagoon Beach	MH 16174	KEFJ-00278	KEFJ 19458
<i>Brachythecium frigidum</i>	Northwest Lagoon Beach	MH 16176	KEFJ-00278	KEFJ 19459
<i>Bryum argenteum</i> var. <i>argenteum</i>	Coleman Bay Cliffs	MH 16346	KEFJ-00278	KEFJ 18841
<i>Buckiella undulata</i>	Crater Bay	MH 16089	KEFJ-00278	KEFJ 19095
<i>Bucklandiella affinis</i>	Harris Peninsula Lakes	MH 16226	KEFJ-00278	KEFJ 19460
<i>Bucklandiella heterosticha</i>	Chance Cove	JKW 17970	KEFJ-00278	KEFJ 19332
<i>Bucklandiella heterosticha</i>	Coleman Bay	MH 16322	KEFJ-00278	KEFJ 19340
<i>Bucklandiella heterosticha</i>	Crater Bay (southside)	MH 16077	KEFJ-00278	KEFJ 19087
<i>Bucklandiella heterosticha</i>	Crater Bay (southside)	MH 16078	KEFJ-00278	KEFJ 19088
<i>Bucklandiella heterosticha</i>	Harris Peninsula Lakes	MH 16217	KEFJ-00278	KEFJ 18991
<i>Bucklandiella heterosticha</i>	McArthur WX	JKW 17830	KEFJ-00278	KEFJ 19263
<i>Bucklandiella heterosticha</i>	Northwest Lagoon Beach	MH 16137	KEFJ-00278	KEFJ 19122
<i>Bucklandiella microcarpa</i>	McArthur WX	JKW 17831	KEFJ-00278	KEFJ 19264
<i>Calliergon cordifolium</i>	Verdant Cove	JKW 18206	KEFJ-00278	KEFJ 19230
<i>Calliergonella cuspidata</i>	Verdant Cove	JKW 18171	KEFJ-00278	KEFJ 19427
<i>Campylopus atrovirens</i>	Coleman Bay	MH 16326	KEFJ-00278	KEFJ 19339
<i>Campylopus atrovirens</i>	Granite Passage Peatland	SKT 2012-7	KEFJ-00278	KEFJ 19171
<i>Campylopus atrovirens</i>	Northwest Lagoon Beach	MH 16139	KEFJ-00278	KEFJ 19186

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
<i>Climacium dendroides</i>	Coleman Bay Cliffs	MH 16343	KEFJ-00278	KEFJ 19062
<i>Climacium dendroides</i>	Crater Bay (southside)	MH 16062	KEFJ-00278	KEFJ 18838
<i>Codriophorus acicularis</i>	Coleman Bay	MH 16318	KEFJ-00278	KEFJ 19049
<i>Codriophorus acicularis</i>	Coleman Bay Cliffs	MH 16341	KEFJ-00278	KEFJ 19061
<i>Codriophorus acicularis</i>	Coleman Bay Cliffs	MH 16344	KEFJ-00278	KEFJ 19063
<i>Codriophorus acicularis</i>	Coleman Bay Cliffs	MH 16347	KEFJ-00278	KEFJ 19182
<i>Codriophorus acicularis</i>	Harris Peninsula Lakes	MH 16257	KEFJ-00278	KEFJ 19178
<i>Codriophorus acicularis</i>	Northwest Lagoon Beach	MH 16170	KEFJ-00278	KEFJ 19362
<i>Codriophorus fascicularis</i>	McArthur WX	JKW 17826	KEFJ-00278	KEFJ 19260
<i>Conostomum tetragonum</i>	McArthur WX	JKW 17816	KEFJ-00278	KEFJ 19389
<i>Conostomum tetragonum</i>	McArthur WX	JKW 17825	KEFJ-00278	KEFJ 19259
<i>Dichodontium pellucidum</i>	Three Hole Bay	MH 16281	KEFJ-00278	KEFJ 19135
<i>Dicranella cerviculata</i>	Crater Bay (northside)	MH 16023	KEFJ-00278	KEFJ 18852
<i>Dicranella heteromallum</i>	Crater Bay (northside)	MH 16031	KEFJ-00278	KEFJ 19586
<i>Dicranella palustris</i>	Harris Peninsula Lakes	MH 16258	KEFJ-00278	KEFJ 19019
<i>Dicranella schreberiana</i>	Harris Peninsula Lakes	MH 16218	KEFJ-00278	KEFJ 18992
<i>Dicranella schreberiana</i>	Three Hole Bay	MH 16272	KEFJ-00278	KEFJ 19448
<i>Dicranodontium denudatum</i>	Crater Bay	MH 16095	KEFJ-00278	KEFJ 19099
<i>Dicranodontium denudatum</i>	Crater Bay (northside)	MH 16022	KEFJ-00278	KEFJ 18853
<i>Dicranodontium denudatum</i>	Northwest Lagoon Beach	MH 16151	KEFJ-00278	KEFJ 19349
<i>Dicranodontium denudatum</i>	Square Bay (southside)	JKW 18003	KEFJ-00278	KEFJ 18864
<i>Dicranodontium denudatum</i>	Three Hole Bay	JKW 18221	KEFJ-00278	KEFJ 19237
<i>Dicranum fuscescens</i> var. <i>fuscescens</i>	Crater Bay (northside)	MH 16033	KEFJ-00278	KEFJ 19461
<i>Dicranum majus</i> var. <i>majus</i>	Crater Bay (northside)	MH 16009	KEFJ-00278	KEFJ 18862
<i>Dicranum majus</i> var. <i>majus</i>	Crater Bay (southside)	MH 16063	KEFJ-00278	KEFJ 19163
<i>Dicranum majus</i> var. <i>majus</i>	Crater Bay (southside)	MH 16081	KEFJ-00278	KEFJ 19091
<i>Dicranum majus</i> var. <i>majus</i>	Northwest Lagoon Beach	MH 16144	KEFJ-00278	KEFJ 19342
<i>Dicranum scoparium</i>	Crater Bay (southside)	MH 16076	KEFJ-00278	KEFJ 19086
<i>Dicranum scoparium</i>	McArthur WX	JKW 17832	KEFJ-00278	KEFJ 19265
<i>Dicranum scoparium</i>	Northwest Lagoon Beach	MH 16119	KEFJ-00278	KEFJ 19114
<i>Dicranum tauricum</i>	North Arm Nuka Bay	JKW 17884	KEFJ-00278	KEFJ 19266
<i>Diphyscium foliosum</i>	Coleman Bay	MH 16324	KEFJ-00278	KEFJ 19053
<i>Distichium capillaceum</i>	Coleman Bay Cliffs	MH 16342	KEFJ-00278	KEFJ 19145
<i>Drepanocladus aduncus</i>	Northwest Lagoon Beach	MH 16156	KEFJ-00278	KEFJ 19351
<i>Fissidens osmundioides</i>	Granite Passage Peatland	SKT 2012-04	KEFJ-00278	KEFJ 19456
<i>Funaria hygrometrica</i> var. <i>hygrometrica</i>	Northwest Lagoon Beach	MH 16155	KEFJ-00278	KEFJ 19350
<i>Grimmia ramondii</i>	Coleman Bay	MH 16323	KEFJ-00278	KEFJ 19052
<i>Grimmia ramondii</i>	Crater Bay	MH 16096	KEFJ-00278	KEFJ 19100
<i>Grimmia ramondii</i>	Granite Passage Peatland	MH 16191	KEFJ-00278	KEFJ 19372
<i>Grimmia ramondii</i>	Harris Peninsula Lakes	MH 16238	KEFJ-00278	KEFJ 19003
<i>Grimmia ramondii</i>	Northwest Lagoon Beach	MH 16167	KEFJ-00278	KEFJ 19359
<i>Grimmia ramondii</i>	Three Hole Bay	JKW 18238	KEFJ-00278	KEFJ 19248
<i>Grimmia torquata</i>	Crater Bay (northside)	MH 16027	KEFJ-00278	KEFJ 18834
<i>Grimmia torquata</i>	McArthur WX	JKW 17813	KEFJ-00278	KEFJ 19387
<i>Herzogiella striatella</i>	Bear Cove Notch	MH 16418	KEFJ-00278	KEFJ 18802
<i>Heterocladium macounii</i>	Coleman Bay Cliffs	MH 16349	KEFJ-00278	KEFJ 19183
<i>Heterocladium macounii</i>	Crater Bay	MH 16099	KEFJ-00278	KEFJ 19166
<i>Heterocladium macounii</i>	Crater Bay (northside)	MH 16013	KEFJ-00278	KEFJ 19160

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
Hygrohypnum ochraceum	Coleman Bay	JKW 18249	KEFJ-00278	KEFJ 19250
Hygrohypnum ochraceum	Coleman Bay	MH 16313	KEFJ-00278	KEFJ 19044
Hygrohypnum ochraceum	Harris Peninsula Lakes	MH 16253	KEFJ-00278	KEFJ 19015
Hygrohypnum ochraceum	Northwest Lagoon Beach	MH 16171	KEFJ-00278	KEFJ 19363
Hylocomiastrum umbratum	Bear Cove Notch	JKW 18280	KEFJ-00278	KEFJ 18916
Hylocomium splendens	Crater Bay (southside)	MH 16059	KEFJ-00278	KEFJ 18833
Hypnum circinale	Crater Bay (northside)	MH 16036	KEFJ-00278	KEFJ 18844
Hypnum circinale	Crater Bay (southside)	MH 16058	KEFJ-00278	KEFJ 18821
Hypnum dieckii	Crater Bay (northside)	MH 16012	KEFJ-00278	KEFJ 18859
Hypnum dieckii	Crater Bay (southside)	MH 16065	KEFJ-00278	KEFJ 18808
Hypnum lindbergii	North Arm Nuka Bay	JKW 17872	KEFJ-00278	KEFJ 19462
Hypnum plicatulum	Crater Bay (southside)	MH 16072	KEFJ-00278	KEFJ 19083
Hypnum plicatulum	Northwest Lagoon Beach	MH 16141	KEFJ-00278	KEFJ 19341
Hypnum subimponens	Northwest Lagoon Beach	MH 16163	KEFJ-00278	KEFJ 19357
Hypnum subimponens	Northwest Lagoon Beach	MH 16169	KEFJ-00278	KEFJ 19361
Isopterygiopsis pulchella	Square Bay (southside)	JKW 18025	KEFJ-00278	KEFJ 18943
Isopterygiopsis pulchella	Square Bay (southside)	JKW 18026	KEFJ-00278	KEFJ 18944
Iwatsukiella leucotricha	Crater Bay (northside)	MH 16041	KEFJ-00278	KEFJ 18832
Iwatsukiella leucotricha	James Lagoon	JKW 17961	KEFJ-00278	KEFJ 19327
Iwatsukiella leucotricha	North Arm Nuka Bay	JKW 17886	KEFJ-00278	KEFJ 19268
Kinbergia oregana	Moonlight Bay	JKW 17930	KEFJ-00278	KEFJ 19302
Kinbergia oregana	Square Bay (southside)	JKW 18012	KEFJ-00278	KEFJ 18865
Kinbergia praelonga	Verdant Cove	JKW 18205	KEFJ-00278	KEFJ 18804
Mnium thomsonii	Bear Cove Notch	MH 16428	KEFJ-00278	KEFJ 18933
Mnium thomsonii	Crater Bay (northside)	MH 16053	KEFJ-00278	KEFJ 18806
Mnium thomsonii	Square Bay (southside)	JKW 18027	KEFJ-00278	KEFJ 18945
Mnium thomsonii	Three Hole Bay	JKW 18237	KEFJ-00278	KEFJ 19247
Myuroclada maximowiczii	Coleman Bay Cliffs	MH 16348	KEFJ-00278	KEFJ 19064
Niphotrichum ericoides	Northwest Lagoon Beach	MH 16120	KEFJ-00278	KEFJ 19115
Oedipodium griffithianum	Three Hole Bay	MH 16293	KEFJ-00278	KEFJ 19025
Oligotrichum hercynicum	Coleman Bay	MH 16314	KEFJ-00278	KEFJ 19045
Oligotrichum hercynicum	Coleman Bay	MH 16315	KEFJ-00278	KEFJ 19046
Oligotrichum hercynicum	Coleman Bay	MH 16316	KEFJ-00278	KEFJ 19047
Oligotrichum parallelum	Crater Bay (northside)	MH 16042	KEFJ-00278	KEFJ 18803
Oligotrichum parallelum	Crater Bay (southside)	MH 16071	KEFJ-00278	KEFJ 19082
Oligotrichum parallelum	Granite Passage Peatland	MH 16194	KEFJ-00278	KEFJ 19374
Oligotrichum parallelum	Verdant Cove	JKW 18199	KEFJ-00278	KEFJ 19225
Oncophorus virens	Bear Cove Notch	MH 16400	KEFJ-00278	KEFJ 18919
Oncophorus wahlenbergii	Coleman Bay	MH 16312	KEFJ-00278	KEFJ 19043
Paraleucobryum enerve	Three Hole Bay	MH 16295	KEFJ-00278	KEFJ 19027
Philonotis fontana var. fontana	Coleman Bay	MH 16319	KEFJ-00278	KEFJ 19050
Philonotis fontana var. fontana	Coleman Bay Cliffs	MH 16339	KEFJ-00278	KEFJ 19059
Philonotis fontana var. fontana	Harris Peninsula Lakes	MH 16256	KEFJ-00278	KEFJ 19018
Philonotis fontana var. fontana	Northwest Lagoon Beach	MH 16173	KEFJ-00278	KEFJ 19365
Plagiomnium medium	Bear Cove Notch	MH 16417	KEFJ-00278	KEFJ 18924
Plagiothecium laetum	Coleman Bay	MH 16317	KEFJ-00278	KEFJ 19048
Plagiothecium piliferum	Crater Bay (northside)	MH 16029	KEFJ-00278	KEFJ 18848
Plagiothecium piliferum	Northwest Lagoon Beach	MH 16113	KEFJ-00278	KEFJ 19112

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
<i>Plagiothecium piliferum</i>	Northwest Lagoon Beach	MH 16138	KEFJ-00278	KEFJ 19123
<i>Pleuroziopsis ruthenica</i>	Bear Cove Notch	MH 16410	KEFJ-00278	KEFJ 18837
<i>Pleurozium schreberi</i>	Crater Bay (southside)	MH 16080	KEFJ-00278	KEFJ 19090
<i>Pogonatum contortum</i>	Crater Bay (northside)	MH 16047	KEFJ-00278	KEFJ 18829
<i>Pogonatum contortum</i>	North Arm Nuka Bay	JKW 17896	KEFJ-00278	KEFJ 19277
<i>Pogonatum dentatum</i>	Crater Bay (northside)	MH 16030	KEFJ-00278	KEFJ 18863
<i>Pogonatum dentatum</i>	McArthur WX	JKW 17829	KEFJ-00278	KEFJ 19262
<i>Pohlia cruda</i>	Crater Bay	MH 16098	KEFJ-00278	KEFJ 19102
<i>Pohlia crudoides</i>	Crater Bay (southside)	MH 16068	KEFJ-00278	KEFJ 19079
<i>Pohlia nutans</i>	Crater Bay (southside)	JKW 18063	KEFJ-00278	KEFJ 18964
<i>Pohlia prolifera</i>	Northwest Lagoon Beach	MH 16145	KEFJ-00278	KEFJ 19343
<i>Pohlia wahlenbergii</i>	Coleman Bay	MH 16327	KEFJ-00278	KEFJ 19055
<i>Pohlia wahlenbergii</i>	Coleman Bay	MH 16330	KEFJ-00278	KEFJ 19056
<i>Pohlia wahlenbergii</i>	Coleman Bay	MH 16331	KEFJ-00278	KEFJ 19057
<i>Polytrichastrum alpinum</i>	Coleman Bay	JKW 18251	KEFJ-00278	KEFJ 19252
<i>Polytrichastrum alpinum</i>	Coleman Bay	MH 16311	KEFJ-00278	KEFJ 19042
<i>Polytrichastrum alpinum</i>	Crater Bay (southside)	MH 16060	KEFJ-00278	KEFJ 18811
<i>Polytrichastrum alpinum</i>	Crater Bay (southside)	MH 16082	KEFJ-00278	KEFJ 19092
<i>Polytrichastrum alpinum</i>	Verdant Cove	JKW 18204	KEFJ-00278	KEFJ 19229
<i>Polytrichastrum formosum</i>	Granite Passage Peatland	MH 16210	KEFJ-00278	KEFJ 19380
<i>Polytrichastrum papillatum</i>	Coleman Bay	MH 16320	KEFJ-00278	KEFJ 19338
<i>Polytrichum juniperinum</i>	Coleman Bay	MH 16321	KEFJ-00278	KEFJ 19051
<i>Polytrichum piliferum</i>	Northwest Lagoon Beach	MH 16114	KEFJ-00278	KEFJ 19113
<i>Pseudoleskea baileyi</i>	Coleman Bay	MH 16310	KEFJ-00278	KEFJ 19041
<i>Pseudoleskea stenophylla</i>	Coleman Bay	MH 18240	KEFJ-00278	KEFJ 19463
<i>Pseudotaxiphyllum elegans</i>	Crater Bay (northside)	JKW 16032	KEFJ-00278	KEFJ 19146
<i>Ptilium crista-castrensis</i>	Crater Bay (southside)	MH 16066	KEFJ-00278	KEFJ 18809
<i>Ptychostomum pseudotriquetrum</i>	Coleman Bay Cliffs	MH 16352	KEFJ-00278	KEFJ 19184
<i>Ptychostomum weigeli</i>	Coleman Bay	MH 16338	KEFJ-00278	KEFJ 19464
<i>Racomitrium lanuginosum</i>	Crater Bay (southside)	JKW 18078	KEFJ-00278	KEFJ 18969
<i>Rhabdoweisia crispata</i>	Crater Bay (northside)	MH 16133	KEFJ-00278	KEFJ 19253
<i>Rhizomnium glabrescens</i>	Crater Bay (northside)	MH 16051	KEFJ-00278	KEFJ 18826
<i>Rhizomnium glabrescens</i>	McArthur WX	JKW 17821	KEFJ-00278	KEFJ 19255
<i>Rhizomnium glabrescens</i>	Moonlight Bay	JKW 17929	KEFJ-00278	KEFJ 19301
<i>Rhizomnium glabrescens</i>	Three Hole Bay	JKW 18233	KEFJ-00278	KEFJ 19243
<i>Rhizomnium magnifolium</i>	Harris Peninsula Lakes	MH 16251	KEFJ-00278	KEFJ 19013
<i>Rhizomnium magnifolium</i>	Northwest Lagoon Beach	MH 16135	KEFJ-00278	KEFJ 19121
<i>Rhizomnium nudum</i>	Coleman Bay	MH 16336	KEFJ-00278	KEFJ 19058
<i>Rhizomnium nudum</i>	Harris Peninsula Lakes	MH 16222	KEFJ-00278	KEFJ 18995
<i>Rhizomnium nudum</i>	Harris Peninsula Lakes	MH 16240	KEFJ-00278	KEFJ 19005
<i>Rhizomnium pseudopunctatum</i>	North Arm Nuka Bay	JKW 17897	KEFJ-00278	KEFJ 19278
<i>Rhytidiadelphus loreus</i>	Crater Bay (southside)	MH 16056	KEFJ-00278	KEFJ 18823
<i>Rhytidiadelphus loreus</i>	McArthur WX	JKW 17822	KEFJ-00278	KEFJ 19256
<i>Rhytidiadelphus squarrosus</i>	Northwest Lagoon Beach	MH 16106	KEFJ-00278	KEFJ 19108
<i>Rhytidiadelphus squarrosus</i>	Square Bay (southside)	JKW 18019	KEFJ-00278	KEFJ 18870
<i>Rhytidiadelphus triquetrus</i>	Northwest Lagoon Beach	MH 16110	KEFJ-00278	KEFJ 19110
<i>Rosulabryum capillare</i>	Northwest Lagoon Beach	MH 16165	KEFJ-00278	KEFJ 19465
<i>Sarmenthyphnum sarmentosum</i>	Coleman Bay Cliffs	MH 16365	KEFJ-00278	KEFJ 19452

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
Sarmenthyphnum sarmentosum	Granite Passage Peatland	MH 16201	KEFJ-00278	KEFJ 18987
Sarmenthyphnum sarmentosum	Harris Peninsula Lakes	MH 16254	KEFJ-00278	KEFJ 19016
Schistidium maritimum	Crater Bay	MH 16091	KEFJ-00278	KEFJ 19096
Schistidium maritimum	North Arm Nuka Bay	JKW 17835	KEFJ-00278	KEFJ 19147
Scouleria aquatica	North Arm Nuka Bay	JKW 17898	KEFJ-00278	KEFJ 19188
Sphagnum compactum	Crater Bay (southside)	JKW 18054	KEFJ-00278	KEFJ 18955
Sphagnum compactum	Crater Bay (southside)	JKW 18065	KEFJ-00278	KEFJ 18966
Sphagnum compactum	Granite Passage Peatland	JKW 18164	KEFJ-00278	KEFJ 19202
Sphagnum compactum	Granite Passage Peatland	JKW 18166	KEFJ-00278	KEFJ 19204
Sphagnum compactum	Granite Passage Peatland	JKW 18167	KEFJ-00278	KEFJ 19205
Sphagnum girgensohnii	Crater Bay (northside)	JKW 18045	KEFJ-00278	KEFJ 19159
Sphagnum linbergii	Crater Bay (northside)	JKW 18029	KEFJ-00278	KEFJ 19466
Sphagnum quinquefarium	Crater Bay (northside)	JKW 18050	KEFJ-00278	KEFJ 18953
Sphagnum russowii	North Arm Nuka Bay	JKW17873	KEFJ-00278	KEFJ 19467
Sphagnum squarrosum	Crater Bay (northside)	JKW 18046	KEFJ-00278	KEFJ 18949
Sphagnum tenellum	Granite Passage Peatland	JKW 18163	KEFJ-00278	KEFJ 19201
Sphagnum tenellum	Granite Passage Peatland	JKW 18165	KEFJ-00278	KEFJ 19203
Sphagnum tenellum	Granite Passage Peatland	JKW 18168	KEFJ-00278	KEFJ 19206
Sphagnum tenellum	Granite Passage Peatland	JKW 18169	KEFJ-00278	KEFJ 19207
Straminergon stramineum	Crater Bay (northside)	MH 16055	KEFJ-00278	KEFJ 18824
Takakia lepidozoides	Three Hole Bay	MH 16282	KEFJ-00278	KEFJ 19136
Takakia lepidozoides	Three Hole Bay	MH 16292	KEFJ-00278	KEFJ 18842
Tayloria serrata	Bear Cove Notch	MH 16409	KEFJ-00278	KEFJ 18922
Tayloria serrata	Bear Cove Notch	MH 16420	KEFJ-00278	KEFJ 18926
Tayloria serrata	Bear Cove Notch	MH 16421	KEFJ-00278	KEFJ 18927
Tayloria serrata	Crater Bay (southside)	MH 16067	KEFJ-00278	KEFJ 18810
Tetraphis geniculata	Bear Cove Notch	MH 16419	KEFJ-00278	KEFJ 18925
Tetraphis geniculata	Verdant Cove	JKW 18189	KEFJ-00278	KEFJ 18840
Tortella tortuosa	Crater Bay (northside)	MH 16015	KEFJ-00278	KEFJ 18857
Tortella tortuosa	Granite Passage Peatland	MH 16180	KEFJ-00278	KEFJ 19367
Ulotia megalospora	Crater Bay (northside)	MH 16039	KEFJ-00278	KEFJ 18849
Ulotia obtusiuscula	Crater Bay (northside)	MH 16038	KEFJ-00278	KEFJ 18843
Ulotia obtusiuscula	Northwest Lagoon Beach	MH 16109	KEFJ-00278	KEFJ 19109
Ulotia phyllantha	Crater Bay	MH 16105	KEFJ-00278	KEFJ 19107
Warnstorfia fluitans	Harris Peninsula Lakes	MH 16230	KEFJ-00278	KEFJ 19000
LIVERWORTS				
Anastrepta orcadensis	Bear Cove Notch	MH 16386	KEFJ-00278	KEFJ 19071
Anastrepta orcadensis	Bear Cove Notch	MH 16387	KEFJ-00278	KEFJ 19072
Anastrepta orcadensis	Bear Cove Notch	MH 16388	KEFJ-00278	KEFJ 19073
Anastrepta orcadensis	Bear Cove Notch	MH 16412	KEFJ-00278	KEFJ 18923
Anastrepta orcadensis	Crater Bay	MH 16102	KEFJ-00278	KEFJ 19104
Anastrepta orcadensis	Crater Bay (northside)	MH 16020	KEFJ-00278	KEFJ 19162
Anastrepta orcadensis	Crater Bay (northside)	MH 16035	KEFJ-00278	KEFJ 18845
Anastrepta orcadensis	Granite Passage Peatland	MH 16190	KEFJ-00278	KEFJ 19371
Anastrophyllum assimile	Crater Bay (southside)	MH 16075	KEFJ-00278	KEFJ 19165
Anastrophyllum assimile	McArthur WX	JKW 17810	KEFJ-00278	KEFJ 19384
Anastrophyllum assimile	McArthur WX	JKW 17815	KEFJ-00278	KEFJ 19388

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
Anastrophyllum assimile	Three Hole Bay	MH 16296	KEFJ-00278	KEFJ 19028
Anastrophyllum minutum	Crater Bay (northside)	MH 16019	KEFJ-00278	KEFJ 19161
Anastrophyllum minutum	Granite Passage Peatland	MH 16196	KEFJ-00278	KEFJ 19176
Anastrophyllum minutum	Northwest Lagoon Beach	MH 16154	KEFJ-00278	KEFJ 19172
Anastrophyllum minutum var. minutum	Bear Cove Notch	MH 16389	KEFJ-00278	KEFJ 19074
Anastrophyllum minutum var. minutum	Bear Cove Notch	MH 16425	KEFJ-00278	KEFJ 18930
Anastrophyllum minutum var. minutum	Three Hole Bay	MH 16287	KEFJ-00278	KEFJ 19141
Anastrophyllum minutum var. minutum	Three Hole Bay	MH 16288	KEFJ-00278	KEFJ 19142
Anastrophyllum minutum var. minutum	Three Hole Bay	MH 16289	KEFJ-00278	KEFJ 19143
Anastrophyllum spheonoloboides	Northwest Lagoon Beach	MH 16128	KEFJ-00278	KEFJ 19170
Anthelia julacea	Crater Bay (southside)	MH 16079	KEFJ-00278	KEFJ 19089
Anthelia julacea	Granite Passage Peatland	MH 16195	KEFJ-00278	KEFJ 19375
Anthelia julacea	Harris Peninsula Lakes	MH 16216	KEFJ-00278	KEFJ 18990
Anthelia julacea	Three Hole Bay	MH 16297	KEFJ-00278	KEFJ 19029
Anthelia juratzkana	McArthur WX	JKW 17809	KEFJ-00278	KEFJ 19468
Barbilophozia floerkei	Harris Peninsula Lakes	MH 16260	KEFJ-00278	KEFJ 19021
Barbilophozia floerkei	Harris Peninsula Lakes	MH 16261	KEFJ-00278	KEFJ 19179
Barbilophozia kunzeana	Three Hole Bay	MH 16266	KEFJ-00278	KEFJ 19125
Barbilophozia lycopodioides	Crater Bay (southside)	MH 16074	KEFJ-00278	KEFJ 19085
Barbilophozia lycopodioides	Harris Peninsula Lakes	MH 16219	KEFJ-00278	KEFJ 18993
Bazzania tricrenata	Bear Cove Notch	MH 16391	KEFJ-00278	KEFJ 19076
Bazzania tricrenata	Bear Cove Notch	MH 16392	KEFJ-00278	KEFJ 19077
Bazzania tricrenata	Crater Bay (northside)	MH 16005	KEFJ-00278	KEFJ 18817
Bazzania tricrenata	Three Hole Bay	MH 16273	KEFJ-00278	KEFJ 19129
Bazzania tricrenata	Three Hole Bay	MH 16276b	KEFJ-00278	KEFJ 19449
Bazzania tricrenata	Three Hole Bay	MH 16277	KEFJ-00278	KEFJ 19133
Bazzania tricrenata	Three Hole Bay	MH 16290	KEFJ-00278	KEFJ 19144
Blepharostoma trichophyllum	Crater Bay (northside)	MH 16002	KEFJ-00278	KEFJ 18814
Calycularia laxa	Crater Bay	MH 16101	KEFJ-00278	KEFJ 19103
Calycularia laxa	Northwest Lagoon Beach	MH 16146	KEFJ-00278	KEFJ 19344
Calycularia laxa	Northwest Lagoon Beach	MH 16147	KEFJ-00278	KEFJ 19345
Calycularia laxa	Northwest Lagoon Beach	MH 16148	KEFJ-00278	KEFJ 19346
Calycularia laxa	Northwest Lagoon Beach	MH 16149	KEFJ-00278	KEFJ 19347
Calypogeia integristipula	Harris Peninsula Lakes	MH 16249	KEFJ-00278	KEFJ 19011
Calypogeia integristipula	Three Hole Bay	MH 16279	KEFJ-00278	KEFJ 19134
Calypogeia muelleriana	Bear Cove Notch	MH 16390	KEFJ-00278	KEFJ 19075
Calypogeia muelleriana	Crater Bay	MH 16097	KEFJ-00278	KEFJ 19101
Calypogeia neesiana	Bear Cove Notch	MH 16427	KEFJ-00278	KEFJ 18932
Calypogeia neesiana	Crater Bay (northside)	JKW 18036	KEFJ-00278	KEFJ 18947
Calypogeia neesiana	Crater Bay (northside)	MH 16018	KEFJ-00278	KEFJ 18805
Cephalozia bicuspidata	McArthur WX	JKW 17808	KEFJ-00278	KEFJ 19383
Cephalozia bicuspidata subsp. ambigua	Harris Peninsula Lakes	MH 16225	KEFJ-00278	KEFJ 19177
Cephalozia bicuspidata subsp. bicuspidata	Northwest Lagoon Beach	MH 16129	KEFJ-00278	KEFJ 19118
Cephalozia bicuspidata subsp. bicuspidata	Three Hole Bay	MH 16298	KEFJ-00278	KEFJ 19030
Cephalozia lunulifolia	Bear Cove Notch	MH 16380	KEFJ-00278	KEFJ 19185
Cephalozia lunulifolia	Crater Bay (southside)	MH 16088	KEFJ-00278	KEFJ 19094
Cephalozia divaricata subsp. divaricata	Bear Cove Notch	MH 16401	KEFJ-00278	KEFJ 18920
Conocephalum salebrosum	Bear Cove	JKW 18282a	KEFJ-00278	KEFJ 19445

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
Conocephalum salebrosum	Bear Cove	JKW 18282b	KEFJ-00278	KEFJ 19446
Diplophyllum albicans	Crater Bay (northside)	MH 16010	KEFJ-00278	KEFJ 18822
Diplophyllum obtusifolium	Crater Bay	MH 16104	KEFJ-00278	KEFJ 19106
Diplophyllum plicatum	Crater Bay (northside)	MH 16049	KEFJ-00278	KEFJ 18827
Diplophyllum taxifolium	Chance Cove	JKW 17967	KEFJ-00278	KEFJ 19469
Douinia ovata	Bear Cove Notch	MH 16384	KEFJ-00278	KEFJ 19069
Douinia ovata	Northwest Lagoon Beach	MH 16112	KEFJ-00278	KEFJ 19111
Frullania nisquallensis	Bear Cove Notch	MH 16385	KEFJ-00278	KEFJ 19070
Frullania nisquallensis	Bear Cove Notch	MH 16402	KEFJ-00278	KEFJ 18921
Gymnocolea inflata	Granite Passage Peatland	MH 16187	KEFJ-00278	KEFJ 19370
Gymnocolea inflata	Harris Peninsula Lakes	MH 16214	KEFJ-00278	KEFJ 18989
Gymnocolea inflata	Three Hole Bay	MH 16267	KEFJ-00278	KEFJ 19126
Gymnocolea inflata	Three Hole Bay	MH 16303	KEFJ-00278	KEFJ 19035
Gymnomitrium concinnatum	Bear Cove Notch	MH 16426	KEFJ-00278	KEFJ 18931
Gymnomitrium concinnatum	Northwest Lagoon Beach	MH 16140	KEFJ-00278	KEFJ 19124
Gymnomitrium concinnatum	Three Hole Bay	MH 16280	KEFJ-00278	KEFJ 19180
Gymnomitrium obtusum	Crater Bay (northside)	MH 16021	KEFJ-00278	KEFJ 18854
Gymnomitrium obtusum	Granite Passage Peatland	MH 16179	KEFJ-00278	KEFJ 19366
Gymnomitrium obtusum	Three Hole Bay	MH 16275	KEFJ-00278	KEFJ 19131
Harpanthus flotovianus	Harris Peninsula Lakes	MH 16221	KEFJ-00278	KEFJ 18994
Herbertus aduncus	Crater Bay (northside)	MH 16014	KEFJ-00278	KEFJ 18858
Herbertus aduncus	Northwest Lagoon Beach	MH 16150	KEFJ-00278	KEFJ 19348
Herbertus dicranus	Crater Bay	MH 16094	KEFJ-00278	KEFJ 19098
Herbertus dicranus	Granite Passage Peatland	MH 16211a	KEFJ-00278	KEFJ 19381
Herbertus dicranus	Granite Passage Peatland	MH 16211b	KEFJ-00278	KEFJ 19455
Herbertus dicranus	Northwest Lagoon Beach	MH 16124	KEFJ-00278	KEFJ 19117
Herbertus dicranus	Three Hole Bay	MH 16299	KEFJ-00278	KEFJ 19031
Herbertus dicranus	Three Hole Bay	MH 16300	KEFJ-00278	KEFJ 19032
Jungermannia exsertifolia	Coleman Bay	MH 16308	KEFJ-00278	KEFJ 19039
Kurzia trichoclados	Bear Cove Notch	MH 16381	KEFJ-00278	KEFJ 19067
Kurzia trichoclados	Crater Bay (southside)	MH 16069	KEFJ-00278	KEFJ 19080
Kurzia trichoclados	Three Hole Bay	MH 16284	KEFJ-00278	KEFJ 19138
Kurzia trichoclados	Three Hole Bay	MH 16285	KEFJ-00278	KEFJ 19139
Lepidozia filamentosa	Bear Cove Notch	MH 16393	KEFJ-00278	KEFJ 19078
Lepidozia filamentosa	Bear Cove Notch	MH 16422	KEFJ-00278	KEFJ 18928
Lepidozia filamentosa	Crater Bay	MH 16103	KEFJ-00278	KEFJ 19105
Lepidozia reptans	Crater Bay (northside)	MH 16008	KEFJ-00278	KEFJ 18847
Lepidozia reptans	East Arm (westside)	JKW 17915	KEFJ-00278	KEFJ 19289
Lophocolea bidentata	Granite Passage Peatland	MH 16189	KEFJ-00278	KEFJ 19175
Lophozia guttulata	Bear Cove Notch	MH 16394	KEFJ-00278	KEFJ 18917
Lophozia guttulata	Harris Peninsula Lakes	MH 16247	KEFJ-00278	KEFJ 19009
Lophozia sudetica	Harris Peninsula Lakes	MH 16228	KEFJ-00278	KEFJ 18999
Lophozia sudetica	Northwest Lagoon Beach	MH 16122	KEFJ-00278	KEFJ 19116
Lophozia sudetica	Three Hole Bay	MH 16269	KEFJ-00278	KEFJ 19128
Lophozia ventricosa	Northwest Lagoon Beach	MH 16117	KEFJ-00278	KEFJ 19167
Lophozia ventricosa	Northwest Lagoon Beach	MH 16118	KEFJ-00278	KEFJ 19168
Lophozia ventricosa	Northwest Lagoon Beach	MH 16126	KEFJ-00278	KEFJ 19169
Lophozia ventricosa var. longiflora	Bear Cove Notch	MH 16399	KEFJ-00278	KEFJ 18918

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
<i>Lophozia wenzelii</i> var. <i>wenzelii</i>	Three Hole Bay	MH 16268	KEFJ-00278	KEFJ 19127
<i>Marsupella emarginata</i>	Coleman Bay	MH 16309	KEFJ-00278	KEFJ 19040
<i>Marsupella emarginata</i>	Granite Passage Peatland	SKT 2012-01	KEFJ-00278	KEFJ 19454
<i>Marsupella emarginata</i>	McArthur WX	JKW 17811	KEFJ-00278	KEFJ 19385
<i>Marsupella emarginata</i>	McArthur WX	JKW 17812	KEFJ-00278	KEFJ 19386
<i>Marsupella emarginata</i> var. <i>emarginata</i>	Crater Bay (northside)	MH 16034	KEFJ-00278	KEFJ 18846
<i>Marsupella emarginata</i> var. <i>emarginata</i>	Granite Passage Peatland	MH 16197	KEFJ-00278	KEFJ 19376
<i>Marsupella emarginata</i> var. <i>emarginata</i>	Three Hole Bay	MH 16305	KEFJ-00278	KEFJ 19036
<i>Marsupella sphacelata</i>	Three Hole Bay	MH 16263	KEFJ-00278	KEFJ 19022
<i>Metzgeria conjugata</i>	Northwest Lagoon Beach	JKW 18118	KEFJ-00278	KEFJ 18883
<i>Moerckia blyttii</i>	Harris Peninsula Lakes	MH 16223	KEFJ-00278	KEFJ 18996
<i>Moerckia blyttii</i>	Harris Peninsula Lakes	MH 16259	KEFJ-00278	KEFJ 19020
<i>Mylia taylorii</i>	Crater Bay (northside)	MH 16007	KEFJ-00278	KEFJ 18820
<i>Mylia taylorii</i>	Granite Passage Peatland	MH 16192	KEFJ-00278	KEFJ 19373
<i>Mylia taylorii</i>	North Arm Nuka Bay	JKW 17885	KEFJ-00278	KEFJ 19267
<i>Mylia taylorii</i>	Three Hole Bay	MH 16294	KEFJ-00278	KEFJ 19026
<i>Mylia taylorii</i>	Three Hole Bay	MH 16301	KEFJ-00278	KEFJ 19033
<i>Nardia scalaris</i>	Coleman Bay	MH 16307	KEFJ-00278	KEFJ 19038
<i>Nardia scalaris</i>	Crater Bay (northside)	MH 16043	KEFJ-00278	KEFJ 18831
<i>Nardia scalaris</i>	Crater Bay (southside)	MH 16070	KEFJ-00278	KEFJ 19081
<i>Nardia scalaris</i>	Harris Peninsula Lakes	MH 16212	KEFJ-00278	KEFJ 18988
<i>Nardia scalaris</i>	Harris Peninsula Lakes	MH 16227	KEFJ-00278	KEFJ 18998
<i>Nardia scalaris</i>	Three Hole Bay	MH 16302	KEFJ-00278	KEFJ 19034
<i>Odontoschisma elongatum</i>	Three Hole Bay	MH 16274	KEFJ-00278	KEFJ 19130
<i>Odontoschisma elongatum</i>	Three Hole Bay	MH 16286	KEFJ-00278	KEFJ 19140
<i>Pellia endiviifolia</i>	McArthur WX	JKW 17805	KEFJ-00278	KEFJ 19382
<i>Pellia epiphylla</i>	Northwest Lagoon Beach	MH 16131	KEFJ-00278	KEFJ 19119
<i>Pellia neesiana</i>	Bear Cove Notch	MH 16424	KEFJ-00278	KEFJ 18929
<i>Plagiochila porelloides</i>	Bear Cove Notch	MH 16411	KEFJ-00278	KEFJ 18836
<i>Plagiochila porelloides</i>	Bear Cove Notch	MH 16423	KEFJ-00278	KEFJ 18861
<i>Plagiochila porelloides</i>	Chance Cove	JKW 17966	KEFJ-00278	KEFJ 19400
<i>Plagiochila porelloides</i>	Granite Passage Peatland	MH 16188	KEFJ-00278	KEFJ 19447
<i>Pleurocladula albescens</i>	Harris Peninsula Lakes	MH 16255	KEFJ-00278	KEFJ 19017
<i>Ptilidium californicum</i>	Crater Bay (northside)	MH 16017	KEFJ-00278	KEFJ 18855
<i>Ptilidium californicum</i>	Crater Bay (southside)	MH 16085	KEFJ-00278	KEFJ 19093
<i>Ptilidium ciliare</i>	Crater Bay (southside)	MH 16073	KEFJ-00278	KEFJ 19084
<i>Ptilidium ciliare</i>	McArthur WX	JKW 17823	KEFJ-00278	KEFJ 19257
<i>Radula bolanderi</i>	Crater Bay (northside)	MH 16001	KEFJ-00278	KEFJ 18812
<i>Riccardia latifrons</i>	Crater Bay (northside)	JKW 18035	KEFJ-00278	KEFJ 18946
<i>Riccardia latifrons</i>	Crater Bay (northside)	MH 16045	KEFJ-00278	KEFJ 18830
<i>Riccardia multifida</i>	Northwest Lagoon Beach	MH 16132	KEFJ-00278	KEFJ 19120
<i>Scapania bolanderi</i>	Crater Bay (northside)	MH 16006	KEFJ-00278	KEFJ 18818
<i>Scapania bolanderi</i>	North Arm Nuka Bay	JKW 17871	KEFJ-00278	KEFJ 19392
<i>Scapania curta</i>	Three Hole Bay	MH 16306	KEFJ-00278	KEFJ 19037
<i>Scapania cuspiduligera</i>	Harris Peninsula Lakes	MH 16252	KEFJ-00278	KEFJ 19014
<i>Scapania paludosa</i>	Coleman Bay Cliffs	MH 16340	KEFJ-00278	KEFJ 19060
<i>Scapania paludosa</i>	Three Hole Bay	MH 16265	KEFJ-00278	KEFJ 19023
<i>Scapania paludosa</i>	Three Hole Bay	MH 16276a	KEFJ-00278	KEFJ 19132

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
Scapania uliginosa	Crater Bay (northside)	MH 16054	KEFJ-00278	KEFJ 18825
Scapania uliginosa	Harris Peninsula Lakes	MH 16250	KEFJ-00278	KEFJ 19012
Scapania umbrosa	Crater Bay (southside)	MH 16057	KEFJ-00278	KEFJ 18813
Scapania umbrosa	Harris Peninsula Lakes	MH 16246	KEFJ-00278	KEFJ 19008
Scapania undulata var. undulata	Crater Bay (northside)	MH 16003	KEFJ-00278	KEFJ 18815
Scapania undulata var. undulata	Verdant Cove	JKW 18181	KEFJ-00278	KEFJ 19209
Schistochilopsis incisa var. incisa	Crater Bay (northside)	MH 16016	KEFJ-00278	KEFJ 18856
Schistochilopsis incisa var. incisa	Harris Peninsula Lakes	MH 16224	KEFJ-00278	KEFJ 18997
Sphenolobopsis pearsonii	Granite Passage Peatland	MH 16181	KEFJ-00278	KEFJ 19368
Tetralophozia setiformis	McArthur WX	JKW 17828	KEFJ-00278	KEFJ 19208
Tritomaria quinquedentata	Crater Bay (southside)	MH 16061	KEFJ-00278	KEFJ 18807
LICHENS				
Alectoria sarmentosa subsp. sarmentosa	Bear Cove	JKW 18266	KEFJ-00278	KEFJ 18902
Alectoria sarmentosa subsp. sarmentosa	Chance Cove	JKW 17978	KEFJ-00278	KEFJ 19337
Alectoria sarmentosa subsp. sarmentosa	Crater Bay	JKW 18099	KEFJ-00278	KEFJ 18982
Alectoria sarmentosa subsp. sarmentosa	East Arm (westside)	JKW 17923	KEFJ-00278	KEFJ 19297
Alectoria sarmentosa subsp. sarmentosa	Fire Cove	JKW 18155	KEFJ-00278	KEFJ 19195
Alectoria sarmentosa subsp. sarmentosa	James Lagoon	JKW 17964	KEFJ-00278	KEFJ 19330
Alectoria sarmentosa subsp. sarmentosa	North Arm Nuka Bay	JKW 17894	KEFJ-00278	KEFJ 19276
Alectoria sarmentosa subsp. sarmentosa	Square Bay (northside)	JKW 17996	KEFJ-00278	KEFJ 19155
Alectoria sarmentosa subsp. sarmentosa	Square Bay (southside)	JKW 18022	KEFJ-00278	KEFJ 18872
Alectoria sarmentosa subsp. sarmentosa	Three Hole Bay	JKW 18222	KEFJ-00278	KEFJ 19238
Alectoria sarmentosa subsp. sarmentosa	Verdant Cove	JKW 18191	KEFJ-00278	KEFJ 19218
Alectoria sarmentosa subsp. sarmentosa	Verdant Cove	JKW 18193	KEFJ-00278	KEFJ 19220
Alectoria sarmentosa subsp. sarmentosa	Verdant Cove	JKW 18200	KEFJ-00278	KEFJ 19226
Baeomyces rufus	Northwest Lagoon Beach	JKW 18126	KEFJ-00278	KEFJ 18891
Bryoria bicolor	Bear Cove Notch	MH 16403	KEFJ-00278	KEFJ 19453
Bryoria bicolor	Fire Cove	JKW 18141	KEFJ-00278	KEFJ 19422
Bryoria bicolor	Three Hole Bay	JKW 18210	KEFJ-00278	KEFJ 19433
Bryoria bicolor	Verdant Cove	JKW 18177	KEFJ-00278	KEFJ 19429
Bryoria cervinula	Crater Bay (northside)	JKW 18032	KEFJ-00278	KEFJ 19470
Bryoria cervinula	Bear Cove	JKW 18261	KEFJ-00278	KEFJ 19471
Bryoria trichodes subsp. americana	Bear Cove Notch	MH 16404	KEFJ-00278	KEFJ 19472
Bryoria trichodes subsp. americana	North Arm Nuka Bay	JKW 17875	KEFJ-00278	KEFJ 19473
Bryoria trichodes subsp. americana	East Arm (westside)	JKW 17912	KEFJ-00278	KEFJ 19474
Bryoria trichodes subsp. americana	East Arm (westside)	JKW 17913	KEFJ-00278	KEFJ 19475
Bryoria trichodes subsp. americana	Moonlight Bay	JKW 17927	KEFJ-00278	KEFJ 19476
Bryoria trichodes subsp. americana	Square Bay (southside)	JKW 18007	KEFJ-00278	KEFJ 19477
Bryoria trichodes subsp. americana	Crater Bay	JKW 18085	KEFJ-00278	KEFJ 19478
Bryoria trichodes subsp. americana	Crater Bay	JKW 18086	KEFJ-00278	KEFJ 19479
Bryoria trichodes subsp. americana	Crater Bay	JKW 18087	KEFJ-00278	KEFJ 19480
Bryoria trichodes subsp. americana	Northwest Lagoon Beach	JKW 18108	KEFJ-00278	KEFJ 19481
Bryoria trichodes subsp. americana	Northwest Lagoon Beach	JKW 18109	KEFJ-00278	KEFJ 19482
Bryoria trichodes subsp. americana	Fire Cove	JKW 18140	KEFJ-00278	KEFJ 19483
Bryoria trichodes subsp. americana	Fire Cove	JKW 18142	KEFJ-00278	KEFJ 19484
Bryoria trichodes subsp. americana	Fire Cove	JKW 18143	KEFJ-00278	KEFJ 19485
Bryoria trichodes subsp. americana	Fire Cove	JKW 18144	KEFJ-00278	KEFJ 19486

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
Bryoria trichodes subsp. americana	Verdant Cove	JKW 18175	KEFJ-00278	KEFJ 19487
Bryoria trichodes subsp. americana	Verdant Cove	JKW 18176	KEFJ-00278	KEFJ 19488
Bryoria trichodes subsp. americana	Three Hole Bay	JKW 18211	KEFJ-00278	KEFJ 19489
Bryoria trichodes subsp. americana	Bear Cove	JKW 18260	KEFJ-00278	KEFJ 19490
Bryoria trichodes subsp. americana	Bear Cove	JKW 18262	KEFJ-00278	KEFJ 19491
Bryoria trichodes subsp. americana	Bear Cove	JKW 18263	KEFJ-00278	KEFJ 19492
Bryoria trichodes subsp. americana	Bear Cove	JKW 18264	KEFJ-00278	KEFJ 19493
Bryoria trichodes subsp. americana	Bear Cove	JKW 18265	KEFJ-00278	KEFJ 19494
Cavernularia hultenii	Bear Cove	JKW 18268	KEFJ-00278	KEFJ 18904
Cavernularia hultenii	Crater Bay	JKW 18093	KEFJ-00278	KEFJ 18976
Cavernularia hultenii	Crater Bay (northside)	JKW 18037	KEFJ-00278	KEFJ 18948
Cavernularia hultenii	East Arm (westside)	JKW 17914	KEFJ-00278	KEFJ 19288
Cavernularia hultenii	Fire Cove	JKW 18149	KEFJ-00278	KEFJ 19190
Cavernularia hultenii	North Arm Nuka Bay	JKW 17901	KEFJ-00278	KEFJ 19281
Cavernularia hultenii	Square Bay (northside)	JKW 17993	KEFJ-00278	KEFJ 19153
Cavernularia hultenii	Square Bay (southside)	JKW 18017	KEFJ-00278	KEFJ 18869
Cavernularia hultenii	Three Hole Bay	JKW 18218	KEFJ-00278	KEFJ 19234
Cavernularia hultenii	Verdant Cove	JKW 18182	KEFJ-00278	KEFJ 19210
Cavernularia lophyrea	Fire Cove	JKW 18148	KEFJ-00278	KEFJ 19189
Cavernularia lophyrea	James Lagoon	JKW 17965	KEFJ-00278	KEFJ 19331
Cavernularia lophyrea	Three Hole Bay	JKW 18216	KEFJ-00278	KEFJ 19232
Cavernularia lophyrea	Three Hole Bay	JKW 18217	KEFJ-00278	KEFJ 19233
Cavernularia lophyrea	Verdant Cove	JKW 18183	KEFJ-00278	KEFJ 19211
Cetraria aculeata	Northwest Lagoon Beach	JKW 18124	KEFJ-00278	KEFJ 18889
Cetraria ericetorum subsp. reticulata	Coleman Bay	JKW 18248	KEFJ-00278	KEFJ 19249
Cetraria ericetorum subsp. reticulata	Northwest Lagoon Beach	JKW 18129	KEFJ-00278	KEFJ 18894
Cetraria islandica subsp. crispiformis	Granite Passage Peatland	JKW 18162	KEFJ-00278	KEFJ 19200
Cetraria islandica subsp. crispiformis	Northwest Lagoon Beach	JKW 18130	KEFJ-00278	KEFJ 18895
Chaenotheca furfuracea	Crater Bay (southside)	JKW 18057	KEFJ-00278	KEFJ 18958
Cladonia amaurocraea	Crater Bay (southside)	JKW 18076	KEFJ-00278	KEFJ 18967
Cladonia amaurocraea	Northwest Lagoon Beach	JKW 18122	KEFJ-00278	KEFJ 18887
Cladonia arbuscula subsp. beringiana	Crater Bay	JKW 18092	KEFJ-00278	KEFJ 18975
Cladonia bellidiflora	East Arm (westside)	JKW 17911	KEFJ-00278	KEFJ 19397
Cladonia bellidiflora	James Lagoon	JKW 17933	KEFJ-00278	KEFJ 19398
Cladonia bellidiflora	North Arm Nuka Bay	JKW 17876	KEFJ-00278	KEFJ 19393
Cladonia bellidiflora	North Arm Nuka Bay	JKW 17883	KEFJ-00278	KEFJ 19396
Cladonia bellidiflora	Northwest Lagoon Beach	JKW 18123	KEFJ-00278	KEFJ 18888
Cladonia bellidiflora	Verdant Cove	JKW 18179	KEFJ-00278	KEFJ 19431
Cladonia carneola	North Arm Nuka Bay	JKW 17882	KEFJ-00278	KEFJ 19495
Cladonia carneola	Northwest Lagoon Beach	JKW 18115	KEFJ-00278	KEFJ 19496
Cladonia carneola	Three Hole Bay	JKW 18214	KEFJ-00278	KEFJ 19497
Cladonia chlorophaea	Fire Cove	JKW 18145	KEFJ-00278	KEFJ 19423
Cladonia chlorophaea	Northwest Lagoon Beach	JKW 18110	KEFJ-00278	KEFJ 19418
Cladonia chlorophaea	Square Bay (southside)	JKW 18009	KEFJ-00278	KEFJ 19405
Cladonia chlorophaea	Square Bay (southside)	JKW 18011	KEFJ-00278	KEFJ 19407
Cladonia chlorophaea	Three Hole Bay	JKW 18230	KEFJ-00278	KEFJ 19437
Cladonia coccifera	Crater Bay (southside)	JKW 18055	KEFJ-00278	KEFJ 18956
Cladonia coccifera	Northwest Lagoon Beach	JKW 18120	KEFJ-00278	KEFJ 18885

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
Cladonia crispata var. crispata	Northwest Lagoon Beach	JKW 18121	KEFJ-00278	KEFJ 18886
Cladonia gracilis subsp. vulnerata	Northwest Lagoon Beach	JKW 18113	KEFJ-00278	KEFJ 19498
Cladonia gracilis subsp. vulnerata	Fire Cove	JKW 19084	KEFJ-00278	KEFJ 19499
Cladonia maxima	Crater Bay (northside)	JKW 18040	KEFJ-00278	KEFJ 18877
Cladonia maxima	Crater Bay (southside)	JKW 18056	KEFJ-00278	KEFJ 18957
Cladonia ochrochlora	Crater Bay	JKW 18089	KEFJ-00278	KEFJ 19415
Cladonia ochrochlora	Crater Bay	JKW 18090	KEFJ-00278	KEFJ 19416
Cladonia ochrochlora	Crater Bay (southside)	JKW 18034	KEFJ-00278	KEFJ 19409
Cladonia ochrochlora	Fire Cove	JKW 18146	KEFJ-00278	KEFJ 19424
Cladonia ochrochlora	James Lagoon	JKW 17934	KEFJ-00278	KEFJ 19399
Cladonia ochrochlora	North Arm Nuka Bay	JKW 17877	KEFJ-00278	KEFJ 19394
Cladonia ochrochlora	Square Bay (southside)	JKW 18002	KEFJ-00278	KEFJ 19403
Cladonia ochrochlora	Square Bay (southside)	JKW 18008	KEFJ-00278	KEFJ 19404
Cladonia ochrochlora	Three Hole Bay	JKW 18212	KEFJ-00278	KEFJ 19434
Cladonia ochrochlora	Verdant Cove	JKW 18178	KEFJ-00278	KEFJ 19430
Cladonia phyllophora	Northwest Lagoon Beach	JKW 18128	KEFJ-00278	KEFJ 18893
Cladonia pyxidata	Granite Passage Peatland	JKW 18160	KEFJ-00278	KEFJ 19198
Cladonia pyxidata	Three Hole Bay	JKW 18231	KEFJ-00278	KEFJ 19438
Cladonia rangiferina	Crater Bay (southside)	JKW 18051	KEFJ-00278	KEFJ 19410
Cladonia rangiferina	Northwest Lagoon Beach	JKW 18119	KEFJ-00278	KEFJ 18884
Cladonia rangiferina	Three Hole Bay	JKW 18235	KEFJ-00278	KEFJ 19245
Cladonia squamosa	Bear Cove	JKW 18256	KEFJ-00278	KEFJ 19441
Cladonia squamosa	Bear Cove	JKW 18257	KEFJ-00278	KEFJ 19442
Cladonia squamosa	Bear Cove	JKW 18258	KEFJ-00278	KEFJ 19443
Cladonia squamosa	Bear Cove	JKW 18259	KEFJ-00278	KEFJ 19444
Cladonia squamosa	Crater Bay (southside)	JKW 18033	KEFJ-00278	KEFJ 19408
Cladonia squamosa	Crater Bay (southside)	JKW 18077	KEFJ-00278	KEFJ 18968
Cladonia squamosa	Fire Cove	JKW 18147	KEFJ-00278	KEFJ 19425
Cladonia squamosa	North Arm Nuka Bay	JKW 17880	KEFJ-00278	KEFJ 19395
Cladonia squamosa	Northwest Lagoon Beach	JKW 18117	KEFJ-00278	KEFJ 19420
Cladonia squamosa	Northwest Lagoon Beach	JKW 18125	KEFJ-00278	KEFJ 18890
Cladonia squamosa	Square Bay (southside)	JKW 18001	KEFJ-00278	KEFJ 19402
Cladonia squamosa	Square Bay (southside)	JKW 18010	KEFJ-00278	KEFJ 19406
Cladonia squamosa	Three Hole Bay	JKW 18213	KEFJ-00278	KEFJ 19435
Cladonia uncialis	Crater Bay (southside)	JKW 18058	KEFJ-00278	KEFJ 18959
Cladonia uncialis	Northwest Lagoon Beach	JKW 18111	KEFJ-00278	KEFJ 19419
Cladonia uncialis	Three Hole Bay	JKW 18232	KEFJ-00278	KEFJ 19439
Cladonia uncialis	Three Hole Bay	JKW 18234	KEFJ-00278	KEFJ 19244
Dibaeis baeomyces	Three Hole Bay	JKW 18215	KEFJ-00278	KEFJ 19231
Dibaeis baeomyces	Three Hole Bay	MH 16283	KEFJ-00278	KEFJ 19137
Ephebe sp.	Coleman Bay	JKW 18239	KEFJ-00278	KEFJ 19588
Ephebe sp.	Coleman Bay	SKT 2012-05	KEFJ-00278	KEFJ 19589
Fuscopannaria confusa	James Lagoon	JKW 17935	KEFJ-00278	KEFJ 19303
Graphis scripta	Three Hole Bay	JKW 18224	KEFJ-00278	KEFJ 19240
Hypogymnia apinnata	Bear Cove	JKW 18267	KEFJ-00278	KEFJ 18903
Hypogymnia apinnata	Crater Bay (northside)	JKW 18049	KEFJ-00278	KEFJ 18952
Hypogymnia apinnata	Fire Cove	JKW 18156	KEFJ-00278	KEFJ 19196
Hypogymnia apinnata	James Lagoon	JKW 17947	KEFJ-00278	KEFJ 19314

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
Hypogymnia apinnata	North Arm Nuka Bay	JKW 17900	KEFJ-00278	KEFJ 19280
Hypogymnia apinnata	Northwest Lagoon Beach	JKW 18131	KEFJ-00278	KEFJ 18896
Hypogymnia apinnata	Square Bay (northside)	JKW 17990	KEFJ-00278	KEFJ 19150
Hypogymnia apinnata	Three Hole Bay	JKW 18223	KEFJ-00278	KEFJ 19239
Hypogymnia apinnata	Verdant Cove	JKW 18192	KEFJ-00278	KEFJ 19219
Hypogymnia duplicata	Fire Cove	JKW 18150	KEFJ-00278	KEFJ 19191
Hypogymnia physodes	Northwest Lagoon Beach	MH 16164	KEFJ-00278	KEFJ 19358
Hypogymnia tubulosa	James Lagoon	JKW 17951	KEFJ-00278	KEFJ 19318
Icmadophila ericetorum	Crater Bay (northside)	JKW 18038	KEFJ-00278	KEFJ 18875
Leptogium lichenoides	Northwest Lagoon Beach	MH 16158	KEFJ-00278	KEFJ 19353
Leptogium saturninum	James Lagoon	JKW 17942	KEFJ-00278	KEFJ 19310
Lichenomphalia umbellifera	Crater Bay (southside)	JKW 18072	KEFJ-00278	KEFJ 19414
Lobaria linita	Crater Bay (southside)	JKW 18059	KEFJ-00278	KEFJ 18960
Lobaria linita	Crater Bay (southside)	JKW 18060	KEFJ-00278	KEFJ 18961
Lobaria linita	Square Bay (southside)	JKW 18016	KEFJ-00278	KEFJ 18868
Lobaria linita	Verdant Cove	JKW 18190	KEFJ-00278	KEFJ 19217
Lobaria oregana	James Lagoon	JKW 17939	KEFJ-00278	KEFJ 19307
Lobaria oregana	James Lagoon	JKW 17956	KEFJ-00278	KEFJ 19323
Lobaria pulmonaria	Bear Cove	JKW 18271	KEFJ-00278	KEFJ 18907
Lobaria pulmonaria	James Lagoon	JKW 17949	KEFJ-00278	KEFJ 19316
Lobaria pulmonaria	North Arm Nuka Bay	JKW 17893	KEFJ-00278	KEFJ 19275
Lobaria scrobiculata	James Lagoon	JKW 17960	KEFJ-00278	KEFJ 19326
Loxosporopsis corallifera	Crater Bay (northside)	JKW 18039	KEFJ-00278	KEFJ 18876
Massalongia carnosa	Northwest Lagoon Beach	MH 16157	KEFJ-00278	KEFJ 19352
Nephroma bellum	Bear Cove	JKW 18270	KEFJ-00278	KEFJ 18906
Nephroma bellum	East Arm (westside)	JKW 17918	KEFJ-00278	KEFJ 19292
Nephroma bellum	Northwest Lagoon Beach	MH 16168	KEFJ-00278	KEFJ 19360
Nephroma helveticum subsp. sipeanum	James Lagoon	JKW 17948	KEFJ-00278	KEFJ 19315
Nephroma helveticum subsp. sipeanum	James Lagoon	JKW 17959	KEFJ-00278	KEFJ 19325
Nephroma helveticum subsp. sipeanum	Northwest Lagoon Beach	JKW 18132	KEFJ-00278	KEFJ 18897
Nephroma helveticum subsp. sipeanum	Verdant Cove	JKW 18185	KEFJ-00278	KEFJ 19213
Nephroma isidiosum	East Arm (westside)	JKW 17954	KEFJ-00278	KEFJ 19321
Nephroma resupinatum	James Lagoon	JKW 17943	KEFJ-00278	KEFJ 19311
Nephroma resupinatum	James Lagoon	JKW 17952	KEFJ-00278	KEFJ 19319
Parmelia pseudosulcata	Bear Cove	JKW 18274	KEFJ-00278	KEFJ 18910
Parmelia pseudosulcata	Crater Bay	JKW 18102	KEFJ-00278	KEFJ 18984
Parmelia saxatilis	Crater Bay	JKW 18101	KEFJ-00278	KEFJ 18881
Parmelia saxatilis	East Arm (westside)	JKW 17920	KEFJ-00278	KEFJ 19294
Parmelia saxatilis	East Arm (westside)	JKW 17922	KEFJ-00278	KEFJ 19296
Parmelia saxatilis	James Lagoon	JKW 17945	KEFJ-00278	KEFJ 19313
Parmelia saxatilis	James Lagoon	JKW 17963	KEFJ-00278	KEFJ 19329
Parmelia saxatilis	North Arm Nuka Bay	JKW 17888	KEFJ-00278	KEFJ 19270
Parmelia saxatilis	Square Bay (northside)	JKW 17994	KEFJ-00278	KEFJ 19154
Parmelia saxatilis	Three Hole Bay	JKW 18226	KEFJ-00278	KEFJ 19242
Parmelia saxatilis	Verdant Cove	JKW 18196	KEFJ-00278	KEFJ 19222
Parmelia saxatilis	Verdant Cove	JKW 18197	KEFJ-00278	KEFJ 19223
Parmelia squarrosa	North Arm Nuka Bay	JKW 17899	KEFJ-00278	KEFJ 19279
Parmeliopsis hyperopta	Harris Peninsula Lakes	MH 16248	KEFJ-00278	KEFJ 19010

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
<i>Peltigera britannica</i>	East Arm (westside)	JKW 17926	KEFJ-00278	KEFJ 19300
<i>Peltigera britannica</i>	Fire Cove	JKW 18151	KEFJ-00278	KEFJ 19192
<i>Peltigera britannica</i>	James Lagoon	JKW 17950	KEFJ-00278	KEFJ 19317
<i>Peltigera collina</i>	James Lagoon	JKW 17936	KEFJ-00278	KEFJ 19304
<i>Peltigera collina</i>	James Lagoon	JKW 17953	KEFJ-00278	KEFJ 19320
<i>Peltigera collina</i>	Northwest Lagoon Beach	JKW 18127	KEFJ-00278	KEFJ 18892
<i>Peltigera membranacea</i>	Crater Bay (southside)	JKW 18064	KEFJ-00278	KEFJ 18965
<i>Peltigera neopolydactyla</i>	Bear Cove	JKW 18253	KEFJ-00278	KEFJ 19440
<i>Peltigera neopolydactyla</i>	Bear Cove Notch	JKW 18275	KEFJ-00278	KEFJ 18911
<i>Peltigera neopolydactyla</i>	Crater Bay (southside)	JKW 18062	KEFJ-00278	KEFJ 18963
<i>Peltigera neopolydactyla</i>	North Arm Nuka Bay	JKW 17902	KEFJ-00278	KEFJ 19282
<i>Peltigera neopolydactyla</i>	North Arm Nuka Bay	JKW 17903	KEFJ-00278	KEFJ 19283
<i>Peltigera neopolydactyla</i>	Three Hole Bay	JKW 18208	KEFJ-00278	KEFJ 19432
<i>Peltigera neopolydactyla</i>	Verdant Cove	JKW 18198	KEFJ-00278	KEFJ 19224
<i>Peltigera praetextata</i>	Harris Peninsula Lakes	MH 16233	KEFJ-00278	KEFJ 19002
<i>Phylliscum demangeonii</i>	Crater Bay	JKW 18103	KEFJ-00278	KEFJ 18882
<i>Physcia caesia</i>	Granite Passage Peatland	SKT 2012-02	KEFJ-00278	KEFJ 19457
<i>Physcia caesia</i>	Northwest Lagoon Beach	MH 16161	KEFJ-00278	KEFJ 19355
<i>Physcia caesia</i>	Three Hole Bay	JKW 18236	KEFJ-00278	KEFJ 19246
<i>Pilophorus acicularis</i>	Chance Cove	JKW 17979	KEFJ-00278	KEFJ 18934
<i>Pilophorus acicularis</i>	Crater Bay (northside)	JKW 18042	KEFJ-00278	KEFJ 18879
<i>Pilophorus clavatus</i>	Crater Bay (northside)	JKW 18043	KEFJ-00278	KEFJ 18880
<i>Pilophorus nigricaulis</i>	Crater Bay (southside)	JKW 18052	KEFJ-00278	KEFJ 19411
<i>Pilophorus nigricaulis</i>	Crater Bay (southside)	JKW 18066	KEFJ-00278	KEFJ 19412
<i>Pilophorus nigricaulis</i>	Three Hole Bay	JKW 18227	KEFJ-00278	KEFJ 19436
<i>Pilophorus nigricaulis</i>	Verdant Cove	JKW 18170	KEFJ-00278	KEFJ 19426
<i>Placopsis gelida</i>	Northwest Lagoon Beach	MH 16166	KEFJ-00278	KEFJ 19500
<i>Platismatia glauca</i>	Bear Cove	JKW 18269	KEFJ-00278	KEFJ 18905
<i>Platismatia glauca</i>	Bear Cove	JKW 18279	KEFJ-00278	KEFJ 18915
<i>Platismatia glauca</i>	Chance Cove	JKW 17972	KEFJ-00278	KEFJ 19334
<i>Platismatia glauca</i>	Chance Cove	JKW 17973	KEFJ-00278	KEFJ 19335
<i>Platismatia glauca</i>	Crater Bay (northside)	MH 18044	KEFJ-00278	KEFJ 19024
<i>Platismatia glauca</i>	East Arm (westside)	JKW 17925	KEFJ-00278	KEFJ 19299
<i>Platismatia glauca</i>	Fire Cove	JKW 18154	KEFJ-00278	KEFJ 19194
<i>Platismatia glauca</i>	North Arm Nuka Bay	JKW 17890	KEFJ-00278	KEFJ 19272
<i>Platismatia glauca</i>	Square Bay (northside)	JKW 17992	KEFJ-00278	KEFJ 19152
<i>Platismatia glauca</i>	Square Bay (northside)	JKW 17997	KEFJ-00278	KEFJ 19156
<i>Platismatia glauca</i>	Square Bay (northside)	JKW 17998	KEFJ-00278	KEFJ 19157
<i>Platismatia glauca</i>	Square Bay (southside)	JKW 18014	KEFJ-00278	KEFJ 18866
<i>Platismatia glauca</i>	Square Bay (southside)	JKW 18015	KEFJ-00278	KEFJ 18867
<i>Platismatia glauca</i>	Three Hole Bay	JKW 18220	KEFJ-00278	KEFJ 19236
<i>Platismatia glauca</i>	Verdant Cove	JKW 18184	KEFJ-00278	KEFJ 19212
<i>Platismatia herrei</i>	Bear Cove	JKW 18278	KEFJ-00278	KEFJ 18914
<i>Platismatia herrei</i>	Crater Bay	JKW 18095	KEFJ-00278	KEFJ 18978
<i>Platismatia herrei</i>	East Arm (westside)	JKW 17924	KEFJ-00278	KEFJ 19298
<i>Platismatia herrei</i>	East Arm (westside)	JKW 17938	KEFJ-00278	KEFJ 19306
<i>Platismatia herrei</i>	Fire Cove	JKW 18137	KEFJ-00278	KEFJ 19421
<i>Platismatia herrei</i>	North Arm Nuka Bay	JKW 17887	KEFJ-00278	KEFJ 19269

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
Platismatia herrei	Verdant Cove	JKW 18186	KEFJ-00278	KEFJ 19214
Platismatia lacunosa	Chance Cove	JKW 17976	KEFJ-00278	KEFJ 19149
Platismatia lacunosa	Crater Bay	JKW 18097	KEFJ-00278	KEFJ 18980
Platismatia lacunosa	Crater Bay	MH 16092	KEFJ-00278	KEFJ 19097
Platismatia lacunosa	Fire Cove	JKW 18152	KEFJ-00278	KEFJ 18835
Platismatia lacunosa	North Arm Nuka Bay	JKW 17895	KEFJ-00278	KEFJ 19148
Platismatia lacunosa	Square Bay (southside)	JKW 18018	KEFJ-00278	KEFJ 19158
Platismatia lacunosa	Verdant Cove	JKW 18173	KEFJ-00278	KEFJ 19428
Platismatia norvegica	Bear Cove	JKW 18272	KEFJ-00278	KEFJ 18908
Platismatia norvegica	Bear Cove	JKW 18273	KEFJ-00278	KEFJ 18909
Platismatia norvegica	Crater Bay	JKW 18094	KEFJ-00278	KEFJ 18977
Platismatia norvegica	Crater Bay	JKW 18096	KEFJ-00278	KEFJ 18979
Platismatia norvegica	Crater Bay	JKW 18098	KEFJ-00278	KEFJ 18981
Platismatia norvegica	Crater Bay (northside)	JKW 18041	KEFJ-00278	KEFJ 18878
Platismatia norvegica	East Arm (westside)	JKW 17917	KEFJ-00278	KEFJ 19291
Platismatia norvegica	Fire Cove	JKW 18153	KEFJ-00278	KEFJ 19193
Platismatia norvegica	James Lagoon	JKW 17937	KEFJ-00278	KEFJ 19305
Platismatia norvegica	James Lagoon	JKW 17944	KEFJ-00278	KEFJ 19312
Platismatia norvegica	North Arm Nuka Bay	JKW 17892	KEFJ-00278	KEFJ 19274
Platismatia norvegica	Northwest Lagoon Beach	JKW 18106	KEFJ-00278	KEFJ 19417
Platismatia norvegica	Square Bay (northside)	JKW 17991	KEFJ-00278	KEFJ 19151
Platismatia norvegica	Square Bay (southside)	JKW 18024	KEFJ-00278	KEFJ 18874
Platismatia norvegica	Three Hole Bay	JKW 18219	KEFJ-00278	KEFJ 19235
Platismatia norvegica	Verdant Cove	JKW 18187	KEFJ-00278	KEFJ 19215
Polychidium muscicola	Harris Peninsula Lakes	MH 16245	KEFJ-00278	KEFJ 19599
Protopannaria pezizoides	East Arm (westside)	JKW 17921	KEFJ-00278	KEFJ 19295
Pseudocyphellaria anomala	Bear Cove	JKW 18276	KEFJ-00278	KEFJ 18912
Pseudocyphellaria anomala	James Lagoon	JKW 17955	KEFJ-00278	KEFJ 19322
Pseudocyphellaria anomala	James Lagoon	JKW 17962	KEFJ-00278	KEFJ 19328
Pseudocyphellaria anomala	Square Bay (southside)	JKW 18021	KEFJ-00278	KEFJ 18871
Pseudocyphellaria crocata	James Lagoon	JKW 17958	KEFJ-00278	KEFJ 19324
Psoroma hypnorum	James Lagoon	JKW 17940	KEFJ-00278	KEFJ 19308
Psoroma hypnorum	North Arm Nuka Bay	JKW 17891	KEFJ-00278	KEFJ 19273
Racodium rupestre	Northwest Lagoon Beach	JKW 18104	KEFJ-00278	KEFJ 19585
Ramalina roesleri	Verdant Cove	JKW 18203	KEFJ-00278	KEFJ 19501
Siphula ceratites	Granite Passage Peatland	MH 16199a	KEFJ-00278	KEFJ 18985
Siphula ceratites	Granite Passage Peatland	MH 16199b	KEFJ-00278	KEFJ 19377
Siphula ceratites	Granite Passage Peatland	MH 16199c	KEFJ-00278	KEFJ 19378
Siphula ceratites	Granite Passage Peatland	MH 16199d	KEFJ-00278	KEFJ 19379
Siphula ceratites	Granite Passage Peatland	MH 16200	KEFJ-00278	KEFJ 18986
Siphula ceratites	McArthur WX	JKW 17817	KEFJ-00278	KEFJ 19390
Siphula ceratites	McArthur WX	JKW 17827	KEFJ-00278	KEFJ 19261
Sphaerophorus tuckermanii	Bear Cove	JKW 18277	KEFJ-00278	KEFJ 18913
Sphaerophorus tuckermanii	Chance Cove	JKW 17971	KEFJ-00278	KEFJ 19333
Sphaerophorus tuckermanii	Crater Bay	JKW 18100	KEFJ-00278	KEFJ 18983
Sphaerophorus tuckermanii	East Arm (westside)	JKW 17916	KEFJ-00278	KEFJ 19290
Sphaerophorus tuckermanii	Fire Cove	JKW 18157	KEFJ-00278	KEFJ 19197
Sphaerophorus tuckermanii	James Lagoon	JKW 17941	KEFJ-00278	KEFJ 19309

Species	Survey Site	Collector & Collection Number	NPS Accession Number	NPS Catalog Number
<i>Sphaerophorus tuckermanii</i>	North Arm Nuka Bay	JKW 17889	KEFJ-00278	KEFJ 19271
<i>Sphaerophorus tuckermanii</i>	Square Bay (northside)	JKW 17995	KEFJ-00278	KEFJ 19254
<i>Sphaerophorus tuckermanii</i>	Square Bay (southside)	JKW 18023	KEFJ-00278	KEFJ 18873
<i>Sphaerophorus tuckermanii</i>	Three Hole Bay	JKW 18225	KEFJ-00278	KEFJ 19241
<i>Sphaerophorus tuckermanii</i>	Verdant Cove	JKW 18195	KEFJ-00278	KEFJ 19221
<i>Stenocybe</i> sp. (see Spribille et al. 2010)	Bear Cove Notch	MH 16406	KEFJ-00278	KEFJ 19588
<i>Stereocaulon paschale</i>	Northwest Lagoon Beach	JKW 18133	KEFJ-00278	KEFJ 18898
<i>Stereocaulon spathuliferum</i>	Crater Bay (northside)	JKW 18047	KEFJ-00278	KEFJ 18950
<i>Stereocaulon spathuliferum</i>	Crater Bay (southside)	JKW 18079	KEFJ-00278	KEFJ 18970
<i>Stereocaulon tomentosum</i>	Northwest Lagoon Beach	JKW 18134	KEFJ-00278	KEFJ 18899
<i>Stereocaulon vesuvianum</i>	Crater Bay (southside)	JKW 18053	KEFJ-00278	KEFJ 18954
<i>Stereocaulon vesuvianum</i>	Crater Bay (southside)	JKW 18070	KEFJ-00278	KEFJ 19413
<i>Stereocaulon vesuvianum</i>	Crater Bay (southside)	JKW 18080	KEFJ-00278	KEFJ 18971
<i>Stereocaulon vesuvianum</i>	Crater Bay (southside)	JKW 18082	KEFJ-00278	KEFJ 18973
<i>Stereocaulon vesuvianum</i>	Granite Passage Peatland	JKW 18161	KEFJ-00278	KEFJ 19199
<i>Tuckermannopsis chlorophylla</i>	Crater Bay (southside)	JKW 18061	KEFJ-00278	KEFJ 18962
<i>Tuckermannopsis chlorophylla</i>	East Arm (westside)	JKW 17919	KEFJ-00278	KEFJ 19293
<i>Tuckermannopsis chlorophylla</i>	Verdant Cove	JKW 18194	KEFJ-00278	KEFJ 18839
<i>Umbilicaria angulata</i>	Crater Bay (northside)	JKW 18048	KEFJ-00278	KEFJ 18951
<i>Umbilicaria angulata</i>	Crater Bay (southside)	JKW 18081	KEFJ-00278	KEFJ 18972
<i>Umbilicaria angulata</i>	Crater Bay (southside)	JKW 18083	KEFJ-00278	KEFJ 18974
<i>Umbilicaria angulata</i>	Northwest Lagoon Beach	JKW 18135	KEFJ-00278	KEFJ 18900
<i>Umbilicaria angulata</i>	Northwest Lagoon Beach	JKW 18136	KEFJ-00278	KEFJ 18901
<i>Umbilicaria deusta</i>	Harris Peninsula Lakes	MH 16231	KEFJ-00278	KEFJ 19001
<i>Umbilicaria deusta</i>	Harris Peninsula Lakes	MH 16244	KEFJ-00278	KEFJ 19007
<i>Umbilicaria hyperborea</i>	Coleman Bay	JKW 18250	KEFJ-00278	KEFJ 19251
<i>Umbilicaria hyperborea</i>	Northwest Lagoon Beach	MH 16159	KEFJ-00278	KEFJ 19354
<i>Umbilicaria torrefacta</i>	Northwest Lagoon Beach	MH 16172	KEFJ-00278	KEFJ 19364
<i>Usnea filipendula</i>	Verdant Cove	JKW 18201	KEFJ-00278	KEFJ 19227
<i>Usnea filipendula</i>	Verdant Cove	JKW 18202	KEFJ-00278	KEFJ 19228
<i>Usnea longissima</i>	Crater Bay (southside)	JKW 19078	KEFJ-00278	KEFJ 19502

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

National Park Service
U.S. Department of the Interior



Natural Resource Stewardship and Science

1201 Oak Ridge Drive, Suite 150

Fort Collins, Colorado 80525

www.nature.nps.gov