



ACADIA NP • APPALACHIAN NST • BOSTON HARBOR ISLANDS NRA • MARSH-BILLINGS-ROCKEFELLER NHP • MINUTE MAN NHP
MORRISTOWN NHP • ROOSEVELT-VANDERBILT NHS • SAINT-GAUDENS NHS • SAUGUS IRON WORKS NHS • SARATOGA NHP • WEIR FARM NHS

Spring Bird Monitoring by Sight and Song

Annual volunteer landbird survey results

Background

The Northeast Temperate Network (NETN) monitors a variety of natural resource indicators, called vital signs, for 12 parks in the northeast and the Appalachian National Scenic Trail. Breeding birds are one of these vital signs because they are a reliable indicator of ecological integrity and a high profile taxonomic group. 2013 was the 8th year of continual landbird monitoring in the forests of Saratoga NHP and many other network parks. NETN and the Vermont Center for Ecostudies accomplished this only with the invaluable help of volunteer birders who crawled out of bed at ungodly early hours, endured hordes of biting mosquitoes and black flies, and contributed their time, efforts, and expert birding skills.

On a broad scale, all 13 NETN parks are located within the temperate deciduous forest biome. At a more refined level, the parks range across four Bird Conservation Regions (BCR) with this park lying in the Lower Great Lakes/St. Lawrence Plain BCR. BCR's, developed by the North American Bird Conservation Initiative, are ecologically defined areas that provide a consistent framework for bird conservation across North America. Each BCR has its own unique list of "priority" species ranked by conservation importance according to a standardized set of criteria.

Methods


A total of six study sites (three grassland sites and three forest sites) consisting of 51 point count stations were established in 2006 and 2007 at Saratoga National Historical Park, and a new forest site (with eight point count stations) was established in 2011 to replace one that was deemed too dangerous by park staff.

The recently published 2013 Landbird report includes an assessment of the ecological integrity of the forest breeding bird community at each park. For this report, instead of combining data from all survey years into a single assessment for the park, they were split into two periods (2006-2009 and 2010-2013) to compare changes over time. The ecological integrity assessments are based on groups of bird species that require similar habitat, food, or other elements for their survival. Over time, these assessments will help to shed light on changes in a broad landscape context and indicate the direction that breeding bird habitat may be going (either towards "highly disturbed" or "pristine").

The forest avian ecological integrity assessment consists of 13 guilds in three ecological integrity categories, with each guild being broadly categorized as "specialist" or "generalist." A specialist is a bird species with



Northern Bobwhites live in agricultural fields, grasslands, open pine or pine-hardwood forests, and grass-brush rangelands. . Virginia State Parks photo.



Most commonly detected species in the park.

very specific habitat needs, or one that has a low rate of population growth. If there are a relatively wide variety of specialist guilds in a park, this indicates a high-integrity habitat condition usually associated with large areas of mature forest. On the other hand, a dominance of generalist species indicates a low-integrity condition. The integrity categories are compositional (indicating species diversity), functional (highlighting ecological processes), and structural (related to the presence of important physical elements, such as tree snags and a shrub layer).

For forest breeding birds, a rank of *Good*, *Caution*, or *Significant Concern* is assigned based on species richness thresholds and ranks. "Good" represents acceptable or desired conditions; "Caution" suggests a problem may exist; "Significant concern" indicates undesired conditions that may be in need of management actions. Project organizers recognize that the assessment is based on ecological criteria, and that especially at historical parks, management goals may not always seek to attain "ecological integrity." A management assessment could be developed by those parks and NETN scientists that would reflect progress towards avian and ecological management goals.

For forest breeding birds, a rank of *Good*, *Caution*, or *Significant Concern* is assigned based on species richness thresholds and ranks. "Good" represents acceptable or desired conditions; "Caution" indicates a problem may exist; "Significant concern" indicates undesired conditions that may be in need of management actions. Project organizers recognize that the assessment is based on ecological criteria, and that especially at historical parks, management goals may not always seek to attain "ecological integrity." A management assessment could be developed by those parks and NETN scientists that would reflect progress towards avian and ecological management goals. For more information about the avian ecological integrity assessment, as well as project methodologies, sampling scheme, etc., see the Breeding Landbird Monitoring Protocol available on NETN's website.

Results and Findings

During 2013, surveys were conducted at 41 point count stations, where observers detected a total of 880 individual birds representing 60 species, including two species that were detected for the first time (Killdeer and Black-throated Green Warbler) during park surveys. Over the 8 survey years an impressive 6,018

Acadian Flycatcher	Hermit Thrush
Alder Flycatcher	House Wren
American Crow	Indigo Bunting
American Goldfinch	Killdeer
American Kestrel	Least Flycatcher
American Redstart	Louisiana Waterthrush
American Robin	Mourning Dove
American Woodcock	Northern Bobwhite
Baltimore Oriole	Northern Cardinal
Barn Swallow	Northern Flicker
Black-and-white Warbler	Northern Harrier
Black-billed Cuckoo	Northern Mockingbird
Black-capped Chickadee	Northern Rough Winged Swallow
Black-throated Green Warbler	Northern Waterthrush
Blue-headed Vireo	Orchard Oriole
Blue Jay	Ovenbird
Blue-gray Gnatcatcher	Pileated Woodpecker
Blue-winged Warbler	Prairie Warbler
Bobolink	Purple Finch
Broad-winged Hawk	Red-bellied Woodpecker
Brown Creeper	Red-breasted Nuthatch
Brown Thrasher	Red-eyed Vireo
Brown-headed Cowbird	Red-tailed Hawk
Canada Goose	Red-winged Blackbird
Cedar Waxwing	Rose-breasted Grosbeak
Chestnut-sided Warbler	Ruby-throated Hummingbird
Chipping Sparrow	Savannah Sparrow
Common Grackle	Scarlet Tanager
Common Raven	Sharp-shinned Hawk
Common Yellowthroat	Song Sparrow
Cooper's Hawk	Tree Swallow
Downy Woodpecker	Tufted Titmouse
Eastern Bluebird	Turkey Vulture
Eastern Kingbird	Veery
Eastern Meadowlark	Vesper Sparrow
Eastern Phoebe	Warbling Vireo
Eastern Towhee	White-breasted Nuthatch
Eastern Wood-Pewee	Willow Flycatcher
European Starling	Wild Turkey
Field Sparrow	Winter Wren
Golden-winged Warbler	Wood Thrush
Gray Catbird	Yellow Warbler
Great Blue Heron	Yellow-bellied Sapsucker
Great Crested Flycatcher	Yellow-billed Cuckoo
Green Heron	Yellow-rumped Warbler
Hairy Woodpecker	Yellow-throated Vireo

All 92 species that have been detected in the park's forest and grasslands since 2006.

individual birds representing 92 species have been detected in the park, with a mean abundance of 21.03 birds per point. During 2013, relative abundance and species richness both dropped slightly from 2012. Of the 10 most commonly detected species across all years, the relative abundances of three increased from 2012, and four species (American Crow, Song Sparrow, Eastern Meadowlark, and American Goldfinch) were below the 8-year mean. During 2013, eight species of regional or USA/Canada conservation concern were recorded (Eastern Wood-Pewee, Veery, Wood Thrush, Blue-winged Warbler, Eastern Towhee, Field Sparrow, Vesper Sparrow, and Baltimore Oriole). A list of species, their relative abundances, and other summary statistics are provided in the full report.

The combined (all forest sites) forest avian ecological integrity assessment comparing the first 4 years of park surveys (2006-2009) to the last 4 years (2010-2013), shows changes to two categories. The proportion of Resident species detected in the sample increased, dropping that guild from "Good" to "Caution," while the proportion of High Canopy Foragers increased slightly, nudging that category from "Significant Concern" to "Caution." In general, the structural and functional metrics were dominated by "Caution" or "Significant Concern" ratings, which continue to suggest that the park's forest bird community may be affected by the generally fragmented landscape consisting of small forest blocks broken up by fields and early successional habitats. In addition, other stressors may be affecting vegetation structure, including deer over-browse of the understory and invasive species, such as European Buckthorn and Japanese Barberry. The forest avian ecological integrity assessment has also been calculated for each forest site and these results are in full report.

The combined grassland avian ecological integrity assessment comparing the first 4 years of park surveys (2006-2009) to the last 4 years (2010-2013), resulted in changes to three categories, all of which declined. Two guilds that evaluate the proportion of grassland-dependent species detected both declined from "Good" to "Caution," the most significant of which was a 25% decline in Partner's-in-Flight (PIF) Grassland Obligate species (e.g. those listed as conservation priorities for this BCR). The other guild that showed a decline was the Edge Generalist metric, which dropped from "Caution" to "Significant Concern" due to a slight increase in the proportion of edge species detected. Grassland communities are considered a high priority in the region, and it would be prudent for the park to continue its efforts to manage grassland habitat to support grassland species. However, because 54% of shrub-dependent birds are of high conservation concern in the region, there is also value in providing habitat for these rare and/or declining species. The grassland avian ecological integrity assessment has also been calculated for each grassland site and these results are in the full report.

More Information

To get a "bird's eye" view of all the park's monitoring sites, visit the Google Earth Park Maps web page (accessible through the leftmost suite of menu choices on NETN's homepage), which also contains bird monitoring routes, Avian Ecological Integrity Assessments and species detection frequencies for all participating NETN park units.

Contacts and Websites

Steve Faccio
Project Lead Scientist

E-mail
sfaccio@vtecostudies.org

Brian Mitchell
NETN Program Manager

Phone/E-mail
802-457-3368 ext. 37
brian_mitchell@nps.gov

Full Report online at:
<https://irma.nps.gov/App/Reference/Profile/2208494>

NETN Website
<http://science.nature.nps.gov/im/units/NETN/index.cfm>



Northeast Temperate Network
54 Elm Street, Woodstock, Vermont 05091
802-457-3368
<http://www.science.nature.nps.gov/im/units/netn/>



Northeast Temperate Network