

Climate Trends, New Orleans Jazz National Historical Park, Louisiana

Patrick Gonzalez

Natural Resource Stewardship and Science, U.S. National Park Service, Washington, DC

February 5, 2014

Historical and projected climate trends for the park (data Daly et al. 2008, IPCC 2013; analysis Wang et al. in preparation, University of Wisconsin and U.S. National Park Service).

	mean	SD	mean	SD
Historical (1950-2010)				
temperature annual average	20.8	0.5 °C	69	1 °F.
temperature linear trend	0.8	N.S. °C/century	1.4	N.S. °F./cent.
precipitation annual average	1600	300 mm/y	62	11 in./year
precipitation linear trend	+21	N.S. %/century	+21	N.S. %/century
Projected (2000-2100)				
Lowest emissions scenario (IPCC RCP 2.6)				
temperature annual average trend	+1.2	0.5 °C/century	+2	1 °F./cent.
precipitation annual average trend	+4	6 %/century	+4	6 %/century
Medium emissions scenario (IPCC RCP 4.5)				
temperature annual average trend	+2.2	0.6 °C/century	+4	1 °F./cent.
precipitation annual average trend	+3	6 %/century	+3	6 %/century
High emissions scenario (IPCC RCP 6.0)				
temperature annual average trend	+2.5	0.6 °C/century	+4.5	1 °F./cent.
precipitation annual average trend	+1	9 %/century	+1	9 %/century
Highest emissions scenario (IPCC RCP 8.5)				
temperature annual average trend	+3.9	0.7 °C/century	+7	1 °F./cent.
precipitation annual average trend	~0	10 %/century	~0	10 %/century

SD = standard deviation; Sig. = statistically significant; N.S. = not statistically significant

Figure 1.

Average annual temperature (1895-2010) for the area within the park boundaries (data Daly et al. 2008; analysis Wang et al. in preparation, University of Wisconsin and U.S. National Park Service).





Figure 2.

Average annual temperature (1950-2010) across the landscape (data Daly et al. 2008; analysis Wang et al. in preparation, University of Wisconsin and U.S. National Park Service).

Linear Trend in Historical Mean Annual Temperature ($^{\circ}\text{C century}^{-1}$)
1950 - 2010

New Orleans Jazz National Historical Park



Data: IPCC 2013, Daly et al. 2008. Analysis: F. Wang, P. Gonzalez, M. Notaro, D. Vimont, J. W. Williams. In preparation.
IPCC AR5 climate projections and exposure assessments for the US National Park System.



Figure 3.

Average annual precipitation (1895-2010) for the area within the park boundaries (data Daly et al. 2008; analysis Wang et al. in preparation, University of Wisconsin and U.S. National Park Service).





Figure 4.

Average annual precipitation (1950-2010) across the landscape (data Daly et al. 2008; analysis Wang et al. in preparation, University of Wisconsin and U.S. National Park Service).

Linear Trend in Historical Total Annual Precipitation (%)
1950 - 2010
New Orleans Jazz National Historical Park



Data: IPCC 2013. Daly et al. 2008. Analysis: F. Wang, P. Gonzalez, M. Notaro, D. Vimont, J. W. Williams. In preparation.
IPCC AR5 climate projections and exposure assessments for the US National Park System.



Figure 5.

Projected change in average annual temperature (difference between 2071-2100 average and 1971-2000 average), for the high emissions scenario (IPCC RCP 6.0) (data Daly et al. 2008, IPCC 2013; analysis Wang et al. in preparation, University of Wisconsin and U.S. National Park Service).

Projected Mean Annual Temperature Change RCP6.0 (°C)
New Orleans Jazz National Historical Park
(Difference between 2071-2100 average and 1971-2000 average)



Data: IPCC 2013. Daly et al. 2008. Analysis: F. Wang, P. Gonzalez, M. Notaro, D. Vimont, J. W. Williams. In preparation.
IPCC AR5 climate projections and exposure assessments for the US National Park System.



Figure 6.

Projected change in average annual precipitation (difference between 2071-2100 average and 1971-2000 average), for the high emissions scenario (IPCC RCP 6.0) (data Daly et al. 2008, IPCC 2013; analysis Wang et al. in preparation, University of Wisconsin and U.S. National Park Service).





Figure 7.

Projected changes in average annual temperature and precipitation for all emissions scenarios (small circle = individual general circulation models, large circle = average of all general circulation models, bars = standard deviation) for the area within the park boundaries (data Daly et al. 2008, IPCC 2013; analysis Wang et al. in preparation, University of Wisconsin and U.S. National Park Service).



References

- Daly, C., M. Halbleib, J.I. Smith, W.P. Gibson, M.K. Doggett, G.H. Taylor, J. Curtis, and P.P. Pasteris. 2008. Physiographically sensitive mapping of climatological temperature and precipitation across the conterminous United States. *International Journal of Climatology* 28: 2031–2064.
- Intergovernmental Panel on Climate Change (IPCC). 2013. *Climate Change 2013: The Physical Science Basis*. Cambridge University Press, Cambridge, UK.
- Wang, F., P. Gonzalez, M. Notaro, D. Vimont, and J.W. Williams. in preparation. IPCC AR5 climate projections and exposure assessments for the US National Park System.