Forest Vulnerability

Project Brief

National Park Service
U.S. Department of the Interior

Natural Resource Stewardship & Science
Climate Change Response Program

Climate, Trees, Pests, and Weeds: Change, Uncertainty, and Biotic Stressors at Gulf Islands National Seashore

Background

Climate change and nonnative tree pests and plants affect forest composition, structure, and function, and the coming decades will likely bring accelerating changes due to these multiple global change factors. We investigated potential forest change in response to climate, differences in projections of change among climate scenarios (uncertainty), and levels of nonnative biotic stressor (tree pests and invasive plants) at 121 national park units in the eastern U.S. (see Fisichelli et al. 2014 for the full article and the project brief for a synthesis). For Gulf Islands National Seashore, we provide here park-specific climate (Table 1) and tree habitat suitability projections (Table 2) for 2100 under two climate scenarios ("least change" and "major change"). These scenarios bracket a range of plausible future conditions based on greenhouse gas emissions and global climate model projections and facilitate assessment of uncertainty in future projections. We also report nonnative tree insects and diseases ("pests") with infestation areas that include the park (Table 3). Park-level nonnative vascular plant ("weeds") information is available from NPSpecies.

Adaptation to ongoing climate change requires revising existing strategies to meet traditional goals and will increasingly require revising goals and developing new strategies. Data presented here is intended to inform managers of potential future change (climate and forests) and exacerbating factors (nonnative species).

More Information

This project is part of ongoing work of the NPS Climate Change Response Program to support park adaptation to changing conditions (websites: for managers, for the public).

Source Publication

Fisichelli, N. A., S. R. Abella, M. P. Peters, and F. J. Krist Jr. 2014. Climate, trees, pests, and weeds: change, uncertainty, and biotic stressors in eastern U.S. national park forests. Forest Ecology and Management 327:31-39.

Contact

Nicholas Fisichelli, Ph.D.
Ecologist, Climate Change Response Program
National Park Service
970-225-3578
Nicholas_Fisichelli@nps.gov

Table 1. Baseline (1961-1990) and future (2070-2099) climates used in tree habitat suitability models for Gulf Islands National Seashore. Future projections for the two climate scenarios ("Least Change" and "Major Change") illustrate a plausible range of future conditions and are shown as departures from baseline values. The Least Change scenario includes a major decrease in future greenhouse gas emissions while the Major Change scenario is a continuation of historical increases in emissions. Climate data are averaged across a roughly 6200 mi^2 (16000 km^2) area centered on the park.

Climate Variable	Baseline (1961-1990)	Least Change (2070-2099)	Major Change (2070-2099)
mean annual temperature	18.9 °C (65.9 °F)	+1.7 °C (+3 °F)	+6.5 °C (+11.6 °F)
mean January temperature	9.1 °C (48.4 °F)	+1.4 °C (+2.6 °F)	+5.3 °C (+9.5 °F)
mean July temperature	27 °C (80.6 °F)	+2 °C (+3.6 °F)	+7.1 °C (+12.8 °F)
seasonality (July-January temp.)	17.9 °C (32.2 °F)	+0.6 °C (+1 °F)	+1.8 °C (+3.2 °F)
mean May-September temp.	25.3 °C (77.6 °F)	+1.9 °C (+3.4 °F)	+7.1 °C (+12.9 °F)
annual precipitation	1653 mm (65.1 in)	+6.7 %	+3.4 %
May-September precipitation	778 mm (30.6 in)	+14 %	-3.1 %

Table 2. Potential changes in habitat suitability (2100 compared with 1990) for tree species in Gulf Islands National Seashore. Species are grouped into decreasing, no change, increasing, mixed results, and new habitat groups based on change class designations for two future climate scenarios (the "least change" scenario represents strong cuts in greenhouse gas emissions and modest climatic changes and the "major change" scenario represents continued increasing greenhouse gas emissions and rapid warming). Change class designations are based on the ratio of future (2100) to baseline (1990) habitat suitability and baseline habitat values, e.g., for common species, large decrease is ≤ 0.5, small decrease is > 0.5 and ≤ 0.8, no change is > 0.8 and ≤ 1.2, small increase is > 1.2 and ≤ 2.0, and large increase is > 2.0. Note: table below uses finer change classes (n=8) than was used in the original article (n=3) to designate uncertainty in habitat projections. Habitat suitability models from the US Forest Service Climate Change Tree Atlas. Contact author for further details on change class definitions.

Decreases in Potential Habitat

Scientific Name	Common Name	Least Change	Major Change
Prunus serotina	black cherry	small decrease	small decrease

No Change in Potential Habitat

Scientific Name	Common Name	Least Change	Major Change
Acer rubrum	red maple	no change	no change
Cornus florida	flowering dogwood	no change	no change
Diospyros virginiana	common persimmon	no change	no change
Ilex opaca	American holly	no change	no change
Liriodendron tulipifera	yellow-poplar	no change	no change
Nyssa sylvatica var. biflora	swamp tupelo	no change	no change
Persea borbonia	redbay	no change	no change
Pinus palustris	longleaf pine	no change	no change
Quercus alba	white oak	no change	no change
Quercus falcata var.falcata	southern red oak	no change	no change
Quercus incana	bluejack oak	no change	no change
Quercus laevis	turkey oak	no change	no change
Quercus marilandica	blackjack oak	no change	no change
Quercus nigra	water oak	no change	no change

Increases in Potential Habitat

Scientific Name	Common Name	Least Change	Major Change
Carya illinoinensis	pecan	large increase	large increase
Carya tomentosa	mockernut hickory	large increase	small increase
Chamaecyparis thyoides	Atlantic white-cedar	small increase	small increase
Fraxinus americana	white ash	large increase	large increase
Fraxinus pennsylvanica	green ash	small increase	small increase
Magnolia grandiflora	southern magnolia	small increase	small increase
Nyssa sylvatica	blackgum	small increase	small increase
Pinus echinata	shortleaf pine	small increase	small increase
Quercus stellata	post oak	small increase	small increase
Quercus virginiana	live oak	small increase	large increase
Taxodium distichum	baldcypress	small increase	small increase
Taxodium distichum var. nutans	pondcypress	small increase	large increase

Mixed Results

Scientific Name	Common Name	Least Change	Major Change
Betula nigra	river birch	small increase	no change
Carpinus caroliniana	American hornbeam	small increase	no change

Mixed Results (continued)

Scientific Name	Common Name	Least Change	Major Change
Liquidambar styraciflua	sweetgum	no change	small increase
Magnolia virginiana	sweetbay	small increase	no change
Nyssa aquatica	water tupelo	no change	small decrease
Pinus clausa	sand pine	no change	small increase
Pinus elliottii	slash pine	no change	small decrease
Pinus taeda	loblolly pine	no change	small increase
Quercus laurifolia	laurel oak	small increase	no change
Quercus phellos	willow oak	no change	small increase

New Potential Habitat

Scientific Name	Common Name	Least Change	Major Change
Carya glabra	pignut hickory	new entry	new entry
Ulmus alata	winged elm	new entry	new entry

Table 3. Nonnative tree insects and diseases ("tree pests") with infestation areas that include Gulf Islands National Seashore. Detection scale is the finest spatial scale at which infestation data were available for the park area. Data derived from the US Forest Service Alien Forest Pest Explorer (AFPE) Database.

Scientific Name	Common Name	Detection Scale
Anarsia lineatella	peach twig borer	state
Aonidiella aurantii	California red scale	state
Blastopsylla occidentalis	eucalyptus psyllid	state
Caliroa cerasi	pear sawfly	state
Carulaspis juniperi	juniper scale	state
Caulocampus acericaulis	maple petiole borer	state
Coleophora serratella	birch casebearer	state
Cryphonectria parasitica	chestnut blight	county
Cyrtepistomus castaneus	Asiatic oak weevil	county
Diaspidiotus perniciosus	San Jose scale	state
Dryocosmus kuriphilus	chestnut gall wasp	state
Eulecanium cerasorum	calico scale	state
Glycaspis brimblecombei	red gum lerp psyllid	state
Homadaula anisocentra	mimosa webworm	county
Icerya purchasi	cottony cushion scale	county
Kaliofenusa ulmi	elm leafminer	state
Lepidosaphes ulmi	oystershell scale	state
Maconellicoccus hirsutus	pink hibiscus mealybug	state
Ophiostoma novo-ulmi	Dutch elm disease	state
Otiorhynchus ovatus	strawberry root weevil	state
Periphyllus lyropictus	Norway maple aphid	state
Phytophthora cinnamomi	littleleaf disease / phytophthora root rot	state
Plagiodera versicolora	imported willow leaf beetle	state
Popillia japonica	Japanese beetle	state
Pseudaonidia duplex	camphor scale	county
Raffaelea spp.	laurel wilt	county

Table 3 (continued). Nonnative tree insects and diseases ("tree pests") with infestation areas that include Gulf Islands National Seashore.

Scientific Name	Common Name	Detection Scale
Scolytus multistriatus	smaller European elm bark beetle	state
Scolytus schevyrewi	banded elm bark beetle	state
Xanthogaleruca luteola	elm leafbeetle	state