



Alagnak

Aniakchak

Katmai

Kenai Fjords

Lake Clark

Black Oystercatchers

Importance

The black oystercatcher is a common and conspicuous member of rocky intertidal marine communities and is thought to be important in structuring nearshore ecosystems. Black oystercatchers are completely dependent on nearshore marine habitats for all critical parts of their life cycle including foraging, breeding, chick-rearing and resting. Because of their complete reliance on nearshore marine habitats, they are also highly susceptible to human disturbance. The black oystercatcher is considered a Management Indicator Species by the Chugach National Forest and a species of concern by the Alaska Shorebird Working Group. It is widely recognized as a species representative of nearshore habitats and therefore particularly amenable to long term monitoring.

Monitoring Approach

Southwest Alaska Network (SWAN) staff and partners have been monitoring black oystercatchers annually in Katmai National Park and Preserve (NPP) since 2006, annually in Kenai Fjords National Park (NP) since 2007 and in western Prince William Sound (WPWS) since 2007, with annual sampling in WPWS starting in 2010. Boat-based surveys are conducted in early summer to determine breeding density. Nests are located on foot and examined for presence of chicks and/or eggs to determine productivity. Prey remains are collected and identified to species to determine chick provisioning habits.



Examining prey remains collected near black oystercatcher nests. NPS photo




Black oystercatcher (top) and a clutch of three black oystercatcher eggs (bottom). NPS photos/K. Chritz

Findings to date

Active nest density continues to be similar across time with little evidence of a trend in all three regions (Figure 1). However, productivity (the number of eggs and chicks per nest) tends to be highly variable across all regions. Because we only visit a nest once per year in our monitoring program, results reflect productivity at the time of the survey, adding to the variability in the results.


Prey remains collected from nests indicate that limpets (predominantly *Lottia* sp.) and mussels (*Mytilus trossulus*) continue to be the dominant food source for black oystercatchers in all three regions. However, we observed a decrease in the proportion of mussels in black oystercatcher diet over time (Figure 2), concurrent with changes in mussel availability (see mussel resource brief). Changes in prey abundance and how those changes potentially affect black oystercatcher populations is a question our long-term monitoring data is designed to answer.

Figure 1



Number of active black oystercatcher nests found per kilometer along shorelines in Kenai Fjords NP, Katmai NP, and western Prince William Sound. Error bars represent +/- 1 standard error.

Figure 2



The average proportion of mussels in black oystercatcher diets through time. Diet remains were collected from nest sites within the study regions. Error bars represent +/- 1 standard error.

For more information:

Contact: Heather Coletti, heather_coletti@nps.gov
Nearshore web site at <http://science.nature.nps.gov/im/units/swan/monitor/nearshore.cfm>

April 2016

Brief designed by: Erin Kunisch
www.nps.gov/rlc/oceanalaska/index.htm