



Weather of Hopewell Furnace National Historic Site and Valley Forge National Historical Park

Mid-Atlantic Network Summary Report for 2016

Natural Resource Data Series NPS/MIDN/NRDS—2017/1101



ON THE COVER

Valley Forge National Historical Park

©Donna Higgins

Weather of Hopewell Furnace National Historic Site and Valley Forge National Historical Park

Mid-Atlantic Network Summary Report for 2016

Natural Resource Data Series NPS/MIDN/NRDS—2017/1101

John E. B. Wofford

Mid-Atlantic Network
National Park Service
120 Chatham Lane
Fredericksburg, VA, 22405

April 2017

U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science
Fort Collins, Colorado

The National Park Service, Natural Resource Stewardship and Science office in Fort Collins, Colorado, publishes a range of reports that address natural resource topics. These reports are of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Data Series is intended for the timely release of basic data sets and data summaries. Care has been taken to assure accuracy of raw data values, but a thorough analysis and interpretation of the data has not been completed. Consequently, the initial analyses of data in this report are provisional and subject to change.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner.

This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data. Data in this report were collected and analyzed using methods based on established, peer-reviewed protocols and were analyzed and interpreted within the guidelines of the protocols.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available in digital format from the [Mid-Atlantic Network website](#) and the [Natural Resource Publications Management website](#). To receive this report in a format that is optimized to be accessible using screen readers for the visually or cognitively impaired, please email irma@nps.gov.

Please cite this publication as:

Wofford, J. E. B. 2017. Weather of Hopewell Furnace National Historic Site and Valley Forge National Historical Park: Mid-Atlantic Network summary report for 2016. Natural Resource Data Series NPS/MIDN/NRDS—2017/1101. National Park Service, Fort Collins, Colorado.

Contents

	Page
Figures.....	iv
Tables.....	v
Abstract.....	vi
List of Acronyms	vii
Introduction.....	1
Methods.....	1
Observing Stations.....	1
Climate Indicators and Climatological Normals	2
Drought Status	4
Data Processing and Quality Assurance	4
Results.....	7
Temperature Summary	7
Precipitation Summary	13
Drought Status	17
Literature Cited	21

Figures

	Page
Figure 1. Location of weather observing stations Hopewell Furnace National Historic Site and Valley Forge National Historical Park selected as most representative of the parks in 2016.....	5
Figure 2. Maps showing departure from average monthly maximum temperature in 2016 compared to the 30-year normal (1981–2010).....	8
Figure 3. Maps showing departure from average monthly minimum temperature in 2016 compared to the 30-year normal (1981–2010).....	9
Figure 4. Maps showing percent of average monthly precipitation in 2016 compared to the 30-year normal (1981–2010).....	14
Figure 5. Palmer Drought Severity Index (PDSI) for the Southeastern Piedmont, 2014–2016.....	17
Figure 6. Mid-month values of the United States Drought Monitor (DM) - Drought Intensity Index for Pennsylvania in 2016.	18
Figure 7. Mid-month values of the United States Drought Monitor (DM) - Drought Intensity Index for the Northeast in 2016.	19

Tables

	Page
Table 1. List of weather observing stations around Hopewell Furnace National Historic Site and Valley Forge National Historical Park selected as most representative of the parks in 2016.....	6
Table 2. Summary of 2016 departure from normal temperature based on the 30-year normal (1981–2010) for selected stations.....	10
Table 3. Summary of 2016 monthly average temperatures for selected stations.....	10
Table 4. Status of 2016 temperature indicators compared to the 30-year normal (1981–2010) for West Chester.....	11
Table 5. Seasonal and annual temperature and precipitation rankings for 2016 over 122 years for the Southeastern Piedmont (NOAA 2017).	12
Table 6. Summary of 2016 percent of normal precipitation based on the 30-year normal (1981–2010) for selected stations.	15
Table 7. Status of 2016 precipitation indicators compared to the 30-year normal (1981–2010) at for selected stations.....	15
Table 8. Summary of 2016 monthly total precipitation for selected stations.....	16
Table 9. Top five wettest days and top five dry spells (consecutive days with a trace or less of liquid precipitation) during 2016 from weather observing stations at West Chester 2 NW (WCHP1) and Glenmoore (GNMP1).....	16

Abstract

The Mid-Atlantic Network (MIDN) weather and climate monitoring protocol supports the documentation of current status and long-term trends in air temperature and precipitation at multiple temporal and spatial scales in MIDN parks. The purpose of this report is to provide a concise weather and climate summary for Hopewell Furnace National Historic Site (HOFU) and Valley Forge National Historical Park (VAFO) for the time period of January 1 to December 31, 2016. HOFU and VAFO are located in the Southeastern Piedmont climate division of Pennsylvania. In 2016, a total of six weather stations were selected to report park weather and climate conditions, and a large suite of precipitation and temperature metrics were summarized. Ultimately, 2016 was a very warm year and ranked as the 4th warmest on record for the Southeastern Piedmont. All of the seasons were warmer than average, and the summer of 2016 was the warmest ever recorded in the Southeastern Piedmont of Pennsylvania. In terms of precipitation, most seasons were near to or below normal. The autumn of 2016 was particularly dry. In total, the year was the 100th wettest (i.e. 23rd driest) since 1895. By the end of 2016, abnormally dry to severe drought conditions were occurring across much of Pennsylvania.

List of Acronyms

ASOS	Automated Surface Observation System
CASTNET	Clean Air Status and Trends Network
COOP	National Weather Service Cooperative Observer Program
HOFU	Hopewell Furnace National Historic Site
MIDN	Mid-Atlantic Network
NCDC	National Climatic Data Center
NHP	National Historical Park
NHS	National Historic Site
NOAA	National Oceanic and Atmospheric Administration
PDSI	Palmer Drought Severity Index
POR	Period of Record
PRISM	Parameter-elevation Regressions on Independent Slopes Model
RAWS	Remote Automated Weather Stations
USDM	United States Drought Monitor
USGS	United States Geological Survey
VAFO	Valley Forge National Historical Park

Introduction

Weather and climate are widely recognized as key drivers of terrestrial and aquatic ecosystems, affecting biotic as well as abiotic ecosystem characteristics and processes. Global and regional scale climatic patterns, trends, and variations are critical to the cycling of elements, nutrients, and minerals through ecosystems and can deliver pollutants from regional and even global sources (National Assessment Synthesis Team 2001). These variations and trends influence the fundamental properties of ecologic systems such as soil-water relationships and plant-soil processes and their disturbance rates and intensity. Information obtained from meteorological monitoring will be useful to interpreting and understanding changes in species composition, community structure, water and soil chemistry, and related landscape processes (Comiskey and Callahan 2008).

The purpose of this report is to provide a concise weather and climate summary for January 1 to December 31, 2016, and to place current patterns and trends in an appropriate historical and regional context (Marshall et al. 2012). It is our intention that this report will satisfy an inherent interest in meteorological phenomena and meet the Mid-Atlantic Network (MIDN) Weather and Climate Monitoring objective:

- Document current status and long-term trends in air temperature and precipitation at multiple temporal scales (e.g., daily, monthly, seasonal, annual, and decadal) and spatial scales (e.g., individual stations and aggregated stations such as climate divisions) utilizing existing weather and climate monitoring programs and datasets.

To accomplish this objective, a variety of atmospheric data streams were evaluated for their quality, longevity, and applicability to the MIDN parks. Since no single weather observing network contains all the pertinent measures of atmospheric phenomena to assess ecosystem health, an objective analysis of the data networks was developed and outlined in the Weather and Climate Monitoring Protocol for the Eastern Rivers and Mountains Network and Mid-Atlantic Network of the National Park Service (Marshall et al. 2012). Through this analysis, a select number of weather/climate observing stations were chosen as representative of each park; these are the primary data sources used in the profile of 2016's climate summary.

In addition to a suite of summary tables, graphs, and narratives, we specifically identify a series of key weather indicators to report status and trends on an annual basis and periodically in separate and more thorough reports (for the most recent trend report, see Knight et al. 2016). These key indicators are further described in the protocol (Marshall et al. 2012) and are summarized in this report.

Methods

Observing Stations

A total of six weather observing stations comprised of three weather observing networks were selected as the most representative weather stations around Hopewell Furnace National Historic Site (NHS) and Valley Forge National Historical Park (NHP) in 2016 (Figure 1, Table 1). Representative stations within a 100-km range of each park were chosen based on several criteria, which include

proximity to the parks, the representativeness of the station to the park elevation profile, the type and frequency of observations, the period of record of the data, and data availability (Marshall et al. 2012). Moreover, the percentage of time a station reports a particular parameter (e.g., temperature) can influence data inclusion.

Hopewell Furnace NHS and Valley Forge NHP are located in Pennsylvania Climate Division 3, known as the “Southeastern Piedmont”. A climate division is a region that is reasonably homogenous with respect to climatic and hydrologic characteristics and is frequently used for compiling climate statistics (<http://www.esrl.noaa.gov/psd/data/usclimdivs/data/map.html>).

Climate Indicators and Climatological Normals

A suite of indicators were identified to provide a summary of weather and climate conditions (Marshall et al. 2012). The following are the climate indicators selected for each of the parks:

- Average Annual Mean Temperature – A measure of the composite mean temperature (averaging the maximum and minimum on a daily basis) for an entire year
- Average Annual Maximum Temperature – A measure of the average of the daily maximum temperature for an entire year
- Highest Annual Maximum Temperature – A measure of the individual highest maximum temperature each year
- Annual Number of Hot Days (maximum temperature equal to or exceeding 90°F (32°C) – A measure of warm season heat using a threshold of days with a 90°F (32°C) maxima
- Average Annual Minimum Temperature – A measure of the average of the daily minimum temperature for an entire year
- Lowest Annual Minimum Temperature – A measure of the individual lowest minimum temperature each year
- Annual Number of Cold Days (maximum temperature less than or equal to 32°F (0°C)) – A measure of cold season chill using a threshold of days with a 32°F (0°C) maxima
- Annual Number of Subfreezing Days (minimum temperature less than or equal to 32°F (0°C)) – A measure of cold season chill using a threshold of days with a 32°F (0°C) minima
- Annual Number of Subzero Days (minimum temperature less than or equal to 0°F (-17.8°C)) – A measure of cold season chill using a threshold of days with a 0°F (-17.8°C) minima
- Growing Season Length (number of days between last freeze in spring and first freeze in fall) – A measure of the local growing season as tallied by the total number of days between the last 32°F (0°C) in the spring and the first 32°F (0°C) in the fall
- Annual Precipitation – A measure of total liquid (and its equivalent melted snow) for all days
- Autumn Precipitation (October 1 – December 31) - The three month total precipitation including rain and melted snow
- Annual Number of Heavy Precipitation Days (liquid equal to or greater than 1.0 in (25mm)) – A measure of the frequency of heavy precipitation events during each year

- Annual Number of Extreme Precipitation Days (liquid equal to or greater than 2.0 in (50mm)) – A measure of frequency of extreme precipitation events during each year
- Annual Number of Micro-droughts – A measure of micro-droughts for the year as the number of periods of 7+ days without rain
- Annual Snowfall – A measure of total solid precipitation (includes sleet and hail; does not include freezing rain)
- Annual Number of Measureable Snow Days – A measure of the severity of winter precipitation as the number of days receiving more than or equal to 0.1 in (0.3 cm) of solid precipitation (includes sleet and hail; does not include freezing rain)
- Annual Number of Moderate Snow Days– A measure of the severity of winter precipitation as the number of days receiving more than or equal to 3 in (7.6 cm) of solid precipitation (includes sleet and hail; does not include freezing rain)
- Annual Number of Heavy Snow Days– A measure of the severity of winter precipitation as the number of days receiving more than or equal to 5 in (12.7 cm) of solid precipitation (includes sleet and hail; does not include freezing rain)

The average value of a climate element over 30 years is defined as a climatological normal, which are calculated and established by NOAA’s National Climatic Data Center (NCDC). Every ten years, NCDC computes new thirty-year climate normals for selected temperature and precipitation elements for a large number of U.S. climate and weather stations. The current (as of 2011) normals cover the period 1981–2010. In this report, the 30-year normals established by NCDC are used as the baseline for comparisons (e.g., departures from normal). In cases where data for the 30-year normal period are not available, we use alternative comparisons such as the new pseudo-normals from NCDC or a recent 10-year period. In some cases, sufficient data may simply not be available to calculate normals. For metrics that NCDC may not routinely calculate a normal, such as the number of days with more than 2 in (55 mm) of rain or liquid equivalent, normals are calculated using the same time period (e.g., 1981–2010) as the current NCDC standard. Throughout the report, descriptions of a station’s values as compared to the normals are described as a difference from the “average”, “mean”, “typical”, “long-term value”, as well as “normal”, to improve the readability of the document. However, all of these terms are comparing a value from one year at that station to that station’s normal, whether it is the 30-year normal or the pseudo-normal calculated on a shorter time frame.

Of the six stations, Glenmoore, West Chester, Reading, and Morgantown have a Period of Record (POR) sufficiently long for a 30-year normal (normal calculated from years 1981–2010) for at least temperature or precipitation. However, Morgantown stopped recording temperature in 1986 and also has a low percent of time reporting precipitation over its POR. Only West Chester was used for the most thorough temperature indicator analysis, as Glenmoore started reporting temperature in 2005. Glenmoore and West Chester are used in the precipitation indicator analysis. Monthly temperature averages are presented for West Chester, Pottstown, Reading, and Glenmore and monthly temperature departures from normal are presented for West Chester, Pottstown, and Reading; all

stations are used for monthly precipitation averages. Due to a short period of record, precipitation departures are not calculated for Valley Creek.

Drought Status

There are a number of drought indices used to estimate the severity of drought in an area using algorithms that incorporate recent temperatures, rainfall, soil moisture, and other information (<http://www.drought.gov>). The main indices we report are the Palmer Drought Severity Index (PDSI) and the United States Drought Monitor (DM) – Drought Intensity Index. While both indices provide excellent summary information on broad-scale conditions, local conditions (such as at the park scale) may vary.

The PDSI is a soil moisture algorithm calibrated for relatively homogeneous regions and is calculated on a monthly basis using precipitation and temperature data, as well as the water content of the soil. The values vary between extremely moist (>4.0) and extreme drought (<-4.0). Positive values indicate periods of above normal precipitation (0 to 2: near normal; 2 to 3: moderate moisture; >3 : very moist) and negative values indicate periods of drought (0 to -2: near normal; -2 to -3: moderate drought; <-3 : severe to extreme drought).

The DM – Drought Intensity Index is a synthesis of multiple indices (including the PDSI) and impacts, which represents a consensus of federal and academic scientists (NIDIS 2017). The DM produces a summary map of drought intensity for the nation and all states each week. It is on a scale ranging from abnormally dry (D0) to exceptional drought (D4).

Data Processing and Quality Assurance

Data from selected stations are downloaded from relevant data streams (ASOS, RAWS, CASTNET, etc.). Variables are calculated using simple statistics (e.g. mean, minimum, maximum, etc.).

Each data stream has varying levels of associated quality assurance (from none to extensive). ASOS, RAWS, and CASTNet data have rigorous and extensive quality assurance beginning at the point of gather (data-logger on site) and at additional levels as described in the respective user's manuals and specification documents.

The COOP data quality control is now much more extensive in real-time as the electronically submitted data passes through several levels of quality assurance and is in a final format within a few days of submission.

Quality assurance of USGS precipitation data is performed manually by comparing rainfall amounts with nearby stations.

Data management procedures are fully outlined in Marshal, et al (2012).




Figure 1. Location of weather observing stations Hopewell Furnace National Historic Site and Valley Forge National Historical Park selected as most representative of the parks in 2016. See Table 1 for station names.

Table 1. List of weather observing stations around Hopewell Furnace National Historic Site and Valley Forge National Historical Park selected as most representative of the parks in 2016.

Station	Observing Network ^A	Station Name	Period of Record (POR)	Percentage of Time Reporting Temperature for 2016	Percentage of Time Reporting Precipitation for 2016	Percentage of Time Reporting Temperature for entire POR	Percentage of Time Reporting Precipitation for entire POR
GNMP1	COOP	Glenmoore	04/01/1959 - Present	100.0	100.0	96.1 ^B	99.2
MGNP1	COOP	Morgantown	06/01/1951 - Present	-	99.5	86.4 ^C	62.9
KPTW	ASOS	Pottstown Heritage Field Airport	01/01/1999 - Present	100.0	100.0	97.4	97.4
KRDG	ASOS	Reading Regional Airport/Carl A. Spaatz Field	01/01/1973 - Present	100.0	100.0	98.0	39.1
WCHP1	COOP	West Chester 2 NW	01/01/1893 - Present	100.0	100.0	90.3	89.1
1473169	USGS	Valley Creek	10/01/2005 - Present	–	100.0	–	98.8

^A Observing networks are: ASOS — Automated Surface Observation System, COOP — National Weather Service Cooperative Observer Program, USGS — United States Geological Survey.

^B Temperature records began at Glenmoore (GNMP1) on October 3, 2005. POR percentages are based off of this start date.

^C Percentage of time reporting temperature for Morgantown is based upon a Period of Record ending on 03/01/1986 because the station stopped reporting temperature after this date.

Results

Temperature Summary

Almost all of the temperature metrics calculated from stations around HOFU and VAFO indicate that 2016 was a warm year. Throughout the year, most months had higher than average temperatures (Figures 2 and 3, Table 2 and 3). Average annual temperatures were also above normal at all of the reporting stations (Tables 2 and 3). In addition, the number of hot days, average annual maximum temperatures, and average annual minimum temperatures were all above average (Table 4). Even so, growing season length was more than a week shorter than normal (Table 4). All of the seasons ranked warmer than average, and the summer of 2016 was the warmest ever recorded in the Southeastern Piedmont of Pennsylvania (Table 5). In all, 2016 was the 4th warmest since the year 1895 (Table 5).




Figure 2. Maps showing departure from average monthly maximum temperature in 2016 compared to the 30-year normal (1981–2010).




Figure 3. Maps showing departure from average monthly minimum temperature in 2016 compared to the 30-year normal (1981–2010).

Table 2. Summary of 2016 departure from normal temperature based on the 30-year normal (1981–2010) for selected stations.

Station name	Station	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
Pottstown Airport*	KPTW	-0.9°F	1.5°F	5.6°F	-1.8°F	-1.2°F	0.2°F	2.4°F	3.1°F	3.8°F	1.5°F	1.1°F	0.1°F	1.2°F
		-0.5°C	0.9°C	3.1°C	-1.0°C	-0.6°C	0.1°C	1.3°C	1.7°C	2.1°C	0.8°C	0.6°C	0.1°C	0.7°C
Reading Airport	KRDG	0.0°F	2.3°F	6.5°F	-0.6°F	0.0°F	1.4°F	3.8°F	4.4°F	4.7°F	3.1°F	2.2°F	1.4°F	1.3°F
		0.0°C	1.2°C	3.6°C	-0.3°C	0.0°C	0.8°C	2.1°C	2.4°C	2.6°C	1.7°C	1.2°C	0.7°C	0.9°C
West Chester 2 NW	WCHP1	0.2°F	2.5°F	5.7°F	-1.3°F	-1.9°F	0.5°F	1.9°F	3.6°F	3.4°F	2.5°F	0.6°F	1.8°F	1.6°F
		0.1°C	1.4°C	3.1°C	-0.7°C	-1.1°C	0.3°C	1.1°C	2.0°C	1.9°C	1.4°C	0.4°C	1.0°C	0.9°C

*Indicates a station's Period of Record does not cover the period of 1981-2010 but is five or more years. In these cases, the departures from normal values were calculated with normals derived from data spanning the length of the station's Period of Record.

Table 3. Summary of 2016 monthly average temperatures for selected stations.

Station name	Station	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
Glenmoore	GNMP1	29.7°F	34.1°F	46.5°F	50.2°F	59.4°F	70.4°F	76.5°F	76.1°F	69.2°F	56.3°F	46.4°F	35.5°F	54.2°F
		-1.3°C	1.2°C	8.1°C	10.1°C	15.2°C	21.3°C	24.7°C	24.5°C	20.7°C	13.5°C	8.0°C	1.9°C	12.3°C
Pottstown Airport*	KPTW	29.2°F	33.9°F	46.3°F	49.5°F	59.5°F	70.0°F	76.8°F	76.0°F	68.7°F	55.4°F	45.4°F	34.5°F	53.8°F
		-1.6°C	1.1°C	7.9°C	9.7°C	15.3°C	21.1°C	24.9°C	24.4°C	20.4°C	13.0°C	7.4°C	1.4°C	12.1°C
Reading Airport	KRDG	29.7°F	34.6°F	47.6°F	51.0°F	61.1°F	71.9°F	78.6°F	77.4°F	70.0°F	56.8°F	46.3°F	35.3°F	55.1°F
		-1.3°C	1.4°C	8.7°C	10.6°C	16.2°C	22.2°C	25.9°C	25.2°C	21.1°C	13.8°C	7.9°C	1.8°C	12.8°C
West Chester 2 NW	WCHP1	29.1°F	33.8°F	45.4°F	49.2°F	58.5°F	70.7°F	76.6°F	76.5°F	68.5°F	55.8°F	44.0°F	35.1°F	53.7°F
		-1.6°C	1.0°C	7.4°C	9.6°C	14.7°C	21.5°C	24.8°C	24.7°C	20.3°C	13.2°C	6.7°C	1.7°C	12.1°C

Table 4. Status of 2016 temperature indicators compared to the 30-year normal (1981–2010) for West Chester.

Temperature Indicator	West Chester 2 NW 2016	West Chester 2 NW 1981–2010
Average Annual Temperature	53.7°F 12.1°C	52.1°F 11.2°C
Average Annual Maximum Temperature	65.6°F 18.7°C	63.8°F 17.7°C
Maximum Temperature	96.0°F 35.6°C	97.0°F 36.1°C
Hot Days (days with Tmax≥90°F/32°C)	36	21
Average Annual Minimum Temperature	41.9°F 5.5°C	40.4°F 4.7°C
Minimum Temperature	4.0°F -15.6 °C	-0.5°F -18.1°C
Cold Days (days with Tmax≤32°F/0°C)	14	16
Sub-freezing Days (days with Tmin≤32°F/0°C)	130	130
Sub-zero Days (days with Tmin≤0°F/-18°C)	0	2
Growing Season Length (days between last spring Tmin 32°F/0°C and first fall Tmin 32°F/0°C)	163	174

Table 5. Seasonal and annual temperature and precipitation rankings for 2016 over 122 years for the Southeastern Piedmont (NOAA 2017).

Parameter	PA Climate Division 3 Rankings — Southeastern Piedmont				
	Jan–Feb–Mar WINTER	Apr–May–Jun SPRING	Jul–Aug–Sep SUMMER	Oct–Nov–Dec AUTUMN	Jan–Dec YEAR
Temperature-2016	9	41-T	1	11	4
Precipitation-2016	60	84-T	67	108-T	100

1 = warmest/wettest year and 122 = coldest/driest year. T = Tie between multiple years

Precipitation Summary

Although monthly precipitation totals were variable, annual totals across all stations were below average (Figure 4 and Table 6). In all, annual precipitation ranged from 3.0 to 7.1 inches (76 to 181 mm) below average (Table 7). Monthly totals ranged from a low of 0.9 inches (23 mm) at Reading Airport in October to a high of 5.9 inches (150 mm) at Glenmore in February (Table 8). The wettest day occurred on September 30th when 2.9 inches (74 mm) of rain was observed. In all, 2016 was the 100th wettest (i.e. 23rd driest) year on record. By then end of 2016, severe drought conditions were occurring across Pennsylvania and the Northeast (Figures 5, 6, and 7).




Figure 4. Maps showing percent of average monthly precipitation in 2016 compared to the 30-year normal (1981–2010).

Table 6. Summary of 2016 percent of normal precipitation based on the 30-year normal (1981–2010) for selected stations.

Station name	Station	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
Glenmoore	GNMP1	100	184	44	54	105	80	110	60	109	33	47	118	85
Morgantown	MGNP1	135	193	42	66	118	57	106	78	98	23	59	117	88
Pottstown Airport*	KPTW	68	164	41	48	93	46	84	66	79	26	80	76	71
Reading Airport*	KRDG	97	132	29	47	105	47	89	131	107	28	74	64	79
West Chester 2 NW	WCHP1	126	156	47	66	126	84	127	92	108	36	45	122	94

*Indicates a station's Period of Record does not cover the period of 1981-2010 but is five years or more. In these cases, the departures from normal values were calculated with normals derived from data spanning the length of the station's Period of Record

Table 7. Status of 2016 precipitation indicators compared to the 30-year normal (1981–2010) at for selected stations.

Precipitation Indicator	West Chester 2 NW 2016	West Chester 2 NW 1981–2010	Glenmoore 2016	Glenmoore 1981–2010
Annual Precipitation	44.8 in 1,138 mm	47.8 in 1,214 mm	41.9 in 1,064 mm	49.0 in 1,245 mm
Autumn (Oct, Nov, Dec) Precipitation	8.3 in 211 mm	12.1 in 307 mm	7.8 in 198 mm	12.0 in 305 mm
Heavy Precipitation Days (days with ≥ 1.0 in (25 mm) rain)	11	14	12	14
Extreme Precipitation Days (days with ≥ 2.0 in (51 mm) rain)	2	1	2	2
Micro-drought (periods of 7+ days without rain)	7	6	9	6
Annual Snowfall	34.0 in 86.4 cm	26.5 in 67.3 cm	34.6 in 87.9 cm	27.8 in 70.6 cm
Measurable Snow Days (days with ≥ 0.1 in (0.3 cm) snow)	11	12	14	13
Moderate Snow Days (days with ≥ 3.0 in (7.6 cm) snow)	3	3	3	3
Heavy Snow Days (days with ≥ 5.0 in (12.7 cm) snow)	2	1	2	1

Table 8. Summary of 2016 monthly total precipitation for selected stations.

Station name	Station	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
Glenmoore	GNMP1	3.6 in	5.9 in	1.8 in	2.2 in	4.4 in	3.3 in	5.5 in	2.4 in	5.0 in	1.4 in	1.8 in	4.7 in	41.9 in
		91 mm	150 mm	46 mm	56 mm	112 mm	84 mm	140 mm	61 mm	127 mm	36 mm	46 mm	119 mm	1,064 mm
Morgantown	MGNP1	4.2 in	5.6 in	1.4 in	2.3 in	4.6 in	2.5 in	5.1 in	3.1 in	4.4 in	1.0 in	2.2 in	4.1 in	40.5 in
		107 mm	142 mm	36 mm	58 mm	117 mm	64 mm	130 mm	79 mm	112 mm	25 mm	56 mm	104 mm	1,029 mm
Pottstown Airport	KPTW	2.1 in	4.6 in	1.5 in	1.9 in	4.0 in	1.8 in	4.2 in	2.7 in	3.7 in	1.1 in	2.8 in	2.8 in	33.1 in
		53 mm	117 mm	38 mm	48 mm	102 mm	46 mm	107 mm	69 mm	94 mm	28 mm	71 mm	71 mm	841 mm
Reading Airport	KRDG	2.9 in	3.3 in	1.0 in	1.8 in	4.4 in	1.8 in	4.0 in	4.7 in	4.6 in	0.9 in	2.6 in	2.1 in	34.2 in
		74 mm	84 mm	25 mm	46 mm	112 mm	46 mm	102 mm	119 mm	117 mm	23 mm	66 mm	53 mm	869 mm
West Chester 2 NW	WCHP1	4.4 in	5.0 in	2.0 in	2.5 in	5.3 in	3.2 in	5.2 in	3.5 in	5.5 in	1.5 in	1.7 in	5.0 in	44.8 in
		112 mm	127 mm	51 mm	64 mm	135 mm	81 mm	132 mm	89 mm	140 mm	38 mm	43 mm	127 mm	1,138 mm
Valley Creek	1473169	3.1 in	5.1 in	1.8 in	2.5 in	3.3 in	1.4 in	4.6 in	3.0 in	4.9 in	1.3 in	3.8 in	3.5 in	38.5 in
		79 mm	130 mm	46 mm	64 mm	84 mm	36 mm	117 mm	76 mm	124 mm	33 mm	97 mm	89 mm	978 mm

Table 9. Top five wettest days and top five dry spells (consecutive days with a trace or less of liquid precipitation) during 2016 from weather observing stations at West Chester 2 NW (WCHP1) and Glenmoore (GNMP1).

Wettest Days in 2016	Dry Spells in 2016
Sept. 30: 2.9 in (74 mm)	Sept. 3 – Sept. 18
Feb. 25: 2.4 in (61 mm)	Aug. 1 – Aug. 12
Dec 1: 2.1 in (53 mm)	Oct. 10 – Oct. 21
Jan 23: 1.6 in (41 mm)	March 5 – March 13
Jan 24: 1.5 in (38 mm)	April 14 – April 22

Drought Status

The monthly Palmer Drought Severity Index (PDSI) indicates that 2016 began somewhat wet but became very dry (Figure 5). According to the PDSI, drought began to occur in October and by November and December, “severe drought” conditions were recorded. Similarly, the Drought Intensity Index shows that from late spring and into fall much of the Commonwealth of Pennsylvania was experiencing “abnormally dry” conditions (Figure 6). Many locations also Pennsylvania experienced moderate to severe drought. Across the Northeast, a number of states experienced severe to extreme droughts during the late summer and fall (Figure 7).




Figure 5. Palmer Drought Severity Index (PDSI) for the Southeastern Piedmont, 2014–2016. Positive values indicate periods of above normal precipitation (0 to 2: near normal; 2 to 3: moderate moisture; >3: very moist) and negative values indicate periods of drought (0 to -2: near normal; -2 to -3: moderate drought; <-3: severe to extreme drought).




Figure 6. Mid-month values of the United States Drought Monitor (DM) - Drought Intensity Index for Pennsylvania in 2016.




Figure 7. Mid-month values of the United States Drought Monitor (DM) - Drought Intensity Index for the Northeast in 2016.

Literature Cited

- Comiskey, J. A., and K. K. Callahan. 2008. Mid-Atlantic Network vital signs monitoring plan. Natural Resource Report NPS/MIDN/NRR—2008/071. National Park Service, Fort Collins, CO.
- Knight P and Others. 2016. Weather of Hopewell Furnace National Historic Site and Valley Forge National Historical Park: Mid-Atlantic Network summary report for 2014. Natural Resource Report. NPS/MIDN/NRR—2016/1113. National Park Service. Fort Collins, Colorado.
- Marshall, M., P. Knight, and J. Comiskey. 2012. Weather and climate monitoring protocol: Eastern Rivers and Mountains Network and Mid-Atlantic Network. Natural Resource Report NPS/ERMN-MIDN/NRR—2012/498. National Park Service, Fort Collins, Colorado.
- National Assessment Synthesis Team. 2001. Climate Change Impacts on United States: The Potential Consequences of Climate Variability and Change. Report for the U.S. Global Change Research Program. Cambridge University Press, Cambridge, UK.
- National Integrated Drought Information System (NIDIS). 2017. National Climate Data Center and United States Department of Agriculture.
<http://droughtmonitor.unl.edu/MapsAndData/MapArchive.aspx> (accessed 07 February 2017).
- National Oceanic and Atmospheric Administration (NOAA). 2017. National Centers for Environmental Information Climatological Rankings. Available at:
<http://www.ncdc.noaa.gov/temp-and-precip/climatological-rankings/> (accessed 23 February 2017).

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS 376/138041, 464/138041, April 2017

National Park Service
U.S. Department of the Interior



Natural Resource Stewardship and Science
1201 Oakridge Drive, Suite 150
Fort Collins, CO 80525

www.nature.nps.gov