Geologic Map of Chiricahua National Monument



NPS Boundary

Geologic Cross Section Lines

Geologic Attitude Observation Localities

- strike and dip of bedding
- strike and dip of foliation
- horizontal foliation
- downthrown side of fault

Geologic Age-Date Localities

- 40Ar/39Ar radiometric sample locality
- K/Ar radiometric sample locality

Linear Geologic Hazard Features

- landslide

High-Angle Fault

- known
- approximate
- concealed

Linear Volcanic Features

- prominent flow layering in Tfre
- trend of bedding in Tfpe
- base of ash cloud deposit in Trcl
- top of surge beds in Trcu

Unconformities

- unconformity, known
- unconformity, concealed

Geologic Point Units

- p — Timp - possible pillow basalt

Linear Geologic Units

- Trcv - Rhyolite Canyon Tuff, virtophyre
- Trca - Rhyolite Canyon Tuff, basal ash
- Tjjf - Jesse James Canyon Tuff, lithic-rich pyroclastic flow
- q - quartz vein

Geologic Contacts

- contact, known
- contact, approximate
- contact, gradational
- map extent

Geologic Units

- Qal - alluvium
- Qaf - alluvial fan deposits
- Qc - colluvium
- QTls - landslide deposits
- Tcg - conglomerate
- Tmt3 - moat deposits, Unit 3 (rhyolite tuff)
- Tmr2 - moat deposits, Unit 2 (rhyolite lava)
- Tmt2 - moat deposits, Unit 2 (tuff)
- Tmr1 - moat deposits, Unit 1 (rhyolite lava)
- Tmt1 - moat deposits, Unit 1 (tuff)
- Tdpl - dacite porphyry lava
- Trco - Outflow facies Rhyolite Canyon Tuff, undivided
- Trcu - Outflow facies Rhyolite Canyon Tuff, upper member
- Trcm - Outflow facies Rhyolite Canyon Tuff, middle member
- Trcl - Outflow facies Rhyolite Canyon Tuff, lower member
- Tjj - Jesse James Canyon Tuff
- Tfre - Rhyolite of Erickson Ridge
- Tfpe - Pyroclastic flow deposits of Erickson Ridge
- Tfrh - Rhyolite of Hands Pass
- Tfph - Pyroclastic flow deposits of Hands Pass
- Tbp - Sedimentary rocks of Bonita Park
- Tjg - Welded tuff of Joe Glen Ranch
- Tim - intermediate and mafic lava flows
- Tims - areas of silicification
- Tr - rhyolite dikes
- Kbls - Bisbee Group, Mural Limestone
- Km - Bisbee Group, Morita Formation
- Kg - Glance Conglomerate
- KJv - volcaniclastic and volcanic rocks
- Pcn - Concha Limestone
- Ps - Scherrer Formation
- Pec - Epitaph Dolomite and Colina Limestone
- Pea - Earp Formation
- Phu - Horquilla Limestone, upper member



Thin surficial units not shown. Thin dotted lines show bedding or flow trends; thicker dotted lines in units Trcu and Trcl are explained in technical description of map units

Figure 12. West to east geologic cross section through Riggs and Sugarloaf Mountains (see figure 11 and map) that shows the distribution of rock layers in the subsurface.

Chiricahua National Monument
Map I-2541, Geologic Section A-A'
Interpretive Map and Guide to the Volcanic Geology of Chiricahua National Monument and Vicinity, Conchise County, Arizona
By John D. Pallister and Edward A. du Bray
U.S. Geological Survey, 1997

MISCELLANEOUS INVESTIGATIONS SERIES
MAP I-2541
Pamphlet accompanies map

EVOLUTION OF THE CHIRICAHUA LANDSCAPE

A. Pre-caldera stage, about 27–32 million years ago



About 28 million years ago, before eruption of the Turkey Creek caldera, the northern Chiricahua Mountains consisted of uplifted masses of Paleozoic and Mesozoic sedimentary rocks (shown in gray and dark green), which were overlain and surrounded by lava and ash flows from two or more active volcanic fields. The volcanoes produced andesite, dacite, and rhyolite of the Faraway Ranch Formation (shown in light green) as well as similar volcanic rocks north of the National Monument (shown in orange). About 27 million years ago, a large body of rhyolite magma (shown in blue) accumulated beneath and uplifted (arrows) the area south of the National Monument.

B. Caldera-forming eruptions, 26.9 million years ago



A series of giant eruptions took place 26.9 million years ago, as the rhyolite magma made its way to the surface and exploded into fragments of pumice and ash. Clouds of ash blanketed a vast area of southern Arizona and New Mexico as hot and dense clouds of pumice, ash, and volcanic gases roared down the flanks of the volcano and filled the Monument valley to a depth of more than 1,600 feet. The eruptions ejected more than 120 cubic miles of rhyolite magma, which is now known as the lower, middle, and upper members of the Rhyolite Canyon Tuff (shown in shades of blue). More than 120 cubic miles of molten rock was erupted, creating a 12-mile-diameter circular depression, or caldera, which also filled with ash and pumice. During the eruption, the caldera floor subsided (arrows). This panel represents the waning stage of the eruptions. During the main phase, the entire region would have been obscured from view by clouds of ash and by storms, which are known to result from volcanic addition of large volumes of ash, heat, and water vapor to the atmosphere.

C. Moat lava eruption, 26.9–26.6 million years ago



Following eruption of the Rhyolite Canyon Tuff, dacite magma (shown in red) uplifted (arrows) the center of the caldera and formed a circular valley, or moat, between the uplifted region and the eroding caldera wall. The dacite magma also erupted and formed lava flows that flowed into the moat, as well as down an ancient channel into the Monument valley. Late during this interval, rhyolite magma erupted again, yielding ash, pumice, and viscous lava flows (shown in yellow), which also accumulated within the caldera moat.

D. Today



Erosion by streams, wind, and landslides has greatly changed the Chiricahua landscape during the past 26 million years. The Turkey Creek caldera has been eroded to the point that it no longer looks like a volcano. The high-standing caldera wall has been eroded away by Pinery Canyon. Several areas that were once valleys or basins, like the National Monument, are now mountains. This "inversion" of topography is the result of more rapid erosion of the older sedimentary rocks than the hard, erosion-resistant rhyolite from the caldera.

The front surface of this block diagram along B-B' represents a cross section through the National Monument along line B-B' on the geologic map, and extending off the map to the south into the adjacent Rustler Park Quadrangle (Pallister and others, 1994). Colors represent the different map units shown on the geologic map, with the following exceptions: all moat rhyolite deposits of the Turkey Creek caldera are generalized in the single color yellow; all Faraway Ranch Formation members are shown as light green, all of the Bisbee Group is shown in dark green, Tertiary volcanic rocks east of the National Monument are shown in orange. Faults are shown in red.

N

0 1 2 Kilometers

0 1 2 Miles

This map graphically presents digital geologic data prepared as part of the NPS Geologic Resources Division's Geologic Resources Inventory. The source map used in creation of the digital geologic data product was:

Pallister, J.S., E. A. Du Bray. 1997. Interpretive Map and Guide to the Volcanic Geology of Chiricahua National Monument and Vicinity, Conchise County, Arizona. Scale 1:24,000. Miscellaneous Investigations Series Map I-2541. U.S. Geological Survey.

Digital geologic data and cross sections for Chiricahua National Monument and all other digital geologic data prepared as part of the Geologic Resources Inventory, are available online at the NPS Data Store: http://science.nature.nps.gov/nrdata/