National Park Service
U.S. Department of the Interior

Natural Resource Stewardship and Science

Grand Canyon National Park

Centennial Paleontological Resource Inventory (Non-Sensitive Version)

Natural Resource Report NPS/GRCA/NRR—2020/2103



ON THE COVER

In situ block of Coconino Sandstone with *Ichniotherium sphaerodactylum* trackways in Grand Canyon National Park. This find was documented in Francischini et al. (2019); it and similar finds are discussed in Chapter 9. Photo by Cassi Knight.

Grand Canyon National Park

Centennial Paleontological Resource Inventory (Non-Sensitive Version)

Natural Resource Report NPS/GRCA/NRR—2020/2103

Vincent L. Santucci[1] and Justin S. Tweet,[2] editors

[1]National Park Service
Geologic Resources Division
1849 "C" Street, NW
Washington, D.C. 20240

[2]National Park Service
9149 79th St. S.
Cottage Grove, Minnesota 55016

March 2020

U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science
Fort Collins, Colorado

The National Park Service, Natural Resource Stewardship and Science office in Fort Collins, Colorado, publishes a range of reports that address natural resource topics. These reports are of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Report Series is used to disseminate comprehensive information and analysis about natural resources and related topics concerning lands managed by the National Park Service. The series supports the advancement of science, informed decision-making, and the achievement of the National Park Service mission. The series also provides a forum for presenting more lengthy results that may not be accepted by publications with page limitations.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner.

This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available in digital format from the Southern Colorado Plateau Inventory & Monitoring Network and the Natural Resource Publications Management website. If you have difficulty accessing information in this publication, particularly if using assistive technology, please email irma@nps.gov.

Please cite this publication as:

Santucci, V. L., and J. S. Tweet, editors. 2020. Grand Canyon National Park: Centennial paleontological resource inventory (non-sensitive version). Natural Resource Report NPS/GRCA/NRR—2020/2103. National Park Service, Fort Collins, Colorado.

NPS 113/167727, March 2020

Contents

Page

Figures xiii

Tables xxv

Appendices xxix

Acknowledgments xxxi

Dedication xxxv

Chapter 1. Introduction and Summary: The Paleontological Heritage of Grand Canyon National Park 1

Literature Cited 10

Chapter 2. History of Paleontological Work at Grand Canyon National Park 11

Up and Down the Long Federal and NGO Trails of Paleontology in Grand Canyon National Park, 1858–2019 11

Introduction 11

Uphill 12

Downhill 16

Laying a Foundation 16

On the Map 17

Making Tracks 18

Digging In 19

Education Flowers 20

The McKeeian Period 24

Super Fossils, Pseudo Fossils 26

Fire 28

About that Ice Age 30

“Toys” Lead to Fossils 31

Fossils Go To School 32

A Final Paleo Note (of a sort) 34

Contents (continued)

Page

Acknowledgments 35

Literature Cited 35

Chapter 3. Stratigraphy of Grand Canyon National Park 45

Introduction 45

Precambrian Stratigraphy of Grand Canyon 50

Grand Canyon Supergroup: Unkar Group 51

Grand Canyon Supergroup: Chuar Group 53

Paleozoic Stratigraphy of Grand Canyon 54

Tonto Group (lower–middle Cambrian) 54

Temple Butte Formation (Middle–Upper Devonian) 57

Redwall Limestone (Lower–Middle Mississippian) 57

Surprise Canyon Formation (Upper Mississippian) 59

Supai Group 60

Pakoon Limestone (lower Permian) 62

Hermit Formation (lower Permian) 62

Coconino Sandstone (lower Permian) 63

Toroweap Formation (lower Permian) 63

Kaibab Formation (lower–middle Permian) 63

Mesozoic Stratigraphy of Grand Canyon 64

Moenkopi Formation (Lower–Middle Triassic) 64

Chinle Formation (Upper Triassic) 65

Conclusions 65

Acknowledgments 66

Literature Cited 66

Chapter 4. Precambrian Paleontology of Grand Canyon National Park 75

Introduction 75

Contents (continued)

Page

History of Research 75
Stratigraphic Distribution of Fossils 77
Paleoproterozoic–Mesoproterozoic basement rocks (upper Paleoproterozoic–lower Neoproterozoic) 78
Unkar Group: Bass Formation (middle–upper Mesoproterozoic) 79
Unkar Group: Hakatai Shale (upper Mesoproterozoic) 80
Unkar Group: Shinumo Quartzite (upper Mesoproterozoic) 81
Unkar Group: Dox Formation (upper Mesoproterozoic) 81
Unkar Group: Cardenas Basalt (upper Mesoproterozoic) 82
Chuar Group: Nankoweap Formation (middle Neoproterozoic: late Tonian age) 82
Chuar Group: Galeros Formation (middle Neoproterozoic: late Tonian age) 83
Chuar Group: Kwagunt Formation (middle Neoproterozoic: late Tonian age) 83
Sixtymile Formation (lower Cambrian) 84
Types of Fossils 84
Stromatolites and Other Microbial Structures 84
Acritarchs and Other Organic-Walled Microfossils 86
Vase-Shaped Microfossils 87
Other Fossils 88
Notable Pseudofossils and Dubiofossils 89
Acknowledgments 91
Literature Cited 91
Appendix 4-A. Precambrian Taxa From GRCA 97
Taxonomic Notes 105
Appendix 4-B. Precambrian Taxa Named From GRCA 106
Chapter 5. Paleozoic Invertebrate Paleontology of Grand Canyon National Park 109
Introduction 109

Contents (continued)

Page

Methods 111

Invertebrate Body Fossils 111

Stratigraphic Distribution of Fossils 113

Cambrian Fauna 113

Devonian Fauna 122

Mississippian Fauna 123

Pennsylvanian Fauna 133

Permian Fauna 136

Summary of the Paleozoic Invertebrate Body Fossils 147

Conclusions 147

Acknowledgments 148

Literature Cited 148

Appendix 5-A. Grand Canyon (GRCA) Paleozoic Invertebrate Specimens List 159

Taxonomic Uncertainty 205

Taxonomic Notes 206

Appendix 5-B. Paleozoic Invertebrate Body Fossils From GRCA From the Literature 209

Appendix 5-C. Paleozoic Invertebrate Body Fossil Taxa Named From GRCA 230

Appendix 5-D. Paleozoic Invertebrate Body Fossil Taxa Potentially Named From GRCA 235

Chapter 6. Paleozoic Vertebrate Paleontology of Grand Canyon National Park: Research History, Resources, And Potential 237

Introduction 237

Devonian 238

Temple Butte Limestone 238

Mississippian 241

Redwall Limestone 241

Contents (continued)

Page

Surprise Canyon Formation....242
Pennsylvanian....245
Supai Group....245
Watahomigi Formation....246
Wescogame Formation....247
Permian....248
Pakoon Limestone....248
Kaibab Formation....248
Conclusions....252
Acknowledgments....253
Literature Cited....253
Chapter 7. Paleozoic Paleobotany of Grand Canyon National Park....257
Introduction....257
History of Research....257
Stratigraphic Distribution of Fossils....259
Surprise Canyon Formation....259
Supai Group....262
Hermit Formation....263
Acknowledgments....270
Literature Cited....270
Appendix 7-A. Paleobotanical Type Specimens....274
Chapter 8. Paleozoic Invertebrate Ichnology of Grand Canyon National Park....277
Introduction....277
Trace Fossil Classifications....278
Stratigraphic Distribution of Trace Fossils....280
Cambrian....283

Contents (continued)

Page

Tonto Group 283
Tapeats Sandstone 284
Bright Angel Shale 288
Muav Limestone 304
Frenchman Mountain Dolostone 305
Devonian 305
Temple Butte Formation 305
Mississippian 306
Redwall Limestone 306
Surprise Canyon Formation 307
Pennsylvanian 307
Supai Group 307
Watahomigi Formation 308
Manakacha Formation 308
Wescogame Formation 308
Permian 309
Esplanade Sandstone 309
Hermit Formation 310
Coconino Sandstone 314
Toroweap Formation 321
Kaibab Formation 321
Conclusions 323
Acknowledgments 324
Literature Cited 324
Chapter 9. Paleozoic Vertebrate Ichnology of Grand Canyon National Park 333
Introduction 333

Contents (continued)

Page

Vertebrate Ichnology 334

History of Track Collection in Grand Canyon and Mogollon Rim Areas 336

Stratigraphic Distribution of Tracks 337

Manakacha Formation 339

Wescogame Formation 341

Hermit Shale 346

Coconino Sandstone 359

Conclusions 372

Acknowledgments 373

Literature Cited 373

Chapter 10. Mesozoic Paleontology of Grand Canyon National Park: Trace Fossils, Stratigraphy, and Regional Correlations 381

Introduction 381

Overview of Triassic Strata at GRCA 383

Moenkopi Formation 383

Chinle Formation (Shinarump Member) 388

Field Methods 390

Results 391

Moenkopi Formation 391

Chinle Formation 395

Conclusions 395

Acknowledgments 396

Literature Cited 396

Chapter 11. Pleistocene/Holocene Cave Fossils From Grand Canyon National Park 403

Ice Age (Pleistocene) Flora, Fauna, Environments, And Climate Of The Grand Canyon, Arizona 403

Contents (continued)

Page

Introduction 403
Preservation Scenarios (Taphonomy) 405
River Corridor Sediments 405
Dry Caves 405
Packrat Middens 406
Taxonomic Groups 408
Plants 408
Invertebrates 409
Vertebrates 409
Select Localities 411
Stanton's Cave and Skull Cave 411
Sandblast Cave and Nearby Caves 412
Little Nankoweap 413
Hance Canyon and Horseshoe Mesa 415
Cremation Creek Caves 416
Double Bopper Cave 417
Rampart, Muav, and Vulture Caves 420
Environments and Climate Discussion 423
Acknowledgments 425
Literature Cited 425
Appendix 11-A. Grand Canyon Ice Age Taxa 433
Plants 433
Invertebrates 448
Vertebrates 452
Ichnofossils and Reproductive Traces 462
Other Fossils 463

Contents (continued)

Page

Chapter 12. Grand Canyon National Park Paleontological Resources Management and Protection 465

Introduction 465

GRCA Paleontological Resource Inventory 465

GRCA Paleontological Resource Monitoring 467

GRCA Paleontological Resource Protection 468

History of Paleontological Resource Protection at GRCA 469

Paleontological Resource Protection Recommendations 472

GRCA Paleontological Resource Research and Permits 473

Internal Permits 473

Previous Research Permits 473

Current Permits 474

GRCA Paleontological Resource Curation and Museum Collections 474

Scope of Paleontological Collections 474

Notable Contributors to Paleontological Collections 475

Paleontological Collections in Outside Repositories 475

Photographic Archives 476

Digital Archives 477

Outreach 477

GRCA Paleontological Resource Interpretation, Education, and Outreach 477

Current Long Range Interpretive Plan 478

Current Paleontological Interpretive Programs 478

Interpretive Signs and Exhibits 488

Interpretive Staff Training and Resources 493

GRCA Paleontological Resource Data Management 494

GRCA Paleontology Archives 495

Contents (continued)

Page

NPS Paleontology Program Archives (WASO) 495

McKee and Walcott Archives 495

Geospatial Database 495

Conclusions 498

Literature Cited 499

Appendix 12-A. GRCA Paleontology Research Permits 1999–2019 501

Appendix 12-B. GRCA Photogrammetry 3D Models 503

Figures

Page

Figure 1-1. Map of Grand Canyon National Park (NPS/MARK NEBEL). 2

Figure 1-2. 2019 National Fossil Day logo featuring Pleistocene fauna from Rampart Cave in Grand Canyon National Park (NPS). 3

Figure 1-3. Notable paleontologists who worked in areas now administered as Grand Canyon National Park, including (from left to right): Charles Doolittle Walcott (USNM photo); John C. Merriam (NPS); Charles Gilmore (USNM); Remington Kellogg (USNM); David White (NPS); and Edwin McKee (USGS). 5

Figure 1-4. Paleontologist Charles Doolittle Walcott explored the Grand Canyon's Precambrian and early Paleozoic strata for traces of early life (NPS). 6

Figure 1-5. Grand Canyon National Park Fossil Trading Card featuring the fossil shark *Amaradontella santuccii* from the Mississippian Surprise Canyon Formation (NPS). 6

Figure 1-6. Civilian Conservation Corps (CCC) constructing the in situ Fossil Fern Exhibit along the South Kaibab Trail during 1937 (NPS). 7

Figure 1-7. In situ block of Coconino Sandstone with *Ichniotherium sphaerodactylum* trackways in Grand Canyon National Park (NPS/CASSI KNIGHT) 8

Figure 1-8. Fossil condor skull from Stevens Cave, Grand Canyon National Park (STEVE EMSLIE). 9

Figure 2-1. The steamboat *Explorer* on the lower Colorado River; the first field trip en route to Grand Canyon (Ives 1861: frontispiece). 13

Figure 2-2. Grand Canyon paleontology begins here, April 1858 (Newberry 1861: Paleontology Plate 2) 15

Figure 2-3. David White (left) and John C. Merriam at a fossil fern quarry in Grand Canyon (note the slab leaning against White's right leg) (NPS). 21

Figure 2-4. Trailside exhibit of fossil ferns, constructed by the Civilian Conservation Corps in the Permian Hermit Shale at Cedar Ridge on South Kaibab Trail, as seen *ca.* 1940s? (NPS). 22

Figure 2-5. Ranger-Naturalist Edwin D. McKee at work in Yavapai Observation Station, *ca.* 1930 (NPS). 23

Figure 2-6. Fossil footprints in a slab of Coconino Sandstone, displayed on the Trail of Time 24

Figures (continued)

Page

Figure 2-7. *Castrocopros martini* Hunt and Lucas, the dung of *Nothrotheriops shastensis* (Sinclair), in Rampart Cave, September 1938 (NPS). 29

Figure 2-8. Skull of the extinct mountain goat, *Oreamnos harringtoni* Stock, from a Grand Canyon cave (NPS). 30

Figure 3-1. Grand Canyon stratigraphy and structural relations (Billingsley et al. 2019: Figure 2) 46

Figure 3-2. Mesoproterozoic rocks of the Unkar Group (Grand Canyon Supergroup) in eastern Grand Canyon 51

Figure 3-3. Mesoproterozoic rocks in contact with lower Tonto Group (Tapeats Sandstone and Bright Angel Shale) in eastern Grand Canyon (Billingsley et al. 2019: Figure 4). 52

Figure 3-4. Upper Tonto Group (Cm=Muav Limestone; Cu="undifferentiated dolomites", now the Frenchman Mountain Dolostone), Temple Butte Formation (Dtb), and Redwall Limestone (Mrw=Whitmore Wash Member; Mrt=Thunder Springs Member; Mrm=Mooney Falls Member) in eastern Grand Canyon (Billingsley et al. 2019: Figure 5). 55

Figure 3-5. Redwall Limestone (Mr), Surprise Canyon Formation (Ms), and overlying Supai Group (Pwa=Watahomigi Formation; Pm=Manakacha Formation; Pwe=Wescogame Formation; Pe=Esplanade Sandstone) (Billingsley et al. 2019: Figure 6). 58

Figure 3-6. Uppermost Supai Group (Pe=Esplanade Sandstone), Hermit Formation (Ph), Coconino Sandstone (Pc), Toroweap Formation (Seligman Member=Pts; Brady Canyon Member=Ptb; Woods Ranch Member=Ptw), and Kaibab Formation (Fossil Mountain Member=Pkf; Harrisburg Member=Pkh) (Billingsley et al. 2019: Figure 7). 62

Figure 3-7. Mesozoic rocks in eastern GRCA at Cedar Mountain (NPS/DIANA BOUDREAU). 65

Figure 4-1. *Chuaria circularis* depicted in Walcott (1899: Plate 27:12). *C. circularis* specimens are up to 5 mm (0.2 in) across, making them the largest of GRCA's Precambrian body fossils. 76

Figure 4-2. An aerial view showing several topographic features west of the Colorado River in eastern GRCA 77

Figure 4-3. Bass Limestone stromatolites (not in life position) (NPS/CASSI KNIGHT). 79

Figures (continued)

Page

Figure 4-4. A mud-cracked surface in the Hakatai Shale 80

Figure 4-5. Left: Dox Formation “fucoids” as shown in McKee (1932); the original caption is “Peculiar casts and moulds in Dox Formation 82

Figure 4-6. Another view of a Bass Limestone stromatolitic horizon just below the Hakatai Shale (NSP/CARL BOWMAN). 85

Figure 4-7. Examples of Alf’s circular structures from the Bass–Hakatai transition interval (RAYMOND M. ALF MUSEUM/DON LOFGREN). 90

Figure 5-1. Grand Canyon view from South Rim Visitors Center with Paleozoic stratigraphic units marked within blue and orange band 110

Figure 5-2. Cambrian trilobite fragments of *Solenopleurella porcata* (USNM 108626; now *Spencella porcata* Rasetti 1963) from undifferentiated Tonto Group, possibly the Muav Limestone. Scale bar 1 cm (0.4 in) (NPS/ANNE MILLER). 114

Figure 5-3. Trilobite fragments including olenellid genal spines of unidentified species, and an olenellid pygidium boxed in red from the Cambrian Bright Angel Shale (GRCA 6598). Scale bar 1 cm (0.4 in) (LINDA S. LASSITER). 115

Figure 5-4. The Cambrian trilobites (GRCA 9529) *Amecephalus althea* (A) and a dolichometopid (B) from the Bright Angel Shale 116

Figure 5-5. Examples of common, well-preserved Bright Angel Shale ptychopariid trilobites 117

Figure 5-6. Bradoriids from the Bright Angel Shale of GRCA. Scale bar 1 cm (0.4 in) (FREDERICK SUNDBERG). 117

Figure 5-7. The phosphatic orthid brachiopod *Wimanella* (GRCA 8423) preserved mostly as outlines of the valves in a slab from the Bright Angel Shale 118

Figure 5-8. An eocrinoid, *Gogia longidactylus* (GRCA 2641), from the Bright Angel Shale 119

Figure 5-9. An illustration of *Tontoia kwaguntensis* (USNM 57660), which is described as a 25 mm (1 in) long animal of trilobite or other arthropod affinity (Walcott 1912b: Plate 24). 120

Figure 5-10. Two hyoliths from the Bright Angel Shale (GRCA numbers uncertain). 121

Figures (continued)

Page

Figure 5-11. An unidentified brachiopod with fine costae like the common *Derbyia* (Strophomenata: Orthotetida) on a fossiliferous slab from the Redwall Limestone (GRCA 11065) 124

Figure 5-12. A coral from the Redwall Limestone of GRCA (GRCA 4470) with poor preservation for clear diagnosis beyond phylum 126

Figure 5-13. Tabulate phaceloid coral (*Syringopora* sp.) from the Redwall Limestone of GRCA showing a massive form with individual corallites (GRCA 11055) 127

Figure 5-14. Gastropods showing poor preservation common in fossils from the Redwall Limestone of the families Euomphalidae (GRCA 7641) **(A)** and Bellerophontidae (GRCA 7628) **(B)** 128

Figure 5-15. A composite sample of unidentified brachiopods, crinoids and coral fragments from the Surprise Canyon Formation (GRCA 57636) 131

Figure 5-16. Fragments of the coral *Michelinia* (Favositida) from the Surprise Canyon Formation (GRCA 7647) 132

Figure 5-17. Examples of brachiopods and bivalves from the Supai Group (GRCA 11326) including prominent examples of *Anthracospirifer tanoensis* (center and right), with the bivalves *Promytilus* and *Permophorus* reported as present as well 134

Figure 5-18. Supai Group brachiopods with reasonable preservation (LINDA S. LASSITER) 135

Figure 5-19. Hermit Formation insect wing of *Tupus whitei* (GRCA 3090) 137

Figure 5-20. Toroweap Formation slab with unidentified bivalves (GRCA 15324) 138

Figure 5-21. Example of impressions of fenestrate-type bryozoan colonies (GRCA 11373) from the Toroweap Formation 139

Figure 5-22. Kaibab Limestone productids from GRCA (LINDA S. LASSITER). **A.** *Peniculauris bassi* (GRCA 21372) 141

Figure 5-23. An unidentified fenestrate bryozoan from the Kaibab Formation 142

Figure 5-24. Conulariid *Conularia kaibabensis* holotype from the Kaibab Formation (USNM 102289) 143

Figure 5-25. Crinoid stems and individual columnals found in the Kaibab Formation (GRCA 11468) 144

Figure 5-26. A large bivalve of the Kaibab Formation, *Allorisma capax* (GRCA 13501) 145

Figures (continued)

Page

Figure 5-27. A common myalinid bivalve, *Aviculopecten* (GRCA 3621), from the Kaibab Formation 145

Figure 5-28. A specimen of the scaphopod *Prodentalium canna* (GRCA 14373) showing the conical shape and small size 146

Figure 5-29. An unidentified sponge from the Kaibab Formation (GRCA 3590) 147

Figure 6-1. Antiarch placoderm, *Bothriolepis*, fossils from the Temple Butte Limestone, Grand Canyon (JOHN-PAUL HODNETT) 240

Figure 6-2. Chondrichthyan fossils from the Mooney Fall Member, Redwall Limestone (JOHN-PAUL HODNETT) 242

Figure 6-3. Chondrichthyans from the Surprise Canyon Formation collected from Grand Canyon National Park (JOHN-PAUL HODNETT) 245

Figure 6-4. Fish fossils from the Watahomigi Formation (JOHN-PAUL HODNETT) 247

Figure 6-5. Chondrichthyan fossils collected from the Kaibab Formation on the North Rim of Grand Canyon National Park (JOHN-PAUL HODNETT) 250

Figure 6-6. Kaibab vertebrate fossils in GRCA collections (JOHN-PAUL HODNETT) 251

Figure 7-1. A. Paleobotanist David White (left) and John C. Merriam working in the Cedar Ridge fossil quarry during field work in the summer of 1927 (NPS) 258

Figure 7-2. *Supaia* specimens (CASSI KNIGHT) 266

Figure 7-3. *Yakia heterophylla* (CASSI KNIGHT) 266

Figure 7-4. Various taxa (CASSI KNIGHT) 268

Figure 8-1. *Rusophycus* in convex hyporelief within the Tapeats Sandstone, central Grand Canyon (NPS/ANNE MILLER). 286

Figure 8-2. *Cruziana* isp. from the Bright Angel Shale preserved in convex hyporelief (NPS/GRCA). 290

Figure 8-3. In situ *Monomorphichnus* aff. *lineatus* var. *giganticus* preserved in convex hyporelief from the lower Bright Angel Shale (NPS/ANNE MILLER) 291

Figure 8-4. *Rusophycus* isp. from the Bright Angel Shale preserved in convex hyporelief (NPS/ANNE MILLER). 292

Figure 8-5. *Palaeophycus* isp. from the Bright Angel Shale preserved in convex hyporelief (NPS/ANNE MILLER). 293

Figures (continued)

Page

Figure 8-6. "*Scalarituba novum*" (GRCA 76901) collected by Eben Rose in September 1998.....294

Figure 8-7. *Treptichnus pedum* (formerly *Phycodes pedum*) from the Bright Angel Shale preserved in convex hyporelief (NPS/ANNE MILLER).....295

Figure 8-8. *Teichichnus* isp. from the Bright Angel Shale showing horizontal spreiten and tapering at one end (NPS/ANNE MILLER).....296

Figure 8-9. In situ *Teichichnus* from the lower Bright Angel Shale (NPS/ANNE MILLER).....297

Figure 8-10. *Bergaueria* isp. from the Bright Angel Shale (NPS/ANNE MILLER).....298

Figure 8-11. *Diplocraterion* isp. from the Bright Angel Shale showing vertical spreiten between limbs indicating downward shifting in the sediment by the tracemaker (NPS/ANNE MILLER).....299

Figure 8-12. Invertebrate traces from the Wescogame Formation (LORENZO MARCHETTI).....309

Figure 8-13. Arthropod trackways from the Hermit Formation of the Mogollon Rim area (LORENZO MARCHETTI).....311

Figure 8-14. *Scoyenia gracilis* (GRCA 8538) from the Hermit Formation (White 1929).....312

Figure 8-15. *Walpia hermitensis* syntypes from the Hermit Formation (White 1929) (SPENCER LUCAS).....313

Figure 8-16. Arthropod trackways from the Coconino Sandstone, attributed to *Paleohelcura tridactyla* (LORENZO MARCHETTI).....316

Figure 8-17. Arthropod trackways from the Coconino Sandstone (LORENZO MARCHETTI).....317

Figure 8-18. Bilobate trails and paired grooves from the Coconino Sandstone.....319

Figure 8-19. Complex, actively filled horizontal structures from the Coconino Sandstone (LORENZO MARCHETTI).....320

Figure 8-20. Worm trail specimen in the upper Kaibab Formation collected by Everette Helman in June 1936 (GRCA 10044).....323

Figure 9-1. Track formation and track measurements (LORENZO MARCHETTI & SPENCER LUCAS).....335

Figures (continued)

Page

Figure 9-2. Historic photos of in situ vertebrate tracks from the Coconino Sandstone at GRCA. 337

Figure 9-3. Localities and geological setting (LORENZO MARCHETTI & SPENCER LUCAS) 338

Figure 9-4. Permian stratigraphy of the Grand Canyon. Panoramic view looking west at the wall of Hermit Canyon showing the track-bearing Permian units (LORENZO MARCHETTI & SPENCER LUCAS). 339

Figure 9-5. Tetrapod footprints from the Manakacha Formation (LORENZO MARCHETTI & SPENCER LUCAS) 340

Figure 9-6. Anamniote footprints from the Wescogame Formation (LORENZO MARCHETTI & SPENCER LUCAS) 342

Figure 9-7. Reptile footprints from the Wescogame Formation (LORENZO MARCHETTI & SPENCER LUCAS) 343

Figure 9-8. Stratigraphic section of the Hermit Shale, measured by Noble (1922) near Bass Point in the Grand Canyon (LORENZO MARCHETTI & SPENCER LUCAS). 347

Figure 9-9. Tetrapod footprints from the Hermit Shale along Hermit Trail in USNM 11707A (LORENZO MARCHETTI & SPENCER LUCAS) 349

Figure 9-10. Anamniote footprints from the Hermit Shale (LORENZO MARCHETTI & SPENCER LUCAS) 350

Figure 9-11. Synapsid footprints from the Hermit Shale (LORENZO MARCHETTI & SPENCER LUCAS) 351

Figure 9-12. Reptile footprints from the Hermit Shale (LORENZO MARCHETTI & SPENCER LUCAS) 352

Figure 9-13. Undetermined footprints from the Hermit Shale (LORENZO MARCHETTI & SPENCER LUCAS) 354

Figure 9-14. Stratigraphic section of the Coconino Sandstone measured along the Hermit Trail (LORENZO MARCHETTI & SPENCER LUCAS). 360

Figure 9-15. Tetrapod footprints from the Coconino Sandstone (LORENZO MARCHETTI & SPENCER LUCAS) 361

Figure 9-16. Anamniote footprints from the Coconino Sandstone (LORENZO MARCHETTI & SPENCER LUCAS) 363

Figures (continued)

Page

Figure 9-17. Synapsid footprints from the Coconino Sandstone, Hermit Trail (LORENZO MARCHETTI & SPENCER LUCAS) 365

Figure 9-18. Reptile footprints from the Coconino Sandstone (LORENZO MARCHETTI & SPENCER LUCAS) 366

Figure 9-19. Stratigraphic distribution of vertebrate tracks in the Paleozoic formations of GRCA (LORENZO MARCHETTI & SPENCER LUCAS) 373

Figure 10-1. Geographic location and stratigraphic section of Cedar Mountain, Grand Canyon National Park (NPS/MARSH ET AL.) 382

Figure 10-2. GRCA 33186, *Rotodactylus* isp. from the Wupatki Member, Moenkopi Formation at Cedar Mountain (NPS/MARSH ET AL.) 387

Figure 10-3. Plant fossils from the Shinarump Member of the Chinle Formation at Cedar Mountain (NPS/MARSH ET AL.) 389

Figure 10-4. Mud cracks superposed on bi-directional ripples in a sandstone from the Moqui Member of the Moenkopi Formation at Cedar Mountain 392

Figure 10-5. Hand sample of orthogonal ripples in alternating layers of laminated sandstone from the Holbrook Member of the Moenkopi Formation at Cedar Mountain (NPS/MARSH ET AL.). 393

Figure 10-6. Internal molds of burrows found ex situ in the Holbrook Member of the Moenkopi Formation at Cedar Mountain (NPS/MARSH ET AL.). 393

Figure 10-7. Hand samples of *Scoyenia* cf. *gracilis* found in the Holbrook Member of the Moenkopi Formation at Cedar Mountain (**A–D**) (NPS/MARSH ET AL.) 395

Figure 11-1. Steve Emslie (right) and Larry Coats (left) excavating a condor (*Gymnogyps*) nest and skeletal remains reclaimed by a packrat and made into a midden, Sandblast Cave, 1984 (EMILEE MEAD). 407

Figure 11-2. A juvenile extinct Harrington's mountain goat (*Oreamnos harringtoni*) skull in a packrat midden, Stevens Cave, 1984 (EMILEE MEAD). 408

Figure 11-3. A mummified myotid bat at its last perch among the gypsum crystals in Double Bopper Cave (NPS/SHAWN THOMAS). 410

Figure 11-4. Ascending into Skylight Cave, 1984 (EMILEE MEAD). 411

Figure 11-5. Close-up of the entrance to Sandblast Cave showing the series of openings, 1984 (EMILEE MEAD). 413

Figures (continued)

Page

Figure 11-6. Stevens Cave entrance located high on the cliff face of the Redwall Limestone, 1984 (EMILEE MEAD).....414

Figure 11-7. An expansive flat region exists above the Redwall Limestone caves in the Little Nankoweap Canyon region 415

Figure 11-8. Entrance to Kaetan Cave, 1980 (JIM MEAD). 417

Figure 11-9. The two main entrances to Double Bopper Cave.....418

Figure 11-10. A mummified ringtail (*Bassariscus astutus*) from Double Bopper Cave 419

Figure 11-11. An articulated gray fox skeleton (not radiocarbon dated) from Double Bopper Cave (NPS/SHAWN THOMAS). 420

Figure 11-12. Rampart Cave in the 1930s 422

Figure 11-13. Inferred elevational distribution of plant communities within and adjacent to the Grand Canyon during the most recent 50,000 years.....425

Figure 12-1. The GRCA paleontology team documenting a new fossil track locality in the Coconino Sandstone during the 2019 PaleoBlitz (Left to Right: Klara Widrig, Anne Miller, Diana Boudreau, and Jered Hansen) (NPS/MARK NEBEL). 467

Figure 12-2. Track blocks have been chiseled out and removed from the Coconino Sandstone along Hermit Trail (left) 469

Figure 12-3. A nail was used by a visitor attempting to pry this brachiopod fossil from the Kaibab Formation (NPS/VINCENT SANTUCCI). 470

Figure 12-4. Scratch marks on the Fossil Fern Exhibit structure along the South Kaibab Trail in 2017 (NPS/ANNE MILLER).....471

Figure 12-5. Photographs from 2013 **(A)** and 2017 **(B)** of the same fossil fern impressions near the exhibit showing vandalism that occurred during Spring Break in 2017 (NPS).....471

Figure 12-6. Rock and fossil specimens found in suspect's backpack after GRCA law enforcement apprehended them removing specimens from park land in 2019 (NPS). 472

Figure 12-7. Entrance portal along the Trail of Time near the Grand Canyon Headquarters building (NPS/DIANA BOUDREAU).....479

Figure 12-8. Crinoid stems, stick and lacy bryozoans, and shell fragments from exposures of the Kaibab Formation can be found along the rim of the canyon (NPS/DIANA BOUDREAU). 480

Figures (continued)

Page

Figure 12-9. Students identify fossils in the Kaibab Formation at the fossil beds during the "Stories in Stone" field trip (NPS/RONNIE COLVIN). 481

Figure 12-10. Park Rangers talk with the public about fossils at the 2016 National Fossil Day celebration at Grand Canyon National Park (NPS). 483

Figure 12-11. NPS paleontologists and educators who assisted with the 10th Annual National Fossil Day activities and programs on the South Rim of Grand Canyon National Park on September 28, 2019 484

Figure 12-12. Advertisement for the four speakers at the Paleontology Symposium on Friday September 27, 2019 at the Shrine of the Ages (NPS). 485

Figure 12-13. Advertisement for the National Fossil Day Celebration event at the South Rim of Grand Canyon National Park (NPS). 486

Figure 12-14. Speakers at the Welcome Ceremony at Mather Amphitheater at the National Fossil Day Celebration on September 28, 2019 487

Figure 12-15. A large pop-up exhibit highlighting Grand Canyon National Park fossils was developed for the 2019 National Fossil Day Celebration at Grand Canyon (NPS/DIANA BOUDREAU). 488

Figure 12-16. Paleozoic exhibit at the Yavapai Geology Museum (NPS/DIANA BOUDREAU). 489

Figure 12-17. Two large chunks of petrified wood used as building materials for the wall along the Rim Trail outside the Verkamp's Visitor Center on the South Rim (NPS/DIANA BOUDREAU). 489

Figure 12-18. A. The geology fireplace in the Bright Angel Lodge designed by Mary Colter exhibits geology and paleontology of the Grand Canyon (NPS/MICHAEL QUINN) 490

Figure 12-19. A coiled nautiloid found in the Kaibab Formation during construction of the Desert View Watchtower on display at Desert View (NPS/SALLY CARTTAR). 491

Figure 12-20. Fossil Fern Exhibit at Cedar Ridge on South Kaibab Trail in the 1930s **(A)** and in the spring of 2019 **(B)** (NPS). 492

Figure 12-21. Teacher workshop at the fossil beds site along the Rim Trail (NPS). 493

Figure 12-22. Screenshot from ArcCatalog showing the structure of the GRCA Paleontology geodatabase (NPS). 496

Figures (continued)

Page

Figure 12-23. Screenshot from ArcMap showing an example of a Paleontology Locality with associated Sites and Photopoints (NPS). 497

Tables

Page

Table 3-1. Grand Canyon area stratigraphy (after Billingsley et al. 2019: 47

Table 3-2. Overview of GRCA stratigraphy and paleontology 48

Table 4-1. Summary of GRCA Precambrian stratigraphy, fossils, and depositional settings in descending order of age, from youngest to oldest; Sixtymile Formation included based on historical classification 78

Appendix Table 4-A-1. Unkar Group fossil taxa 99

Appendix Table 4-A-2. Chuar Group fossil taxa 100

Appendix Table 4-B-1. Precambrian taxa named from specimens found within GRCA. 106

Table 5-1. The number of Paleozoic invertebrate body fossils of GRCA by phylum as reported from the literature or as listed in the museum collections included in this report 112

Table 5-2. Foraminifera from the Redwall Limestone of the Grand Canyon in USNM collections. 128

Appendix Table 5-A-1. Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other) 160

Appendix Table 5-A-2. Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other) 173

Appendix Table 5-A-3. Specimens of invertebrate body fossils from the Supai Group of GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other) 186

Appendix Table 5-A-4. Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other) 190

Appendix Table 5-B-1. Cambrian invertebrate taxa reported from GRCA localities in the literature 210

Appendix Table 5-B-2. Devonian and Mississippian invertebrate taxa reported from GRCA localities in the literature. 216

Tables (continued)

Page

Appendix Table 5-B-3. Supai Group invertebrate taxa reported from GRCA localities in the literature 222

Appendix Table 5-B-4. Post-Supai Group Permian invertebrate taxa reported from GRCA localities in the literature 225

Appendix Table 5-C-1. Fossil invertebrate taxa named from specimens collected from GRCA 231

Appendix Table 5-D-1. Fossil invertebrate taxa named from specimens possibly found within GRCA 236

Table 7-1. List of all pteridosperm species reported in White (1929a), including locality information 265

Table 7-2. List of all conifer species reported in White (1929a), including locality information 267

Table 7-3. Fruits of uncertain affinities reported by David White (1929a) with locality information. 269

Appendix Table 7-A-1. Table of all known paleobotanical type specimens from GRCA, including name and specimen numbers 274

Table 8-1. List of valid ichnogenera and their respective architectural designs (sensu Buatois et al. 2017), ordered by geological unit (from lower to upper) 281

Table 8-2. List of invalid ichnotaxa, ordered by geological unit (from lower to upper) 282

Table 8-3. List of collected trace fossil specimens from the Bright Angel Shale in Grand Canyon, housed in the GRCA Museum Collections 300

Table 9-1. Museum collections, Wescogame Formation. 345

Table 9-2. Museum collections, Hermit Shale. 356

Table 9-3. Museum collections, Coconino Sandstone. 368

Appendix Table 11-A-1. Ice Age plant taxa reported from the Grand Canyon 433

Appendix Table 11-A-2. Ice Age invertebrate taxa reported from the Grand Canyon. 448

Appendix Table 11-A-3. Ice Age vertebrate taxa reported from the Grand Canyon. 452

Appendix Table 11-A-4. Ice Age ichnofossils and reproductive traces reported from the Grand Canyon 462

Tables (continued)

Page

Appendix Table 11-A-5. Other Ice Age fossil taxa reported from the Grand Canyon 463

Appendix Table 12-A-1. List of paleontological research permits issued from 1999 to 2019..... 501

Appendix Table 12-B-1. GRCA specimens which have been documented as 3D photogrammetry models. 503

Appendix Table A-1. Fossil taxa named from specimens found within GRCA. 506

Appendix Table B-1. Fossil taxa named from specimens possibly found within GRCA. 521

Appendix Table C-1. Cambrian fossil taxa 527

Appendix Table C-2. Devonian and Mississippian fossil taxa 538

Appendix Table C-3. Supai Group fossil taxa 551

Appendix Table C-4. Post-Supai Permian fossil taxa 557

Appendices

Page

Appendix A. Fossil Taxa Named From GRCA 505
Appendix B. Fossil Taxa Potentially Named From GRCA 521
Appendix C. Stratigraphic Tables of GRCA Paleozoic Taxa 525
Appendix D. Outside Repositories of GRCA Fossils 591
Appendix E. Paleontological Resource Law and Policy 595
Appendix F. Geologic Time Scale 603

Acknowledgments

The Grand Canyon National Park Centennial Paleontological Resource Inventory was made possible through the collaboration, contributions and support of an extensive multidisciplinary team. Although the 100th anniversary of GRCA was an exceptionally busy year at the park, we received exceptional support from the park's leadership and staff throughout this project. Central to the decision-making in undertaking the various paleontology-focused projects and events during 2019 at GRCA was Jeanne Calhoun, Chief of Science & Resource Management. Planning for the GRCA Paleontological Resource Inventory and associated activities was initiated between Jeanne Calhoun and Vincent L. Santucci, NPS Senior Paleontologist during the summer of 2018. We are grateful for the enthusiastic support for this project from Jeanne, Deputy Chief of Science & Resource Management Sarah Haas, Acting Superintendent Woody Smeck, Deputy Superintendent Lisa Carrico, and Jan Balsom, Senior Adviser for Stewardship and Tribal Programs.

A long list of GRCA staff played a variety of roles during the planning and implementation of the paleontology-focused work at the park, especially the National Fossil Day Celebration. We extend our appreciation to members of the GRCA Interpretation Staff, Education and Creative Media, including: Laurel Brierly, Sharon Cawley, Vanessa Ceja-Cervantes, Jeremy Childs, Lillian Daniels, Celia Dubin, Jennifer Eberlein, Kevin Garcia, Joel Kane, Grace Lilly, Bryan Maul, Michael Quinn, Warren Rudkin, Todd Stoeberl, Chris Symons, and John Wishart. A number of GRCA staff provided assistance, technical support and information to the paleontological resource inventory, including: Joelle Baird, Santiago Garcia, Jered Hansen, Edward "Ted" McClure, and Eugenia Sullivan.

This project benefited through the work of the previous GRCA Physical Science Program Manager Deanna Greco and previous GRCA Deputy Chief of Science and Resource Management Jane Rodgers, who helped secure funding for a number of paleontology interns, including: Jeff Dobbins, Taylor Hartman, Cassi Knight, and James Super. Each of the interns assisted with compiling baseline paleontological resource data for GRCA during their assignments at the park and shared information that is presented in this report.

Important behind-the-scenes work for this project was put forth by staff from the NPS Geologic Resources Division (GRD), and we extend our thanks to Hal Pranger, Lima Soto, Paige Lambert, Jim Wood, Jack Wood, Jason Kenworthy and John Graham. Additional thanks is extended to staff from the NPS Office of Public Affairs, including Jeff Olson, Kathy Kupper, and Victoria Stauffenberg for helping to share information about this project with the public and media.

One of the components of the GRCA Paleontological Resource Inventory was the GRCA PaleoBlitz. The GRCA PaleoBlitz was a field-based inventory for paleontological resources by teams of specialist and subject matter experts in various disciplines of paleontology (e.g. fossil invertebrates, vertebrates, plants and trace fossils). We extend our appreciation to staff from the NPS Biological Resources Division (BRD) for awarding financial support to host the GRCA PaleoBlitz, including Elaine Leslie (Chief of BRD, now retired), and BRD staff Kiersten Jarvis and Kriston Barnes. We would like to extend our thanks to James "Max" Bovis and Tom Olson for the volunteer assistance and participation with the GRCA PaleoBlitz.

A National Fossil Day (NFD) Celebration was organized as a two-day event hosted on the South Rim of GRCA on Friday, September 27 and Saturday, September 28, 2019, as part of the paleontology-focused activities in conjunction with the park's centennial. On Friday evening, a Grand Canyon National Park Paleontology Symposium was organized at the Shrine of Ages and consisted of four fossil-themed presentations open to the public. The speakers included paleontologists Jim Mead (The Mammoth Site), John-Paul Hodnett (Maryland-National Capital Park and Planning Commission), Linda Sue Lassiter (Northern Arizona University) and Spencer Lucas (New Mexico Museum of Natural History and Science). On Saturday, the National Fossil Day Celebration was hosted outside of the GRCA South Rim Visitor Center for the public, and an evening program for the public was presented by NPS Senior Paleontologist Vincent L. Santucci. We wish to thank the many partners that participated in this event, meeting with the public and promoting the scientific and educational values of fossils. Thank you to Bruce MacFadden (Paleontological Society, Florida Museum of Natural History); Jim Mead (The Mammoth Site); Ed Robeck, Geoff Camphire, Adam Blankenbicker, and Sequoyah McGee (American Geosciences Institute); David and Janet Gillette, Jennifer Glennon, and Richard McMichael (Museum of Northern Arizona); Doug and Hazel Wolfe (Zuni Dinosaur Institute for Geosciences); Andy Grass and Sherman Mohler (Arizona Museum of Natural History, Southwest Paleontological Society); John-Paul Hodnett (Maryland-National Capital Park and Planning Commission); Joel Despain (National Cave and Karst Research Institute); Sandy Croteau (Protectors of Tule Springs); Tom Olson (Geology for Kids); and Anne Scott and Eleanour Snow (US Geological Survey). We are also grateful to our colleagues from the NPS for their participation in the Grand Canyon Centennial National Fossil Day Celebration, including: Jason Kenworthy and Justin Tweet (NPS Geologic Resources Division); Mary Carpenter (Badlands National Park); Maria Rodriguez (Glen Canyon National Recreation Area); Erin Eichenberg and Mary Ontiveros (Tule Springs Fossil Beds National Monument); and Robyn Henderek (Zion National Park).

One of the main components of the 2019 GRCA Centennial Paleontological Resource Inventory was the compilation of the extensive paleontological resource information for the park. The park preserves an extraordinary record of Paleozoic flora and fauna which have been the focus of scientific collection and study for approximately 150 years. A team of notable paleontologists were selected to develop original manuscripts documenting the scope, significance and distribution of fossils throughout GRCA. The final report contains thematic chapters presenting a comprehensive overview of the paleontological resource of GRCA. A sincere thank you to each of the lead authors and co-authors who prepared original peer-reviewed manuscripts which are included in this publication, including: P. J. Bergman (Northern Arizona University); Kim Besom (Grand Canyon National Park); Diana Boudreau (Geoscientists-in-the-Parks, American Conservation Experience); Mary Carpenter (Badlands National Park); Carol Chambers (Northern Arizona University); Ronnie Colvin (Grand Canyon National Park); Tim Connors (NPS Geologic Resources Division); Dave Elliott (Northern Arizona University); John Foster (Utah Field House of Natural History State Park Museum); Heitor Francischini (Federal University of Rio Grande do Sul); John-Paul Hodnett (Maryland-National Capital Park and Planning Commission); Adrian Hunt (Flying Heritage and Combat Armor Museum); Colleen Hyde (Grand Canyon National Park); Cassi Knight (Paleoworks Consulting); Linda Lassiter (Northern Arizona University); Spencer Lucas (New Mexico Museum of

Natural History and Science); Lorenzo Marchetti (Prehistoric Museum Geoskop); Adam Marsh (Petrified Forest National Park); Jim Mead (The Mammoth Site); Anne Miller (Northern Arizona University); Mark Nebel (Grand Canyon National Park); William Parker (Petrified Forest National Park); Vincent Santucci (NPS Geologic Resources Division); Earle Spamer (retired—American Philosophical Society); Fred Sundberg (retired—Show Low High School); Shawn Thomas (Bat Conservation International); Benjamin Tobin (University of Kentucky); Justin Tweet (NPS Geological Resources Division); Sebastian Voigt (Prehistoric Museum Geoskop); and Klara Widrig (Geoscientists-in-the-Parks).

Each of the chapters within the GRCA Centennial Paleontological Resource Inventory report was peer-reviewed and will be published in both NPS internal reports and as an open-access publication in the journal *Geology of the Intermountain West*. First, we want to extend our appreciation and thank you to Don Weeks (DOI Regions 6–8 [formerly NPS Intermountain Region]), who served as the peer review coordinator for the GRCA Paleontological Resource Inventory Report. The peer review process for so many individual chapters was an involved process, and Don did a great job in completing the peer review for the many sections of this large report. Additional thanks are extended to the many peer reviewers who help us to improve the content, quality and scientific integrity of information contained in this publication. Thanks are extended to: Harry Butowsky (retired—NPS); Don Corrick (Big Bend National Park); Carol Dehler (Utah State University); Bill DiMichele (Smithsonian Institute); Erin Eichenberg (Tule Springs Fossil Beds National Monument); David Elliott (Northern Arizona University); Matt Friedman (Museum of Paleontology, University of Michigan); Hendrik Klein (Saurierwelt Paläontologisches Museum Neumarkt); Cindy Looy (Smithsonian Institute); Spencer Lucas (New Mexico Museum of Natural History and Science); Jeff Martz (University of Houston); Andrew Milner (St. George Dinosaur Discovery Site); Nicholas Minter (University of Portsmouth); Gary Morgan (New Mexico Museum of Natural History); Sterling Nesbitt (Virginia Polytechnic Institute and State University); Susannah Porter (University of California: Santa Barbara); Wayne Ranney (Grand Canyon river guide, writer, and educator); and Rick Toomey (Mammoth Cave National Park).

Finally, we want to acknowledge the Grand Canyon National Park Paleontology Project Core Team. The individuals below contributed to all aspects of planning, implementation, research, field work, museum collections review, specimen photography, and development of interpretive media. They also reviewed draft manuscripts, issued research and collecting permits, and coordinated communication with park staff, cooperating paleontologists, partners and other individuals involved with the GRCA Centennial Paleontological Resource Inventory. An After Action Report for this project was prepared by paleontology intern Diana Boudreau and is available to download at the following website: https://irma.nps.gov/DataStore/Reference/Profile/2268299.

Two paleontology interns, Diana Boudreau and Klara Widrig, were hired for this project and served six months helping to coordinate the many components of the paleontology-focused activities at GRCA. Colleen Hyde and Kim Besom of the GRCA Museum have been long-term advocates for GRCA paleontology and were extremely helpful with the many museum-related requests during this project. We dedicate this report to Colleen and Kim in appreciation for their years of service

supporting GRCA paleontology. GRCA Interpretive Ranger Ronnie Colvin inspired our team with her creative energy, wonderful ideas, and the experiences she has gained by hosting National Fossil Day events at GRCA in the past. Our National Fossil Day Coordinator John-Paul Hodnett (Maryland-National Capital Park and Planning Commission) first suggested hosting a NFD event at GRCA and was instrumental in the success of this special event during the park's centennial. We were fortunate to share in the successful completion of a Master's Thesis from Northern Arizona University by paleontologist Anne Miller. Anne's contributions to this project were many, and her previous work at GRCA prepared her well to help plan field logistics to ensure that all our team remained safe during their ventures for fossils in the canyon. Our scientific advisor for the project, Ronda Newton, helped our team to navigate important discussions and requirements, including our research and collecting permit, to support our planning and project activities. GRCA GIS Program Manager Mark Nebel served many key roles during the 16 months of work involved with the early planning, developing of funding requests, and providing cutting-edge technical geospatial support that we can benefit from in our paleontology work in other NPS areas. Vincent Santucci and Justin Tweet from the NPS Paleontology Program provided coordination of project planning, weekly assignments and meeting, communication, and development of the GRCA Paleontological Resource Inventory Report.

The GRCA Centennial Paleontological Resource Inventory represents the largest and most comprehensive park-specific paleontological resource project ever undertaken by the NPS. Our hope is that this effort will not only contribute to the understanding of the extensive fossil record at GRCA, but to also comprehensively assemble important baseline paleontological resource data to support and inform park management. This report has benefited from the work of those who have come before us, and during this 100th Anniversary we offer the GRCA Paleontological Resource Inventory Report to serve and inspire rangers, visitors and those others who will be the stewards of the GRCA fossil record in the future.

Dedication

On behalf of the Grand Canyon National Park Centennial Paleontological Resource Inventory team we would like to dedicate this report to Colleen Hyde and Kim Besom (GRCA Museum) and to all National Park Service curators, museum specialists, museum technicians, collections managers and archivists.



A photo of Kim Besom (left) and Colleen Hyde (right) standing at an overlook on the Grand Canyon rim

First, we recognize that the work we are involved with in the NPS Paleontology Program and at GRCA has consistently benefited by the dedicated professionalism of park museum staff. For paleontology, maintaining fossil collections in museums is a fundamental and core value of the science. The meticulous detail and care provided for fossil collections in the NPS, not only contributes to the science of paleontology, but it is fundamentally important to our preservation/conservation mission and goals in the NPS. This is especially true for our non-renewable remains of ancient organisms preserved in our national parks.

Moreover, much of this important and often behind the scenes work undertaken by museum professionals, is the foundation for our scientific research and resource management related to non-renewable fossil resources. We have clearly benefited by the support from many museum professionals in the NPS, who through their work contribute to the scientific integrity of the important legacy data associated with park museum fossil collections. We are reminded continuously

during our work in the NPS Paleontology Program, how we have benefited from the meticulous work of museum professionals at all levels (WASO, regions and parks).

Both Colleen and Kim exemplify the best of the best in professional museum curators. Not only do they maintain the huge database of knowledge related to what is now more than 100 years of GRCA history and museum objects that preserve the rich stories of the park, but they do so with passion and dedication. Colleen and Kim have been so very helpful to each one of our GRCA Paleontology Team with our many requests for assistance. As long as we have worked together with Colleen and Kim, they have always been so helpful and friendly in support of our research. They have accommodated a long list of researchers that were sent their way to view collections and obtain curatorial records to support their research. A number of current and past paleontology interns praise Colleen and Kim on a regular basis.

This GRCA Paleontological Resource Inventory project is an important one for us all—and it is equally important that we are able to extend our thanks and tribute to the museum professionals who have supported our work and we have all come to call our friends and colleagues.

Chapter 1. Introduction and Summary: The Paleontological Heritage of Grand Canyon National Park

By Vincent L. Santucci[1]

[1]National Park Service
Geologic Resources Division
1849 “C” Street, NW
Washington, D.C. 20240

Throughout my life I have been bestowed the privilege of experiencing the world-renowned landscape and resources of the Grand Canyon from many perspectives and viewsheds (Figure 1-1). My first views were standing and taking photos from the many vantage points and overlooks along the North and South rims. I have enjoyed many hikes into the canyon with colleagues from the National Park Service (NPS) or with academic geologists and paleontologists. On a few occasions I ventured down and then back up the trails of the canyon with my children Sarah, Bethany, Luke, Jacob, Brianna and Abigail, often carrying one or more in my arms on the climb against gravity. I traversed by foot to the base of the canyon at Phantom Ranch and gained a greater appreciation for the geologic story preserved in the park strata. I have gazed intensely out the window of many commercial aircraft from above this geologic wonder of Earth, contemplating the geomorphic “grandeur” created over "Deep Time" and the artistry of processes perfected by “Mother Nature.” I pinch myself when I recall the opportunity when my friend Justin Tweet and I were granted permission to fly into the western portion of the Grand Canyon on a small NPS plane operated by a pilot from Lake Mead National Recreation Area. As we meandered above the Colorado River and below the canyon rim, we celebrated when we located one of the paleontological crown jewels of the NPS: the entrance to Rampart Cave. I have studied the Grand Canyon through many scientific articles and geologic/topographic maps, and in my contemplation and dreams, I always ask myself, “What is still out there? What fossilized remains from our past are yet to be discovered?” I love this Grand Canyon!

The stars were aligned in 2019 to assemble a team, with the support of the Grand Canyon National Park leadership, to undertake a comprehensive paleontological resource inventory for the park. One of the primary catalysts to this ambitious task was tied to the centennial celebration for Grand Canyon National Park, in recognition of the 100th anniversary of the park’s establishment on February 26, 1919. The discussions for the paleontological resource inventory began between the NPS Paleontology Program and Grand Canyon National Park staff in 2018, which led to tremendous support to move forward with a park-specific paleontological resource inventory in conjunction with the centennial. A multidisciplinary team of subject matter experts were assembled to assist in the development of the thematic manuscripts which make up this publication and to assist with a number of paleontology-focused tasks and activities related to the paleontological resources of Grand Canyon National Park.




Figure 1-1. Map of Grand Canyon National Park (NPS/MARK NEBEL).

In addition to this publication focused on the paleontology of Grand Canyon National Park, two other major tasks were undertaken during the park's centennial. One consisted of hosting the Grand Canyon National Park PaleoBlitz, only the second such undertaking sponsored by the NPS Paleontology Program (the first was hosted at Chickasaw National Recreation Area in 2016). The other was the hosting of the 2019 National Fossil Day Celebration at Grand Canyon National Park (Figure 1-2). As part of the Grand Canyon National Park Centennial events, in September 2019 we gathered together at Mather Point and celebrated National Fossil Day with our friends, partners and the public. Diana Boudreau (2020) summarizes the various components and accomplishments in an after-action report available through IRMA at https://irma.nps.gov/DataStore/Reference/Profile/2268299.




Figure 1-2. 2019 National Fossil Day logo featuring Pleistocene fauna from Rampart Cave in Grand Canyon National Park (NPS).

The principal goals and objectives for the Grand Canyon National Park Centennial Paleontological Resource Inventory were to identify the scope, significance, distribution and management issues associated with the fossils of the park. Through this effort we compiled baseline paleontological resource information for park managers and staff to better understand the non-renewable resources under their stewardship, and in turn help to inform park planning and decision-making which may relate to park fossils. This publication is written for park managers, in the language and content most useful for considering the many aspects of management, protection, interpretation and scientific study of National Park Service fossils.

It is also important to recognize the role of the Paleontological Resource Preservation Act of 2009 (PRPA) in defining the need for undertaking this paleontological resource inventory for Grand Canyon National Park. Specifically, Section 6302 of PRPA (16 USC 470aaa-1) mandates for the NPS and other Department of Interior bureaus that "The Secretary shall manage and protect paleontological resources using scientific principles and expertise. The Secretary shall develop appropriate plans for inventory, monitoring, and the scientific and educational use of paleontological resources." The work undertaken in conjunction with the 2019 Grand Canyon National Park Centennial Paleontological Resource Inventory exemplifies this legal standard and scientific principle.

The Grand Canyon National Park Centennial Paleontological Resource Inventory represents the largest such inventory in National Park Service history. The work required the collaboration of a team of paleontologists, NPS staff, and a number of partners. Subject matter experts with previous research and experience involving Grand Canyon fossils were involved in various aspects of this inventory including drafting or peer review of the manuscripts presented in this undertaking. The collective work is represented by this publication, by new fossil specimens cataloged in the park's museum collection, and through new interpretive media focused on Grand Canyon paleontology. This work represents the compilation of historical and current research and fossil collections associated with paleontology of Grand Canyon National Park. In addition, fossils documented from outside the administrative boundaries of the park, from the greater Grand Canyon region, have also been considered as part of this inventory.

The first technical chapter (2) in the Grand Canyon National Park Centennial Paleontological Resource Inventory is *History of Paleontological Work at Grand Canyon National Park: Up and Down the Long Federal and NGO Trails of Paleontology in Grand Canyon National Park, 1858–2019* by Earle E. Spamer. This chapter presents a historic overview of the park's paleontology, including fieldwork and research. This rich history begins with the first discovery of a fossil in the Grand Canyon in April 1858 in the Diamond Creek area by members of the Ives Expedition and published in 1861 by John Strong Newberry (Newberry 1861). A long list of notable paleontologists have worked in Grand Canyon and are summarized in Spamer's chapter (Figure 1-3).




Figure 1-3. Notable paleontologists who worked in areas now administered as Grand Canyon National Park, including (from left to right): Charles Doolittle Walcott (USNM photo); John C. Merriam (NPS); Charles Gilmore (USNM); Remington Kellogg (USNM); David White (NPS); and Edwin McKee (USGS).

A chapter (3) devoted to a basic overview on the *Stratigraphy of Grand Canyon National Park* was prepared by Tim Connors and others. Given the paleontological focus of this publication, there was a need to provide a consistent stratigraphic context for the individual chapters. This chapter was not intended to present a comprehensive guide to the geology of Grand Canyon National Park, nor to address unresolved or controversial interpretations or perspectives on the park's geology. The focus on the stratigraphy was viewed as important given the close relationship to the park's paleontology and paleoenvironments represented by the various stratigraphic units.

Grand Canyon National Park preserves important and fossiliferous exposures of Precambrian strata. Prior to the area being initially proclaimed a national monument in 1908, paleontologist Charles Doolittle Walcott ventured into the Grand Canyon on his quest for early fossils (Figure 1-4). In the chapter (4) titled *Precambrian Paleontology of Grand Canyon National Park* by Justin Tweet, he reports on the oldest known fossils preserved at the park and in some cases the earliest reported occurrences of unusual fossil forms such as *Chuaria circularis* and vase-shaped microfossils.

The invertebrate fossil record from the Paleozoic strata of Grand Canyon National Park represents by far the most abundant and diverse category of park fossils. This includes both invertebrate body fossils and trace fossils largely preserved in marine sedimentary rocks, and specific chapters are dedicated to each. Linda Lassiter and others have prepared a detailed summary of the invertebrate body fossils (Chapter 5), titled *Paleozoic Invertebrate Paleontology of Grand Canyon National Park*. Anne Miller and others prepared a comparable work reporting on the invertebrate trace fossils (Chapter 8) which is titled *Paleozoic Invertebrate Ichnology of Grand Canyon National Park.*

Fossil vertebrates were first reported from the Devonian Temple Butte Limestone in Grand Canyon by Charles Walcott (1880) during the initial surveys undertaken by the U.S. Geological Survey. The Mississippian Surprise Canyon Formation has yielded a rich fossil chondrichthyan fauna which has recently been described, with a number of new fossil shark taxa being named (Hodnett and Elliott 2018) (Figure 1-5). The authors who published on the Surprise Canyon Formation sharks, JP Hodnett and David Elliott, are also the authors for the chapter (6) in this publication titled *Paleozoic Vertebrate Paleontology of Grand Canyon National Park: Research History, Resources, and Potential.*




Figure 1-4. Paleontologist Charles Doolittle Walcott explored the Grand Canyon's Precambrian and early Paleozoic strata for traces of early life (NPS).



Figure 1-5. Grand Canyon National Park Fossil Trading Card featuring the fossil shark *Amaradontella santuccii* from the Mississippian Surprise Canyon Formation (NPS).

Fossil plants are summarized by Cassi Knight in her chapter (7) *Paleozoic Paleobotany of Grand Canyon National Park*. This chapter reviews the known Paleozoic plant macrofossil resources in Grand Canyon National Park, detailing previously published plant fossil occurrences from the Surprise Canyon Formation, the Supai Group formations, and the Hermit Formation (Figure 1-6). Many of the taxonomic names used by David White in his 1929 monograph on the plant fossils of the Hermit Formation needed to be reviewed in light of more recent publications and taxonomic considerations.




Figure 1-6. Civilian Conservation Corps (CCC) constructing the in situ Fossil Fern Exhibit along the South Kaibab Trail during 1937 (NPS).

An important Late Paleozoic vertebrate ichnofossil record has been documented from the Hermit Formation and Coconino Sandstone from within Grand Canyon National Park (Figure 1-7). Smithsonian paleontologist Charles Gilmore made large collections of fossil vertebrate tracks from the park during the 1920s. The fossil tracks from the park include some of the oldest occurrences of tetrapod tracks preserved in eolian strata. Lorenzo Marchetti and co-authors review these and other finds in their chapter (9) titled *Paleozoic Vertebrate Ichnology of Grand Canyon National Park.*

The limited exposure of Mesozoic strata in Grand Canyon National Park is represented by the Triassic Moenkopi Formation and the Shinarump Member of the Chinle Formation. Petrified Forest National Park paleontologists Adam Marsh and Bill Parker coordinated field inventories of the Triassic units in the park during 2019. Marsh and others prepared a chapter (10) titled *Mesozoic Paleontology of Grand Canyon National Park: Trace Fossils, Stratigraphy, and Regional*

Correlations. Marsh and his team documented both invertebrate and vertebrate trace fossils from the Moenkopi Formation in the park, which are the first examples of either reported from the Mesozoic rocks of Grand Canyon National Park.



Figure 1-7. In situ block of Coconino Sandstone with *Ichniotherium sphaerodactylum* trackways in Grand Canyon National Park (NPS/CASSI KNIGHT). This find was documented in Francischini et al. (2019).

The area in and around Grand Canyon National Park contains more than 650 documented caves. Some of these dry caves preserve Pleistocene/Holocene paleontological resources including Ice Age mammals and birds (Figure 1-8). Among these fossils, rare mummified remains of Pleistocene bats and other mammals are preserved with soft tissues and hair in a few of these caves within the canyon. One of the more notable caves in Grand Canyon is Rampart Cave, which preserved stratified sequences of ground sloth dung, but much of the material was lost to an accidental anthropogenic fire in the 1970s. Jim Mead and his team have contributed a chapter (11) titled *Pleistocene/Holocene cave fossils from Grand Canyon National Park* to summarize these discoveries.



Figure 1-8. Fossil condor skull from Stevens Cave, Grand Canyon National Park (STEVE EMSLIE).

The final chapter of this publication is specifically dedicated to the management of Grand Canyon National Park's rich and diverse paleontological resources. Diana Boudreau and others developed a chapter (12) titled *Grand Canyon National Park Paleontological Resources Management and Protection* to help park managers and staff understand the range of activities and issues associated with the management of non-renewable resources. This chapter focuses not only on the management of fossils found within the rock strata of the park, but also presents information related to fossils in the park's museum collection and other outside repositories which maintain park fossil collections. This chapter also presents information related to a Geospatial Database developed and piloted by park staff to manage fossil locality data for Grand Canyon National Park.

In addition to chapter-specific appendices, six general appendices are included at the end of this document: fossil taxa named from specimens found within GRCA (Appendix A); fossil taxa named from specimens possibly found within GRCA (Appendix B); summary tables of Paleozoic taxa reported from GRCA (Appendix C); contact information for outside repositories of GRCA fossils (Appendix D); paleontological resource law and policy information (Appendix E); and a geologic time scale (Appendix F).

The study of the paleontological resources at Grand Canyon are brief, spanning slightly more than 160 years since the first fossils were collected from Diamond Creek during the Ives Expedition in 1858. The team contributing to this publication is dedicated to the paleontology of Grand Canyon National Park and all recognize that we have just scratched the surface in terms of the potential for future fossil discovery. This report is intended to provide a baseline of paleontological resource information as of the 100th anniversary of Grand Canyon National Park in 2019. We collectively admire and benefit from the work of those paleontologists who have come before us and hope that

our work will inspire and support the work by those paleontologists and park managers who will come in the future.

Literature Cited

Boudreau, D. M. 2020. Grand Canyon National Park: 2019 paleontology project after action report. Natural Resource Report NPS/GRCA/NRR—2020/2054. National Park Service, Fort Collins, Colorado.

Francischini, H., S. G. Lucas, S. Voight, L. Marchetti, V. L. Santucci, C. L. Knight, J. R. Wood, P. Dentzien-Dias, and C. L. Schultz. 2019. On the presence of *Ichniotherium* in the Coconino Sandstone (Cisuralian) of the Grand Canyon and remarks on the occupation of deserts by non-amniote tetrapods. Paläontologische Zeitschrift online:1–19.

Hodnett, J.-P. M., and D. K. Elliott. 2018. Carboniferous chondrichthyan assemblages from the Surprise Canyon and Watahomigi formations (latest Mississippian–Early Pennsylvanian) of the western Grand Canyon, northern Arizona. Journal of Paleontology 92, Memoir 77.

Newberry, J. S. 1861. Geological report. *In* Ives, J. C., Report upon the Colorado River of the West, explored in 1857 and 1858. U.S. 36th Congress, 1st Session, House Executive Document 90, Serial 1058, Part 3.

Chapter 2. History of Paleontological Work at Grand Canyon National Park

Up and Down the Long Federal and NGO Trails of Paleontology in Grand Canyon National Park, 1858–2019

By Earle E. Spamer[1]

[1]Academy of Natural Sciences
Research Associate
Philadelphia, Pennsylvania

Introduction

The Grand Canyon! Anywhere in the world the name rouses recognition. Monumentally impossible to describe (or so have said thousands of writers who then effused their own descriptions), it has been a lure to geologists since 1858. From the start, the rocks were read for the clues of their relative ages. It has been the draw of government agencies and non-government organizations (NGO) alike. The national park is a century old now; the canyon six to 70-some million years (depending upon with whom you argue, and about which parts of the canyon you consider); and fossils in the canyon have awaited the hammer and scanning electron microscope for even more than a billion years. So, to avoid the traps of superlatives and the gulping periods of time, this is a fast trot through "the best of" Grand Canyon paleontology, refreshed with bits of human history, with a few pauses on peculiar details—a 100th birthday present to the national park. Here, beginning with the first Grand Canyon field trip in 1858, is an accounting of how the first explorers, and scientists and educators over the years, have fashioned our understanding and encouraged our participation in the story of ancient life presented in Grand Canyon's strata and secluded deposits. With this long look backward, we also may gain an appreciation for how paleontologists, federal administrators, and NGO champions built up the scientific and educational programs that modern resource managers receive as a legacy.

There is a lot of "business" behind this paper, in the form of stratigraphic and taxonomic summaries, which I provided in a lengthy overview of Grand Canyon paleontology (Spamer 1984). It still stands as a fairly reliable history of research, with an extensive list of references, and some handy species lists. For those so interested, there is also a corresponding general-audience overview from the perspective of the geologists in Grand Canyon (Spamer 1995; see also Ranney 2013, 2014). The present paper, in celebration of the national park's centennial anniversary and saluting National Fossil Day 2019, looks at the history from the perspective of the people and the agencies responsible for fossil studies in the canyon, and some of the educational projects that came about due to these studies. To this is added a brief rundown of research handed in since 1984.

So much is available in Grand Canyon publications that deal with paleontology that to supplement this paper I have prepared a *Bibliography of Paleontology in the Grand Canyon Region and in the Stratigraphic Continuity of Grand Canyon Formations*. This PDF with more than a thousand citations also complements the commemorative events of 2019. Occasionally updated, it can be

downloaded from the web at https://ravensperch.org/bibliography-of-paleontology-of-the-grand-canyon-region/.

In this day and age it has become so easy to search online for reliable additional basic information—such as photos of the many people who are mentioned here, as well as maps and beautifully detailed representations of Grand Canyon's stratigraphic column (which even this year has been revised)—that it seems needless to fill out this paper with numerous asides and portraits. Instead, the illustrations here show a few places, things, and events of interest that are mentioned.

And a note of thanks to Park Service managers: *These* are the very kinds of work that rely on the lands and resources you oversee, even if much of it never comes to your attention.

Uphill

"Not really on the way to anywhere" pretty much describes the Grand Canyon; even the earliest trail-blazing expeditions in the neighborhood did not go there. In those days the canyon was known only vaguely and unimaginatively as "Big Cañon"; big and impassable (so it was heard). Although the name "Grand Canyon" was first used by physical scientist and statistician Lorin Blodget [1823–1901] (1857:90, 97), it was overlooked until John Wesley Powell [1834–1902] popularized it a decade later. The first field trip that *did* go to Big Cañon was quite out of the ordinary.

John Strong Newberry [1822–1892] was the first geologist to arrive, on mule, back in 1858. He had come to the region aboard a 50-foot sternwheel steamboat, made of iron and wood, commissioned by the U.S. Army in Philadelphia. It was little more than a bucket and a boiler. The mission was to explore the Colorado River from the sea upstream in order to determine the river's navigable reach and thus the possibilities of communicating with the interior West from the sea. (There also was a bit of secret political intrigue, too, only more recently understood, that indicates that this was a reconnaissance in the face of a potential civil war with Utah's Mormons; but that's a whole separate story having nothing to do with science.)

Three steamers already on the lower Colorado River were deemed too expensive to hire. *Explorer* (Figure 2-1), built in a hurry during the summer of 1857 and tested by a spin on the Delaware River between Pennsylvania and New Jersey, was seen as more economical even with shipping costs. It was sent in eight pieces first to New York, then by ship to the two-year-old railroad across the Isthmus of Panama, and, after two more voyages by steamer and schooner in the Pacific, unloaded and wrestled together at Robinson's Landing, an improvised shipyard on the mudflats of the Colorado River delta in Mexico. There, during December, it was decided that the three-ton boiler could make groundings on sandbars a dangerous affair for the low-sided, deckless boat, so two heavy timbers were bolted lengthwise beneath the hull. Unfortunately, that increased the vessel's draft, a serious concern on such a shallow river. The slot in the transom, through which the engine's drive rod passed to the paddlewheel, was too deep and would have allowed water into the boat underway on the Colorado, a river not as placid as the tidewater Delaware. The engine was raised and an iron patch was fitted over the lower part of the slot.




Figure 2-1. The steamboat *Explorer* on the lower Colorado River; the first field trip en route to Grand Canyon (Ives 1861: frontispiece).

The expedition commander, veteran western explorer Lt. Joseph C. Ives [1829–1868], engaged there at Robinson's Landing steamer captain David C. Robinson [1833–1874] to pilot the craft on the river. With Robinson at the tiller perched atop *Explorer*'s small aft cabin (the boat did not even have a wheel), Andrew Carroll [*ca*. 1830–?], the engineer who had accompanied the boat from Philadelphia, and a skeleton crew, *Explorer* beat its way 150 miles to Fort Yuma, constantly challenged by the deceptive, wiggly channels of the delta and shifting river bottoms. At Fort Yuma on the California side of the Colorado and Gila River confluence, the rest of the small crew, who had traveled through the desert from San Diego, came aboard. Ives' (1861) formal report provides a detailed record of all of these happenings, as well as a fascinating travelogue of the remainder of the expedition.

Explorer's crew was an unlikely bunch: Captain Robinson and engineer Carroll; Lt. Ives and soldiers; Newberry as the mission's physician, geologist, and general naturalist; Friedrich von Egloffstein [1824–1885], a German baron who served as cartographer and illustrator and who was a veteran of western explorations elsewhere; and Balduin Möllhausen [1825–1905], a burly German draftsman, illustrator, and naturalist's field assistant, another western exploration veteran who had already visited the lower Colorado River. The expedition was cheerfully waved on its way by the fort's soldiers and smirking townspeople. *Explorer* promptly ran aground within sight of the fort. One might be forgiven if it brings to mind the campy 1960 Jack Lemmon film, *The Wackiest Ship in the Army*.

Still they did make their way upstream to find the head of navigation. They reached a suitable point in Black Canyon, not far downstream from where Hoover Dam was built eight decades later—by violently crashing into a rock, casting Ives, Newberry and Möllhausen from the cabin top, not to overlook "the fireman, who was pitching a log into the fire [and] went half-way in with it" (Ives 1861, Pt. 1: 82). Three days later, after a brief survey farther upstream in a skiff and *Explorer* patched, they glided back down to meet up with a long army packtrain of mules. Ives divided his command, sent the boat back to Yuma and set out overland to the east en route to Fort Defiance with

soldiers, Native Americans, a hundred-odd mules, Mexican packers, the Germans, our faithful Newberry, and a dog, "Grizzly."

The direction of travel was different, amidst this age of explorers who blazed wagon roads and railroad routes westward. However, the region they entered was not wholly unknown to Ives and Möllhausen. They had already traveled the general route near the 35th parallel a few years earlier when they were attached to a westbound expedition led by Lt. Amiel Whipple [1817–1863]. Laboring up from the Basin and Range to the Colorado Plateau, the now landward expedition approached the Grand Canyon—and thus, finally, the first geologist arrived. Somewhere in the vicinity, Newberry (or perhaps Möllhausen, Newberry's field assistant) gathered fossils that established a "Carboniferous" age (Permian today) for the rim rocks (Figure 2-2). The Hualapai people they had met might have been amused by men who hammered on rocks. A small party rode down to Diamond Creek's confluence with the Colorado River, the first non-Indigenous people to reach the bottom of the canyon.

While Möllhausen sketched, Newberry made geological observations and drew up an accurate stratigraphic column from the rim to the Colorado River. In his measured section (Newberry 1861: 55; and see his figure 12, page 42), he noted "*Euomphalus, Spirifer*, &c.", "imperfect corals," "corals (*Chaetetes*?)" (a genus now known as a sponge), and "green shales with mud furrows, resembling casts of worm holes." Even though the fossils of the rim rocks were helpful in establishing the relative age of those strata, Newberry bemoaned in his written report (page 55), "In the absence of [index] fossils it is impossible to determine the precise geological age of any of the strata composing the … section below the limestone which forms the summit of the cliffs." When all was said and done, though, Newberry drew up a detailed collection list for the whole expedition, itemizing nearly a thousand specimens of rocks, including fossils galore. Newberry also made his one most astute observation for the ages, that even the dry side canyons were created by running water rather than by cataclysm.

The Ives party pressed on to Cataract Creek, its lower end now known as Havasu Creek, where the mules could barely be turned around when the way became too sketchy for animals. The teams were sent back twenty miles to the last known water while some men went ahead on foot. The weighty Egloffstein managed to wreck a precarious Havasupai ladder, stranding himself. He wandered off to the Havasupai village of Supai, the first "foreigner" to go there since Spanish padre Francisco Garcés [1738–1781] arrived in July 1776. (Egloffstein was rescued when the soldiers lowered gun slings to hoist him up.)

Thus was the first brief, but productive, field trip to the Grand Canyon. Regrouped, Ives continued eastward, sending most of the party off toward Fort Defiance (Möllhausen had traveled the route with Whipple) while he, Egloffstein, and a few men made a side trip to visit the Hopi. After the expedition disbanded, Ives returned to Yuma, where he sold *Explorer* six months after its maiden (and only government) voyage. It served anonymous local companies for a while, broke loose in a flood, and drifted away toward the sea.




Figure 2-2. Grand Canyon paleontology begins here, April 1858 (Newberry 1861: Paleontology Plate 2). All are from the undifferentiated Permian Kaibab Limestone–Toroweap Formation. **Figures 1–8.** *Productus ivesi*, a new species; **Figures 9 and 10.** *P. occidentalis*, a new species; from "Middle Carboniferous limestone, banks of the Colorado, near mouth of Diamond river" [Peach Springs Canyon between Peach Springs and the confluence of Diamond Creek]; **Figures 11–13.** *Streptorhynchus pyramidalis*, a new species, from "Carboniferous limestone over red cross-stratified sandstone at Camp 70, on high mesa west of Little Colorado" [western part of the Coconino Plateau not far east from Diamond Creek]. These localities today are a part of the Hualapai Indian Reservation.

Downhill

A decade later, another government-sponsored expedition arrived in the canyon—this time from the north, by river from Wyoming and Utah. In August 1869, Civil War veteran Maj. John Wesley Powell, now a civilian and with eight crew members aboard three boats, passed into Grand Canyon having charted the Green and Colorado Rivers to the mouth of the Little Colorado River. They had support from the Smithsonian Institution and some military favors in the form of supplies. Although they had passed sequentially down through all of the Colorado Plateau's strata, there hadn't been much time for collecting. From this point onward was Powell's "Great Unknown." They were effectively off the map now, with only Egloffstein's generalized physiographic map (in Ives 1861) as an informer, a map good for the lower Colorado River region but quite inaccurately portraying the Grand Canyon region.

Powell, though geologically trained, reported little on fossils in his expositions on the strata; and anyway by this time he was more concerned with where the river was leading him, and getting his crew through on rapidly diminishing rations. He led a second trip in 1871–1872, pausing for a winter layover in Kanab, Utah, but he cut that journey short in the middle of Grand Canyon when he suddenly decided at Kanab Creek that they had achieved their objectives. Thereafter, the emphasis was on mapping the region on the plateaus. Powell's (1875, 1895) renditions of his river exploits remain classic tales of Grand Canyon and Colorado River running. The plateau surveys under his direction produced in 1882 what is probably the greatest (and most readable) scientific book on Grand Canyon physiography and geological history, *Tertiary History of the Grand Cañon District*, with its magnificent double-folio *Atlas*, by Clarence Edward Dutton [1841–1912].

Laying a Foundation

The first expeditions to and through the Grand Canyon were enough to establish that *this* was an interesting place, worthy of further study. By 1881, Powell had become the second director of the newly organized U.S. Geological Survey, and his ongoing surveys of the Colorado Plateau country at last brought a refined geologist's eye to the whole Grand Canyon. Powell, in the field, directed the construction of a trail into the northeastern corner of the canyon—today this is the tortuous, unmaintained Nankoweap Trail. He assigned the up-and-coming geologist, Charles Doolittle Walcott [1850–1927] (who would himself succeed Powell as USGS director), to spend the winter of 1882–1883 in the depths of the canyon, where with a collector, a cook, and a packer accompanying him, he studied for the first time the structural and stratigraphic relationships of the Proterozoic and lower Paleozoic rocks there—a remarkably instructive study to this day. Powell had recognized the basics of these relationships during his river expeditions, but now was the time to do some real field work.

Walcott saw quite a lot that piqued his interest in "Precambrian" fossils. He saw the stromatolites in these strata, which he called "an obscure Stromatopora-like group of forms" and "concretionary limestones." He also reported pteropods referred to *Hyolithes*, a trilobite fragment, a *Lingula*-like brachiopod, and a brachiopod related to *Acrothele*. This led Walcott to assume, with a Cambrian bias, that perhaps complex biological forms had evolved earlier than previously known. His research included ongoing observations, even if it was published in a paper on a wholly different geographic (and geologic) area. For example, see Walcott's (1884: 432) footnote in a paper on the Paleozoic of

Texas: “Since the paper on the Pre-Carboniferous strata of the Grand Cañon was published [Walcott 1883], a fragment of a trilobite, probably of the genus *Ptychoparia*, has been detected in a bit of shale from the Chuar group.” Alas, all of these reports of body fossils have subsequently been reinterpreted as pseudofossils, the “fool’s gold” of paleontology.

Yet in the process, Walcott also found a remarkable Neoproterozoic fossil, a megascopic alga he named in 1899 *Chuaria circularis*, which he believed to be a primitive brachiopod. Even though *Chuaria* has since been found around the world, so problematic this organism has been that it has been variously assigned to other phyla in the major kingdoms (it even once was considered to be a trilobite egg) and as a pseudofossil. Now *Chuaria*, definitely a fossil, is thought to be represented at different life stages also by *Tawuia* and *Longfengshania*. In a bizarre sort of way its form- and function-based taxonomies act like those of ichnofossils! Studies in China and India have been particularly robust; but the well-known, well-studied enigma persists. It all began in the Grand Canyon: Walcott’s original, small shale chips are in the U.S. National Museum (the Smithsonian Museum of Natural History).

Walcott was hardly a Grand Canyon novice. A couple of years earlier, the young man had had his first hand at geologizing in the west when he worked along Kanab Creek, the large tributary to the Colorado coming down to Grand Canyon from Utah, getting his and our bearings on the canyon’s entire Paleozoic sequence (Walcott 1880, 1883). Fossils were not lacking. These collections and others contributed decades later to Walcott’s very detailed series of publications on brachiopods and trilobites, produced during his 20-year position as the Secretary (director) of the Smithsonian Institution (see the paleontology bibliography online). The usefulness of the canyon’s fossil record was proven. Now geologists had a proper first analysis of the principal formations of the canyon.

On the Map

Geologists need maps. By the late 1800s, rudimentary 1:250,000-scale maps were available for the Grand Canyon region, products of Powell’s surveys. In summer 1902, the USGS’s François Matthes [1874–1948] and crew began a heart-stopping mission—using plane tables and alidades to create 15-minute quadrangles of the eastern portion of Grand Canyon with contour intervals of 80 ft (24 m); a project completed the following summer. They traveled into the canyon as well, and blazed a trail for their horses up through Bright Angel Creek to the North Rim. The resulting quads still are masterful examples of scientific art, but at the time they provided geologists with important tools for detailing more of the Grand Canyon’s lithology in the field.

While in the early 20th century some geologists were beginning to scrutinize the physiographic history of the canyon, others were heading into the chasm on foot and hoof.

Paleontology again took the back seat, serving only to assist in identifying relative ages, with no purposefully comprehensive collections being made. First on the ground now were Frederick Leslie Ransome [1868–1935], who in 1908 published a study of the Neoproterozoic sedimentary rocks and in 1916 produced correlations of Paleozoic strata in Arizona; Levi Fatzinger Noble [1882–1965], whose 1909 doctoral dissertation from Yale was based on his studies in the central Grand Canyon where Paleozoic and Neoproterozoic rocks are well exposed, with publications on this area in 1910,

1914, and 1922; Nelson Horatio Darton [1865–1948], who published in 1910 a reconnaissance of northern Arizona and in 1925 produced a very comprehensive résumé of Arizona geology; and Charles Schuchert [1858–1942], who published on the Grand Canyon Cambrian and "Carboniferous" (which included Permian strata) in 1918. This laid the groundwork for studies that, finally, began to look seriously at the fossil fauna and flora.

Making Tracks

Paleontological studies in the Grand Canyon have in fact been vigorous. Valuable collections of acritarchs, invertebrates, vertebrates, plants, and cryptic remains have been taken from Neoproterozoic to Permian strata and Pleistocene–Holocene deposits. Some of these collections contributed to the development of new concepts in stratigraphy that since have become a part of basic geological principles. However, the findings of Grand Canyon paleontology began modestly, sometimes peculiarly.

A variety of trace fossils had been known for years. The first were Newberry's observations of green-shale "worm holes" that now are well recognized and beautifully exposed in much of the Cambrian Muav Limestone. (In 1892 Joseph Francis James [1857–1897] suggested that these casts should be named *Scolithus arizonicus*, but without a description they fall into the taxonomic wastebasket of *nomina nuda*, unfounded names.) Walcott, too, widely recognized these abundant "fucoids," which he attributed to annelid worm trails and trilobite burrows. It is the ichnofauna of the canyon that first brought Grand Canyon paleontology to the fore, both professionally and popularly. While we can dismiss an anonymous 1884 report from two miners who found footprints 5.5 m (18 ft) apart made by a human some 4.0 m (13 ft) tall, who thus was shown to have been "fairly bounding along," it nonetheless demonstrates the things that can capture the public's interest. This report was interesting enough to have been picked up in France, too, so public attention was there. It's a shame that the miners never fulfilled their promise to return to remove the footprints to "exhibit them to public gaze."

Richard Swann Lull [1867–1957] had first written about "Nature's hieroglyphics" in 1904, including passing mention of a trackway in the Grand Canyon. In 1918 he produced the first dedicated study of vertebrate tracks from the canyon, describing two new ichnospecies from the Pennsylvanian–Permian Supai Group and four new ichnospecies and the now well-known ichnogenus *Laoporus* from the Permian Coconino Sandstone. What made the Coconino fossils all the more remarkable, Lull noted, as had Schuchert who collected them, was that they all were preserved on the foresets of fossil sand dunes, always advancing upslope. This apparently peculiar trait would in later years become the subject of much study.

By this time, although the Grand Canyon was widely known and was a popular tourist destination, for a variety of political reasons it still had not been designated a national park. (Since 1893 portions of the canyon had been a national forest preserve [1893], national game preserve [1906], and national monument [1908].) Finally, with Congress making Grand Canyon a national park in February 1919, even more public attention was drawn to it. In response, the National Park Service, itself only three years old, was faced with the increasing pressures of "what to do with it." This entailed promoting

the scenic, historic, and educational values the landscape holds, in addition to managing and entertaining visitors from around the world.

As might be expected, these programs came with the concerns of cost, promotion, and maintenance. Fortunately, the Santa Fe Railway had for more than two decades already been self-servingly advertising the Grand Canyon. Before government oversight of authorized concessionaires fully took root, the railroad independently completed a line to the rim in 1901, and by 1905 had built El Tovar Hotel and the adjacent Hopi House that drew visitors to buy American Indian crafts by the basketful. (In 1899 the rail line had been built from Williams, Arizona, to a site named Anita, to service copper mines established there. This station was just 19 rail miles short of Grand Canyon [Richmond 1998, 2017].) While the Santa Fe had thus also made it possible for scientists of various sorts (such as botanists and entomologists) to visit the canyon in pursuit of gatherings in their fields, they had no vested interest in "science" *per se*. That fell to the Park Service, and it would have to entail what we today call "outside funding" and "outreach."

Digging In

The industrialist and philanthropist Andrew Carnegie [1835–1919] used his billions of dollars (in today's figures) for educational and scientific projects after he retired from his business ventures in 1901. This included the founding of the Carnegie Institution of Washington, in 1902. Today, as the Carnegie Institution for Science, it is a leader in scientific discovery, in widely diverse fields. From 1920 to 1938, paleontologist and educator John Campbell Merriam [1869–1945] served as president of the institution (in 1919 he had also been president of the Geological Society of America), and Charles D. Walcott (also a GSA president, 1901) had served as an institution trustee 1902–1928 even while he directed the Geological Survey (to 1907) and the Smithsonian Institution (1907–1927). Not surprisingly, they encouraged programs of study in the earth sciences at the fledgling national park.

On recommendation, the National Park Service asked the Smithsonian's vertebrate paleontologist Charles Whitney Gilmore [1874–1945] to visit Schuchert's trackway site in the Coconino Sandstone along Hermit Trail. The purpose was to prepare a public educational exhibit. At the time, Hermit Trail had been improved by the Santa Fe so that tourists could reach Hermit Camp, the railway's tourist camp that deflected visitors from using the Bright Angel Trail that by dint of dozens of spurious "mining claims" was still largely (and illegally) "owned" by Ralph Henry Cameron [1863–1953], U.S. Senator from Arizona [1921–1927]. Gilmore went on to spend the next several field seasons in the canyon, studying and collecting from the Permian and Pennsylvanian strata along several trails. In the process, he described numerous new species and genera of vertebrate and invertebrate tracks (Gilmore 1926, 1927, 1928) and reported the first trackway found on the north side of the canyon (Gilmore and Sturdevant 1928).

More than relegated to stodgy scientific reports, Gilmore's work was noted in popular literature; after all, footprints are something that all can relate to, even more so that they were very old and were *in the Grand Canyon*. The prodigiously productive Merriam, too, promoted this work in his own educational publications.

Knowing that the Coconino Sandstone tracks were seen to face upslope on the fossil dunes, Gilmore (1926: 4) suggested they represented “an old trail leading to water, or possibly recording a great migration of animal life.” These were additional concepts with which an engaged public could easily identify. As time went on, interest in this peculiar trait did not wane. During the 1930s and 40s, experiments on modern sand dunes and in the laboratory, sometimes using living animals, showed that the answer was more a simple process of mechanics: essentially (and a bit simplistically), upslope trackways required sure planting of feet, while downslope travel incurred more sliding and tracks were readily erased.

Education Flowers

Many kinds of botanical fossils are found in the Grand Canyon. Some of these early finds were based on physical forms, which proved really to be sedimentary features and thus not fossils at all; for example, a new seaweed-like species *Rivularites permiensis* was described by David White [1862–1935] in 1929, from the Permian Hermit Shale, although various species of this genus have been reassessed as microbial sedimentary features. The higher systematic plants, found in abundance particularly in the Hermit Shale, are obviously fossils; after all, if it looks like a fern it probably is a fern (Figure 2-3). White’s magnificent 1929 monograph (published by the Carnegie Institution) remains a primary guide to Grand Canyon fossil plants. (Suitably, White’s grave in the Grand Canyon Cemetery is marked by a plaque in which are molded three of his new species; the figures are reproduced from his Hermit Shale monograph.) Some of White’s paleobotanical taxa, such as *Supaia*, even have significance as biomarkers for the early Permian, having been found in other strata around the world. Regretfully, paleobotanical work on the vascular plants of the Grand Canyon has been sparse since White’s work there.

All of this field work—ichnofossils and plants in particular—did not come cheaply. It was the Carnegie Institution that had underwritten the costs. There were the purely scientific objectives, of course, but the findings could be turned into educational projects. By the late 1920s the newly founded Grand Canyon Natural History Association (today the Grand Canyon Conservancy) was producing a monthly newsletter, *Nature Notes*, which frequently presented brief write-ups of the latest paleontological finds from the canyon. Glen Sturdevant [1895–1929], the only ranger–naturalist at Grand Canyon, almost single-handedly wrote the articles and news notes. After his untimely death in the Colorado River, a geology student who had been working under Merriam’s programs in the canyon during recent summers, Edwin “Eddie” Dinwiddie McKee [1906–1984], stayed on as the next ranger–naturalist. As we shall see, McKee exploded into the earth sciences in the canyon and around the world.



Figure 2-3. David White (left) and John C. Merriam at a fossil fern quarry in Grand Canyon (note the slab leaning against White's right leg) (NPS).

Through the 1930s, the Carnegie Institution supported research and funded publications about many of the Grand Canyon's strata; sedimentological, stratigraphic, and paleontological studies alike. The institution's *Year Book* series dutifully recorded each year's support and results from its underwritten programs. In 1928, the Yavapai Observation Station was dedicated as a place where visitors could observe and learn about the geology of the canyon in a museum setting, staffed by Park Service rangers, with an observation area overlooking the canyon and displays of rocks and fossils. In 1937, David White's work was recognized by the development of a trailside exhibit in the Hermit Shale at Cedar Ridge on the South Kaibab Trail, a locale which still displays a surface with ferns (Figure 2-4). This small glassed-over exhibit, built by the Civilian Conservation Corps, was re-dedicated in 2008 during a CCC symposium at Grand Canyon; and it is now undergoing further restoration by Grand Canyon interpretive and Volunteers-In-Parks staff.



Figure 2-4. Trailside exhibit of fossil ferns, constructed by the Civilian Conservation Corps in the Permian Hermit Shale at Cedar Ridge on South Kaibab Trail, as seen *ca.* 1940s? (NPS).

E. D. McKee was instrumental in the development of Yavapai, staffing it himself from time to time during the late 1920s and 1930s (Figure 2-5); that is, when he wasn't doing other naturalist duties in the park, or on one of his far-ranging geology field trips. Over the years, geological interpretation at Yavapai became rather static. The usual wear and tear on the hardware, and changing public perspectives on educational presentations, eventually called for a remodeling of Yavapai by 1980. It was a more sleek—some said "dumbed down"—exhibit now partly commandeered as a tourists' gift shop. McKee, long since having had left the canyon and by now a senior geologist in the USGS, was dismayed over this conversion. Fortunately, in the early 2000s, Yavapai was completely refitted again, this time bringing back detailed and informative modern exhibits on Grand Canyon geology, restoring the building to meet its original mission and brightening its original architecture. It was rededicated in 2007 as the Yavapai Geology Museum.



Figure 2-5. Ranger-Naturalist Edwin D. McKee at work in Yavapai Observation Station, *ca.* 1930 (NPS).

McKee would have been even more ecstatic over the most recent addition to the park's educational offerings in geology and paleontology. In 1994, geologist Karl Karlstrom from the University of New Mexico, and others, proposed the construction of a "Trail of Time" along the already existing paved Rim Trail. After many concept studies and arrangements with the park, a 4.6-km (3-mi) trail was laid with markers and rock samples from every stratigraphic formation and member in the canyon. It can be entered at several points and be informative in either direction. The trail is accessible to wheelchairs and people with strollers and is convenient to shuttle bus stops and parking areas. Best of all, trail users are encouraged to "please touch" the rock specimens.

The trail is inlaid with small bronze markers, 1 m (3 ft) apart, each "giant step" representing one million years in the history of the earth from "Today" by the Yavapai museum westward to the origin of the earth at Maricopa Point, 4,560 Ma (with one more, lone marker miles farther to the west at Pima Point that denotes the comparable "distance" to the origin of the universe, 13,750 Ma). The Trail of Time rock samples include fossils where appropriate (Figure 2-6). They are interpreted by striking informational signs at the appropriate "ages" along the trail. Most of the rocks and slabs, far larger than geologists' usual "hand specimens," were collected along the Colorado River and taken

out of the canyon by raft. In a few cases where strata are not exposed along the river, the rocks were flown out strung together like a necklace beneath a heavy-lift helicopter.



Figure 2-6. Fossil footprints in a slab of Coconino Sandstone, displayed on the Trail of Time. This slab was retrieved from a flagstone quarry near Ash Fork, Arizona (NPS).

As might be expected, the project was very complex, involving dozens of people and calling for substantial funding for concept preparation, planning and on-site testing, logistics of making rock collections, preparation of specimens, designing and manufacturing signage and ground markers, and installation, not to mention administrative assistance and permitting through the park. The recently published guide to the trail by Karl Karlstrom and Laura Crossey (2019) has an engaging, exhaustive text and is richly illustrated. Though written for the interested casual audience, the guide will be seen by professionals as informative to them as well. With it, even those who cannot visit the canyon can experience and learn from the Trail of Time.

The McKeeian Period

In 1927, Edwin D. McKee had been in the U.S. Naval Academy, but he was influenced by his former Boy Scout leader at Boy Scout Troop 1 in Washington, D.C., to take advantage of a summer internship at the Grand Canyon with John C. Merriam. There he was the field assistant to David

White and C. W. Gilmore. (Incidentally, his former troop leader was none other than Grand Canyon's plane-table cartographer, François Matthes.) He enrolled in the geology program at Cornell University and spent subsequent summers at Grand Canyon. The tragedy of Glen Sturdevant's drowning in the Colorado moved McKee into a permanent position as the park's ranger–naturalist. He never looked back.

McKee privately published *Ancient Landscapes of the Grand Canyon Region* in 1931, which of course prominently featured the fossil life found in the area. This booklet went through 30 variously revised printings until 1985. His first geological attentions were given to the large-scale cross-bedding of the Permian desert Coconino Sandstone, so prominently displayed in the canyon. The Carnegie Institution published his first major work in 1933, *The Coconino Sandstone—Its History and Origin*, that included the ichnofauna of invertebrates and vertebrates.

The canyon's other strata beckoned. McKee had become a master of minutiae, looking at sediments and fossils from one end of the canyon to the other. He became the first geologist to systematically trace key beds from one area to another, which was possible by measuring stratigraphic sections at different places in the canyon. His first monograph was yet another Carnegie publication, *The Environment and History of the Toroweap and Kaibab Formations of Northern Arizona and Southern Utah* (1938), on the Permian units that form the rim of the canyon.

Field work had been going apace in the canyon's other Paleozoic strata, too, which would produce important monographs in coming years. First up were the Cambrian formations of the Tonto Group, analyzed in another Carnegie publication in 1945 (delayed partly by the war) with paleontologist Charles Elmer Resser [1889–1943]: *Cambrian History of the Grand Canyon Region*. It was this monograph that for the first time clearly demonstrated the on- and off-lap sequencing of advancing and retreating seas on a landscape, made possible by tracing key beds. This varying, periodically reversing action of the sea, and the changing sedimentology and fossil assemblages produced by this activity, has since become the stuff of elementary geological education; diagrams printed in the book have been widely reprinted in textbooks, too.

McKee left the Grand Canyon in 1938 rather than be moved to be a ranger elsewhere in the park system. He first went to the Museum of Northern Arizona, then on to the University of Arizona where he eventually became geology department chair, and later on to the U.S. Geological Survey as a research geologist in sedimentology. During this time, in addition to publishing the occasional paper, he produced two more major monographs, effectively rounding out the Paleozoic sequence for Grand Canyon. First was the 1969 volume with Raymond Charles Gutschick [1913–2002], *History of the Redwall Limestone of Northern Arizona*, followed in 1982 by McKee's *The Supai Group of Grand Canyon*.

The Redwall and Supai yielded to McKee's quintessential methods of observation and reporting. The corresponding monographs, following on the two-author Cambrian monograph of 1945, are composed of separately authored chapters by authorities in their fields, with observations on lithologies, paleoenvironments, and fossil systematics. In paleontology, the Redwall monograph has separate chapters for crinoids (J. C. Brower), bryozoans (Helen Duncan), blastoids (Donald B.

Macurda, Jr.), corals (William J. Sando), and foraminifera (Betty Skipp), with McKee and Gutschick reserving chapters on the Redwall faunas in general, brachiopods, and "miscellaneous fossil groups: algae and stromatolites, holothurians, trilobites, ostracodes, and fish."

McKee's Supai monograph of 1982 was even more comprehensive, though focusing more on lithologic analyses, including such esoteric subjects as "insoluble residue patterns" (Walter H. Peirce et al.) and "stable isotope analyses" (McKee). Still, fossils were documented in the chapters on "Biostratigraphy of the Watahomigi Formation" (Mackenzie Gordon, Jr.) and "Distribution of Age and Flora" (McKee). There also was a chapter by George H. Billingsley and McKee on "Pre-Supai Buried Valleys" that announced fossil-dated Chesterian-age deposits of paleochannels in the top of the Redwall Limestone. These intermittently exposed valleys were described in 1985 as a new formation, the Surprise Canyon Formation, by Billingsley and Stanley S. Beus. (McKee had earlier named the Permian Toroweap Formation, members of the three formations in the Cambrian Tonto Group, members of the Mississippian Redwall Limestone, and divided the Permian–Pennsylvanian Supai Formation into four formations, elevating the Supai to group rank.) The Surprise Canyon Formation was afforded its own monograph (Billingsley and Beus 1999) that followed the McKee design of separately authored chapters, which included "Megafossil Paleontology" (Beus) and "Conodont Biostratigraphy" (Harriet Martin and James E. Barrick).

Placing all of this Grand Canyon work in broader perspective, McKee focused on studying modern sediments as analogs of paleo-sediments, work that took him first to the Colorado River delta in Mexico. There, his wife, Barbara, satisfactorily explained to quizzical Yaqui Indians in whose boat they were guided through the channels that he "gets paid" to fill bags with mud. His field travels took him around the world, including to Saudi Arabia where Bedouin tribesmen appeared "out of nowhere" to watch the astonishing act of trickling a tanker truck of water into desert sand dunes (so that the dunes could be cut open to allow study of their internal structure). The tribesmen left after having had their waterbags filled. At the USGS in Denver, McKee had a laboratory where he conducted flume-sediment experiments, which he continued even in retirement. He used remote-sensing technology and Skylab observations in his studies of world sand seas as well. Not long before he died, McKee had been on South America's Rio Orinoco, surveying its sediments as an analog to the Supai's Watahomigi Formation. All this, and much more, was a remarkable series of achievements for a man with only a bachelor's degree. He and his wife are buried in the Grand Canyon Cemetery, near David White. The headstone, a water-sculpted piece of Cambrian Tapeats Sandstone, was collected by USGS geologist George Billingsley near Pumpkin Spring in western Grand Canyon.

Super Fossils, Pseudo Fossils

The Proterozoic sequence of sedimentary rocks beneath the Great Unconformity that separates them from the Paleozoic strata—the Unkar and Chuar Groups, comprising the Grand Canyon Supergroup—were studied by geologists as far back as Walcott in the early 1880s, with renewed interest in the early 20th century. A flurry of additional interest came about in the late 1930s as a result of the Carnegie Institution's battery of field studies in the canyon. But first, some museum drawer-cleaning.

For years, the presumed "Precambrian" fossils that Walcott had gathered in the 1880s had been accepted without question by later workers. In the 1930s and 40s their biogenic origins were questioned, and all but *Chuaria* were relegated to the bin of pseudofossils. Even so, the Carnegie-sponsored explorations yielded more. In 1934, Clarence Edgar Van Gundy [1908–1985] examined some of the Supergroup strata exposed in easternmost Grand Canyon. In 1934 he described a new formation that he called the Nankoweap Group, stratigraphically atop the Unkar Group. (For years the Nankoweap Formation was a loner, lying unconformably between the Unkar and Chuar Groups, but more recently it has been reassigned as the basal unit of the Chuar.) He also reported finding a jellyfish (Van Gundy 1937), which may have misled another geologist on Carnegie-sponsored canyon work, Norman Ethan Allen Hinds [1893–1961], to declare the entire Supergroup to be lower Paleozoic in age. Hinds had been working throughout the 1930s on "Algonkian" and other "Precambrian" relationships in the American West. (It should be pointed out that Carnegie was not the sole, though continuously generous, benefactor over the years. For example, in 1933 the Marsh Fund of the National Academy of Sciences provided funds to Edwin McKee for work on the Paleozoic strata; Gregory et al. 1933.)

The jellyfish was sensational news for its implications on the timing of the appearance of multicellular life. In 1941, Ray Smith Bassler [1878–1961] named Van Gundy's jellyfish *Brooksella canyonensis*. Although the title of his paper read "a supposed jellyfish" he assigned it, ironically, to a genus that had been named by C. D. Walcott (he did, though, acknowledge that the Grand Canyon specimen could be inorganic). In 1960, Preston Ercelle Cloud, Jr. [1912–1991], having examined the "problematicum" *B. canyonensis*, took Bassler to task for publishing a retouched photograph and said that the specimen is on an under, rather than upper, surface of a slab; thus it "seems to be the reverse imprint of a subradial fracture system of unknown origin in an underlying shale, bits of which are preserved in the fractures" (Cloud 1960:44).

Van Gundy's jellyfish holds to a life of its own. Over the years it was reinterpreted—if it was organic, that is—as a starlike burrow of a filter-feeding benthic worm (a trace fossil), and in 1969 was accordingly redescribed by Martin Fritz Glaessner [1906–1989] as *Asterosoma*? *canyonensis*, which did nothing more than remove the fossil from among the medusoid taxa. It also was thought to be something possibly akin to the novel late Proterozoic Ediacaran fauna of Australia. By the 1980s, it was suggested to be a complex metazoan trace fossil, after which interest in this fossil waned, although more recently a study of phylogenetic affinities and taphonomy of the genus *Brooksella* (though from the Cambrian of Georgia and Alabama) by Ciampaglio et al. (2006) "tentatively" sustained the Grand Canyon jellyfish as Glaessner's trace, *Asterosoma*? *canyonensis*. It still has not died the death of pseudofossils.

Proterozoic jellyfish in the Grand Canyon have had other, even older, representatives. In 1959 Raymond Manfred Alf [1905–1999] reported "possible fossils" from the Bass Limestone (basal unit of the Unkar Group and the oldest of Grand Canyon's sedimentary formations), which he described as jellyfish impressions. These soon were swept into the pseudofossil bin by Cloud (1960: 43, footnote) who interpreted them as "gas blisters or their impressions."

Pseudojellyfish and other wonders notwithstanding, in the Chuar Group there is a dazzling array of fossils, which even Walcott's notoriously keen eye could never have seen because scanning electron microscopy is needed to study them. In 1977, Bonnie Bloeser, while working on her Master's thesis at UCLA, published with three co-authors a paper on microfossils that she had found in the Chuar Group on Nankoweap Butte. These millimeter-sized, vase-shaped, organic-walled objects of unclear origin occur worldwide, but what she had found was a new genus, which she named *Melanocyrillium* in 1985. The known "vase-shaped microfossils" at Grand Canyon now include at least 16 species and appear to be the fossils of amoebas (Porter et al. 2003; Morais et al. 2019). Many other kinds of microfossils were observed as well, including spheroids and carbonaceous filaments; and the Grand Canyon Supergroup yields traces of various algal-like remains in many strata. Were it not so difficult to gain physical access to these strata, probably many more studies would be possible. Nonetheless, as we shall see, that does not mean that nothing has been done.

Fire

The youngest of Grand Canyon's paleomenageries and paleoherbaria—Pleistocene faunas and flora—were completely unknown until well into the 20th century. In summer 1936, National Park Service employee Willis Evans was in far western Grand Canyon, where Hoover Dam's Lake Mead was rapidly drowning out the last of the Colorado River's rapids. There, he discovered Rampart Cave, finding inside what was called a sloth stable (Anonymous 1936). Soon the discovery was widely reported even in popular public literature, including *Life* magazine (Anonymous 1937).

Willis had found a lair used by now-extinct giant ground sloths; specifically, the Shasta ground sloth, *Nothrotheriops shastensis* (Sinclair). They had left their bones in there, but by far the greatest trove was the abundance of fossil dung (Figure 2-7), in layers up to 2 m (6 ft) deep. The discovery was fortuitous, because at that time there was plenty of young, able, and nearly free labor nearby: the Civilian Conservation Corps. Young men were pressed into service making excavations in the cave, gathering bones (even pieces of fur), and retrieving barrels full of dung that preserved the plant material that once grew in the western Grand Canyon—the stuff of paleoecologists' dreams.

Losing no time on the opportunity, the National Park Service thought that Rampart Cave would make an ideal public exhibit. Later they reported (Anonymous 1936), "Plans for showing the cave to the public have not been completed, but it is hoped that an exhibit in situ can be made. Tentative plans include running a trench through the cave and lining the walls of the trench with glass so that bones or any possible human artifacts may be seen." While humans seem never to have inhabited the cave, the amount of useful material there was enough to ride the same Carnegie education wagon that was making the rounds in the eastern part of the canyon, and Carnegie underwrote some of the research. The exhibit never came to be, even though the possibility of tourism was there. After Hoover Dam was built, tourist concessions evolved in Lake Mead National Recreation Area, which included long boat rides that stopped at Rampart Cave and ran 19 km (12 mi) up into the western part of Grand Canyon. However, the gradual silting in of the lower end of the canyon, as the "red river" slowed and dropped its sediment load in the head of the lake, ended the tours.




Figure 2-7. *Castrocopros martini* Hunt and Lucas, the dung of *Nothrotheriops shastensis* (Sinclair), in Rampart Cave, September 1938 (NPS).

Even though no public exhibition ever came to Rampart Cave, studies of its contents proved quite valuable to paleontologists and paleoenvironmentalists. Archaeologist Mark Raymond Harrington [1882–1971] paid a visit soon after the discovery, commenting in part (1936: 226–227), "I have often thought the critter must have been pretty smelly for his aroma to last ten thousand years!" Soon afterward, in 1938 Jerome Douglass Laudermilk [1893–1956] and Philip Alexander Munz [1892–1974] published the first scientific studies of the flora from Rampart Cave and the nearby Muav Cave, and in 1942 Robert Warren Wilson [1909–2006] published the first comprehensive (though "preliminary") faunal study itemizing 11 vertebrate species, including the ground sloth and the mountain goat *Oreamnos harringtoni* Stock (Figure 2-8). Both publications were Carnegie productions. In 1946 Gordon Cortis Baldwin [1908–1983] briefly updated the findings based on a 1942 survey he had done under Arthur Remington Kellogg [1892–1969]. In 1960, Loye Holmes Miller [1874–1970] reported a variety of fossil birds from Rampart Cave, including the California Condor, although the highly fragmented condition of the material was likely due to "trampling by the ponderous ground sloths." In 1961, Paul Schultz Martin [1928–2010] and colleagues produced the first paleoecological survey of the cave, extending studies to include pollen analyses of the dung.



Figure 2-8. Skull of the extinct mountain goat, *Oreamnos harringtoni* Stock, from a Grand Canyon cave (NPS).

Unfortunately, the cave was not far above a very convenient lake surface, and from time to time it attracted visitors even after it had been gated. In 1977, apparently by accident, the rich, dry dung in the cave was set on fire, probably from a home-made torch used for illumination. It was almost impossible to put out despite valiant efforts at suffocation and dumping tons of water, destroying the better part of the cave's contents; it smoldered for a year (Anonymous 1977, 1978; U.S. National Park Service 1977). The Park Service has since put stronger safeguards in place to keep unauthorized visitors out of the cave. (Due to years of drought in the American West, the level of Lake Mead has fallen dramatically. Today, the Colorado River has reestablished itself in the lake-bottom sediments covering its original channel, where steep banks and heavy vegetation make even stopping here difficult. The lake itself is presently reached some 32 km or 20 mi farther downstream.)

About that Ice Age

Despite the destruction of the valuable Rampart Cave remains, Grand Canyon is nonetheless full of caves. Many were the lairs of animals, perhaps most notably the persistent packrat, *Neotoma*. To these, one may add many more secluded, sheltered rock areas in which packrat middens have been found in abundance. Paleoenvironmental data are locked up in these middens, from which investigators have teased out a story of changing climates in the Grand Canyon since the time of the last glaciation in North America. The dateable and identifiable twigs, leaves, and pollen packed away

and indurated by urine, show the coming and going of desert and woodland environments at various elevations as the advancing and receding glaciers influenced the world's atmosphere. When cooler, wetter conditions arrive, so do more plants and animals. Things are on the warmer, dry side now.

The first systematic approach to analyzing late Pleistocene environments as revealed in Grand Canyon's packrat middens was a 1974 study published by Arthur M. Phillips, III and Thomas R. Van Devender, a preamble to Phillips' 1977 doctoral dissertation on the Grand Canyon Pleistocene. This was expanded upon by Van Devender and Jim I. Mead (1976); and Van Devender (1977) identified four stages of plant community dominance in the western part of the canyon between 24,000 and 8,500 years ago. Mead's 1981 "The Last 30,000 Years of Faunal History Within the Grand Canyon" was a foundational part of work on his 1983 doctoral dissertation on the extinct Harrington's mountain goat (*Oreamnos harringtoni*) in Grand Canyon. The data from nine caves and 50 packrat middens displayed a gradual early Holocene change from woodland to desert faunal and floral communities, a change completed 8,500 years ago. Kenneth Lee Cole and Samantha T. Arundel (2006) pinpointed the Younger Dryas in Grand Canyon, an anomalous cold period in the midst of the warm-up following the last major glaciation. Most early data were from western Grand Canyon sites, but Kenneth Lee Cole's 1981 doctoral dissertation took care of the eastern part of the canyon. Cole and Mead (1981) described the animal fractions of 53 packrat middens in eastern Grand Canyon. Many regional studies corroborate and suggest refinements to the findings made in Grand Canyon; too many to practically discuss here.

"Toys" Lead to Fossils

Stanton's Cave in the Marble Canyon section of Grand Canyon National Park is well-known to river runners. Historically, it is an important site that relates to Robert Brewster Stanton's [1846–1922] flash-in-the-pan 1889–1890 project to survey a railroad route through the canyons from Colorado to California. Scientifically, it is even more important, because it contains not only abundant fossils, but evidence of human use as well. Split-twig figurines, fashioned from a single twig (like willow) split down the middle, with its two fronds twisted and wrapped to create an animal effigy, some with stick spears in the body, were first discovered in 1933 by early river runners who thought they were aboriginal Indians' toys. They are much more than that; considering that many of them had been secreted beneath rock cairns and contain fossil dung pellets, we might see them as a sort of hunting charm. They have been dated here in the canyon to between 2,000 and 4,000 years.

Robert Clark Euler [1924–2002], archaeologist and ethnologist at Grand Canyon National Park, edited in 1978 what still stands as the comprehensive summary of studies at Stanton's Cave. The volume contains separately authored chapters; those of which pertain to paleontological studies are on ungulate remains (C. R. Harrington), zooarchaeological analysis of small vertebrates (John W. Olsen and Stanley J. Olsen), fish remains (Robert Rush Miller and Gerald R. Smith), macroscopic plant materials (Richard H. Hevly), and bird bones (Amadeo M. Rea and Lyndon L. Hargrave), with a paleoecological chapter about the cave by Eleanora I. Robbins, Paul Martin, and Austin Long, and a chapter on the cave "during and after the last Ice Age" by Paul Martin.

Unlike the disaster at Rampart Cave, Stanton's Cave has been well preserved and protected. A "bat gate" at the entrance prohibits unauthorized human access, while allowing the bat colony that lives in the cave easy exit and entrance.

Fossils Go To School

One might notice in this overview that federal involvement was a mainstay of geological research in the late 1800s and early 1900s. At first, geologists were a part of nationally directed surveys in the field. Some independent researchers took the field in the 1910s when the canyon was still very much an unread book. Once the National Park Service had been organized in 1916, and Grand Canyon was finally made a national park in 1919, the Park Service took the reins for a while by encouraging the involvement of the (Washington-based) Carnegie Institution to get researchers into the canyon and to develop outreach programs through museum and trailside exhibits. Almost by brute force, the Grand Canyon's geology and paleontology was being nailed down.

Wartime came, with a reduction in field work, and in the post-war period researchers were striking out on their own. The deep pockets of the Carnegie were not as open to them as had been during the end of the Roaring Twenties and into the Depression, as perspectives on the nature of field work and where it was conducted shifted. More researchers were affiliated with academic institutions that sometimes offered their own funding sources for these kinds of activities, and for publication. Others, though they were working for federal agencies like the USGS, were in the same boat. Occasional research was still going on in the canyon, but the Park Service was out of the picture except in the administrative context of managing resources and issuing permits (once that became a more rigorously applied procedure).

Only a few federal geologists were studying the Grand Canyon, and it fell mostly to the universities and museums to undertake that work. Once the Baby Boom generation reached college age, there was no lack for people, though; and in some measure with the concomitant "age of environmentalism," interest in the fantastic Grand Canyon blossomed anew. Still, this was a time when computers didn't do the things they do today, and "poster sessions" at academic conferences looked like taped-together science projects on easels. Now, in just the past couple of decades, laboratory methodologies, publishing and printing techniques, and the topics of applied studies have so dramatically matured that our forebears in science surely would be as appreciatively dumbfounded as are some of us older geologists.

I take the opportunity here to update my 1984 systematic overview with a few notes that document the kinds of research conducted in the canyon since then. These are a mix of systematic, taxonomic, and paleoecological publications, selected only as a representative set rather than a complete list. (See the online bibliography for a more complete list.) From an administrative viewpoint, it's important to realize that these investigations demonstrate the value and use of resources overseen by federal agencies, and the need of these agencies to gather in data about elements of those resources that have over the past century gone into collections beyond their purview. The "bullet point"-type notes that follow are grouped taxonomically and chronologically, and they further demonstrate the shift to academic sources as well as the implementation of new technologies.

Proterozoic: Microbial mat structures in the Chuar Group have been examined by Bottjer and Hagadorn (2007) and Bohacs et al. (2007). A Master's thesis on *Microfossils from the Neoproterozoic Chuar Group, Grand Canyon, Arizona: Taxonomy, paleoecological analysis and implications for life during the onset of Neoproterozoic glaciation* (Nagy 2008). A study of systematics of organic-walled microfossils of the Chuar Group, including new taxa (Porter and Riedman 2016). Algae in Cryogenean oceans are discussed by Brocks et al. (2017). Perspectives on Ediacaran metazoan ecosystems include comparisons to the Chuar Group (Bowyer et al. 2017).

Paleozoic: An important field conference on Cambrian stratigraphy and paleontology in northern Arizona and southern Nevada (Hollingsworth et al. 2011). Brachiopods of the Redwall Limestone (Carter et al. 2014). Chondrichthyan assemblages of the Surprise Canyon and Watahomigi Formations (Hodnett and Elliott 2018).

Ichnofossils: A new trace fossil, *Angulichnus alternipes*, described from the Bright Angel Shale of Grand Canyon (Elliott and Martin 1987). Coconino Sandstone invertebrate trackways (Braddy 1995). Permian tetrapod tracks from Grand Canyon (Hunt and Santucci 1998). *Bicavichnites martini*, new ichnogenus and ichnospecies, described from the Bright Angel Shale (Lane et al. 2003). Discovery of the "oldest vertebrate trackway in Grand Canyon" (*Chelichnus*?) (Rowland 2019, a paper presented in a symposium convened in honor of Grand Canyon National Park's 2019 centennial, Earth Day 2019, and the 150th anniversary of John Wesley Powell's 1869 pioneering Colorado River expedition). *Ichniotherium* in the Coconino Sandstone of Grand Canyon (Francischini et al. 2019).

Cave fossils: Age and diet of fossil California condors in Grand Canyon (Emslie 1987). Mammalian biogeography (Harris 1990; study sites include several Grand Canyon caves). Parasites in sloth dung from Rampart Cave (Schmidt et al. 1992). A Master's thesis on *Late Pleistocene Aves, Chiroptera, Perissodactyla, and Artiodactyla from Rampart Cave, Grand Canyon, Arizona* (Carpenter 2003). An overview of late Pleistocene Grand Canyon cave faunas (Mead et al. 2003). A late Pleistocene bat mummy (apparently from an eastern Grand Canyon cave) (Mikesic and Chambers 2004). An overview of 50,000 years of vegetation and climate history on the Colorado Plateau in Utah and Arizona (Coats et al. 2008). Molecular identification of the extinct mountain goat, *Oreamnos harringtoni* (Campos et al. 2010). A stable-isotope analysis of subfossil bat guano serving as a long-term environmental archive, based on material from Bat Cave in western Grand Canyon (Wurster et al. 2010). Paleoenvironment of the extinct shrub-ox *Euceratherium collinum* on the Colorado Plateau (Kropf et al. 2007, which includes references to "eastern Grand Canyon caves" that are identified only by archaeological site numbers). Larsen et al. (2018) made a DNA study of "27,000-year-old papillomavirus infection and long-term codivergence with rodents" based on packrat midden collections made on Poston Butte, Grand Canyon, in 1979. Hunt and Lucas (2018) described a new ichnogenus and ichnospecies of Shasta ground sloth dung, *Castrocopros martini*, based on latrinite specimens from Rampart Cave. Delsuc et al. (2019) included Rampart Cave data in DNA sequencing of fossil sloths, revealing an evolutionary history and biogeography of these animals.

If we briefly step across the boundary from Park Service to Bureau of Land Management jurisdiction, we can include a couple of interesting lines of research on the diet and health of aboriginal people living in Antelope Cave on the Uinkaret Plateau to the north of Grand Canyon. A

tick found in a human coprolite was identified by Johnson et al. (2008) as an adult or nymph of *Dermacentor andersoni* (Stiles), probably having passed through a digestive tract; and there is a case study by Reinhard et al. (2012) on determining the pathoecological relationship between ancient diet and modern diabetes based on analyses of human coprolites from the cave. Many studies have also been made of human-made artifacts and of faunal remains here; see particularly the comprehensive overview by Janetski et al. (2013).

Administrative needs: In recent years, there have been various important management projects from the National Park Service, embracing Grand Canyon, including: Vertebrate trackways in National Park Service units (Santucci et al. 1998, 2006). Paleontological resources associated with Park Service caves (Santucci et al. 2001). Paleontology as a tool for conservation (Chure 2002). An inventory of packrat middens in Park Service lands (Tweet et al. 2012). A Master's thesis on *Late Pleistocene and Holocene* Bison *of Grand Canyon and Colorado Plateau: Implications from the use of paleobiology for natural resource management policy* (Martin 2014), followed by the publication by Martin et al. (2017). Application of new technologies in resource inventory and monitoring (Santucci and Wood 2015). Concomitant archaeological and paleontological work in Grand Canyon caves (Conservation Legacy 2016). An inventory of trilobites from Park Service lands (Norr et al. 2016). A register of name-bearing fossil specimens and taxa from National Park Service areas (Tweet et al. 2016). A history of preserving fossils in the national parks (Santucci 2017), which includes Grand Canyon National Park and Grand Canyon–Parashant National Monument.

As these highlights of the last three decades show, paleontology, the science of the long-dead, is very much alive at Grand Canyon. It is important to realize that it was quite convenient that most of the early fossil collections from the canyon went to the Smithsonian Institution, but as the 20th century progressed more collections began to arrive in universities and independent museums. It has become all the more urgent for federal resource managers to keep up with the ever increasing, widely dispersed, collections that are made on federal lands. The means are at hand—through the diligent work of individuals, of course, but also through the use of such documentary records as digital databases.

A Final Paleo Note (of a sort)

The last we had heard of Joseph C. Ives' steamboat, *Explorer*, it was unmanned, adrift in a flood, heading for the Gulf of California. A survey party on the Colorado River delta during the winter of 1929–30 discovered a partially buried iron boat hull, which was investigated more closely during the following July (Sykes 1937:90–92). The wreck was found in Sonora, Mexico, about 48 straight-line km (30 mi) from Yuma and far from an active river channel, clearly aground for a long time. Although it was nothing but a skeleton of ribs and some iron hull panels (Sykes 1937: Figure 10), its dimensions were close to those described by Ives (1861). All the wooden parts of the boat were long gone, as was its boiler, though when that had been salvaged is unknown. What made its identification possible was that there were bolts where the two massive lengthwise timbers had been affixed when *Explorer* was reassembled in December 1857, which were meant to stiffen the unnervingly flexible hull. More importantly, an iron patch on a surviving portion of the transom (Sykes 1937: Figure 11),

made when the engine was raised and the drive-arm opening reconfigured, effectively confirmed the identity of *Explorer*.

Just when *Explorer* was lost is uncertain, though likely before 1865 according to Sykes. Apparently, the boat went out of control during a flood coming from the mouth of the Gila River at Yuma and was secured to the Colorado bank farther downstream. Later it broke free and disappeared. The constantly shifting channels of the Colorado delta, especially during floods, eventually stranded it in a since-abandoned slough, where it was rediscovered four decades later. There's been no report since 1930—an unceremonious end to the vessel that set in motion the first geology field trip to the Grand Canyon. Perhaps, someday, it will be a fragmentary ichnofossil of the Anthropocene.

Acknowledgments

I thank Harry Butowsky and Wayne Ranney for their formal reviews; Diana Boudreau, Justin Tweet, and Klara Widrig for helpful preliminary readings and comments; and Vince Santucci for tapping me on the shoulder for this paper. Photos credited to NPS are courtesy of Grand Canyon National Park Museum Collections.

Literature Cited

Anonymous. 1884a. ["Two miners report finding a series of footprints in Grand Cañon, Arizona"]. Harper's Weekly 28, January 26:63.

Anonymous. 1884b. Ancienne race humaine. L'Exploration (Paris) 17:728. [Rewritten from information in the *Champion* newspaper (Peach Springs, Arizona), which reported that "MM. Spencer et Ridenour" (Charles Spencer and William "Bill" B. Ridenour, miners) had found large footprints attributed by local Indians to an ancestral race.]

Anonymous. 1936. Find well preserved sloths in cave resembling stable. Science News Letter 30(804):157.

Anonymous. 1937. Carnegie expedition tracks prehistoric giant sloth to its lair. Life 3(12):48–50.

Anonymous. 1977. Fossil fire. National Park Service Newsletter 12(4):3.

Anonymous. 1978. Grand Canyon National Park (in Park Briefs section). Courier (U.S. National Park Service) 1(4):8. [Rampart Cave fire extinguished; Paul Martin assesses damage.]

Alf, R. M. 1959. Possible fossils from the early Proterozoic Bass Formation, Grand Canyon, Arizona. Plateau 31:60–63.

Baldwin, G. C. 1946. Notes on Rampart Cave. The Masterkey 20:94–96.

Bassler, R. S. 1941. A supposed jellyfish from the pre-Cambrian of the Grand Canyon. Proceedings of the U.S. National Museum 89:519–522.

Billingsley, G. H., and S. S. Beus. 1985. The Surprise Canyon Formation, an Upper Mississippian and Lower Pennsylvanian (?) rock unit in the Grand Canyon, Arizona. U.S. Geological Survey, Washington, D.C. Bulletin 1605:A27–A33.

Billingsley, G. H., and S. S. Beus, editors. 1999. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona Bulletin 61.

Blodget, L. 1857. Climatology of the United States, and of the temperate latitudes of the North American continent. J. B. Lippincott and Co., Philadelphia, and Trübner and Co., London.

Bloeser, B. 1985. *Melanocyrillium*, a new genus of structurally complex late Proterozoic microfossils from the Kwagunt Formation (Chuar Group), Grand Canyon, Arizona. Journal of Paleontology 59:741–765.

Bloeser, B., J. W. Schopf, R. J. Horodyski, and W. J. Breed. 1977. Chitinozoans from the late Precambrian Chuar Group of the Grand Canyon, Arizona. Science 195:676–679.

Bohacs, K. M., and C. K. Junium. 2007. Microbial mat sedimentary structures and their relation to organic-carbon burial in the Middle Neoproterozoic Chuar Group, Grand Canyon, Arizona, USA. Pages 208–213 *in* J. Scheiber, P. Bose, P. G. Eriksson, S. Sarkar, W. Altermann, and O. Cataneanu, editors. Atlas of microbial mat features preserved within the clastic rock record. Elsevier, Amsterdam and New York. Atlases in Geoscience 2.

Bottjer, D., and J. W. Hagadorn. 2007. Mat growth features. Pages 53–71 *in* J. Scheiber, P. Bose, P. G. Eriksson, S. Sarkar, W. Altermann, and O. Cataneanu, editors. Atlas of microbial mat features preserved within the clastic rock record. Elsevier, Amsterdam and New York. Atlases in Geoscience 2.

Bowyer, F., R. A. Wood, and S. W. Poulton. 2017. Controls on the evolution of Ediacaran metazoan ecosystems: A redox perspective. Geobiology 15(4):516–551.

Braddy, S. J. 1995. The ichnotaxonomy of the invertebrate trackways of the Coconino Sandstone (Lower Permian), northern Arizona. New Mexico Museum of Natural History and Science Bulletin 6:219–224.

Brocks, J. J., A. J. M. Jarrett, E. Sirantoine, C. Hallmann, Y. Hoshino, and T. Liyanage. 2017. The rise of algae in Cryogenian oceans and the emergence of animals. Nature 548:578–581.

Campos, P. F., E. Willersley, J. I. Mead, M. Hofreiter, and M. T. Gilbert. 2010. Molecular identification of the extinct mountain goat, *Oreamnos harringtoni* (Bovidae). Boreas 39:18–23.

Carpenter, M. C. 2003. Late Pleistocene Aves, Chiroptera, Perissodactyla, and Artiodactyla from Rampart Cave, Grand Canyon, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Carter, J. L., D. K. Brezinski, A. D. Kollar, and J. T. Dutro, Jr. 2014. Brachiopoda taxonomy and biostratigraphy of the Redwall Limestone (Lower Mississippian) of Arizona. Annals of the Carnegie Museum 82:257–290.

Chure, D. 2002. Hindsight makes better foresight: Paleontology as a new tool for conservation. Park Science 21(2):43–46.

Coats, L. L., K. L. Cole, and J. I. Mead. 2008. 50,000 years of vegetation and climate history on the Colorado Plateau, Utah and Arizona, USA. Quaternary Research 70:322–338.

Ciampaglio, C. N., L. E. Babcock, C. L. Wellman, A. R. York, and H. K. Brunswick. 2006. Phylogenetic affinities and taphonomy of *Brooksella* from the Cambrian of Georgia and Alabama, USA. Palaeoworld 15:256–265.

Cloud, P. E., Jr. 1960. Gas as a sedimentary and diagenetic agent. American Journal of Science 258-A:35–45.

Cole, K. L. 1981. Late Quaternary environments in the eastern Grand Canyon: Vegetational gradients over the last 25,000 years. Dissertation. University of Arizona, Tucson, Arizona.

Cole, K. L., and J. I. Mead. 1981. Late Quaternary animal remains from packrat middens in the eastern Grand Canyon, Arizona. Journal of the Arizona-Nevada Academy of Science 16:24–25.

Conservation Legacy. 2016. Stewards Individual Placement Program. Pages 9–10 *in* Conservation Legacy annual report: 2016. Conservation Legacy, Durango, Colorado.

Darton, N. H. 1910. A reconnaissance of parts of northwestern New Mexico and northern Arizona. U.S. Geological Survey, Washington, D.C. Bulletin 435.

Darton, N. H. 1925. A résumé of Arizona geology. Arizona Bureau of Mines Bulletin 119, Geological Series 3. [Also an errata sheet.]

Delsuc, F., M. Kuch, G. C. Gibb, E. Karpinski, D. Hackenberger, P. Szpak, J. G. Martínez, J. I. Mead, H. G. McDonald, R. D. E. MacPhee, G. Billet, L. Hautier, and H. N. Poinar. 2019. Ancient mitogenomes reveal the evolutionary history and biogeography of sloths. Current Biology 29:2031–2042.

Dutton, C. E. 1882. Tertiary history of the Grand Cañon district, with atlas. U.S. Geological Survey, Washington, D.C. Monograph 2.

Elliott, D. K., and D. L. Martin. 1987. A new trace fossil from the Cambrian Bright Angel Shale, Grand Canyon, Arizona. Journal of Paleontology 61:641–648.

Emslie, S. D. 1987. Age and diet of fossil California condors in Grand Canyon, Arizona. Science 237:768–770.

Euler, R. C., editor. 1978. The archaeology, geology, and paleobiology of Stanton's Cave, Grand Canyon National Park, Arizona. Grand Canyon Natural History Association Monograph 6.

Francischini, H., S. G. Lucas, S. Voight, L. Marchetti, V. L. Santucci, C. L. Knight, J. R. Wood, P. Dentzien-Dias, and C. L. Schultz. 2019. On the presence of *Ichniotherium* in the Coconino Sandstone (Cisuralian) of the Grand Canyon and remarks on the occupation of deserts by non-amniote tetrapods. Paläontologische Zeitschrift online:1–19.

Gilmore, C. W. 1926. Fossil footprints from the Grand Canyon. Smithsonian Miscellaneous Collections 77(9).

Gilmore, C. W. 1927. Fossil footprints from the Grand Canyon: Second contribution. Smithsonian Miscellaneous Collections 80(3).

Gilmore, C. W. 1928. Fossil footprints from the Grand Canyon: Third contribution. Smithsonian Miscellaneous Collections 80(8).

Gilmore, C. W., and G. E. Sturdevant. 1928. Discovery of fossil tracks on the North Rim of the Grand Canyon. Science 67:216.

Glaessner, M. F. 1969. Trace fossils from the Precambrian and basal Cambrian. Lethaia 2:369–393.

Gregory, W. K., E. D. Merrill, T. W. Vaughan, D. White, and W. H. Howell. 1933. Marsh Fund. Pages 17–18 *in* Report of the National Academy of Sciences: Fiscal Year 1931-1932. U.S. Government Printing Office, Washington, D.C.

Harrington, M. R. 1936. A new ground-sloth den. The Masterkey 10:225–227.

Harris, A. H. 1990. Fossil evidence bearing on southwestern mammalian biogeography. Journal of Mammalogy 71:219–229.

Hodnett, J.-P. M., and D. K. Elliott. 2018. Carboniferous chondrichthyan assemblages from the Surprise Canyon and Watahomigi formations (latest Mississippian–Early Pennsylvanian) of the western Grand Canyon, northern Arizona. Journal of Paleontology 92, Memoir 77.

Hollingsworth, J. S., F. A. Sundberg, and J. R. Foster, editors. 2011. Cambrian stratigraphy and paleontology of northern Arizona and southern Nevada; the 16th Field Conference of the Cambrian Stage Subdivision Working Group, International Subcommission on Cambrian Stratigraphy, Flagstaff, Arizona, and southern Nevada, United States. Museum of Northern Arizona Bulletin 67 and "ISCS 16 Guidebook Errata Sheet".

Hunt, A. P., and S. G. Lucas. 2018. The record of sloth coprolites in North and South America: Implications for terminal Pleistocene extinctions. New Mexico Museum of Natural History and Science Bulletin 79:277–298.

Hunt, A. P., and V. L. Santucci. 1998. Taxonomy and ichnofacies of Permian tetrapod tracks from Grand Canyon National Park, Arizona. Pages 94–96 *in* V. L. Santucci, and L. McClelland,

editors. National Park Service paleontological research. U.S. National Park Service, Geologic Resources Division Technical Report NPS/NRGRD/GRDTR-98/01.

Ives, J. C. 1861. Report upon the Colorado River of the West, explored in 1857 and 1858. U.S. 36th Congress, 1st Session, Senate Executive Document [no number], 5 separately paginated parts in 1 volume. [Also as House Document 90.]

James, J. F. 1892. Studies in problematic organisms—the genus *Scolithus*. Geological Society of America Bulletin 3:32–44.

Janetski, J. C., D. E. Newman, and J. D. Wilde. 2013. A report of archaeological excavations at Antelope Cave and Rock Canyon Shelter, northwestern Arizona. Brigham Young University, Provo, Utah. Museum of Peoples and Cultures Occasional Paper 19.

Johnson, K. L., K. J. Reinhard, L. Sianto, A. Araújo, S. L. Gardner, and J. Janovy, Jr. 2008. A tick from a prehistoric Arizona coprolite. Journal of Parasitology 94:296–298.

Karlstrom, K., and L. Crossey. 2019. The Grand Canyon Trail of Time companion: Geology essentials for your canyon adventure. [no imprint] ("Design: Mary Williams Design, Flagstaff, AZ", "Printing: Four Colour Printing/Printed in Korea")

Kropf, M., J. I. Mead, and R. S. Anderson. 2007. Dung, diet, and the paleoenvironment of the extinct shrub-ox (*Euceratherium collinum*) on the Colorado Plateau, USA. Quaternary Research 67:143–151.

Lane, A. A., S. J. Braddy, D. E. G. Briggs, and D. K. Elliott. 2003. A new trace fossil from the Middle Cambrian of the Grand Canyon, Arizona, USA. Palaeontology 46:987–997.

Laudermilk, J. D., and P. A. Munz. 1938. Plants in the dung of *Nothrotherium* from Rampart and Muav Caves, Arizona. Carnegie Institution of Washington Publication 487:271–281.

Larsen, B. B., K. L. Cole, and M. Worobey. 2018. Ancient DNA provides evidence of 27,000-year-old papillomavirus infection and long-term codivergence with rodents. Virus Evolution 4(1):vey014.

Lull, R. S. 1904. Nature's hieroglyphics. Popular Science Monthly, December:139–149.

Lull, R. S. 1918. Fossil footprints from the Grand Canyon of the Colorado. American Journal of Science (4th series) 45:337–346.

Martin, J. M. 2014. Late Pleistocene and Holocene *Bison* of Grand Canyon and Colorado Plateau: Implications from the use of paleobiology for natural resource management policy. Thesis. East Tennessee State University, Johnson City, Tennessee.

Martin, J. M., R. A. Martin, and J. I. Mead. 2017. Late Pleistocene and Holocene *Bison* of the Colorado Plateau. Southwestern Naturalist 62:14–28.

Martin, P. S., B. E. Sabels, and D. Shutler, Jr. 1961. Rampart Cave coprolite and ecology of the Shasta ground sloth. American Journal of Science 259:102–127.

McKee, E. D. 1931. Ancient landscapes of the Grand Canyon region: The geology of Grand Canyon, Zion, Bryce, Petrified Forest, and Painted Desert. [No imprint] (printed by Lockwood-Hazel Co., Atchison, Kansas].

McKee, E. D. 1933. The Coconino sandstone—its history and origin. Carnegie Institution of Washington Publication 440:77–115.

McKee, E. D. 1938. The environment and history of the Toroweap and Kaibab formations of northern Arizona and southern Utah. Carnegie Institution of Washington Publication 492.

McKee, E. D. 1982, The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D., and R. C. Gutschick. 1969. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and C. E. Resser. 1945. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

Mead, J. I. 1981. The last 30,000 years of faunal history within the Grand Canyon, Arizona. Quaternary Research 15:311–326.

Mead, J. I. 1983. Harrington's extinct mountain goat (*Oreamnos harringtoni*) and its environment in the Grand Canyon, Arizona. Dissertation. University of Arizona, Tucson, Arizona.

Mead, J. I., Coats, L. L., and Schubert, B. W. 2003. Late Pleistocene faunas from caves in the eastern Grand Canyon, Arizona. Pages 64–68 *in* B. W. Schubert, J. I. Mead, and R. W. Graham, editors. Ice age cave faunas of North America. Indiana University Press, Bloomington and Indianapolis, Indiana.

Mikesic, D. G., and C. Chambers. 2004. Return of the bat mummy; a desert cave preserves a 10,000-year-old spotted bat. Bats 22(4): cover, inside front cover, 8–11.

Miller, L. 1960. Condor remains from Rampart Cave, Arizona. The Condor 62(1):70. [See also "Correction" 62(4):298.]

Morais, L., D. J. G. Lahr, I. D. Rudnitzki, B. T. Freitas, G. R. Romero, S. M. Porter, A. H. Knoll, and T. R. Fairchild. 2019. Insights into vase-shaped microfossil diversity and Neoproterozoic biostratigraphy in light of recent Brazilian discoveries. Journal of Paleontology 93(4):612–627.

Nagy, R. M. 2008. Microfossils from the Neoproterozoic Chuar Group, Grand Canyon, Arizona: Taxonomy, paleoecological analysis and implications for life during the onset of Neoproterozoic glaciation. Thesis. University of California at Santa Barbara, Santa Barbara, California.

Newberry, J. S. 1861. Geological report; *in* J. C. Ives. Report upon the Colorado River of the West, explored in 1857 and 1858. U.S. 36th Congress, 1st Session, Senate Executive Document [no number], Part 3, 154 p. [separately paginated part in volume], with four maps in volume pocket.

Noble, L. F. 1909. The geology of the Shinumo area, Grand Canyon, Arizona. Dissertation. Yale University, New Haven, Connecticut.

Noble, L. F. 1910. Contributions to the geology of the Grand Canyon, Arizona.—The geology of the Shinumo area. American Journal of Science (4th series) 29:369–386, 497–528.

Noble, L. F. 1914. The Shinumo quadrangle, Grand Canyon district, Arizona. U.S. Geological Survey, Washington, D.C. Bulletin 549.

Noble, L. F. 1922. A section of the Paleozoic formations of the Grand Canyon at Bass Trail. Pages 23–73 *in* Shorter contributions to general geology, 1922. U.S. Geological Survey, Washington, D.C. Professional Paper 131-B.

Norr, M. R., V. L. Santucci, and J. S. Tweet. 2016. An inventory of trilobites from National Park Service areas. New Mexico Museum of Natural History and Science Bulletin 74:179–203.

Phillips, A. M., III. 1977. Packrats, plants and the Pleistocene in the lower Grand Canyon. Dissertation. University of Arizona, Tucson, Arizona.

Phillips, A. M., III, and T. R. Van Devender. 1974. Pleistocene packrat middens from the lower Grand Canyon of Arizona. Journal of the Arizona Academy of Science 9:117–119.

Porter, S. M., and L. A. Riedman. 2016. Systematics of organic-walled microfossils from the ca. 780-740 Ma Chuar Group, Grand Canyon, Arizona. Journal of Paleontology 90:815–853.

Porter, S. M., R. Meisterfeld, and A. H. Knoll. 2003. Vase-shaped microfossils from the Neoproterozoic Chuar Group, Grand Canyon: A classification guided by modern testate amoebae. Journal of Paleontology 77:409–429.

Powell, J. W. 1875. Exploration of the Colorado River of the West and its tributaries; explored in 1869, 1870, 1871, and 1872, under the direction of the Secretary of the Smithsonian Institution. Washington, U.S. Government Printing Office, 291 p., map and river profile on two sheets laid in. [U.S. 43rd Congress, 1st Session, House Miscellaneous Document 300, Serial 1622.]

Powell, J. W. 1895. Canyons of the Colorado. Flood and Vincent, The Chatauqua-Century Press, Meadville, Pennsylvania.

Ranney, W. 2013. Geologists through time in the Grand Canyon: From Newberry to a new century. Pages 113–118 *in* R. D. Quartaroli, compiler and editor. A rendezvous of Grand Canyon historians: Ideas, arguments, and first-person accounts: Proceedings of the Third Grand Canyon History Symposium, January 2012. Grand Canyon Historical Society, Flagstaff, Arizona.

Ranney, W. 2014. A pre-21st century history of ideas on the origin of the Grand Canyon, Geosphere 10(2):233–242.

Ransome, F. L. 1908. Pre-Cambrian sediments and faults in the Grand Canyon of the Colorado. Science 27:667–669.

Ransome, F. L. 1916. Some Paleozoic sections in Arizona and their correlations. U.S. Geological Survey, Washington, D.C. Professional Paper 98-K:133–166.

Reinhard, K. J., K. L. Johnson, S. LeRoy-Toren, K. Wieseman, I. Teixeira-Santos, and M. Vieira. 2012. Understanding the pathoecological relationship between ancient diet and modern diabetes through coprolite analysis; a case example from Antelope Cave, Mojave [*sic*] County, Arizona. Current Anthropology 53:506–512.

Richmond, A. 1998. Rails to the rim: Milepost guide to the Grand Canyon Railway (4th edition). Grand Canyon Railway, Williams, Arizona.

Richmond, A. 2017. The story of Grand Canyon Railway: Cowboys, miners, presidents and kings (new edition). Grand Canyon Association, Grand Canyon, Arizona.

Rowland, S. M. 2019. The oldest vertebrate trackway in Grand Canyon and the dawn of reptiles. Pages 14–15 *in* K. E. Karlstrom, L. J. Crossey, S. Semken, T. Stoeberl, and J. Calhoun, convenors. Grand Canyon Geology and Geoscience Education Public Symposium, April 18-20, 2019; in honor of Grand Canyon National Park's 2019 centennial celebration, Earth Day 2019, and the 150th anniversary of John Wesley Powell's 1869 pioneering Colorado River expedition. [No imprint].

Santucci, V. L. 2017. Preserving fossils in the national parks: A history. Earth Sciences History 36:245–285.

Santucci, V. L., and J. R. Wood. 2015. Application of new technologies supporting paleontological resource inventory and monitoring in Intermountain Region parks. Crossroads in Science 3:73–81.

Santucci, V. L., A. P. Hunt, and M. G. Lockley. 1998. Fossil vertebrate tracks in National Park Service areas. Dakoterra 5:107–114.

Santucci, V. L., A. P. Hunt, T. Nyborg, and J. P. Kenworthy. 2006. Additional fossil vertebrate tracks in National Park Service areas. New Mexico Museum of Natural History and Science, Bulletin 34:152–158.

Santucci, V. L., J. Kenworthy, and R. Kerbo. 2001. An inventory of paleontological resources associated with National Park Service caves. NPS Geological Resources Division, Denver, Colorado. Technical Report NPS/NRGRD/GRDTR-01/02.

Schmidt, G. D., D. W. Duszynski, and P. S. Martin. 1992. Parasites of the extinct Shasta ground sloth *Nothrotheriops shastensis*, in Rampart Cave, Arizona. Journal of Parasitology 78:811–816.

Schuchert, C. 1918a. On the Carboniferous of the Grand Canyon. American Journal of Science (4th series) 45:347–361.

Schuchert, C. 1918b. The Cambrian of the Grand Canyon of Arizona. American Journal of Science (4th series) 45:362–369.

Spamer, E. E. 1984. Paleontology in the Grand Canyon of Arizona: 125 years of lessons and enigmas from the late Precambrian to the present. The Mosasaur 2:45–128.

Spamer, E. E. 1995. Sublime stone: Grand Canyon geology after Powell. Canyon Legacy 24:16–26, back cover.

Sykes, G. 1937. The Colorado delta. Carnegie Institution of Washington Publication 460.

Tweet, J. S., V. L. Santucci, and A. P. Hunt. 2012. An inventory of packrat (*Neotoma* spp.) middens in National Park Service areas. New Mexico Museum of Natural History and Science Bulletin 57:355–368.

Tweet, J. S., V. L. Santucci, and H. G. McDonald. 2016. Name-bearing fossil type specimens and taxa named from National Park Service areas. New Mexico Museum of Natural History and Science Bulletin 73:277–288.

U.S. National Park Service. 1977. Rampart Cave sloth dung fire, 1976-1977. Grand Canyon National Park.

Van Devender, T. R., and J. I. Mead. 1976. Late Pleistocene and modern plant communities of Shinumo Creek and Peach Springs Wash, lower Grand Canyon, Arizona. Journal of the Arizona Academy of Science 11:16–22.

Van Gundy, C. E. 1934. Some observations of the Unkar group of the Grand Canyon Algonkian. Grand Canyon Nature Notes 9(8):338–349.

Van Gundy, C. E. 1937. Jellyfish from the Grand Canyon Algonkian. Science 85:314.

Walcott, C. D. 1880. The Permian and other Paleozoic groups of the Kanab Valley, Arizona. American Journal of Science (3rd series) 20:221–225.

Walcott, C. D. 1883. Pre-Carboniferous strata in the Grand Cañon of the Colorado, Arizona. American Journal of Science (3rd series) 26:437–442, 484.

Walcott, C. D. 1884. Note on Paleozoic rocks of central Texas. American Journal of Science (3rd series) 28:431–433.

Walcott, C. D. 1899. Pre-Cambrian fossiliferous formations. Geological Society of America Bulletin 10:199–244.

White, D. 1929. Flora of the Hermit Shale, Grand Canyon, Arizona. Carnegie Institution of Washington Publication 405.

Wilson, R. W. 1942. Preliminary study of the fauna of Rampart Cave, Arizona. Carnegie Institution of Washington Publication 530:169–185.

Wurster, C. M., D. A. MacFarlane, M. I. Bird, P. Ascough, and N. B. Athfield. 2010. Stable isotopes of subfossil bat guano as a long-term environmental archive: Insights from a Grand Canyon cave deposit. Journal of Cave and Karst Studies 72:111–121.

Chapter 3. Stratigraphy of Grand Canyon National Park

By Timothy B. Connors[1], Justin S. Tweet[2] and Vincent L. Santucci[3]

[1]National Park Service
Geologic Resources Division
12795 W. Alameda Parkway
Lakewood, Colorado 80225

[2]National Park Service
9149 79th Street S.
Cottage Grove, Minnesota 55016
[3]National Park Service

Geologic Resources Division
1849 "C" Street, NW
Washington, D.C. 20240

Introduction

The story that Grand Canyon tells is a spectacle of approximately two billion years of earth history (approximately one-half of the age of the earth) in its rock record, with an equally extensive paleontological component. There is no other place on Earth where the pages of Earth's story can be read so easily by the observer to reveal such a long, rich, geologic history of events that are recorded in the layers. Dr. John Strong Newberry said it best in the mid-19th century: "the most splendid exposure of stratified rocks that there is in the world" (Beus 2003).

Grand Canyon rocks can be simplified into three main packages: Vishnu Basement rocks, Grand Canyon Supergroup rocks, and layered Paleozoic rocks. These are each separated by major unconformities and indicate formation under differing geologic conditions and during different time intervals (Mathis 2006). Colorado Plateau uplift and recent downcutting in the canyon and volcanic activity are also responsible for younger geologic materials as well.

Grand Canyon National Park (GRCA) hosts extensive exposures of many Precambrian and Phanerozoic units ranging in age from Proterozoic to Triassic (Figure 3-1, Table 3-1 and 3-2). These units consist of igneous and metamorphic rocks and numerous sedimentary lithologies (siltstones, sandstones, conglomerates, limestones, and dolostones), many of which are extremely fossiliferous. Paleozoic sedimentary rocks are responsible for approximately 900 m (3,000 ft) of the stairstep topography and viewshed in the Grand Canyon. Mesozoic sedimentary rocks likely once covered the Paleozoic section, but these rocks are now only seen in rare isolated outcrops in GRCA (Billingsley et al. 2019).

This summary presents a focused overview of the stratigraphy of GRCA and does not delve into the broader and complex geologic topics and themes associated with the origin and geologic history of the Grand Canyon itself. It is a brief overview focused on the stratigraphic framework for Grand

Canyon to provide a context for the rich and diverse paleontological resources presented in this report and establishes consistency for the other chapters.




Figure 3-1. Grand Canyon stratigraphy and structural relations (Billingsley et al. 2019: Figure 2). Recently, the Sixtymile Formation was proposed to be Cambrian (not Proterozoic) (Karlstrom et al. 2018, 2020) and the Nankoweap was moved into the Chuar Group (Dehler et al. 2017). Mesozoic rocks younger than the Chinle Formation are not found within the boundaries of GRCA, but are present in the immediate vicinity.

Table 3-1. Grand Canyon area stratigraphy (after Billingsley et al. 2019: Table 2, with updates). Lower case denotes informal names. Mesozoic rocks younger than the Chinle Formation are not found within the boundaries of GRCA, so are omitted, but are present in the immediate vicinity.

Era	Period/Subperiod	Formation	Member
Mesozoic (Mz)	Triassic (Tr)	Chinle Formation	Shinarump Member
	Triassic (Tr)	Moenkopi Formation	Holbrook Member
	Triassic (Tr)	Moenkopi Formation	Moqui Member
	Triassic (Tr)	Moenkopi Formation	Wupatki Member
Paleozoic (Pz)	Permian (P)	Kaibab Formation	Harrisburg Member
	Permian (P)	Kaibab Formation	Fossil Mountain Member
	Permian (P)	Toroweap Formation	Woods Ranch Member
	Permian (P)	Toroweap Formation	Brady Canyon Member
	Permian (P)	Toroweap Formation	Seligman Member
	Permian (P)	Coconino Sandstone	–
	Permian (P)	Hermit Formation	–
	Permian (P)	Esplanade Sandstone	–
	Permian (P)	Pakoon Limestone	–
	Pennsylvanian (IP)	Wescogame Formation	–
	Pennsylvanian (IP)	Manakacha Formation	–
	Pennsylvanian (IP)	Watahomigi Formation	–
	Mississippian (M)	Surprise Canyon Formation	–
	Mississippian (M)	Redwall Limestone	Horseshoe Mesa Member
	Mississippian (M)	Redwall Limestone	Mooney Falls Member
	Mississippian (M)	Redwall Limestone	Thunder Springs Member
	Mississippian (M)	Redwall Limestone	Whitmore Wash Member
	Devonian (D)	Temple Butte Formation	–
	Cambrian (Є)	Frenchman Mountain Dolostone	–
	Cambrian (Є)	Muav Limestone	Havasu Member
	Cambrian (Є)	Muav Limestone	Gateway Canyon Member
	Cambrian (Є)	Muav Limestone	Kanab Canyon Member
	Cambrian (Є)	Muav Limestone	Peach Springs Canyon Member
	Cambrian (Є)	Muav Limestone	Rampart Cave Member
	Cambrian (Є)	Bright Angel Shale	Flour Sack Member
	Cambrian (Є)	Bright Angel Shale	red-brown member
	Cambrian (Є)	Tapeats Sandstone	–
	Cambrian (Є)	Sixtymile Formation	–
Neoproterozoic (Z)	–	Kwagunt Formation	Walcott Member
	–	Kwagunt Formation	Awatubi Member
	–	Kwagunt Formation	Carbon Butte Member
	–	Galeros Formation	Carbon Canyon Member
	–	Galeros Formation	Jupiter Member
	–	Galeros Formation	Tanner Member
	–	Nankoweap Formation	–
	–	Cardenas Basalt	–

Table 3-1 (continued). Grand Canyon area stratigraphy (after Billingsley et al. 2019: Table 2, with updates). Lower case denotes informal names. Mesozoic rocks younger than the Chinle Formation are not found within the boundaries of GRCA, so are omitted, but are present in the immediate vicinity.

Era	Period/Subperiod	Formation	Member
Mesoproterozoic (Y)	–	Dox Formation	Ochoa Point Member
	–	Dox Formation	Comanche Point Member
	–	Dox Formation	Solomon Temple Member
	–	Dox Formation	Escalante Creek Member
	–	Shinumo Sandstone	–
	–	Hakatai Shale	–
	–	Bass Formation	Hotauta Conglomerate Member
Paleoproterozoic (X)	–	Zoroaster Granite	–
	–	Vishnu Schist	–

Table 3-2. Overview of GRCA stratigraphy and paleontology. See the various chapters for more paleontological information.

Formation	Age	Paleontological Resources
Upper Cenozoic sediments	Pleistocene–Holocene	Almost entirely late Pleistocene–Holocene fossils, predominantly from dry cave and crevice deposits; horsetails, ferns, gnetales, conifers, and angiosperms (macrobotanical), driftwood, pollen, nematodes and their eggs (in dung), bivalves, aquatic and terrestrial gastropods, ostracodes, arthropods (ticks, scorpions, millipedes, beetles, flies, hemipterans, cicadas, hymenopterans, lepidopterans, antlions, grasshoppers), osteichthyans, frogs, salamanders, turtles, lizards, snakes, birds (accipitriforms, anseriforms, apodiforms, cathartiforms, charadriiforms, columbiforms, falconiforms, galliforms, gruiforms, passeriforms, pelecaniforms, piciforms, podocipediforms, strigiforms), mammals (sloths, shrews, rodents, rabbits, bats, carnivorans, proboscidean, horses, artiodactyls), dung (lizard, mammal), bird regurgitation pellets, packrat middens, ringtail middens, and bird eggshell and nests
Chinle Formation	Late Triassic	Petrified wood
Moenkopi Formation	Early–?Middle Triassic	Invertebrate trace fossils and vertebrate tracks (*Rotodactylus*)
Kaibab Formation	early Permian	Dasycladacean algae, sponges, rugose corals, conulariids, bryozoans, brachiopods, bivalves, nautiloids, gastropods, scaphopods, trilobites, crinoids, echinoids, chondrichthyans (ctenacanthiforms, hybodontiforms, euselachians, petalodontiforms, and holocephalans), platysomid actinopterygians, indeterminate actinopterygian teeth and scales, and invertebrate burrows and trails
Toroweap Formation	early Permian	Bryozoans, brachiopods, bivalves, nautiloids, gastropods, scaphopods, ostracodes, crinoids, echinoids, and stromatolites
Coconino Sandstone	early Permian	Invertebrate burrows, trails, and tracks, anamniote tracks (cf. *Amphisauropus* and *Ichniotherium*), reptile tracks (cf. *Dromopus*, *Erpetopus*, and *Varanopus*), synapsid tracks (cf. *Tambachichnium*), and undetermined tetrapod tracks

Table 3-2 (continued). Overview of GRCA stratigraphy and paleontology. See the various chapters for more paleontological information.

Formation	Age	Paleontological Resources
Hermit Formation	early Permian	Horsetails, "seed ferns", ginkgoes, conifers, undetermined plants, eurypterids, insects, invertebrate burrows, trails, and tracks, anamniote tracks (*Amphisauropus*, *Batrachichnus*, and *Ichniotherium*), reptile tracks (*Dromopus*, *Erpetopus*, and *Hyloidichnus*), synapsid tracks (*Dimetropus*), undetermined tetrapod tracks, and possible microbial features
Esplanade Sandstone (in west transitions to Pakoon Limestone)	early Permian	Conifers (*Walchia*), undetermined plants, bioclasts of marine invertebrates (corals, bryozoans, pelmatozoans, and brachiopods or bivalves), invertebrate burrows and trails, and foraminifers
Wescogame Formation	Late Pennsylvanian	Undetermined plants, bioclasts of invertebrate fossils (bryozoans, pelmatozoans, and brachiopods or bivalves), holocephalan chondrichthyans, invertebrate burrows, trails, and tracks, anamniote tracks (cf. *Amphisauropus*, *Batrachichnus*, and cf. *Limnopus*), of reptiles (*Varanopus*), undetermined tetrapod tracks, foraminifers, and microbial features
Manakacha Formation	Middle Pennsylvanian	Undetermined ferns and other plants, bioclasts of invertebrate fossils (bryozoans, ostracodes, pelmatozoans, and brachiopods or bivalves), microbial trace fossils (stromatolites), invertebrate burrows, trails, and tracks, undetermined tetrapod tracks, foraminifers, and "algal" bioclasts (calcispheres and *Girvanella*)
Watahomigi Formation	Early–Middle Pennsylvanian	Equisetopsids (*Calamites*), "seed ferns" (*Neuropteris*), conifers (*Cordaites* and *Walchia*), *Taeniopteris*, undetermined plants, corals including tabulates, conulariids, bryozoans, brachiopods, bivalves, gastropods, trilobites, crinoids, echinoids, conodonts, chondrichthyans (holocephalan and indeterminate dermal denticles), undetermined fish teeth, microbial trace fossils (stromatolites), invertebrate burrows and trails, foraminifers, and "algae"
Surprise Canyon Formation	Late Mississippian	*Calamites*, *Lepidodendron*, *Lepidostrobophyllum*, undetermined wood and other plant fossils, rugose and tabulate corals, bryozoans, brachiopods, bivalves, gastropods, trilobites, ostracodes, asteroids, blastoids, crinoids, echinoids, conodonts, chondrichthyans (thrinacodontids, xenacanthiforms, symmoridforms, ctenacanthiforms, hybodontiforms, euselachians, indeterminate elasmobranchs, paraselachians, orodontiforms, eugenodontiforms, petalodontiforms, and holocephalans), indeterminate actinopterygians, indeterminate tetrapods, microbial trace fossils ("algal" laminations, oncolites, stromatolites), invertebrate burrows and trails, foraminifers, and "algae"
Redwall Limestone	Early–Middle Mississippian	Rugose and tabulate corals, bryozoans, brachiopods, nautiloids, gastropods, trilobites, blastoids, crinoids, holocephalan chondrichthyans, undetermined fish teeth, invertebrate burrows and trails, foraminifers, "algae", and calcispheres
Temple Butte Formation	Middle–Late Devonian	Rugose corals, brachiopods, gastropods, conodonts, placoderms, sarcopterygians, indeterminate fish, invertebrate burrows and trails, and trace fossils or stromatoporoid sponges
Frenchman Mountain Dolostone	middle–late Cambrian	Invertebrate burrows and trails

Table 3-2 (continued). Overview of GRCA stratigraphy and paleontology. See the various chapters for more paleontological information.

Formation	Age	Paleontological Resources
Muav Limestone	middle Cambrian	Sponges, brachiopods, hyoliths, helcionelloids, trilobites, eocrinoids, enigmatic invertebrates (*Chancelloria*, *Scenella*), invertebrate burrows and trails, and *Girvanella*-like structures (oncolites)
Bright Angel Shale	middle Cambrian	Cryptogam spores, brachiopods, hyoliths, trilobites, bradoriids, eocrinoids, enigmatic invertebrates (*Chancelloria*, *Tontoia*), microbial wrinkle structures, invertebrate burrows and trails, leiospheres, filament mats resembling *Nematothallus*, non-marine cryptospores, terrestrial algal cell clusters, enigmatic fossils (*Margaretia*), and possibly sponges
Tapeats Sandstone	early–middle Cambrian	Brachiopods, trilobites, and invertebrate burrows and trails
Sixtymile Formation	early Cambrian	Potential undetermined fragment
Kwagunt Formation	middle Neoproterozoic (late Tonian)	Stromatolites and other microbial features, acritarchs and colonial organic-walled microfossils, microbial filaments, vase-shaped microfossils, various unspecified microfossils, “vampire traces” on microfossils, chemical evidence for possible sponges, and possible meiofaunal traces
Galeros Formation	middle Neoproterozoic (late Tonian)	Stromatolites and other microbial features, acritarchs and colonial organic-walled microfossils, microbial filaments, various unspecified microfossils, and “vampire traces”
Nankoweap Formation	middle Neoproterozoic (late Tonian)	None to date, unless *Brooksella canyonensis* is organic
Cardenas Basalt	late Mesoproterozoic	None to date; fossils are unlikely but not impossible
Dox Formation	late Mesoproterozoic	Stromatolites; also dubiofossils
Shinumo Quartzite	late Mesoproterozoic	None confirmed; also dubiofossils
Hakatai Shale	late Mesoproterozoic	Stromatolites and other microbial features in the Bass–Hakatai transition zone; also dubiofossils
Bass Formation	middle–late Mesoproterozoic	Stromatolites and other microbial structures, possible microfossils, and possible microbial filaments; also dubiofossils
Paleoproterozoic–Mesoproterozoic basement	late Paleoproterozoic–early Mesoproterozoic	Unfossiliferous igneous and high-grade metamorphic rocks

Precambrian Stratigraphy of Grand Canyon

The Precambrian rocks of GRCA consist of igneous, metamorphic, and sedimentary rocks. Precambrian sedimentary and igneous rocks are generally only exposed in the eastern and central Grand Canyon regions along the canyon depths, while Proterozoic crystalline rocks are only exposed along the Colorado River and tributaries in eastern and western Grand Canyon (Billingsley et al. 2019).

The base of the Precambrian section is composed of various igneous and metamorphic bodies of Paleoproterozoic age, overlaid by a series of primarily sedimentary units. The “Vishnu Basement rocks” (consisting of generically the Elves Chasm Gneiss, and granites and schists) will not be

treated here as they do not contain paleontological resources. The “Grand Canyon Supergroup rocks” are divided into the Mesoproterozoic Unkar Group (consisting of the Bass Formation, Hakatai Shale, Shinumo Sandstone, Dox Formation, and Cardenas Basalt), and the Neoproterozoic Chuar Group (Nankoweap Formation, Galeros Formation, and Kwagunt Formation). These sedimentary rocks are discussed in further detail in the Precambrian paleontology chapter; capsule descriptions are included here.

Grand Canyon Supergroup: Unkar Group

The Unkar Group consists of the Mesoproterozoic Bass Formation, Hakatai Shale, Shinumo Sandstone, Dox Formation, and Cardenas Basalt (Figures 3-2 and 3-3).




Figure 3-2. Mesoproterozoic rocks of the Unkar Group (Grand Canyon Supergroup) in eastern Grand Canyon. Yo=Hotauta Conglomerate Member; Yb=Bass Formation; Yc=Cardenas Basalt; Yh=Hakatai Shale; Ys=Shinumo Sandstone (Billingsley et al. 2019: Figure 3).




Figure 3-3. Mesoproterozoic rocks in contact with lower Tonto Group (Tapeats Sandstone and Bright Angel Shale) in eastern Grand Canyon (Billingsley et al. 2019: Figure 4).

Unkar Group: Bass Formation (Mesoproterozoic)

The Bass Formation is primarily composed of dolomite, with some interbedded sandstone, mudstone, and pebble conglomerate, about 60 to 100 m (200 to 330 ft) thick. The basal part of the formation is a cobble conglomerate known as the Hotauta Member. The Bass Formation is interpreted as mostly shallow to restricted marine, with increasing clastic input over time. It grades into the overlying Hakatai Shale. The base of the formation dates to approximately 1254 Ma (million years ago) (Timmons et al. 2005, 2012). This formation is significant for preserving the oldest evidence of life in GRCA.

Unkar Group: Hakatai Shale (Mesoproterozoic)

The Hakatai Shale is a clastic unit consisting of primarily siltstone and fine-grained sandstone, with lithologies ranging from mudstone to conglomerate, varying from 137 to 300 m (450 to 980 ft) thick. The upper contact with the Shinumo Sandstone is unconformable. It is interpreted as a shallow water unit from marginal marine, tidal flat and deltaic settings, deposited at least in part after 1187 Ma (Timmons et al. 2005, 2012).

Unkar Group: Shinumo Sandstone (Mesoproterozoic)

The Shinumo Sandstone is a mostly quartzitic sandstone interpreted as a high-energy shoreface unit. It is approximately 355 to 410 m (1,160 to 1,350 ft) thick and has a gradational contact with the overlying Dox Formation (Timmons et al. 2005, 2012). It may be as old as ca. 1170 Ma (Timmons et al. 2012) or as young as 1140 Ma (Mulder et al. 2017).

Unkar Group: Dox Formation (Mesoproterozoic)

The Dox Formation is predominantly composed of red sandstone. It is interpreted as initially a fluvial to deltaic unit, becoming more marine over time (Timmons et al. 2012; Mulder et al. 2017). It has been divided into four members, in ascending order: the Escalante Creek, Solomon Temple, Comanche Point, and Ochoa Point Members, with a combined thickness of approximately 920 m (3,020 ft) (Elston 1989a). Deposition occurred between approximately 1140 and 1104 Ma (Timmons et al. 2012; Mulder et al. 2017).

Unkar Group: Cardenas Basalt (Mesoproterozoic)

The Cardenas Basalt is an unfossiliferous basalt unit formed by eruptions that began near the end of Dox Formation deposition, as shown by interfingering Dox beds and Cardenas lava flows. It is about 300 m (980 ft) thick and dates to approximately 1104 Ma. Its upper contact with the Nankoweap Formation is unconformable (Timmons et al. 2005, 2012).

Grand Canyon Supergroup: Chuar Group

The Chuar Group consists of the Neoproterozoic Nankoweap, Galeros and Kwagunt Formations, each with their own members.

Chuar Group: Nankoweap Formation (Neoproterozoic)

The Nankoweap Formation can be divided into a lower red unit of hematite-cemented sandstone and mudstone, and an upper white unit of siltstone and sandstone (Timmons et al. 2012). These two informal members have an unconformable contact, and the overall thickness of the formation varies greatly from 113 to more than 250 m (370 to more than 820 ft) (Elston 1989a). This unit was recently found to be much younger than previously inferred by dating detrital zircons, at less than approximately 782 Ma, and has been added to the Chuar Group (Dehler et al. 2017).

Chuar Group: Galeros Formation (Neoproterozoic)

The Galeros Formation is a dominantly clastic unit, mostly mudstones with some sandstone and dolomite beds. It is divided into four members, in ascending order the Tanner, Jupiter, Carbon Canyon, and Duppa Members. Like the similar overlying Kwagunt Formation, it is interpreted as representing primarily wave- and tidal-influenced marine deposition and supratidal. The upper contact with the Kwagunt Formation is gradational, and the two together are about 1,600 m (5,250 ft) thick (Dehler et al. 2001, 2012). It dates from after 782 Ma to approximately 751 ± 7.6 Ma (Rooney et al. 2018).

Chuar Group: Kwagunt Formation (Neoproterozoic)

The Kwagunt Formation is lithologically similar to the Galeros Formation and is also divided into several members (in ascending order the Carbon Butte, Awatubi, and Walcott Members). It was also primarily deposited in shallow subtidal to intertidal settings, with more frequent episodes of subaerial exposure than the Galeros Formation (Dehler et al. 2001, 2012). Deposition occurred after approximately 751 Ma to about 729 ± 0.9 Ma (Rooney et al. 2018).

Paleozoic Stratigraphy of Grand Canyon

Outcrops of 17 distinct Paleozoic formations have been reported in GRCA, ranging in age from the Cambrian to the Triassic. These rocks vary greatly in depositional environments ranging from open marine to eolian terrestrial. As discussed in following chapters, they also preserve a broad array of fossils, from Cambrian invertebrate burrows and trails, to Devonian fish, to Mississippian crinoids, to Pennsylvanian vertebrate tracks, to Permian plants and insects. GRCA boasts one of the most complete Paleozoic records in the National Park System, particularly from the Late Devonian through the end of the Permian.

Tonto Group (lower–middle Cambrian)

The Tonto Group (Figures 3-3 and 3-4) consists of the Sixtymile Formation, Tapeats Sandstone, Bright Angel Shale (or Formation), Muav Limestone (or Formation), and Frenchman Mountain Dolostone (Karlstrom et al. 2020). Historically it included only the Tapeats, Bright Angel, and Muav Formations. It is misleading to consider these three units as simple "layer cake" beds. The formations are defined by lithology and because deposition occurred over many small-scale marine regressions and transgressions during the overall marine transgression, the lithologies intertongue extensively, making mapping complicated (Beus and Billingsley 1989; Huntoon 1989).

Tonto Group: Sixtymile Formation (lower Cambrian)

The Sixtymile Formation was thought to be Precambrian in age until recently, when dating of detrital zircons established it as Cambrian in age (Karlstrom et al. 2018). It is only found in a few areas of eastern GRCA and is composed of red- to white sandstone and siltstone with chert and interformational breccia (Elston 1979). What had previously been described as the lowest part of the formation has been transferred to the upper Kwagunt Formation (Timmons et al. 2001). The Sixtymile Formation was deposited in lacustrine, fluvial, and shallow marine settings in fault-controlled basins. Detrital zircons indicate it was deposited between 520 and 509 Ma, making it contemporaneous in part with rocks of the lower Tonto Group in the western Grand Canyon and Lake Mead regions (Karlstrom et al. 2018). There is an angular unconformity between the Sixtymile Formation and the overlying Tapeats (Tonto Group) (A. Mathis, pers. comm., December 2019).




Figure 3-4. Upper Tonto Group (Cm=Muav Limestone; Cu="undifferentiated dolomites", now the Frenchman Mountain Dolostone), Temple Butte Formation (Dtb), and Redwall Limestone (Mrw=Whitmore Wash Member; Mrt=Thunder Springs Member; Mrm=Mooney Falls Member) in eastern Grand Canyon (Billingsley et al. 2019: Figure 5).

Tonto Group: Tapeats Sandstone (lower–middle Cambrian)

The Tapeats Sandstone is a medium- to coarse-grained, cliff-forming conglomeratic sandstone (Beus and Billingsley 1989). At GRCA, this unit is deposited on what had been the hilly terrain of weathered Precambrian rocks (the Grand Canyon Supergroup in eastern GRCA, the older Vishnu Basement in western GRCA) (Middleton and Elliott 2003). The unconformity with all underlying Precambrian rocks is known as the Great Unconformity. The base of the Tapeats Sandstone is locally conglomeratic, with mudstone and fine sandstone becoming common toward the top, where the Tapeats Sandstone forms a transition zone with the overlying Bright Angel Shale (Middleton and Elliott 2003). Three members may be apparent in the western part of the canyon, with a shale (mudstone)-rich member sandwiched between sandstone members (Elston 1989d).

Historically, the Tapeats Sandstone and the rest of the Tonto Group were considered to span much of the Cambrian and were interpreted as a classic example of a gradual marine transgression in which the nearshore sands of the Tapeats Sandstone were replaced by successively deeper marine deposits of the Bright Angel Shale and Muav Limestone (McKee and Resser 1945). More recent study indicates that the marine transgression responsible for the Tonto Group took place over a much shorter time frame (Karlstrom et al. 2018). West of GRCA, the upper Tapeats Sandstone includes

rocks deposited approximately 508 to 504 Ma, while in eastern GRCA, the Tapeats Sandstone has a maximum depositional age of 505.4 ± 8.0 Ma (Karlstrom et al. 2018).

The Tapeats Sandstone is typically interpreted as representing shallow marine sand deposition under significant tidal influence, with more terrestrial environments toward the base (Hereford 1977; Middleton 1989; Middleton and Elliott 2003). However, the formation may have been more continental overall, perhaps a fluvial braidplain (Baldwin et al. 2004). The thickness of the formation varies from very thin or absent where deposited over prominent paleotopographic highs, to 90 m (300 ft), 12 to 15 m (40 to 50 ft) of which are part of a transition zone (Beus and Billingsley 1989).

Tonto Group: Bright Angel Shale (Middle Cambrian)

The Bright Angel Shale is a mixed formation mostly composed of shale (mudstone) to fine-grained sandstone (Middleton and Elliott 2003). The rocks are sometimes divided into numerous members (McKee 1945; Spamer 1984; Beus and Billingsley 1989). It appears to have been deposited between approximately 505 to 501 Ma in Grand Canyon (Karlstrom et al. 2018). It has a complex gradational and intertonguing relationship with the overlying Muav Limestone (Middleton 1989). To simplify matters, Elston (1989d) has suggested transferring the lower portion of the Muav Limestone to the Bright Angel Shale. The Bright Angel Shale is about 107 to 150 m (350 to 500 ft) thick (Billingsley 2000).

The Bright Angel Shale is generally interpreted as a shallow marine shelf unit (Middleton and Elliott 2003). The various members correspond to minor transgressions and regressions (Elston 1989d; Beus and Billingsley 1989). When interpreted as more continental, the rocks are instead seen as representing estuary and tidal flat settings (Baldwin et al. 2004) influenced by storm events (Elliott and Martin 1987). The lack of acritarchs in the mudstones, the dominant lithology of the formation, may be evidence for minimal marine influence in those rocks (Baldwin et al. 2004).

Tonto Group: Muav Limestone (middle Cambrian)

The Muav Limestone is composed of limestone, dolomite, thin shale (mudstone) and siltstone, and conglomerate (Spamer 1984; Middleton and Elliott 2003), and forms cliffs at GRCA (Middleton and Elliott 2003). Like the Bright Angel Shale, it can be divided into multiple members (Spamer 1984; Middleton 1989; Middleton and Elliott 2003). Trilobites of the Muav Limestone can be attributed to the same part of the Cambrian as the Bright Angel Shale of eastern GRCA (Karlstrom et al. 2018), so it is likely not substantially younger. It is between 45 and 245 m (150 and 800 ft) thick (Spamer 1984). Its upper contact is an unconformity with the unnamed dolomite unit (Beus and Billingsley 1989).

The Muav Limestone is interpreted as representing subtidal to supratidal offshore deposits (Middleton and Elliott 2003). The various members correspond to minor transgressions and regressions (Elston 1989d; Beus and Billingsley 1989). There are also some tidal flat deposits, particularly in the western part of GRCA (Wanless 1973; Baldwin et al. 2004).

Tonto Group: Frenchman Mountain Dolostone (middle–?upper Cambrian)

Above the Muav Limestone at GRCA is a unit historically known as the "undifferentiated dolomites", now assigned to the Frenchman Mountain Dolostone by Karlstrom et al. (2020). It consists of white to gray dolomite unit with thin layers of shale (mudstone) between beds, especially in the lower part of the unit. Its exact age is uncertain, due to the paucity of fossils. The thickness varies from 60 to 140 m (200 to 450 ft) (Beus and Billingsley 1989). This unit is found in western GRCA (Middleton 1989). It is also sometimes called the "Supra-Muav" or "Grand Wash Dolomite" in the literature (Middleton 1989), although the latter name is precluded from formal usage because "Grand Wash" is already in use for a different unit in the area (Elston 1989d). This unit is interpreted as shallow subtidal to possibly intertidal in depositional setting (Middleton and Elliott 2003), deposited in a regressing sea (Spamer 1984).

Temple Butte Formation (Middle–Upper Devonian)

The Temple Butte Formation (Figure 3-4) is a dolomite (dolostone) and sandstone unit, becoming mostly dolomitic in western GRCA (Beus 1989). In eastern GRCA, it is discontinuous, filling channels cut into the underlying Cambrian rocks. It becomes a thicker and continuous layer in the western part of the park, with dolomite over the channel fill (Spamer 1984). Some descriptions have combined part of the unnamed Cambrian dolomite with the formation (Beus 2003a). Conodont fossils have been used to date the Temple Butte Formation to the late Middle and early Late Devonian (Beus 1980). Where present, it is up to 135 m (450 ft) thick in western GRCA (Beus and Billingsley 1989). Both the lower contact with Cambrian rocks and the upper contact with the Redwall Limestone are unconformities (Spamer 1984).

Most of the Temple Butte Formation is interpreted as representing shallow, subtidal, open marine settings in western Grand Canyon, although some of the dolomite may be supratidal and the channel fill could correspond to tidal channels in eastern Grand Canyon (Beus 2003a). A transgression occurred during the deposition of this unit, moving west to east (Beus 1989).

Redwall Limestone (Lower–Middle Mississippian)

The Redwall Limestone (Figures 3-4 and 3-5) is made up mostly of limestone, with some dolomite, chert, and mudstone (Beus et al. 1989). The most detailed description of the unit is McKee and Gutschick (1969a), which includes data from a number of GRCA localities. This cliff-forming unit is actually gray, but in the canyon it has been stained red on the surface by iron oxides washed from the overlying Supai Group (McKee and Gutschick 1969b). There are four members, all present at GRCA; from oldest to youngest, they are the Whitmore Wash, Thunder Springs, Mooney Falls, and Horseshoe Mesa Members (McKee 1963). The Whitmore Wash Member is mostly limestone and dolomite, thickening from 15 m (50 ft) in eastern GRCA to 36 m (120 ft) in western GRCA, which dates to the Early Mississippian. The Thunder Springs Member is a distinctively banded unit, due to alternating carbonate and chert beds. It is 30 m (100 ft) thick in eastern GRCA, increasing to 43 m (140 ft) in western GRCA. It is slightly younger than the Whitmore Wash Member. The Mooney Falls Member is a massive cliff-forming limestone, and spans from 76 m (250 ft) thick in eastern GRCA to 104 m (340 ft) thick in western GRCA. It dates to the early Middle Mississippian. Finally, the Horseshoe Mesa Member, composed of limestone ledges, is thinnest, ranging from 0 to 30 m (0

to 100 ft) thick. It is absent where the overlying channel-filling Surprise Canyon Formation is present. It is slightly younger than the Mooney Falls Member (Beus and Billingsley 1989). Within the formation, there is a depositional hiatus between the Thunder Springs and Mooney Falls Members that becomes progressively greater from west to east (Beus 1989). Shortly after its deposition, the upper part of the Redwall Limestone eroded to form a karst terrain (McKee and Gutschick 1969d) and erodes into overhangs and caves today (McKee and Gutschick 1969b).



Figure 3-5. Redwall Limestone (Mr), Surprise Canyon Formation (Ms), and overlying Supai Group (Pwa=Watahomigi Formation; Pm=Manakacha Formation; Pwe=Wescogame Formation; Pe=Esplanade Sandstone) (Billingsley et al. 2019: Figure 6).

The Redwall Limestone records two marine transgression-regression cycles. The older and larger cycle is represented by the transgressional Whitmore Wash Member and the regressional Thunder Springs Member, and the second cycle is represented by the transgressional Mooney Falls Member and the regressional Horseshoe Mesa Member (Beus 1989). The marine body transgressed from west to east, forming a shallow sea (Beus 2003b). Several types of limestone and other rocks are found throughout the members and correspond to different environments on the shelf (McKee and Gutschick 1969c). Distinct fossil assemblages are found from these different settings. For example, featureless limestone appears to represent lime mud deposits that were not conducive to life, with the only numerous fossils being massive colonial corals. Oolitic limestone (limestone composed of small spherical particles) is probably from warm shallow water with moderate energy, and has a faunal

assemblage of foraminifera, corals, ostracodes, and sea cucumbers, with algal structures (McKee and Gutschick 1969d).

Fossil preservation in the Redwall Limestone is quite variable, depending on the matrix, organisms, and environment. Fossils found in chert are often the best, though usually preserved as molds (McKee and Gutschick 1969d). Fossils in GRCA's Whitmore Wash Member were often destroyed when limestone was altered to dolomite (Beus 2003b), a common phenomenon in southeastern GRCA (McKee and Gutschick 1969e). Fossils are common in the Thunder Springs and Mooney Falls members (Beus and Billingsley 1989). The limestone beds of the Thunder Springs Member are crinoid-rich, while the chert beds are formed by silicified bryozoan limestones and mudstones (Beus and Billingsley 1989). The best fossils are found in the chert beds (Beus 2003b). Bryozoans dominate the Thunder Springs Member fossil assemblages in eastern GRCA, while crinoids dominate central GRCA, and a mixed bryozoan-brachiopod-gastropod-crinoid fauna is present in western GRCA. Fossils of the Mooney Falls Member are only well-preserved in a few scattered zones. Otherwise, specimens are fragmentary (McKee and Gutschick 1969e). Fossils are rare again in the Horseshoe Mesa Member (Beus and Billingsley 1989); however, when present, Horseshoe Mesa Member fossils are well-preserved (Beus 2003b).

Surprise Canyon Formation (Upper Mississippian)

The Surprise Canyon Formation (Figure 3-5) is a discontinuous unit found filling paleo-valleys and other karst features eroded in the upper Redwall Limestone. The Surprise Canyon Formation also occurs in caves in the Redwall Limestone's Mooney Falls and Horseshoe Mesa Members (Billingsley and Beus 1985). It is found only in the Grand Canyon region. Formally named in 1985, its outcrops were first thought to be part of the Redwall Limestone or the Watahomigi Formation (Billingsley and Beus 1985). After it was recognized as a distinct unit, but before it was formally described, it was known as the pre-Supai buried valleys or canyons (Billingsley and McKee 1982; Spamer 1984). The lower portion is composed of fluvial conglomerate and sandstone with some mudstone and siltstone (Beus 2003b). The coarsest material is found near the base, grading up into sandstone (Beus and Billingsley 1989). The middle portion is a cliff-forming marine limestone. Finally, the upper part includes marine slope-forming siltstone, sandstone, and silty to sandy limestone (Beus 2003b). Most of the limestone of the upper unit is at the top, so there is a siltstone-sandstone slope above the middle unit's cliff leading to a cliff higher in the upper unit (Billingsley and McKee 1982). The valleys filled by the Surprise Canyon Formation are as much as 120 m (400 ft) deep. It was deposited a few million years after the Redwall Limestone, and dates to the end of the Mississippian (Beus 2003b). The upper contact is an unconformity with the Watahomigi Formation (Beus 1989).

The Surprise Canyon Formation forms a dendritic drainage system that can be traced through GRCA (Billingsley and Beus 1999). Flow moved from east to west (Beus 2003b). The three parts of the formation formed under different conditions. In general, the lower sandstone/conglomerate portion is interpreted as fluvial, the middle limestone portion is interpreted as marine, and the upper silty portion is interpreted as estuary. The eastern depositional area may have been fluvial during its entire deposition (Beus 2003b). An alternate paleoenvironmental interpretation for the entire formation is as

a more widespread shallow sea. This interpretation would be more consistent with the distribution of some of the marine fossils, but is not favored (Beus 2003b).

Supai Group

The Supai Group (Figure 3-5) was recognized for many years as the Supai Formation in the Grand Canyon. It was designated as a group and was divided into four formations in 1975. In ascending order, these are the Watahomigi Formation, Manakacha Formation, Wescogame Formation, and Esplanade Sandstone (McKee 1975). The coeval Pakoon Limestone intertongues with the Esplanade Sandstone in western GRCA (Blakey and Knepp 1989). The Supai Group as a whole is thought of as a broad coastal plain, over which the sea advanced from the west and retreated several times. The four formations represent different stages of several transgressive-regressive cycles, with the depositional setting oscillating between continental (particularly eolian) and shallow marine environments (Blakey 2003).

Supai Group: Watahomigi Formation (Lower–Middle Pennsylvanian)

The Watahomigi Formation (Figure 3-5) is composed of mudstone, siltstone, limestone, and dolostone. The lower and upper portions are slope-forming red beds, and the middle is a ledge-forming carbonate (Blakey 2003). These parts can be recognized throughout the Grand Canyon (Beus and Billingsley 1989). Carbonates dominate western GRCA and mudstone dominates the eastern outcrops in the park, with very little of the middle unit present (McKee 1982b). It is 24 to 91 m (80 to 300 ft) thick at GRCA, becoming thicker from east to west (Beus and Billingsley 1989). The formation mostly dates to the Early Pennsylvanian. An erosional horizon represented by a conglomerate marks both the base of the upper section and the Early–Middle Pennsylvanian boundary (McKee 1982b). It was deposited after a short hiatus following the deposition of the Surprise Canyon Formation (Beus 1989). The upper contact with the Manakacha Formation may be another unconformity (Blakey and Knepp 1989), or conformable (Blakey 2003).

The Watahomigi Formation is interpreted as a shoreline unit, deposited in shallow marine to coastal plain settings (Blakey and Knepp 1989). It is part of a marine transgression (McKee 1982c). The upper portion had more marine influence than the lower portion (Blakey 2003). During the Early Pennsylvanian, a sea was present west of the modern Little Colorado River, which expanded to the east during the early middle Pennsylvanian (McKee 1982a). Fossils in the Watahomigi Formation suggest low energy conditions (McKee 1982d), but possibly too energetic or with too much sand and silt for extensive coral growth (Gordon 1982).

Supai Group: Manakacha Formation (Middle Pennsylvanian)

The Manakacha Formation (Figure 3-5) is primarily a mix of sandstone and limestone, with some mudstone, conglomerate, and dolostone (Blakey and Knepp 1989). It is usually exposed as a lower cliff and upper slope, with a conglomeratic zone between the two. Unlike other Supai Group formations, there is not a basal conglomerate (Beus and Billingsley 1989). Carbonates are prominent in western GRCA, grading to sandstone and mudstone in central GRCA, and then mudstone and sandstone in eastern GRCA. The top of the unit is a widely recognized channeled surface that marks an unconformity (McKee 1982b). Its thickness is relatively consistent throughout the park, ranging

from 61 to 84 m (200 to 275 ft) thick (Beus and Billingsley 1989). The Manakacha Formation dates to the early Middle Pennsylvanian.

The Manakacha Formation was initially interpreted as a dominantly marine formation (McKee 1982c), representing marine shelf to open marine environments, with mudstone limited to restricted marine environments and the dominant sandstone and limestone deposited under high energy (Blakey and Knepp 1989). More recently, it has been interpreted as dominantly eolian. Eolian deposition began encroaching from the north into the area that had been submerged by the marine transgression of the Watahomigi Formation (Blakey 2003).

Supai Group: Wescogame Formation (Upper Pennsylvanian)

The Wescogame Formation (Figure 3-5) is a mixed unit, with limestones prominent in extreme western GRCA, sandstones dominant in central GRCA, and mudstones increasing in prominence in eastern GRCA. It is exposed as a lower cliff and upper slope (Blakey 2003). It is the most complex of the Supai Group formations, with rapidly shifting rock types (Blakey and Knepp 1989). The thickness is between 30 and 69 m (100 and 225 ft) at GRCA (Beus and Billingsley 1989). Both the upper and lower contact are unconformities (Blakey 2003). The Wescogame Formation dates to the end of the Late Pennsylvanian.

The Wescogame Formation is interpreted as predominately eolian, representing one or more large dune fields (Blakey 2003). Fluvial, coastal plain, shoreline, shelf, and open marine settings are also likely represented in its various rock types (Blakey and Knepp 1989).

Supai Group: Esplanade Sandstone (lower Permian)

The Esplanade Sandstone (Figures 3-5 and 3-6) is a quartz-rich sandstone, with basal and upper slope-forming beds of finer sediments (McKee 1982c). It was deposited during the early Permian (McKee 1982d). The lower portion of the Esplanade Sandstone intertongues with the Pakoon Limestone in western GRCA (Blakey and Knepp 1989). The combined Esplanade Sandstone–Pakoon Limestone thickens from east to west, going from 91 m (300 ft) thick in eastern GRCA to more than 137 m (450 ft) in the western part of the park (Billingsley 1997). The lower contact with the Wescogame Formation and the upper contact with the Hermit Formation are unconformable. The base of the unit in eastern and central GRCA is a conglomerate that fills paleochannels in the Wescogame Formation (Beus and Billingsley 1989).

The depositional environment of this formation has been interpreted in multiple ways. The marine interpretation sees the Esplanade Sandstone as mostly high-energy marine sandstone with more terrestrial beds at the top and bottom (McKee 1982c; Blakey and Knepp 1989). The more current interpretation is that it is an eolian unit (Beus and Billingsley 1989; Blakey 2003), or part of a large coastal plain (Blakey 2003). The base was probably less eolian than the rest of the unit (Blakey 2003). Marine influence increased to the west, as evidenced by the change into the Pakoon Limestone (McKee 1982c). Some gypsum is also present (Blakey 2003).




Figure 3-6. Uppermost Supai Group (Pe=Esplanade Sandstone), Hermit Formation (Ph), Coconino Sandstone (Pc), Toroweap Formation (Seligman Member=Pts; Brady Canyon Member=Ptb; Woods Ranch Member=Ptw), and Kaibab Formation (Fossil Mountain Member=Pkf; Harrisburg Member=Pkh) (Billingsley et al. 2019: Figure 7).

Pakoon Limestone (lower Permian)

The Pakoon Limestone is a heterogeneous unit including dolomite, limestone, sandstone, mudstone, and gypsum (Blakey and Knepp 1989). It is mostly dolomite and limestone in the Grand Canyon region (Blakey 2003). It intertongues with the lower Esplanade Sandstone in western GRCA (Blakey and Knepp 1989), and dates to the earliest Permian (Blakey 2003). The Pakoon Limestone is interpreted as a clear water, shallow marine unit (Blakey 2003). It is not mapped separately from the Esplanade Sandstone within GRCA (Billingsley and Wellmeyer 2004; Billingsley et al. 2006a).

Hermit Formation (lower Permian)

The Hermit Formation (Figure 3-6; formerly known as Hermit Shale) is a mixed red bed unit composed of very fine grained sandstone, siltstone, and minor mudstone. At GRCA, it is known as a reddish-brown, slope-forming unit (Blakey 2003). Its common alternate name is a misnomer, as it includes very little true shale. The thickness varies greatly from 49 m (160 ft) in eastern GRCA to 244 m (800 ft) in western GRCA (Beus and Billingsley 1989). It dates to the late early Permian (Blakey 2003). Although there is an unconformity between the Hermit Formation and the underlying Esplanade Sandstone with deep channel cuts, there was probably little time between the two (White 1927). The upper contact, with the Coconino Sandstone, is also disconformable, but is sharp (Beus

and Billingsley 1989). The Hermit Formation is interpreted as a broad coastal plain and fluvial, but was also deposited as loess and scattered eolian dunes (Blakey and Middleton 2012). The climate was probably semi-arid, with long hot, dry seasons (White 1929).

Coconino Sandstone (lower Permian)

The Coconino Sandstone (Figure 3-6) is a fine-grained eolian sandstone, changing from white and tan, to brown or red in western GRCA (Beus and Billingsley 1989). This unit is bracketed by other units dated to the late early Permian (Blakey and Knepp 1989). From east to west, the unit thickens rapidly in eastern GRCA to 210 m (700 ft) and then thins to practically nothing in western GRCA. Its base forms a sharp unconformity with the Hermit Formation (Beus and Billingsley 1989). The upper contact with the Toroweap Formation intertongues (Blakey and Knepp 1989). The Coconino Sandstone is interpreted as an eolian unit formed as an erg (Hunt et al. 2005). Sand was deposited by wind action (Blakey and Knepp 1989).

Toroweap Formation (lower Permian)

The Toroweap Formation (Figure 3-6) has been studied extensively and offers striking lateral and vertical changes in lithofacies over a relatively small area. Members with carbonate and evaporite lithologies are more easily discerned in western outcrops and these distinctions become absent in the eastern phase that is mostly cross-bedded sandstone (Turner 2003).

In the west it can be divided into three members in the GRCA area; in ascending order these are the Seligman, Brady Canyon, and Woods Ranch Members The relatively thin Seligman Member appears to intertongue and be conformable with the underlying Coconino Sandstone and is no thicker than 15 m (45 ft) at GRCA (Turner 2003). Above the Seligman Member is the overlying Brady Canyon Member, a cliff-forming carbonate unit composed of limestone and mixed dolostone in western GRCA. The Brady Canyon Member is thickest in western GRCA, up to 93 m (280 ft) thick. The Brady Canyon Member thins uniformly to the east to its depositional edge near Marble Canyon and grades into the overlying Woods Ranch Member, mostly made of repetitive evaporites, limestone, and sandstone. The Woods Ranch Member forms distinctive slopes and attains a maximum thickness of about 60 m (180 ft) (Turner 2003). The Woods Ranch Member is interpreted as a shallow evaporitic marine shelf. The climate during deposition of the Toroweap Formation is thought to have been semi-arid to arid (Turner 2003). At GRCA, gypsum and/or contorted sandstones of the Woods Ranch Member always underlie the Kaibab Formation. (Hopkins and Thompson 2003).

Most fossils in the Toroweap Formation are from the Brady Canyon Member, with fossils in the Woods Ranch Member limited to an unusual *Schizodus* bed near the top of the member (McKee 1938; Rawson and Turner 1974).

Kaibab Formation (lower–middle Permian)

The Kaibab Formation (Figure 3-6) is a complex sedimentary package of numerous lithologies. At GRCA, it forms the canyon rim and is 90 to 120 m (300 to 400 ft) thick (Hopkins and Thompson 2003). Early workers divided the Kaibab Formation into the Gamma, Beta, and Alpha Members (McKee 1938), which have since been subsumed into the Fossil Mountain and overlying Harrisburg members. The Fossil Mountain Member is the equivalent of the Gamma and Beta Members, and the

Harrisburg Member is the equivalent of the Alpha Member (Blakey and Knepp 1989). Chert is a major feature of the Fossil Mountain Member at GRCA, and it is quite voluminous and varied in character and weathers to form distinct recesses along cliff faces. It is mostly attributed to the original distribution and abundance of siliceous sponges and spicules. In the west, the Fossil Mountain Member is more carbonate-rich (fossiliferous limestone) but becomes more siliciclastic eastward (sandstone, sandy carbonate, and dolomite) (Hopkins and Thompson 2003). It thickens westward and ranges from 75 to 205 m (250 to 300 ft) thick, to approximately 60 m (200 ft) at the type section at Fossil Mountain along the south rim. The Harrisburg Member constitutes the uppermost cliffs and ledges at GRCA and is a mixed unit including gypsum, dolostones, sandstone, redbeds, chert, and minor limestone. Thicknesses range from 25 to 90 m (80 to 300 ft) at GRCA, and numerous subunits are discernable in its overall extent (Hopkins and Thompson 2003).

The Kaibab Formation is evidence of an ancient seaway covering the GRCA area in the Permian. A complex depositional history is evidenced by the mixing of carbonates and siliciclastics with numerous variations of subtidal to shallow-marine settings. The Fossil Mountain Member documents a west to east shift of fossiliferous open-marine limestones to restricted-marine sandy dolostones and the Harrisburg records retreat of the Kaibab Sea (Hopkins and Thompson 2003).

Mesozoic Stratigraphy of Grand Canyon

Limited exposures of Mesozoic formations are found at Cedar Mountain near Desert View in GRCA, including the Lower–Middle Triassic Moenkopi Formation and the Upper Triassic Chinle Formation. These units were evaluated for paleontological resources during the 2019 GRCA PaleoBlitz and are discussed in Chapter 10 of this volume. Additionally, the Lower Jurassic Wingate Sandstone, Moenave Formation, Kayenta Formation (and Springdale Sandstone Member), and Navajo Sandstone are known in the surrounding Grand Canyon region (Billingsley et al. 2019) but not within GRCA.

Moenkopi Formation (Lower–Middle Triassic)

The Moenkopi Formation (Figure 3-7) is a continental red-bed unit found across the American Southwest (McKee 1954; Stewart et al. 1972a) that includes marginal marine depositional facies in its western exposures (Nevada and Utah) and regressive freshwater fluvial and lacustrine facies in its eastern exposures (Arizona and New Mexico). The only complete section of Moenkopi Formation exposed within GRCA occurs at Cedar Mountain, adjacent to the far eastern boundary near Desert View. This 2 km (1.2 mi) wide feature is largely covered by loose talus and juniper trees, but includes the Wupatki, Moqui, and Holbrook Members, all of which are also exposed along the nearby Little Colorado River Valley from Cameron to Holbrook, Arizona. Noble (1922) determined that the Moenkopi Formation at Cedar Mountain is nearly 150 m (490 ft) thick. The Wupatki Member at Cedar Mountain is characterized by low mounds of ripple-laminated sandstone, the Moqui Member is a slope-former with interbedded evaporite/channel complexes, and the Holbrook Member includes the cliff-forming “upper massive sandstone”. Fossils have been reported from the Moenkopi Formation at GRCA (Marsh et al. this report), and similar sections nearby are known for producing actinopterygian fish, mastodonsauroid, trematosaurian, and brachyopid temnospondyl amphibians, tanystropheid reptiles, and pseudosuchian archosaurs (Welles 1947, 1969; Nesbitt 2000,

2005a, 2005b). Terrestrial vertebrate (Lucas 2010; Martz and Parker 2017), ichnological biochronology (McKee 1954; Klein and Lucas 2010; Henderek et al. 2017), and U-Pb detrital zircon geochronology (Dickinson and Gehrels 2009) suggest that at least the uppermost part of the Moenkopi Formation is Middle Triassic in age.



Figure 3-7. Mesozoic rocks in eastern GRCA at Cedar Mountain (NPS/DIANA BOUDREAU).

Chinle Formation (Upper Triassic)

The only exposure of the Chinle Formation (Figure 3-7) within eastern GRCA caps the Moenkopi Formation section at Cedar Mountain and is represented by the Shinarump Member (formerly the "Shinarump Conglomerate"; Noble 1922; Repenning et al. 1969; Stewart et al. 1972b). It is approximately 8 m (26 ft) thick here and is characterized by well-cemented channel conglomerates with mud rip-up clasts and pieces of (or entire) petrified conifer trees. No vertebrate fossils are known from the Chinle Formation (Shinarump Member) at GRCA, but terrestrial vertebrate biochronology and U-Pb detrital zircon geochronology of overlying and/or equivalent units constrain the entire Chinle Formation to the Late Triassic (Lucas 2010; Atchley et al. 2013; Riggs et al. 2016; Martz and Parker 2017; Kent et al. 2019).

Conclusions

The Grand Canyon serves as a geologic and paleontologic window into the past. The park contains colorful, awe-inspiring rocks and traces of life that showcase spectacular stratigraphy and tell a vast story of almost two billion years of earth history and organism evolution, making it one of the geologic wonders on Earth. The Vishnu Basement rocks, Grand Canyon Supergroup rocks, and Layered Paleozoic rocks combine to present a story like no place on Earth. These old rocks contrast nicely with the geologically "young" age of the canyon. Magmatism, volcanism, metamorphism, deposition, and erosion are all visible on a grand scale at Grand Canyon, leaving their evidence for

the viewer to decipher this planet's rich geologic history and record of organism evolution over the eons.

Acknowledgments

We wish to acknowledge the thoughtful and thorough reviews of George Billingsley (USGS, Flagstaff, Arizona), for all of his geologic mapping and stratigraphic expertise in GRCA, for being a great friend to the NPS for so many years, for being such an expert on GRCA geology, and for sharing that expertise through his numerous publications. Allyson Mathis has been another excellent source on the stratigraphy for GRCA from her years as an interpretive ranger there and her publishing of the "Yardstick of Geologic Time"; her review of this manuscript led to using more definitive sources for sure and for that we are very grateful. Additional thanks to Monica Erdman (USGS, Menlo Park, California) for the high-resolution images borrowed from Billingsley's photo collection.

Literature Cited

Atchley, S. C., L. C. Nordt, S. I. Dworkin, J. Ramezani, W. G. Parker, S. R. Ash, and S. A. Bowring. 2013. A linkage among Pangean tectonism, cyclic alluviation, climate change, and biologic turnover in the Late Triassic: The record from the Chinle Formation, southwestern United States. Journal of Sedimentary Research 83:1147–1161.

Baldwin, C. T., P. K. Strother, J. H. Beck, and E. Rose. 2004. Palaeoecology of the Bright Angel Shale in the eastern Grand Canyon, Arizona, USA, incorporating sedimentological, ichnological and palynological data. Pages 213–236 *in* D. McIlroy, editor. The application of ichnology to palaeoenvironmental and stratigraphic analysis. Geological Society, London, United Kingdom. Special Publication 228.

Beus, S. S. 1980. Late Devonian (Frasnian) paleogeography and paleoenvironments in northern Arizona. Pages 55–69 *in* T. D. Fouch and E. R. Magathan, editors. Paleozoic paleogeography of the west-central United States: Rocky Mountain paleogeography symposium 1. Society of Economic Paleontologists, Rocky Mountain Section, Denver, Colorado.

Beus, S. S. 1989. Devonian and Mississippian geology of Arizona. Pages 287–311 *in* J. P. Jenney and S. J. Reynolds, editors. Geologic evolution of Arizona. Arizona Geological Society Digest 17.

Beus, S. S., and M. Morales, editors. 2003. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Beus, S. S., and M. Morales. 2003a. Introducing the Grand Canyon. Pages 1–8 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Beus, S. S. 2003a. Temple Butte Formation. Pages 107–114 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Beus, S. S. 2003b. Redwall Limestone and Surprise Canyon Formation. Pages 115–135 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Beus, S. S., and G. H. Billingsley. 1989. Paleozoic strata of the Grand Canyon, Arizona. Pages 122–127 *in* D. P. Elston, G. H. Billingsley, and R. A. Young, editors. Geology of Grand Canyon, northern Arizona (with Colorado River guides): Lees Ferry to Pierce Ferry, Arizona. P. M. Hanshaw. Field trips for the 28th international geological congress. American Geophysical Union, Washington, D.C.

Billingsley, G. H. 1997. The Permian clastic sedimentary rocks of northwestern Arizona. Pages 106–124 *in* F. Maldonado and L. D. Nealey, editors. Geologic studies in the Basin and Range-Colorado Plateau transition in southeastern Nevada, southwestern Utah, and northwestern Arizona, 1995. U.S. Geological Survey, Reston, Virginia. Bulletin 2153.

Billingsley, G. H., and H. M. Hampton. 2000. Geologic map of the Grand Canyon 30’ X 60’ Quadrangle, Coconino and Mohave Counties, northwestern Arizona. U.S. Geological Survey, Reston, Virginia. Geologic Investigations Series Map I-2688. Scale 1:100000.

Billingsley, G. H., and E. D. McKee. 1982. Pre-Supai buried valleys. Pages 137–153 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

Billingsley, G. H., and S. S. Priest. 2013. Geologic map of the Glen Canyon Dam 30’ x 60’ Quadrangle, Coconino County, Arizona. U.S. Geological Survey, Reston, Virginia. Scientific Investigations Map SIM-3268. Scale 1:100,000.

Billingsley, G. H., and J. L. Wellmeyer. 2004. Geologic map of the Mount Trumbull 30’ X 60’ Quadrangle, Mohave and Coconino Counties, northwestern Arizona. U.S. Geological Survey, Reston, Virginia. Geologic Investigations Series Map I-2766. Scale 1:100,000.

Billingsley, G. H, S. S. Beus, and P. Grover. 1999. Stratigraphy of the Surprise Canyon Formation; Chapter B. Pages 9–16 *in* G. H. Billingsley and S. S. Beus, editors. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona Bulletin 61.

Billingsley, G. H., D. L. Block, and H. C. Dyer. 2006a. Geologic map of the Peach Springs 30’ x 60’ Quadrangle, Mohave and Coconino Counties, northwestern Arizona. U.S. Geological Survey, Reston, Virginia. Scientific Investigations Map SIM-2900. Scale 1:100,000.

Billingsley, G. H., T. J. Felger, and S. S. Priest. 2006. Geologic map of the Valle 30’ x 60’ Quadrangle, Coconino County, northern Arizona. U.S. Geological Survey, Reston, Virginia. Scientific Investigations Map SIM-2895. Scale 1:100,000.

Billingsley, G. H., S. S. Priest, and T. J. Felger. 2007. Geologic map of Wupatki National Monument and vicinity, Coconino County, northern Arizona. U.S. Geological Survey, Reston, Virginia. Scientific Investigations Map 2958. Scale 1:24,000.

Billingsley, G. H., S. S. Priest, and T. J. Felger. 2007. Geologic map of the Cameron 30’ x 60’ Quadrangle, Coconino County, northern Arizona. U.S. Geological Survey, Reston, Virginia. Scientific Investigations Map SIM-2977. Scale 1:100,000.

Billingsley, G. H., S. S. Priest, and T. J. Felger. 2008. Geologic map of the Fredonia 30’ x 60’ Quadrangle, Mohave and Coconino Counties, northern Arizona. U.S. Geological Survey, Reston, Virginia. Scientific Investigations Map SIM-3035. Scale 1:100,000.

Billingsley, G. H., P. W. Stoffer, and S. S. Priest. 2012. Geologic map of the Tuba City 30’ x 60’ Quadrangle, Coconino County, Arizona. U.S. Geological Survey, Reston, Virginia. Scientific Investigations Map SIM-3227. Scale 1:50,000.

Billingsley, G. H., G. Goodwin, S. E. Nagorsen, M. E. Erdman, and J. T. Sherba. 2019. Geologic field photograph map of the Grand Canyon region, 1967–2010. U.S. Geological Survey, Reston, Virginia. General Information Product 189.

Blakey, R. C. 2003. Supai Group and Hermit Formation. Pages 136–162 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Blakey, R. C., and R. Knepp. 1989. Pennsylvanian and Permian geology of Arizona. Pages 313–347 *in* J. P. Jenney and S. J. Reynolds, editors. Geologic evolution of Arizona. Arizona Geological Society Digest 17.

Blakey, R. C., and L. T. Middleton. 2012. Geologic history and paleogeography of Paleozoic and early Mesozoic sedimentary rocks, eastern Grand Canyon, Arizona. Pages 81–92 *in* J. M. Timmons and K. E. Karlstrom, editors. Grand Canyon geology; two billion years of Earth’s history. Geological Society of America, Boulder, Colorado. Special Paper 489.

Blakey, R. C., and L. T. Middleton. 1998. Permian rocks in north-central Arizona: A comparison of the sections at Grand Canyon and Sedona. Pages 97–125 *in* E. D. Deubendorfer, editor. Geological excursions in northern and central Arizona. Geological Society of America, Rocky Mountain Section, Flagstaff, Arizona.

Chronic, H. 1952. Molluscan fauna from the Permian Kaibab Formation, Walnut Canyon, Arizona. Geological Society of America Bulletin 63(2):95–165.

Dehler, C. M., M. E. Elrick, K. E. Karlstrom, G. A. Smith, L. J. Crossey, and M. J. Timmons. 2001. Neoproterozoic Chuar Group (~800–742 Ma), Grand Canyon: A record of cyclic marine deposition during global climatic and tectonic transitions. Sedimentary Geology 141–142:465–499.

Dehler, C. M., S. M. Porter, and J. M. Timmons. 2012. The Neoproterozoic Earth system revealed from the Chuar Group of Grand Canyon. Pages 49–72 *in* J. M. Timmons and K. E. Karlstrom, editors. Grand Canyon geology; two billion years of Earth’s history. Geological Society of America, Boulder, Colorado. Special Paper 489.

Dehler, C., G. Gehrels, S. Porter, M. Heizler, K. Karlstrom, G. Cox, L. Crossey, and M. Timmons. 2017. Synthesis of the 780-740 Ma Chuar, Uinta Mountain, and Pahrump (ChUMP) groups, western USA; implications for Laurentia-wide cratonic marine basins. Geological Society of America Bulletin 129(5–6):607–624.

Dickinson, W. R., and G. E. Gehrels. 2009. Insights into North American paleogeography and paleotectonics from U-Pb ages of detrital zircons in Mesozoic strata of the Colorado Plateau, USA. International Journal of Earth Sciences 99:1247–1265.

Elston, D. P. 1979. Late Precambrian Sixtymile Formation and orogeny at top of the Grand Canyon Supergroup, northern Arizona. U.S. Geological Survey, Washington, D.C. Professional Paper 1092.

Elston, D. P. 1989a. Middle and late Proterozoic Grand Canyon Supergroup, Arizona. Pages 94–105 *in* D. P. Elston, G. H. Billingsley, and R. A. Young, editors. Geology of Grand Canyon, northern Arizona (with Colorado River guides): Lees Ferry to Pierce Ferry, Arizona. P. M. Hanshaw. Field trips for the 28th international geological congress. American Geophysical Union, Washington, D.C.

Elston, D. P. 1989b. Correlations and facies changes in lower and middle Cambrian Tonto Group, Grand Canyon, Arizona. P. Pages 131–136 *in* D. P. Elston, G. H. Billingsley, and R. A. Young, editors. Geology of Grand Canyon, northern Arizona (with Colorado River guides): Lees Ferry to Pierce Ferry, Arizona. P. M. Hanshaw. Field trips for the 28th international geological congress. American Geophysical Union, Washington, D.C.

Ford, T. D., and C. M. Dehler. 2003. Grand Canyon Supergroup: Nankoweap Formation, Chuar Group, and Sixtymile Formation. Pages 53–75 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Gordon, M., Jr. 1982. Biostratigraphy of the Watahomigi Formation. Pages 113–135 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

Henderek, R. L., V. L. Santucci, B. Rizner, J. Tweet, and J. R. Wood. 2017. Wupatki National Monument paleontological resources inventory. Natural Resource Report NPS/WUPA/NRR-2017/1524.

Hendricks, J. D., and G. M. Stevenson. 2003. Grand Canyon Group Supergroup: Unkar Group. Pages 39–52 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Hereford, R. 1977. Deposition of the Tapeats Sandstone (Cambrian) in central Arizona. Geological Society of America Bulletin 88(2):199–211.

Hopkins, R. L., and K. L. Thompson. 2003. Kaibab Formation. Pages 196–211 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Hunt, A. P., S. G. Lucas, V. L. Santucci, and D. K. Elliot. 2005. Permian vertebrates of Arizona. New Mexico Museum of Natural History and Science Bulletin 29:10–15.

Huntoon, P. W. 1989. Cambrian stratigraphic nomenclature, Grand Canyon, Arizona: Mappers nightmare. Pages 128–129 *in* D. P. Elston, G. H. Billingsley, and R. A. Young, editors. Geology of Grand Canyon, northern Arizona (with Colorado River guides): Lees Ferry to Pierce Ferry, Arizona. P. M. Hanshaw. Field trips for the 28th international geological congress. American Geophysical Union, Washington, D.C.

Jenson, J. 1986. Stratigraphy and facies analysis of the upper Kaibab and lower Moenkopi formations in Southwest Washington County, Utah. Geology Studies 33(1):21–43.

Karlstrom, K. E., B. R. Ilg, M. L. Williams, D. P. Hawkins, S. A. Bowring, and S. J. Seaman. 2003. Paleoproterozoic rocks of the granitic gorges. Pages 9–38 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Karlstrom, K., and L. Crossey. 2019. The Grand Canyon Trail of Time companion: Geology essentials for your canyon adventure. Four Colour Printing, Korea.

Karlstrom, K., J. Hagadorn, G. Gehrels, W. Matthews, M. Schmitz, L. Madronich, J. Mulder, M. Pecha, D. Giesler, and L. Crossey. 2018. Cambrian Sauk transgression in the Grand Canyon region redefined by detrital zircons. Nature Geoscience 11:438–443.

Karlstrom, K. E., M. T. Mohr, M. D. Schmitz, F. A. Sundberg, S. M. Rowland, R. Blakey, J. R. Foster, L. J. Crossey, C. M. Dehler, and J. W. Hagadorn. 2020. Redefining the Tonto Group of Grand Canyon and recalibrating the Cambrian timescale. Geology in press. doi: https://doi.org/10.1130/G46755.1.

Kent, D. V., P. E. Olsen, C. Lepre, C. Rasmussen, R. Mundil, G. E. Gehrels, D. Giesler, R. B. Irmis, J. W. Geissman, and W. G. Parker. 2019. Magnetochronology of the entire Chinle Formation (Norian age) in a scientific drill core from Petrified Forest National Park (Arizona, U.S.A.) and implications for regional and global correlations in the Late Triassic. Geochemistry, Geophysics, Geosystems 20(11):4654–4664.

Klein, H., and S. G. Lucas. 2010. Tetrapod footprints – their use in biostratigraphy and biochronology of the Triassic. Pages 419–446 *in* S. G. Lucas, editor. The Triassic timescale. Geological Society of London, London, United Kingdom. Special Publication 334.

Lucas, S. G. 2010. The Triassic timescale based on nonmarine tetrapod biostratigraphy and biochronology. Pages 447–500 *in* S. G. Lucas, editor. The Triassic timescale. Geological Society of London, London, United Kingdom. Special Publication 334.

Martz, J. W, and W. G. Parker. 2017. Revised formulation of the Late Triassic Land Vertebrate "Faunachrons" of western North America: Recommendations for codifying nascent systems of vertebrate biochronology. Pages 39–124 *in* K. E. Zeigler and W. G. Parker, editors. Terrestrial depositional systems: Deciphering complexities through multiple stratigraphic methods. Elsevier, Oxford, United Kingdom.

Mather, T. J. 1970. Stratigraphy and paleontology of Kaibab Formation (Permian), Mogollon rim region, Arizona. Journal of Sedimentary Petrology 40(2):770–771.

Mathis, A., and C. Bowman. 2007. Telling time at Grand Canyon National Park. Park Science 24(2):78–83.

Mathis, A. 2006. Grand Canyon yardstick of geologic time: A guide to the canyon's geologic history and origin. Grand Canyon Association, Grand Canyon, Arizona.

McKee, E. D. 1938. The environment and history of the Toroweap and Kaibab formations of northern Arizona and southern Utah. Carnegie Institution of Washington Publication 492.

McKee, E. D. 1945. Stratigraphy and ecology of the Grand Canyon Cambrian. Pages 5–168 *in* E. D. McKee and C. E. Resser. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

McKee, E. D. 1954. Stratigraphy and history of the Moenkopi Formation of Triassic age. Geological Society of America Memoir 61.

McKee, E. D. 1963. Nomenclature for lithologic subdivisions of the Mississippian Redwall Limestone, Arizona: Article 65. Pages C21–C22 *in* Geological Survey Research 1963. U.S. Geological Survey, Washington, D.C. Professional Paper 475-C.

McKee, E. D. 1975. The Supai Group: Subdivision and nomenclature. U.S. Geological Survey, Washington, D.C. Bulletin 1395-J.

McKee, E. D. 1982a. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D. 1982b. Stratigraphy and lithology. Pages 29–50 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D. 1982c. Introductory summary. Pages 21–27 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D. 1982d. Distribution and age of flora and fauna. Pages 75–112 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D., and R. C. Gutschick. 1969a. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969b. Introduction. Pages 1–11 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969c. Interpretation of environments. Pages 553–570 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969d. Sequence of sediments and unconformities. Pages 13–95 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969e. The Redwall faunas. Pages 125–172 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and C. E. Resser. 1945. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

Middleton, L. T. 1989. Cambrian and Ordovician depositional systems in Arizona. Pages 273–286 *in* J. P. Jenney and S. J. Reynolds, editors. Geologic evolution of Arizona. Arizona Geological Society Digest 17.

Middleton, L. T., and D. K. Elliott. 2003a. Tonto Group. Pages 90–106 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Middleton, L. T., D. K. Elliott, and M. Morales. 2003b. Coconino Sandstone. Pages 163–179 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Morales, M. 2003. Mesozoic and Cenozoic Strata of the Colorado Plateau near the Grand Canyon. Pages 212–221 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Mulder, J. A., K. E. Karlstrom, K. Fletcher, M. T. Heizler, J. M. Timmons, L. J. Crossey, G. E. Gehrels, and M. Pecha. 2017. The syn-orogenic sedimentary record of the Grenville Orogeny in southwest Laurentia. Precambrian Research 294:33–52.

Nesbitt, S. J. 2000. A preliminary report on new vertebrate fossil sites, including a microsite from the Holbrook Member of the Moenkopi Formation, Holbrook, Arizona. Mesa Southwest Museum Bulletin 7:17–30.

Nesbitt, S. J. 2005a. Stratigraphy and tetrapod fauna of major quarries in the Moenkopi Formation (Early-Middle Triassic) along the Little Colorado River of northern Arizona. Mesa Southwest Bulletin 11:18–33.

Nesbitt, S. J. 2005b. Osteology of the Middle Triassic pseudosuchian archosaur *Arizonasaurus babbitti*. Historical Biology 17:19–47.

Noble, L. F. 1922. A section of the Paleozoic formations of the Grand Canyon at Bass Trail. Pages 23–73 *in* Shorter contributions to general geology, 1922. U.S. Geological Survey, Washington, D.C. Professional Paper 131-B.

Rawson, R. R., and C. E. Turner. 1974. The Toroweap Formation: A new look. Pages 155–190 *in* T. N. V. Karlstrom, G. A. Swann, and R. L. Eastwood, editors. Geology of northern Arizona, with notes on archaeology and paleoclimate. Geological Society of America, Rocky Mountain Section, Flagstaff, Arizona. Guidebook 27, Part 1, regional studies.

Repenning, C. A., M. E. Cooley, and J. P. Akers. 1969. Stratigraphy of the Chinle and Moenkopi Formations, Navajo and Hopi Indian Reservations. U.S. Geological Survey, Washington, D.C. Professional Paper 521-B.

Riggs, N. R., Z. A. Oberling, E. R. Howell, W. G. Parker, A. P. Barth, M. R. Cecil, and J. W. Martz. 2016. Sources of volcanic detritus in the basal Chinle Formation, southwestern Laurentia, and implications for the Early Mesozoic magmatic arc. Geosphere 12:439–463.

Rooney, A. D., J. Austermann, E. F. Smith, Y. Li, D. Selby, C. M. Dehler, M. D. Schmitz, K. E. Karlstrom, and F. A. Macdonald. 2018. Coupled Re-Os and U-Pb geochronology of the Tonian Chuar Group, Grand Canyon. Geological Society of America Bulletin 130(7–8):1085–1098.

Rowland, S. M. 1987. Paleozoic stratigraphy of Frenchman Mountain, Clark County, Nevada. Pages 53–56 *in* Hill, M. L., editor. Cordilleran section of the Geological Society of America. Geological Society of America, Boulder, Colorado. Centennial Field Guide 6.

Sorauf, J. E., and G. H. Billingsley. 1991. Members of the Toroweap and Kaibab formations, Lower Permian, northern Arizona and southwestern Utah. The Mountain Geologist 28(1):9–24.

Spamer, E. E. 1984. Paleontology in the Grand Canyon of Arizona: 125 years of lessons and enigmas from the late Precambrian to the present. The Mosasaur 2:45–128.

Stewart, J. H., F. G. Poole, and R. F. Wilson. 1972a. Stratigraphy and origin of the Moenkopi Formation and related strata in the Colorado Plateau region. U.S. Geological Survey, Washington, D.C. Professional Paper 691.

Stewart, J. H., F. G. Poole, and R. F. Wilson. 1972b. Stratigraphy and origin of the Chinle Formation and related Upper Triassic strata in the Colorado Plateau region. U.S. Geological Survey, Washington, D.C. Professional Paper 690.

Timmons, J. M., K. E. Karlstrom, C. M. Dehler, J. W. Geissman, and M. T. Heizler. 2001. Proterozoic multistage (ca. 1.1 and 0.8 Ga) extension recorded in the Grand Canyon Supergroup and establishment of northwest- and north-trending tectonic grains in the southwestern United States. Geological Society of America Bulletin 113(1):163–181.

Timmons, J. M., K. E. Karlstrom, M. T. Heizler, S. A. Bowring, G. E. Gehrels, and L. J. Crossey. 2005. Tectonic inferences from the ca. 1255-1100 Ma Unkar Group and Nankoweap Formation, Grand Canyon: Intracratonic deformation and basin formation during protracted Grenville orogenesis. Geological Society of America Bulletin 117(11–12):1573–1595.

Timmons, J. M., J. Bloch, K. Fletcher, K. E. Karlstrom, M. Heizler, and L. J. Crossey. 2012. The Grand Canyon Unkar Group; Mesoproterozoic basin formation in the continental interior during supercontinent assembly. Pages 25–47 *in* J. M. Timmons and K. E. Karlstrom, editors. Grand Canyon geology; two billion years of Earth's history. Geological Society of America, Boulder, Colorado. Special Paper 489.

Turner, C. E. 2003. Toroweap Formation. Pages 180–195 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Tweet, J. S., V. L. Santucci, J. P. Kenworthy, and A. L. Mims. 2009. Paleontological resource inventory and monitoring—Southern Colorado Plateau Network. Natural Resource Technical Report. NPS/NRPC/NRTR—2009/245. Natural Resource Program Center. Fort Collins, Colorado.

Wanless, H. R. 1973. Cambrian of the Grand Canyon: A reevaluation. The American Association of Petroleum Geologists Bulletin 57(4):810–811.

Welles, S. P. 1947. Vertebrates from the Upper Moenkopi Formation of northern Arizona. University of California Publication 27:241–294.

Welles. S. P. 1969. Collecting Triassic vertebrates in the Plateau Province. Journal of the West 8:231–246.

White, C D. 1927. Study of the fossil floras in the Grand Canyon, Arizona. Carnegie Institution of Washington Year Book 26:366–369.

White, C. D. 1929. Flora of the Hermit shale, Grand Canyon, Arizona. Carnegie Institution of Washington Publication 405.

Chapter 4. Precambrian Paleontology of Grand Canyon National Park

By Justin Tweet[1]

[1]National Park Service
9149 79th Street S.
Cottage Grove, Minnesota 55016

Introduction

The Precambrian paleontology of Grand Canyon National Park (GRCA) is fundamentally unlike the paleontology of any other portion of the park's substantial stratigraphic column. There are no shells, teeth, bones, footprints, leaves, or similar large fossils. The only fossils visible to the naked eye are layered structures left by microbial mats and the very largest of a diverse assemblage of microfossils. The tools of choice are not shovels and picks, but chemical treatments and powerful microscopes. Very few visitors would be able to spot a Precambrian fossil. Notwithstanding the humble, inconspicuous nature of these fossils, they offer important glimpses at two stages in the development of life, approximately 1,250 to 1,100 million years ago (Ma) and 780 to 730 Ma, long before the appearance of familiar multicellular organisms with hard parts. The Precambrian fossils of GRCA are the oldest fossils of the park and include type specimens for 18 taxa (Appendix 4-B), such as the organic-walled microfossil *Chuaria circularis* and species of *Melanocyrillium*, the first described vase-shaped microfossil, as well as the notable pseudofossil (resembling a fossil but inorganic in origin) *Brooksella canyonensis*, first described as a possible jellyfish. They also have a place in the history of Precambrian paleontology, from Charles Walcott's early explorations to the present.

History of Research

An excellent summary of the early Precambrian paleontology of GRCA can be found in Spamer (1984). Although Powell (1876) and White (1876) briefly mentioned potential fossils in the Grand Canyon Supergroup (at that time defined as a group), the study of Precambrian fossils at GRCA was pioneered by Charles Doolittle Walcott in the 1880s and 1890s. The study of Precambrian fossils in general was just beginning, and Walcott made several misidentifications among his more lasting discoveries (see also "Notable Pseudofossils and Dubiofossils" below). For example, his erroneous identifications of several "fossils" led him to initially regard the Chuar Group as Cambrian in age (Walcott 1883). Of his initial assemblage, only the "stromatoporoids" (stromatolites) and some of the "brachiopods" (*Chuaria*) proved to be legitimate fossils, and he retreated on several of the identifications later (Walcott 1899). Walcott (1899) includes stratigraphic descriptions of the Grand Canyon Supergroup and the description of *Chuaria circularis* (Figure 4-1). Dawson (1897) also gave the name *Cryptozoon occidentale* to an example of Walcott's "stromatoporoids" during this time frame.

After the 1890s, Walcott's attention turned to other topics, including the overlying Cambrian strata at GRCA, and little research on Precambrian fossils at GRCA was conducted until the 1920s. Toward the end of the 1920s, David White briefly mentioned some fossils and potential fossils from the

Precambrian rocks of GRCA alongside his research on later Paleozoic plant fossils from the canyon (White 1927, 1928a, 1928b, 1929). Because of White's paleobotanical focus, the Precambrian fossils and pseudofossils he described were primarily stromatolites, due to their "algal" origin, and "fucoids", a now-obsolete term for what were then thought to be seaweeds. "Fucoids" as a whole are now known to include various invertebrate burrows and burrow-like sedimentary features such as mud cracks, and the Precambrian "fucoids" appear to be inorganic features.



Figure 4-1. *Chuaria circularis* depicted in Walcott (1899: Plate 27:12). *C. circularis* specimens are up to 5 mm (0.2 in) across, making them the largest of GRCA's Precambrian body fossils.

Between the 1920s and 1970s, most of the work on GRCA Precambrian paleontology focused on objects now generally regarded as pseudofossils. The most significant of these is the long-controversial *Brooksella canyonensis* from the Nankoweap Formation, which was first interpreted as a potential jellyfish (Van Gundy 1937, 1951; Bassler 1941). A running dialog on the interpretation of various enigmatic features took place in the literature (e.g., Seilacher 1956; Alf 1959; Cloud 1968; Glaessner 1969; Nitecki 1971). Toward the end of the 1960s, Ford et al. (1969) published initial work toward a complete stratigraphic re-evaluation of the Chuar Group, and Downie (1969) published a brief discussion of microfossils from these rocks, an area of research that would become increasingly significant.

Beginning with the late 1960s publications, paleontological work in the Precambrian of GRCA has focused on the microfossils of the Kwagunt and Galeros formations of the Chuar Group. Many advances and discoveries were published during the 1970s, including: the formal division of the Chuar Group (Ford and Breed 1973a); the redescription of *Chuaria circularis* (Ford and Breed 1973b, 1977); the first report of GRCA filamentous fossils (Schopf et al. 1973); and the initial description of vase-shaped microfossils (Bloeser et al. 1977). Since the 1970s, Precambrian research at GRCA has documented fossils including acritarchs and other organic-walled microfossils (Vidal and Ford 1985; Nagy et al. 2009; Porter and Riedman 2016), filaments (Horodyski and Bloeser 1983), vase-shaped microfossils (Bloeser 1985; Porter and Knoll 2000; Porter et al. 2003), and apparent puncture traces (Porter 2016a). Chemical biomarkers have also been studied (Summons et al. 1988; Brocks et al. 2016). In recent years there have also been re-assessments of the Nankoweap Formation (Dehler et al. 2017) and Sixtymile Formation (Karlstrom et al. 2018), finding both to be younger than originally thought. In the case of the Sixtymile Formation, it is now known to be

Cambrian in age, but it will be included briefly in this chapter due to the recency of its reassessment and long historical placement in the Proterozoic.

Stratigraphic Distribution of Fossils

The Precambrian rocks of GRCA include crystalline Paleoproterozoic basement rocks and a sequence of Mesoproterozoic and Neoproterozoic sedimentary rocks known as the Grand Canyon Supergroup. The Supergroup is divided into the Mesoproterozoic Unkar Group and the Neoproterozoic Chuar Group, which in turn are both divided into several formations. The Unkar Group includes, in ascending order (oldest first), the Bass Formation, Hakatai Shale, Shinumo Quartzite, Dox Formation, and Cardenas Basalt. The Chuar Group includes the Nankoweap, Galeros, and Kwagunt formations. As mentioned above, the Sixtymile Formation was previously thought to be a Precambrian formation, but is now known to be Cambrian in age.

Although the crystalline rocks beneath the Grand Canyon Supergroup can be found near the western end of GRCA, the rocks of the Supergroup are exposed only in eastern GRCA, from roughly Nankoweap Canyon in the east to Deer Creek Falls in the west; all of their outcrops are limited to GRCA. The Unkar Group is more widely exposed, being found from a short distance below the Little Colorado River to Deer Creek Falls. The Chuar Group is limited to the stretch of the canyon between approximately Nankoweap Canyon and Unkar Creek, and except for outcrops of the Nankoweap Formation on the south/east side of the Colorado River within a few km or mi of the Tanner Trail, is only exposed west or north of the river (Figure 4-2). Outcrops are discontinuous, and it is not uncommon for rocks of the overlying Tonto Group to overlie the ancient crystalline rocks without the Supergroup between them, indicating erosion between the Paleoproterozoic and Cambrian. Fossils are confirmed from the Bass Formation, Dox Formation, Galeros Formation, and Kwagunt Formation. More controversial reports have come from the Hakatai Shale, Shinumo Quartzite, and Nankoweap Formation (Table 4-1).




Figure 4-2. An aerial view showing several topographic features west of the Colorado River in eastern GRCA. The distant mesa is Nankoweap Mesa. Nankoweap Butte is to the left. Kwagunt Butte is near the center of the photo. Awatubi Crest is in the foreground (NPS).

Formations are discussed below in ascending order. See Table 4-1 for a brief summary. The following information is also included in tabular form in Appendix 4-A.

Table 4-1. Summary of GRCA Precambrian stratigraphy, fossils, and depositional settings in descending order of age, from youngest to oldest; Sixtymile Formation included based on historical classification. Details and references can be found in the text

Formation	Age	Fossils Within GRCA	Depositional Environment
Sixtymile Formation	early Cambrian	Potential undetermined fragment	Lacustrine, fluvial, and shallow marine
Kwagunt Formation	middle Neoproterozoic (late Tonian)	Stromatolites and other microbial features, acritarchs and colonial organic-walled microfossils, microbial filaments, vase-shaped microfossils, various unspecified microfossils, “vampire traces” on microfossils, chemical evidence for possible sponges, possible meiofaunal traces	Primarily subtidal to intertidal with active tectonism
Galeros Formation	middle Neoproterozoic (late Tonian)	Stromatolites and other microbial features, acritarchs and colonial organic-walled microfossils, microbial filaments, various unspecified microfossils, “vampire traces”	Primarily subtidal to supratidal, intermittently restricted with active tectonism
Nankoweap Formation	middle Neoproterozoic (late Tonian)	None to date, unless *Brooksella canyonensis* is organic	Primarily marine shoreface with active tectonism
Cardenas Basalt	late Mesoproterozoic	None to date; fossils are unlikely but not impossible	Extrusive igneous
Dox Formation	late Mesoproterozoic	Stromatolites; dubiofossils	Fluvial and deltaic to estuarine to marine over time
Shinumo Quartzite	late Mesoproterozoic	None confirmed; dubiofossils	Marine shoreface
Hakatai Shale	late Mesoproterozoic	Stromatolites and other microbial features in the Bass–Hakatai transition zone; dubiofossils	Marginal marine, tidal, and deltaic
Bass Formation	middle–late Mesoproterozoic	Stromatolites and other microbial structures, possible microfossils and microbial filaments; dubiofossils	Subtidal to supratidal, intermittently restricted with active tectonism
Paleoproterozoic–Mesoproterozoic basement	late Paleoproterozoic–early Mesoproterozoic	Unfossiliferous igneous and high-grade metamorphic rocks	Various; igneous rocks that formed at depth, sedimentary rocks with origins obscured by metamorphism

Paleoproterozoic–Mesoproterozoic basement rocks (upper Paleoproterozoic–lower Neoproterozoic)

The sedimentary rocks of the Grand Canyon Supergroup are deposited on igneous and high-grade metamorphic rocks that include the oldest known rocks in southwestern North America (the Elves Chasm gneiss of 1,840 ± 1 Ma; Holland et al. 2015). These crystalline basement rocks are unfossiliferous and included here for geologic completeness.

Unkar Group: Bass Formation (middle–upper Mesoproterozoic)

The Bass Formation is a mixed unit dominated by dolomite, interpreted as altered original limestone, with interbedded sandstone, mudstone, and pebble conglomerate, and a basal cobble conglomerate known as the Hotauta Member. The depositional environment is interpreted as shallow and sometimes restricted marine with fluvio-tidal clastic input from active tectonism, becoming more clastic in the transition to the Hakatai Shale (Timmons et al. 2005, 2012). The base of the formation dates to approximately 1254.8 ± 1.6 Ma (Timmons et al. 2005), although some contested detrital zircons date to around 1200 Ma (Timmons et al. 2012).

The Bass Formation preserves the oldest evidence of life in GRCA. The most abundant and obvious fossils in the formation are various forms of stromatolites (Figure 4-3). “Algal reefs” were reported in the formation by White (1927, 1928a), who identified tabular stromatolites, lenticular stromatolites, winding or knobby masses, and *Collenia*-type columnar forms (White 1928a). The *Collenia* forms can be divided into *C. undosa*, *C. symmetrica*, and *C. frequens* (Beus et al. 1974). More recently, Lathrop (2018) observed crinkly, domal, small domal, conical, and pseudo-conical stromatolites, and noted that stromatolites were sometimes ripped up and redeposited by mass flows. Stromatolites persist into the transition to the Hakatai Shale; stromatolitic carbonate beds typical of the Bass Formation are found interbedded with clastic beds typical of the Hakatai Shale, to the extent that sometimes Hakatai-type clasts of sandstone and conglomerate are encrusted by stromatolites (Timmons et al. 2012). Possible microbial filaments have been observed in thin sections of Bass Formation stromatolites (Dalton 1972; Beus et al. 1974; Ford and Breed 1977), as well as calcispheres (calcitic microfossils) (Dalton 1972; Lathrop 2018). Lathrop’s calcispheres were found in conical stromatolites and have been replaced by silica.



Figure 4-3. Bass Limestone stromatolites (not in life position) (NPS/CASSI KNIGHT).

Other types of potential fossils from the Bass Formation are tantalizing but unconfirmed or considered to be pseudofossils. In the former category, there are: “algal” scales and vertical plates (White 1928a); mysterious *Chuaria*-like objects first mentioned by White (1928a); objects first reported as potential bivalved animals (Smith 1966) that may be microbial oncolites (Ford and Breed 1977); branching structures resembling brown algae (Nitecki 1971); ovoid pellets resembling fecal pellets, associated with potential calcispheres, stromatolites, and potential filaments (Ford and Breed 1977); and organic detritus (Vidal and Ford 1985). In the latter category are: possible animal traces (White 1927); “fusiform remains” in dark red beds transitional into the Hakatai Shale (White 1929); and potential sponges and medusae (Alf 1959), now widely regarded as pseudofossils (Cloud 1968; Glaessner 1969; Nitecki 1971; Ford and Breed 1977). The *Chuaria*-like objects reported by White (1928a) cannot be evaluated, because no locality information or specimens exist (Ford and Breed 1977), but Nitecki (1971) did report finding *Chuaria*-like microfossils after dissolving an apparently inorganic siliceous “biscuit” concretion.

Unkar Group: Hakatai Shale (upper Mesoproterozoic)

The Hakatai Shale is primarily composed of siltstone and fine-grained sandstone, although beds range from mudstone to conglomerate (Figure 4-4). It is interpreted as the result of shallow water deposition in marginal marine, tidal flat, and deltaic settings. Detrital zircons indicate at least some deposition occurred after 1187 Ma (Timmons et al. 2012).



Figure 4-4. A mud-cracked surface in the Hakatai Shale. Mud cracks are sometimes mistaken for burrows (NPS/CARL BOWMAN).

Fossils are unconfirmed from the Hakatai Shale. Aside from stromatolites found in the transition with the Bass Formation, mentioned above, most records appear to be pseudofossils. The GRCA museum has records of Hakatai “algal” structures, although in the absence of more detailed provenance information, these could be from Bass Formation beds in the transitional zone. White (1929) reported potential “algal thalli” impressions in this formation. Seilacher (1956) reported features he compared

to the trilobite trace fossil known as *Rusophycus*, but they were regarded as inorganic in Cloud (1968). Other "fucoids" were reported by White (1928a). Nitecki (1971) noticed "worm burrows" in the Hakatai Shale, which he regarded as possibly sedimentary.

Unkar Group: Shinumo Quartzite (upper Mesoproterozoic)

The Shinumo Quartzite or Sandstone is primarily composed of quartz sandstone, with some feldspar-rich sandstone in the lower part of the formation and a lag deposit at the unconformable contact with the underlying Hakatai Shale. It is interpreted as a high-energy shoreface unit. Laterally continuous contorted beds, resulting from soft-sediment deformation, are found in this unit (Timmons et al. 2012). Detrital zircons indicate it was deposited ca. 1170 Ma (Timmons et al. 2012), but it could be as young as 1140 Ma (Mulder et al. 2017).

Fossils have not been confirmed from the Shinumo Quartzite. White (1929) mentioned features that could be "fucoids" or inorganic near the base of the formation. Miller and Byers (1984) described the Shinumo Quartzite as "heavily burrowed" in a table based on a personal communication, without further discussion. As noted by Spamer (1984), burrows in the Shinumo Quartzite would be very significant, but to date this observation has not been replicated. Many of the features interpreted to be potentially biogenic in origin could be explained by the dynamic soft-sediment deformation in the unit (C. Dehler, pers. comm., May 2019).

Unkar Group: Dox Formation (upper Mesoproterozoic)

The Dox Formation is a clastic unit dominated by red sandstone, with lesser quantities of siltstone and other mudrocks. Deposition began in fluvial and deltaic settings, becoming more marine over time, with rivers becoming estuaries and nearshore marine to possibly intertidal deposition setting in. Deposition occurred between approximately 1140 and 1104 Ma (Timmons et al. 2012; Mulder et al. 2017).

The Dox Formation, like the Bass Formation, is noted for stromatolites. Stromatolites were first noted in passing by Ford and Breed (1977), and are found in the Comanche Point Member (Stevenson and Beus 1982; Hendricks and Stevenson 1990; Timmons et al. 2012), the second highest of four members recognized in the formation. Dox Formation stromatolites have not been described in detail to date. McKee (1932) pointed out Dox Formation "fucoids" that are more like mud cracks based on the photograph (Figure 4-5). Park collections attribute one of the specimens (GRCA 11442, the right object) to the ichnogenus *Teichichnus*, but it is not clear why this ichnogenus was invoked because the features do not have the distinctive "stacked" appearance of *Teichichnus*.



Figure 4-5. Left: Dox Formation “fucoids” as shown in McKee (1932); the original caption is “Peculiar casts and moulds in Dox Formation. Photo by Grant”. No scale was provided. Right: The specimens as relocated in 2019, placed in a similar setting and photographed with a scale. The two pieces are actually part and counterpart, with the left piece having concave equivalents of the convex features on the right piece (the large light gray redox spot is a tie point) (NPS/JUSTIN TWEET).

Unkar Group: Cardenas Basalt (upper Mesoproterozoic)

The volcanic Cardenas Basalt began erupting before the end of Dox Formation deposition, as shown by some interfingering beds of the two units. Eruption occurred approximately 1104 Ma (Timmons et al. 2012). Although extrusive igneous units sometimes include fossils, such as tree molds in volcanic flows and organisms buried by tuffs and other pyroclastic deposits, no fossils have been reported from the Cardenas Basalt.

Chuar Group: Nankoweap Formation (middle Neoproterozoic: late Tonian age)

The Nankoweap Formation includes a lower red unit of hematite-cemented sandstone and mudstone, and an upper white unit of siltstone and thin-bedded sandstone, separated by a local lag deposit (Timmons et al. 2012). It is interpreted as indicating shoreface and mudflat deposition (C. Dehler, pers. comm., May 2019), perhaps marine or lacustrine (Timmons et al. 2005). The Nankoweap Formation was recently found to be younger than previously suspected, approximately at most 782 Ma, making it very similar in age to the established formations of the Chuar Group (Dehler et al. 2017). Because of this, and the apparent absence of a pronounced unconformity with the overlying Galeros Formation, Dehler et al. (2017) moved the Nankoweap Formation into the Chuar Group as the lowest unit.

No uncontroversial fossils have been reported from the Nankoweap Formation. A famous likely pseudofossil, *Brooksella canyonensis*, was named from a Nankoweap Formation specimen (Bassler 1941); its lengthy history is discussed under “Notable Pseudofossils and Dubiofossils” below. As

with other Grand Canyon Supergroup units, the Nankoweap Formation has features that resemble sinuous trace fossils, but which are more likely truncated mud cracks (Ford 1990).

Chuar Group: Galeros Formation (middle Neoproterozoic: late Tonian age)

The Galeros Formation is primarily clastic, with variegated and organic-rich mudstones to sandstones represented, but a small percentage of dolomite is also present. Four members are recognized, in ascending order the Tanner, Jupiter, Carbon Canyon, and Duppa members. Deposition occurred in subtidal to supratidal environments, and possibly terrestrial environments as well (Dehler et al. 2001, 2012, 2017). Until recently, the formation had not been directly dated. The lower part of the formation is younger than the <782 Ma Nankoweap Formation (Dehler et al. 2017). An organic-rich dolomite in the lower Carbon Canyon Member has been dated using the Re-Os system to 757 ± 6.8 Ma, and a marcasite nodule in the basal Awatubi Member of the Kwagunt Formation has been dated to 751 ± 7.6 Ma (Rooney et al. 2018). Therefore, the Galeros Formation spans no more than about 25 million years, and likely less.

All four members of the Galeros Formation are fossiliferous (Porter and Riedman 2016). Most fossils are microfossils, but there are two prominent stromatolitic horizons, not only notable for their fossils but also for being carbonate beds in a primarily clastic formation. The lower horizon, the *Stratifera*/*Inzeria* horizon, is at the base of the Jupiter Member. The upper horizon, the *Baicalia* horizon, is near the top of the Carbon Canyon Member (Ford and Breed 1973a; Dehler et al. 2001, 2012, 2017). The Galeros Formation has also yielded significant microfossils, primarily acritarchs (Ford and Breed 1973b; Bloeser et al. 1977; Vidal and Ford 1985; Nagy et al. 2009; Dehler et al. 2012, 2017; Porter and Riedman 2016). Filaments and sheaths (Vidal and Ford 1985; Porter and Riedman 2016) and organic detritus (Downie 1969; Vidal and Ford 1985) have also been reported. Puncture marks, described as “vampire traces”, have been identified on some microfossils (Porter 2016a, 2016b, 2017).

Chuar Group: Kwagunt Formation (middle Neoproterozoic: late Tonian age)

Similar to the underlying Galeros Formation, the Kwagunt Formation is primarily a fine-grained organic-rich and variegated clastic unit with occasional dolomite beds, and is divided into several members. In ascending order, the members of the Kwagunt Formation are the Carbon Butte, Awatubi, and Walcott members. The Kwagunt Formation was primarily deposited in subtidal to intertidal environments, although with more episodes of subaerial exposure, especially in the upper Walcott Member (Dehler et al. 2001, 2012, 2017). The majority of the formation is younger than a ca. 751 Ma marcasite nodule in the basal Awatubi Member (Rooney et al. 2018). A tuff at the top of the formation, formerly dated to 742 ± 6 Ma (Karlstrom et al. 2000), has been re-dated via a higher precision U-Pb method to 729 ± 0.9 Ma (Rooney et al. 2018).

The Awatubi and Walcott members of the Kwagunt Formation are notably fossiliferous, but no fossils have yet been reported from the Carbon Butte Member. There is a prominent stromatolitic horizon, the *Baicalia–Boxonia* horizon, at the base of the Awatubi Member. At the base of the Walcott Member is the “flaky dolomite bed”, which is thought to be composed of microbial laminations that have been reworked and altered by soft-sediment deformation (Ford and Breed 1973a; Dehler et al. 2001, 2012, 2017). As in the Galeros Formation, most fossils are microfossils,

including acritarchs and colonial organic-walled microfossils (Bloeser et al. 1977; Vidal and Ford 1985; Horodyski 1993; Nagy et al. 2009; Dehler et al. 2012, 2017; Porter and Riedman 2016), filaments and sheaths (Schopf et al. 1973; Horodyski and Bloeser 1983; Vidal and Ford 1985; Horodyski 1993; Porter and Knoll 2000; Dehler et al. 2012, 2017; Porter and Riedman 2016), and vase-shaped microfossils (Bloeser et al. 1977; Bloeser 1985; Vidal and Ford 1985; Porter and Knoll 2000; Porter et al. 2003). Acritarchs are abundant and diverse in the Awatubi Member but disappear within the Walcott Member, while vase-shaped microfossils are primarily found in the Walcott Member (Porter and Riedman 2016; Dehler et al. 2017).

The Kwagunt Formation has also yielded more obscure fossils in GRCA. "Vampire traces" continue into this formation, and larger circular or semi-circular punctures have been found on vase-shaped microfossils (Porter 2016a, 2016b, 2017). Chemical analysis of Kwagunt Formation rocks has yielded steranes, a type of biomarker (Summons et al. 1988; Dehler et al. 2012; Brocks et al. 2016), including a sterane profile that is typical of some groups of sponges (Brocks et al. 2016). Possible traces smaller than a millimeter (0.04 in) have also been reported by Horodyski (1988, 1993), who interpreted them as trace fossils made by meiofauna.

Sixtymile Formation (lower Cambrian)

The Sixtymile Formation has recently been shown to be Cambrian in age (Karlstrom et al. 2018), but it is included here because of its traditional Precambrian placement. It is only found in a few areas of eastern GRCA, and is composed of red sandstone with chert and breccia (Elston 1979). What had previously been described as the lowest part of the formation has been transferred to the upper Kwagunt Formation (Timmons et al. 2001; Kenny 2017). The Sixtymile Formation was deposited in lacustrine, fluvial, and shallow marine settings. Detrital zircons indicate it was deposited between 520 and 509 Ma, making it contemporaneous in part with rocks of the lower Tonto Group in the western Grand Canyon–Lake Mead region (Karlstrom et al. 2018). Karlstrom et al. (2020) have included it in the Tonto Group as the basalmost formation. The supplementary information for Karlstrom et al. (2018) included a figure of an object in a chert nodule, described as "akin to a lithistid demosponge or perhaps a coralomorph", representing the first published report of a potential fossil in the Sixtymile Formation

Types of Fossils

The Precambrian fossil record of GRCA can be divided into two major categories: stromatolites and other microbial structures, which are apparent to the naked eye; and microfossils, which have only been studied since the late 1960s, with the notable exception of *Chuaria*. The Galeros and Kwagunt formations have significant microfossil assemblages. The search for fossils in the park's Precambrian rocks has also led to the description of inorganic features as fossils; some of these pseudofossils (non-fossils) or dubiofossils (unclear if fossils or not) have been the subjects of lengthy debate in the literature and are documented here for completeness.

Stromatolites and Other Microbial Structures

The most visible uncontroversial fossils in the Precambrian rocks of GRCA are stromatolites, which at this time are known from four formations (Bass, Dox, Galeros, and Kwagunt). The stromatolites

from the Bass Formation are the oldest fossils known from the park (Figures 4-3 and 4-6), between 1.25 and 1.2 billion years old.




Figure 4-6. Another view of a Bass Limestone stromatolitic horizon just below the Hakatai Shale (NSP/CARL BOWMAN).

Stromatolites and related structures such as oncolites and thrombolites are trace fossils produced by microbial colonies. In the past, they were frequently described as “algal”, but with the refinement of the definition of algae, it is more appropriate to describe the producers under the generalized term “microbes”. Stromatolites form from cycles of sediment being trapped on the surface of a microbial mat; the colony grows over the layer of sediment to have access to the outside environment, and a new layer of sediment begins to form. The colonies can have a variety of forms, related to environmental conditions and the type of microbes. Morphologies include flat layers, crinkly layers, groups of columns, domes, and cones. Oncolites form concentric layers around objects, and thrombolites form “clotted” structures instead of layers. There may also be microbial laminations not divisible into distinct colonies. Brecciated and folded microbial laminations are thought to be the source of the “flaky dolomite bed” of the Kwagunt Formation (Ford and Breed 1973a).

Some researchers have applied scientific names to stromatolites, but this is controversial; describing them in terms of morphology is more common. Taxonomic names applied to GRCA stromatolites include the following: stromatolites from the Bass Formation have been identified as *Collenia* (White 1928a; Beus et al. 1974; Ford and Breed 1977); *Baicalia*-type stromatolites have been identified from the Galeros and Kwagunt formations (Ford and Breed 1973a; Dehler et al. 2001, 2012); *Inzeria*-type and *Stratifera*-type stromatolites have been identified from the Galeros Formation (Ford and Breed

1973a; Dehler et al. 2001, 2012, 2017); *Boxonia*-type stromatolites have been reported from the Kwagunt Formation (Ford and Breed 1973a; Dehler et al. 2001, 2012); and the stromatolite taxon *Cryptozoon occidentale* was named from what was later identified as most likely the Awatubi Member of the Kwagunt Formation (Dawson 1897; Ford and Breed 1973a). As mentioned above, stromatolites form distinct horizons in the Galeros and Kwagunt formations. White (1928a) and Lathrop (2018) identified diverse morphologies in the Bass Formation stromatolites, but in general it has been more typical to report the presence of stromatolites in passing.

Acritarchs and Other Organic-Walled Microfossils

Many of the Precambrian microfossils of GRCA are organic-walled structures. They include acritarchs, colonial forms, and filaments and sheaths. "Acritarch" is a term for a non-colonial microfossil of uncertain origin with a central cavity and organic walls rather than mineralized walls. The term should be considered a category rather than a taxonomic group, because many different organisms produce structures that could be described as acritarchs (Evitt 1963). GRCA's Precambrian acritarchs include forms with smooth or ornamented walls, generally spherical to elliptical, mostly ranging in size from a few tens of μm to a few hundred μm in diameter (Porter and Riedman 2016). Aside from the *Chuaria*-like Bass Formation microfossils described by White (1928a), which have not been subsequently relocated, and Nitecki's microfossils from the same formation (Nitecki 1971), Precambrian acritarchs at GRCA are limited to the Galeros and Kwagunt formations. They are most diverse in the Galeros Formation (Nagy et al. 2009). A number of taxa have been recognized (Vidal and Ford 1985; Nagy et al. 2009; Porter and Riedman 2016; Dehler et al. 2017), five of which were named from GRCA fossils. The taxonomy differs in each report, so following specific finds from report to report can be confusing. Leiosphaerids are another distinct type of acritarch. They have simple smooth walls and are found in abundance higher in section than most of the acritarchs, in the Awatubi Member of the Kwagunt Formation. There are also colonial forms, such as aggregates of small round cells known as *Bavlinella* (*Sphaerocongregus* of older reports), also abundant in the Awatubi and overlying Walcott members (Nagy et al. 2009; as Nagy et al. noted, not all occurrences of *Bavlinella* are necessarily fossils).

Chuaria circularis is the most historically notable Precambrian microfossil from GRCA, and like many Precambrian fossils described in the 19th century or early 20th century has had a long history of controversy. It is unusual for its great size, being visible to the naked eye: *C. circularis* fossils are generally 2 to 3 mm (0.08 to 0.1 in) across, up to 5 mm (0.2 in) (Ford 1990). It has also had an outsized impact: the work that led to the division of the Chuar Group and helped initiate the study of microfossils in these rocks was begun at the request of the National Museum of Natural History to relocate Walcott's type locality for *Chuaria circularis* (Ford and Breed 1973a). Aside from size, *C. circularis* can be described as "flattened carbonaceous spheroids, now discs," lacking pores or other openings, or regular surficial ornamentation, but with wrinkles or cracks (Figure 4-1). In the field, they can be seen as black shiny discs on bedding laminae, although when prepared they look yellowish and translucent (Ford and Breed 1973b).

Although the report in White (1876) of "*Lingulella* and *Obolella*" from Kwagunt Valley sounds superficially like *Chuaria*, it is unlikely that these objects were *Chuaria* (Spamer 1988). The first

definite report of *Chuaria* is the "small Discinoid shell" of Walcott (1883). Walcott formally described it some years later as *Chuaria circularis* (Walcott 1899). Over the next few decades *Chuaria* was the subject of numerous attempts at classification and was reported from many other places across the world (see summaries in Spamer 1984 and 1988), although many of these reports are questionable (Vidal and Ford 1985). Authors suggested everything from brachiopods, to hyolith operculae, to gastropods, to ostracodes, to foraminifera, to algae, to trilobite eggs, to pseudofossils (Ford and Breed 1973b; Spamer 1984). By the 1960s, opinion had crystallized around an algal identification (Ford and Breed 1973b). Ford and Breed (1973a, 1973b, 1977) redescribed *Chuaria circularis* from GRCA material. Because of the diversity of fossils from numerous locations that have been described as *Chuaria*, caution should be exerted when applying interpretations to GRCA *Chuaria* that are based on fossils from other locations: they may not represent the same organisms. At some locations outside of GRCA, such as in India, fossils described as *Chuaria* are linked to fossils known as *Tawuia*, perhaps as two parts of the same structure (e.g., Kumar 2001; Sharma et al. 2009). Although *Tawuia* is apparently only found with *Chuaria* (Xiao and Dong 2006), the opposite is not the case, as *Tawuia* has never been reported from GRCA. Therefore, *Chuaria circularis* of GRCA is not necessarily the same kind of fossil as the "*Chuaria-Tawuia* complex".

Filaments and Sheaths

Potential filamentous structures have been reported from the Bass Formation (Dalton 1972; Beus et al. 1974; Ford and Breed 1977), but uncontroversial filaments are only known from the Galeros and Kwagunt formations. Filaments and sheaths were first reported by Schopf et al. (1973) from a cherty pisolite bed in the Walcott Member of the Kwagunt Formation. Several taxa have since been reported from shale beds (Vidal and Ford 1985; Horodyski 1993; Porter and Riedman 2016).

GRCA Precambrian filaments can be divided into several groups. The most abundant forms are narrow, non-branching, non-tapering, and non-tubular, between one and several μm across (Horodyski 1993). This group was divided by Porter and Riedman (2016) into several species of *Siphonophycus* based on size. They are found both isolated and as parts of fragments of filamentous mats. Similar but larger tubular filaments up to several tens of μm across have been found, but are rarer. There are also rare non-tubular filaments from 10 to >100 μm across. One carbonaceous form is apparently limited to a 0.1-mm-thick (0.004 in) interval near the top of the Awatubi Member of the Kwagunt Formation, in which it is very abundant. They are most likely from eukaryotic algae, but could be prokaryotic (Horodyski 1993).

Vase-Shaped Microfossils

Vase-shaped microfossils (VSMs) are non-colonial fossils shaped like a simple vase with a wider "base" and narrower "upper" end featuring an opening. They were first reported from GRCA and are now known from a number of formations across the world, but flourished only between approximately 760 Ma and the onset of the first Snowball Earth glaciation at 717 Ma (MacDonald et al. 2010). They are frequently preserved as siliceous or calcareous casts, sometimes coated by organic material or oxides, probably replacing an original organic material. Some feature a "honeycomb" appearance, derived from an original scaled surface. Most are straight, but some have curved necks. In some forms there is a distinct "neck", while in others the shape tapers without an

obvious break. Some have a “collar” or other distinct apertural feature, while others do not. They range in size from approximately 20 to 300 μm long and 10 to 200 μm wide (Porter and Knoll 2000). A few features previously reported in the literature appear to have been mistaken. Bloeser (1985) interpreted some features as opercula, but these appear to be sediment plugs instead (Porter and Knoll 2000). The internal vesicles of Horodyski (1993) appear to be crystalline precipitates (Porter and Knoll 2000). As of this writing, a dozen species of VSMs have been named from GRCA fossils, all from the Kwagunt Formation (Porter et al. 2003). They can be exceedingly abundant in certain beds, up to 4000/mm^3 in some nodules (Porter and Knoll 2000). They are interpreted as the fossils of testate amoebas (“testate” meaning they form a structure, a test, around the body) (Porter and Knoll 2000), apparently ancient examples of arcellinid amoebas (Lahr et al. 2019).

VSMs are first found in the uppermost Awatubi Member, where they overlap with *Chuaria*, leiosphaerids, and *Sphaerocongregus* (=*Bavlinella*). These other groups disappear higher in the Awatubi Member or in the lower half of the Walcott Member, leaving the VSMs as the dominant group. This pattern has been interpreted as the result of prolonged eutrophication, in which nutrient-rich water promotes “algal” blooms, and the increased productivity depletes oxygen from the water, disrupting the existing phytoplankton ecosystem while providing an opportunity for protozoans to thrive (Nagy et al. 2009).

VSMs were first reported in Bloeser et al. (1977) as “tear- and flask-shaped microfossils”, attributed to chitinozoans (a type of flask-shaped organic-walled microfossil of uncertain origin otherwise known from Paleozoic rocks). At that time two forms were distinguished: flask-shaped with a short “neck” with pronounced “collar”; and slightly longer, thinner tear-shaped forms lacking a distinct neck. The initial genus, *Melanocyrillium*, was mentioned several times before formal publication (e.g., Spamer 1984; Vidal and Ford 1985). Bloeser (1985) formally described *Melanocyrillium* and separated the fossils into three species: *M. fimbriatum* for the tear-shaped form, and *M. hexodiadema* and *M. horodyskii* for flask-shaped forms with two different “necks” and “collars”. She also removed them from the chitinozoans, suggesting that they were cysts of an unknown alga. Porter and Knoll (2000) first proposed the testate amoeba hypothesis, which is now widely accepted. Porter et al. (2003) provided a description of the GRCA VSM assemblage, naming a number of new taxa and transferring *Melanocyrillium fimbriata* and *M. horodyskii* to a new genus, *Trigonocyrillium*. Additional taxa were reported from GRCA in Morais et al. (2019).

Other Fossils

Some researchers have explored hydrocarbons in the Precambrian rocks. This line of research was initiated by Summons et al. (1988), who identified biomarkers including steranes and neohopanes that indicate eukaryotes. Brocks et al. (2016), continuing this work, identified a new sterane they called cryostane, which they linked to a group of sponges known as demosponges (with the caveat that other eukaryotes could have produced it). They interpreted this as a type of sterol evolved to protect cells against toxins released by protist predators.

Holes have been observed in some of the microfossils, such as semi-circular and circular holes in VSMs. They are thought to represent predation and/or scavenging (Porter et al. 2003; Porter 2016a). One unusual class of these perforations consists of minute circular holes, 0.1 to 3.4 μm across, found

in the walls of at least seven species of GRCA organic-walled microfossils. They are believed to represent "vampire-like" predation behavior as exhibited today by several groups of eukaryotes and are, to date, the oldest direct evidence for predation on eukaryotes (Porter 2016a).

In addition, some other types of fossils have been reported, but generally in passing or otherwise in such little detail that it is not possible to evaluate the report or to be sure that they do not belong to one of the other categories. They include various unspecified cellular fossils in the Chuar Group (Schopf et al. 1973; Bloeser et al. 1977; Vidal and Ford 1985; Ford 1990; Dehler et al. 2012), Bass Formation calcispheres (Dalton 1972; Ford and Breed 1977; Lathrop 2018), and the possible Kwagunt meiofaunal traces mentioned by Horodyski (1988, 1993). Undefined organic detritus has also been reported from the Bass Formation (Vidal and Ford 1985) and Chuar Group (Downie 1969).

Notable Pseudofossils and Dubiofossils

Reports of pseudofossils (resembling a fossil but inorganic in origin) and dubiofossils (potentially a fossil, but origin unclear) in the Precambrian rocks of GRCA go back to the "*Lingulella* and *Obolella*" of White (1876). Walcott (1883) reported finding a "small Discinoid shell, a couple of specimens of a Pteropod allied to *Hyolithes triangularis* and an obscure Stromatopora-like group of forms" in rocks of the Chuar Group. A few years later, he added a possible trilobite fragment and a *Lingula*-like shell to the lot (Walcott 1886). By 1899, he'd reevaluated the "hyolith" as potentially of mechanical origin and decided that the trilobite fragment could have been misinterpreted, but had also added an *Acrothele*-like brachiopod (Walcott 1899). After observing some of Walcott's specimens, Horodyski (1993) interpreted the *Acrothele*-like brachiopod (USNM 33801) as an "apparently mineralic precipitate along a fracture or bedding surface", and the trilobite fragment (USNM 33802) as "probably a fortuitously exposed portion of a carbonaceous layer". Ford and Breed (1977) suspected that the lingulid shell could also be dismissed in a similar way. The only fossils from Walcott's assemblage are the "discinoid shell", which he named *Chuaria circularis* in 1899, and the "Stromatopora", actually stromatolites. "Stromatopora", now used for certain layered sponges, is a term that in the 19th century was not firmly distinguished from the various terms then used for what are now recognized as stromatolites.

The most significant of the various Precambrian pseudofossils and dubiofossils is *Brooksella canyonensis*. It first came to attention in the 1930s, a putative jellyfish uncovered in the course of work on what became named the Nankoweap Formation (Van Gundy 1937). Bassler (1941) formally created the name *Brooksella canyonensis* for Van Gundy's specimen (USNM 99438), although he was hesitant to declare it a jellyfish, arguing that it could also be algal or inorganic. Although the jellyfish hypothesis has long been out of favor, there have been decades of arguments and counter-arguments, split between organic (e.g., Glaessner 1969; Kauffman and Steidtmann 1981; Kauffman and Fursich 1983) and inorganic (e.g., Cloud 1968; Ford and Breed 1977; Ford 1990) camps. The organic camp favors a trace-fossil origin, such as a worm-like burrowing sediment feeder (Glaessner 1969) or an advanced metazoan burrow produced by complex feeding behavior (Kauffman and Fursich 1983). The inorganic camp has suggested such origins as gas-escape structures or compactions (Cloud 1968), similar to "sand volcanoes" formed by fluid escape (Ford and Breed 1977; Ford 1990), or an aggregation of mud rolls (Fedonkin and Runnegar 1992).

Another category of Precambrian pseudofossils is the “fucoids”. White (1928a, 1929) and McKee (1932) reported “fucoids” in several of the Grand Canyon Supergroup units. “Fucoid”, from Fucaceae, an order of brown algae, is an obsolete term applied to what were once thought to be fossil seaweeds, but which are now known to be invertebrate burrows or various burrow-like sedimentary features such as mud cracks (Figure 4-4). McKee (1932) recognized that most “fucoids” were not seaweed fossils and used the term in a general fashion to encompass anything with that general shape, including a photographed slab of Dox Formation “fucoids” (Figure 4-5). Mud cracks are thought to be responsible for putative “trace fossils” in the Nankoweap Formation (Ford 1990).

Several other pseudofossils and dubiofossils are worth noting. Seilacher (1956) reported the presence of bilobed features 1 mm (0.04 in) across in the Hakatai Shale, which he compared to the trilobite resting trace *Rusophycus*. Cloud (1968) considered them inorganic. Alf (1959) reported several potential fossils from near the Bass–Hakatai transition interval, including vermiform markings, more putative jellyfish (Figure 4-7), and a possible sponge. Cloud (1968) regarded one of the “jellyfish” varieties as more likely marks left by water drops and the other variety as gas-escape structures or compactions, and the “sponge” as a silica nodule. Glaessner (1969) disagreed with the water drop interpretation of some of Alf’s “jellyfish”, which he thought might be algal colonies instead, but agreed with the silica nodule identification of the “sponge”, and regarded the vermiform markings as filled mud cracks. Ford and Breed (1973a, 1977) disagreed with Glaessner’s algal interpretation, and returned to a gas escape structure explanation. Nitecki (1971) reported similar nodules to the “sponge” and considered them inorganic, but was able to produce apparent microfossils by dissolving one of them in hydrofluoric acid.




Figure 4-7. Examples of Alf’s circular structures from the Bass–Hakatai transition interval (RAYMOND M. ALF MUSEUM/DON LOFGREN).

Acknowledgments

I wish to thank Diana Boudreau, Anne Miller, Vince Santucci, and Klara Widrig for their preliminary reviews and comments; Don Weeks for his assistance in organizing peer review; Carol Dehler (Utah State University) and Susannah Porter (University of California Santa Barbara) for their detailed reviews; Colleen Hyde (GRCA Museum) for her assistance with McKee's specimens; Spencer Lucas (New Mexico Museum of Natural History and Science) for his observations regarding McKee's Dox "fucoids"; and Andrew Farke and Don Lofgren of the Raymond M. Alf Museum for their assistance concerning Raymond Alf's Precambrian collections from GRCA.

Literature Cited

Alf, R. M. 1959. Possible fossils from the early Proterozoic Bass Formation, Grand Canyon, Arizona. Plateau 31(3):60–63.

Bassler, R. S. 1941. A supposed jellyfish from the pre-Cambrian of the Grand Canyon. Proceedings of the United States National Museum 89(3104):519–522.

Beus, S. S., R. R. Rawson, R. O. Dalton, G. M. Stevenson, V. S. Reed, and T. M. Daneker. 1974. Preliminary report on the Unkar Group (Precambrian) in Grand Canyon, Arizona. Pages 34–53 *in* T. N. V. Karlstrom, G. A. Swann, and R. L. Eastwood, editors. Geology of northern Arizona; with notes on archaeology and paleoclimate, part 1, regional studies. Geological Society of America, Rocky Mountain Section Meeting, Flagstaff, Arizona.

Bloeser, B. 1985. *Melanocyrillium*, a new genus of structurally complex late Proterozoic microfossils from the Kwagunt Formation (Chuar Group), Grand Canyon, Arizona. Journal of Paleontology 59(3):741–765.

Bloeser, B., J. W. Schopf, R. J. Horodyski, and W. J. Breed. 1977. Chitinozoans from the late Precambrian Chuar Group of the Grand Canyon, Arizona. Science 195(4279):676–679.

Brocks, J. J., A. J. M. Jarrett, E. Sirantoine, F. Kenig, M. Moczydlowska, S. Porter, and J. Hope. 2016. Early sponges and toxic protists: Possible sources of cryostane, an age diagnostic biomarker antedating Sturtian snowball Earth. Geobiology 14(2):129–149.

Cloud, P. E., Jr. 1960. Gas as a sedimentary and diagenetic agent. American Journal of Science 258-A:35–45.

Cloud, P. E., Jr. 1968. Pre-metazoan evolution and the origins of the Metazoa. Pages 1–72 *in* E. T. Drake, editor. Evolution and environment. Yale University Press, New Haven, Connecticut.

Dalton, R. O. 1972. Stratigraphy of the Bass Formation. Thesis. Northern Arizona University, Flagstaff, Arizona.

Dawson, J. W. 1897. Note on *Cryptozoon* and other ancient fossils. Canadian Record of Science 7:203–219.

Dehler, C. M., M. E. Elrick, K. E. Karlstrom, G. A. Smith, L. J. Crossey, and M. J. Timmons. 2001. Neoproterozoic Chuar Group (~800–742 Ma), Grand Canyon: A record of cyclic marine deposition during global climatic and tectonic transitions. Sedimentary Geology 141–142:465–499.

Dehler, C. M., S. M. Porter, and J. M. Timmons. 2012. The Neoproterozoic Earth system revealed from the Chuar Group of Grand Canyon. Pages 49–72 *in* J. M. Timmons and K. E. Karlstrom, editors. Grand Canyon geology; two billion years of Earth's history. Geological Society of America, Boulder, Colorado. Special Paper 489.

Dehler, C., G. Gehrels, S. Porter, M. Heizler, K. Karlstrom, G. Cox, L. Crossey, and M. Timmons. 2017. Synthesis of the 780-740 Ma Chuar, Uinta Mountain, and Pahrump (ChUMP) groups, western USA; implications for Laurentia-wide cratonic marine basins. Geological Society of America Bulletin 129(5–6):607–624.

Downie, C. 1969. Palynology of the *Chuaria* shales of the Grand Canyon. Pages 121–122 *in* D. L. Bass, editor. Geology and natural history of the Grand Canyon region. Four Corners Geological Society, Flagstaff, Arizona. Field Conference 5.

Elston, D. P. 1979. Late Precambrian Sixtymile Formation and orogeny at top of the Grand Canyon Supergroup, northern Arizona. U.S. Geological Survey, Washington, D.C. Professional Paper 1092.

Evitt, W. R. 1963. A discussion and proposals concerning fossil dinoflagellates, hystrichospheres, and acritarchs, II. Proceedings of the National Academy of Sciences of the United States of America 49(3):298–302.

Fedonkin, M. A., and B. N. Runnegar. 1992. Proterozoic metazoan trace fossils. Pages 389–395 *in* J. W. Schopf and C. Klein, editors. The Proterozoic biosphere: A multidisciplinary study. Cambridge University Press, Cambridge, United Kingdom.

Ford, T. D. 1990. Grand Canyon Supergroup: Nankoweap Formation, Chuar Group, and Sixtymile Formation. Pages 49–70 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology. Oxford University Press, New York, New York.

Ford, T. D., and W. J. Breed. 1973a. Late Precambrian Chuar Group, Grand Canyon, Arizona. Geological Society of America Bulletin 84(4):1243–1260.

Ford, T. D., and W. J. Breed. 1973b. The problematic Precambrian fossil *Chuaria*. Palaeontology 16:535–550.

Ford, T. D., and W. J. Breed. 1977. *Chuaria circularis* Walcott and other Precambrian fossils from the Grand Canyon. Journal of the Palaeontological Society of India 20:170–177.

Ford, T. D., W. J. Breed, and C. Downie. 1969. Preliminary geologic report of the Chuar Group, Grand Canyon, Arizona. Pages 114–122 *in* D. L. Bass, editor. Geology and natural history of the Grand Canyon region. Four Corners Geological Society, Flagstaff, Arizona. Field Conference 5.

Glaessner, M. F. 1969. Trace fossils from the Precambrian and basal Cambrian. Lethaia 2(4):369–393.

Hendricks, J. D., and G. M. Stevenson. 1990. Grand Canyon Supergroup: Unkar Group. Pages 29–47 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology. Oxford University Press, New York, New York.

Holland, M. E., K. E. Karlstrom, M. F. Doe, G. E. Gehrels, M. Pecha, O. P. Shufeldt, G. Begg, W. L. Griffin, and E. Belousova. 2015. An imbricate mid-crustal suture zone; the Mojave-Yavapai Province boundary in Grand Canyon, Arizona. Geological Society of America Bulletin 127(9–10):1391–1410.

Horodyski, R. J. 1988. Late Proterozoic fossils from the western U.S.: Could they represent the oldest skeletal metaphytes and the oldest meiofaunal traces? Abstracts with Programs - Geological Society of America 20(7):A256.

Horodyski, R. J. 1993. Paleontology of Proterozoic shales and mudstones: Examples from the Belt Supergroup, Chuar Group and Pahrump Group, western USA. Precambrian Research 61(3–4):241–278.

Horodyski, R. J., and B. Bloeser. 1983. Possible eukaryotic algal filaments from the late Proterozoic Chuar Group, Grand Canyon, Arizona. Journal of Paleontology 57(2):321–326.

Karlstrom, K. E., C. M. Dehler, Z. D. Sharp, J. W. Geissman, M. E. Elrick, J. M. Timmons, L. J. Crossey, S. A. Bowring, K. Keefe, A. H. Knoll, S. M. Porter, D. J. Des Marais, and A. B. Weil. 2000. The Chuar Group of the Grand Canyon: Record of breakup of Rodinia, associated change in the global carbon cycle, and ecosystem expansion by 740 Ma. Geology 28:619–622.

Karlstrom, K., J. Hagadorn, G. Gehrels, W. Matthews, M. Schmitz, L. Madronich, J. Mulder, M. Pecha, D. Giesler, and L. Crossey. 2018. Cambrian Sauk transgression in the Grand Canyon region redefined by detrital zircons. Nature Geoscience 11:438–443.

Karlstrom, K. E., M. T. Mohr, M. D. Schmitz, F. A. Sundberg, S. M. Rowland, R. Blakey, J. R. Foster, L. J. Crossey, C. M. Dehler, and J. W. Hagadorn. 2020. Redefining the Tonto Group of Grand Canyon and recalibrating the Cambrian timescale. Geology in press. doi: https://doi.org/10.1130/G46755.1.

Kauffman, E. G., and F. Fursich. 1983. *Brooksella canyonensis*: A billion year old complex metazoan trace fossil from the Grand Canyon. Abstracts with Programs - Geological Society of America 15(6):608.

Kauffman, E. G., and J. R. Steidtmann. 1981. Are these the oldest metazoan trace fossils? Journal of Paleontology 55:923–947.

Kenny, R. 2017. Oxygen and hydrogen isotope ratios of chert from the Sixtymile Formation in Grand Canyon National Park, USA: A warm palaeoclimate, freshwater deposit. Acta Geochimica 36(2):250–259.

Kumar, S. 2001. Mesoproterozoic megafossil *Chuaria–Tawuia* association may represent parts of a multicellular plant, Vindhyan Supergroup, central India. Precambrian Research 106(3–4):187–211.

Lahr, D. J. G., A. Kosakyan, E. Lara, E. A. D. Mitchell, L. Morais, A. L. Porfirio-Sousa, G. M. Ribeiro, A. K. Tice, T. Pánek, S. Kang, M. W. Brown. 2019. Phylogenomics and morphological reconstruction of Arcellinida testate amoebae highlight diversity of microbial eukaryotes in the Neoproterozoic. Current Biology 29(6):991–1001.

Lathrop, E. C. 2018. Understanding the Late Mesoproterozoic Earth system from the oldest strata in Grand Canyon: C-isotope stratigraphy and facies analysis of the 1254 Ma Bass Formation, Grand Canyon Supergroup, AZ., USA. Dissertation. Utah State University, Logan, Utah.

Macdonald, F. A., M. D. Schmitz, J. L. Crowley, C. F. Roots, D. S. Jones, A. C. Maloof, J. V. Strauss, P. A. Cohen, D. T. Johnston, and D. P. Schrag. 2010. Calibrating the Cryogenian. Science 327(5970):1241–1243.

McKee, E. D. 1932. Some Fucoides from Grand Canyon. Grand Canyon Nature Notes 7(8):77–81.

Miller, M. F., and C. W. Byers. 1984. Abundant and diverse early Paleozoic infauna indicated by the stratigraphic record. Geology 12(1):40–43.

Morais, L., D. J. G. Lahr, I. D. Rudnitzki, B. T. Freitas, G. R. Romero, S. M. Porter, A. H. Knoll, and T. R. Fairchild. 2019. Insights into vase-shaped microfossil diversity and Neoproterozoic biostratigraphy in light of recent Brazilian discoveries. Journal of Paleontology 93(4):612–627.

Mulder, J. A., K. E. Karlstrom, K. Fletcher, M. T. Heizler, J. M. Timmons, L. J. Crossey, G. E. Gehrels, and M. Pecha. 2017. The syn-orogenic sedimentary record of the Grenville Orogeny in southwest Laurentia. Precambrian Research 294:33–52.

Nagy, R. M., S. M. Porter, C. M. Dehler, and Y. Shan. 2009. Biotic turnover driven by eutrophication before the Sturtian low-latitude glaciation. Nature Geoscience 2:415–418.

Nitecki, M. H. 1971. Pseudo-organic structures from the Precambrian Bass Limestone in Arizona. Fieldiana: Geology (New Series) 23(1):1–9.

Porter, S. M. 2016a. Tiny vampires in ancient seas: Evidence for predation via perforation in fossils from the 780-740 million-year-old Chuar Group, Grand Canyon, USA. Proceedings of the Royal Society, Biological Sciences 283(1831):20160221.

Porter, S. M. 2016b. Data from: Tiny vampires in ancient seas: Evidence for predation via perforation in fossils from the 780–740 million-year-old Chuar Group, Grand Canyon, USA. Dryad Digital Repository. Available at: https://doi.org/10.5061/dryad.6pr6c (accessed May 2019).

Porter, S. M. 2017. A view of microbial ecosystems and oxygen from the mid-Neoproterozoic (780-730 Ma) Chuar Group, Grand Canyon supergroup, Arizona. Abstracts with Programs - Geological Society of America 49(6):abstract 11-7.

Porter, S. M., and A. H. Knoll. 2000. Testate amoebae in the Neoproterozoic Era: Evidence from vase-shaped microfossils in the Chuar Group, Grand Canyon. Paleobiology 26(3):360–385.

Porter, S. M., and L. A. Riedman. 2016. Systematics of organic-walled microfossils from the ca. 780–740 Ma Chuar Group, Grand Canyon, Arizona. Journal of Paleontology 90(5):815–853.

Porter, S. M., R. Meisterfeld, and A. H. Knoll. 2003. Vase-shaped microfossils from the Neoproterozoic Chuar Group, Grand Canyon: A classification guided by modern testate amoebae. Journal of Paleontology 77(3):409–429.

Powell, J. W. 1876. Report on the geology of the eastern portion of the Uinta Mountains and a region of country adjacent thereto. U.S. Geological and Geographical Survey of the Territories, Government Printing Office, Washington, D.C.

Rooney, A. D., J. Austermann, E. F. Smith, Y. Li, D. Selby, C. M. Dehler, M. D. Schmitz, K. E. Karlstrom, and F. A. Macdonald. 2018. Coupled Re-Os and U-Pb geochronology of the Tonian Chuar Group, Grand Canyon. Geological Society of America Bulletin 130(7–8):1085–1098.

Schopf, J. W., T. D. Ford, and W. J. Breed. 1973. Microorganisms from the Late Precambrian of the Grand Canyon, Arizona. Science 179(4080):1319–1321.

Seilacher, A. 1956. Der Beginn des Kambriums als biologische Wende. Neues Jahrbuch für Geologie und Palaeontologie, Abhandlungen 103:155–180.

Sharma, M., S. Mishra, S. Dutta, S. Banerjee, and Y. Shukla. 2009. On the affinity of *Chuaria-Tawuia* complex: A multidisciplinary study. Precambrian Research 173(1–4):123–136.

Smith, A. R. 1969. Precambrian fossils from the Bass Limestone of the Grand Canyon Series. Special Paper - Geological Society of America 121:639.

Spamer, E. E. 1984. Paleontology in the Grand Canyon of Arizona: 125 years of lessons and enigmas from the late Precambrian to the present. The Mosasaur 2:45–128.

Spamer, E. E. 1988. Geology of the Grand Canyon: An annotated bibliography, with an annotated catalogue of Grand Canyon type fossils. Volume 3. Second supplement (to 1987) with an annotated bibliography of the world literature on the Grand Canyon type fossil *Chuaria*

circularis Walcott, 1899, an index fossil for the late Proterozoic. Geological Society of America, Boulder, Colorado. Microform Publication 17.

Stevenson, G. M., and S. S. Beus. 1982. Stratigraphy and depositional setting of the upper Precambrian Dox Formation in Grand Canyon. Geological Society of America Bulletin 93(2):163–173.

Summons, R. E., S. C. Brassell, G. Eglinton, E. Evans, R. J. Horodyski, N. Robinson, and D. M. Ward. 1988. Distinctive hydrocarbon biomarkers from fossiliferous sediment of the late Proterozoic Walcott Member, Chuar Group, Grand Canyon, Arizona. Geochimica et Cosmochimica Acta 52(11):2625–2637.

Timmons, J. M., K. E. Karlstrom, C. M. Dehler, J. W. Geissman, and M. T. Heizler. 2001. Proterozoic multistage (ca. 1.1 and 0.8 Ga) extension recorded in the Grand Canyon Supergroup and establishment of northwest- and north-trending tectonic grains in the southwestern United States. Geological Society of America Bulletin 113(1):163–181.

Timmons, J. M., K. E. Karlstrom, M. T. Heizler, S. A. Bowring, G. E. Gehrels, and L. J. Crossey. 2005. Tectonic inferences from the ca. 1255-1100 Ma Unkar Group and Nankoweap Formation, Grand Canyon: Intracratonic deformation and basin formation during protracted Grenville orogenesis. Geological Society of America Bulletin 117(11–12):1573–1595.

Timmons, J. M., J. Bloch, K. Fletcher, K. E. Karlstrom, M. Heizler, and L. J. Crossey. 2012. The Grand Canyon Unkar Group: Mesoproterozoic basin formation in the continental interior during supercontinent assembly. Pages 25–47 *in* J. M. Timmons and K. E. Karlstrom, editors. Grand Canyon geology; two billion years of Earth's history. Geological Society of America, Boulder, Colorado. Special Paper 489.

Van Gundy, C. E. 1937. Jellyfish from Grand Canyon Algonkian. Science 85(2204):314.

Van Gundy, C. E. 1951. Nankoweap group of the Grand Canyon Algonkian of Arizona. Geological Society of America Bulletin 62(8):953–959.

Vidal, G., and T. D. Ford. 1985. Microbiotas from the late Proterozoic Chuar Group (northern Arizona) and Uinta Mountain Group (Utah) and their chronostratigraphic implications. Precambrian Research 28(3–4):349–389.

Walcott, C. D. 1883. Pre-Carboniferous strata in the Grand Cañon of the Colorado, Arizona. American Journal of Science (3rd series) 26:437–442.

Walcott, C. D. 1886. Second contribution to the studies on the Cambrian faunas of North America. U.S. Geological Survey, Washington, D.C. Bulletin 30.

Walcott, C. D. 1899. Pre-Cambrian fossiliferous formations. Geological Society of America Bulletin 10:199–244.

White, C. A. 1876. Invertebrate paleontology of the Plateau province, together with notice of a few species from localities beyond its limits in Colorado. Pages 74–135 *in* J. W. Powell. Report on the geology of the eastern portion of the Uinta Mountains and a region of country adjacent thereto. U.S. Geological and Geographical Survey of the Territories, Government Printing Office, Washington, D.C.

White, C. D. 1927. Study of the fossil floras in the Grand Canyon, Arizona. Carnegie Institution of Washington Year Book 26:366–369.

White, C. D. 1928a. Algal deposits of Unkar Proterozoic age in the Grand Canyon, Arizona. Proceedings of the National Academy of Sciences of the United States of America 14(7):597–600.

White, C. D. 1928b. Study of the fossil floras in the Grand Canyon, Arizona. Carnegie Institution of Washington Year Book 27:389–390.

White, C. D. 1929. Study of the fossil floras in the Grand Canyon, Arizona. Carnegie Institution of Washington Year Book 28:392–393.

Xiao, S., and L. Dong. 2006. Morphological and ecological history of Proterozoic macroalgae. Pages 57–90 *in* S. Xiao and A. J. Kaufman, editors. Neoproterozoic geobiology and paleobiology. Springer, Dordrecht, the Netherlands.

Appendix 4-A. Precambrian Taxa From GRCA

It was not possible to critically evaluate every cited occurrence. There are undoubtedly instances below where the same taxon is listed more than once due to differing interpretations or general changes in taxonomic usage. This is not limited to choices of genera and species; one author may describe a find as [*Genus*] [*species*], where another may prefer [*Genus*] sp. In addition, it was not within the scope of the project to evaluate every potential synonymy and genus-species combination. The most recent name is generally used, with other combinations included in notes following the tables to allow translation between different references. Some cited taxa have outdated names, but the author did not provide enough information to determine more appropriate names. In these cases, the taxonomy is left as the author gave it. Records are marked stratigraphically with "Y" in the corresponding formation column; questionable records are marked with "?".

GCM = Grand Canyon museum collections (only unique results unreported in the literature are included)

Due to the number of taxa and formations, entries have been split into two tables, incorporating the following formations in ascending order:

Appendix Table 4-A-1: Unkar Group

- Yud = Dox Formation
- Yus = Shinumo Quartzite

- Yuh = Hakatai Shale
- Yub = Bass Formation

Appendix Table 4-A-2: Chuar Group

- Zck = Kwagunt Formation
- Zcg = Galeros Formation
- Zcn = Nankoweap Formation

Appendix Table 4-A-1. Unkar Group fossil taxa. Yud = Dox Formation, Yus = Shinumo Quartzite, Yuh = Hakatai Shale, and Yub = Bass Formation.

Category	Taxon	Yub	Yuh	Yus	Yud	References
Structures of microbial origin	Various “algal” structures	Y	Y	–	–	White 1928a, GCM
	Possible microbial laminations	–	–	–	Y	Stevenson and Beus 1982
	Stromatolites: *Collenia frequens*	Y	–	–	–	Beus et al. 1974, Ford and Breed 1977
	Stromatolites: *Collenia symmetrica*	Y	–	–	–	Beus et al. 1974, Ford and Breed 1977
	Stromatolites: *Collenia undosa*	Y	–	–	–	Beus et al. 1974, Ford and Breed 1977
	Stromatolites: *Collenia*-type	Y	–	–	–	White 1928a
	Stromatolites: general	Y	?*	–	Y	Nitecki 1971, Ford and Breed 1977, Stevenson and Beus 1982, Beus 1987, Hendricks and Stevenson 1990, Timmons et al. 2012, Lathrop 2018
Acritarchs	*Chuaria*-like microfossils	Y	–	–	–	White 1928a, Nitecki 1971
Filaments and sheaths	Possible “algal” filaments	Y	–	–	–	Dalton 1972, Beus et al. 1974, Ford and Breed 1977
	Enigmatic branching structures	Y	–	–	–	Nitecki 1971
Miscellaneous fossils	Calcispheres (Calcitarcha)	Y	–	–	–	Dalton 1972, Ford and Breed 1977, Lathrop 2018
	Organic detritus	Y	–	–	–	Vidal and Ford 1985
	Ovoid pellets	Y	–	–	–	Ford and Breed 1977
Select pseudofossils and dubiofossils	Apparent algal thalli impressions	–	Y	–	–	White 1929
	Apparent bivalved fossils (?oncolites)	Y	–	–	–	Smith 1969, Ford and Breed 1977
	“Jellyfish”	Y	–	–	–	Seilacher 1956, Alf 1959, Cloud 1968, Glaessner 1969, Ford and Breed 1977
	“*Rusophycus*”	–	Y	–	–	Seilacher 1956, Cloud 1968, Glaessner 1969, Webby 1970
	“Sponge”	Y	–	–	–	Alf 1959, Cloud 1968, Ford and Breed 1977
	“*Teichichnus*”	–	–	–	Y	GCM
	Various “fucoids” and other putative invertebrate trace fossils	Y	Y	Y	Y	White 1927, 1928a, 1929; McKee 1932; Alf 1959; Cloud 1968; Glaessner 1969; Nitecki 1971; Ford and Breed 1977

* Stromatolites are noted in the Bass–Hakatai transition zone, but are limited to beds of the Bass facies (Timmons et al. 2012).

Appendix Table 4-A-2. Chuar Group fossil taxa. Zck = Kwagunt Formation, Zcg = Galeros Formation, and Zcn = Nankoweap Formation.

Category	Taxon	Zcn	Zcg	Zck	Zcu	References
Invertebrate trace fossils	Possible meiofaunal traces	–	–	Y	–	Horodyski 1988, 1993
Fossils and structures of microbial origin	Circular and semi-circular holes in vase-shaped microfossils	–	–	Y	–	Porter et al. 2003, Porter 2016a
	"Cryptalgal" features	–	Y	–	–	Ford 1990
	"Microbial mats"	–	–	Y	–	Bohacs and Junium 2007
	Stromatolites: *Baicalia*-type	–	Y	Y	–	Ford and Breed 1973a; Bloeser et al. 1977; Ford 1990; Dehler et al. 2001, 2012, 2017
	Stromatolites: *Boxonia*-type	–	–	Y	–	Ford and Breed 1973a; Bloeser et al. 1977; Ford 1990; Dehler et al. 2001, 2012
	Stromatolites: *Cryptozoon occidentale*	–	–	Y	–	Dawson 1897, Ford and Breed 1973a
	Stromatolites: *Inzeria*-type	–	Y	–	–	Ford and Breed 1973a; Bloeser et al. 1977; Ford 1990; Dehler et al. 2001, 2012, 2017
	Stromatolites: *Stratifera*-type	–	Y	–	–	Ford and Breed 1973a; Bloeser et al. 1977; Ford 1990; Dehler et al. 2001, 2012, 2017
	Stromatolites: general	–	Y	Y	–	Ford and Breed 1973a; Horodyski and Bloeser 1983; Ford 1990; Porter and Knoll 2000; Dehler et al. 2001, 2017; Porter and Riedman 2016
	"Vampire traces"	–	Y	Y	Y	Porter 2016a, 2016b, 2017
Acritarchs and colonial organic-walled microfossils	**Overall acritarchs and colonial organic-walled microfossils**	**–**	**Y**	**Y**	**Y**	**–**
	Bavlinella faveolata	–	–	Y	–	Nagy et al. 2009, Dehler et al. 2012
	Caelatimurus foveolatus	–	Y	–	–	Porter and Riedman 2016
	Cerebrosphaera globosa	–	Y	Y	Y	Nagy et al. 2009, Porter and Riedman 2016, Dehler et al. 2017
	Cerebrosphaera sp.	–	Y	–	–	Dehler et al. 2017
	Chuaria circularis	–	Y	Y	–	Ford and Breed 1973a, 1973b; Bloeser et al. 1977; Vidal and Ford 1985; Ford 1990; Porter and Knoll 2000; Nagy et al. 2009; Dehler et al. 2012, 2017; Porter and Riedman 2016
	Culcitulisphaera revelata	–	Y	Y	–	Nagy et al. 2009, Porter and Riedman 2016

Appendix Table 4-A-2 (continued). Chuar Group fossil taxa. Zck = Kwagunt Formation, Zcg = Galeros Formation, and Zcn = Nankoweap Formation.

Category	Taxon	Zcn	Zcg	Zck	Zcu	References
Acritarchs and colonial organic-walled microfossils (continued)	*Galerosphaera walcottii*	–	Y	Y	Y	Vidal and Ford 1985, Nagy et al. 2009, Porter and Riedman 2016
	Kaibabia gemmulella	–	Y	Y	–	Vidal and Ford 1985, Nagy et al. 2009, Porter and Riedman 2016
	Kildinosphaera chagrinata	–	Y	Y	Y	Vidal and Ford 1985, Nagy et al. 2009
	Lanulatisphaera laufeldii	–	Y	Y	Y	Vidal and Ford 1985, Nagy et al. 2009, Porter and Riedman 2016, Dehler et al. 2017
	Leiosphaeridia asperata	–	Y	Y	–	Vidal and Ford 1985
	Leiosphaeridia crassa	–	Y	Y	–	Porter and Riedman 2016
	Leiosphaeridia jacutica	–	Y	Y	–	Porter and Riedman 2016
	Leiosphaeridia minutissima	–	Y	Y	–	Porter and Riedman 2016
	Leiosphaeridia tenuissima	–	Y	Y	–	Porter and Riedman 2016
	Leiosphaeridia sp.	–	Y	Y	–	Vidal and Ford 1985, Porter and Riedman 2016
	Unspecified leiosphaerids	–	Y	Y	–	Dehler et al. 2017
	Microlepidopalla mira	–	Y	–	–	Porter and Riedman 2016
	Navifusa majensis	–	Y	–	–	Porter and Riedman 2016
	Palaeastrum dyptocranum	–	Y	–	–	Porter and Riedman 2016
	Squamosphaera colonialica	–	Y	Y	Y	Vidal and Ford 1985, Nagy et al. 2009, Porter and Riedman 2016
	Cf. *Stictosphaeridium* sp.	–	Y	Y	–	Vidal and Ford 1985, Dehler et al. 2012
	Clustered cf. *Stictosphaeridium* sp.	–	Y	–	–	Vidal and Ford 1985
	Synsphaeridium sp.	–	Y	Y	Y	Vidal and Ford 1985, Nagy et al. 2009, Porter and Riedman 2016, Dehler et al. 2017
	Tasmanites rifejicus	–	Y	Y	Y	Vidal and Ford 1985, Nagy et al. 2009
	Trachysphaeridium timofeevi	–	Y	Y	Y	Vidal and Ford 1985, Nagy et al. 2009
	Corroded *Trachysphaeridium* spp.	–	Y	–	–	Vidal and Ford 1985
	Valeria lophostriata	–	Y	Y	Y	Vidal and Ford 1985, Nagy et al. 2009, Porter and Riedman 2016
	Vidalopalla cf. *verrucata*	–	Y	Y	–	Vidal and Ford 1985, Porter and Riedman 2016

Appendix Table 4-A-2 (continued). Chuar Group fossil taxa. Zck = Kwagunt Formation, Zcg = Galeros Formation, and Zcn = Nankoweap Formation.

Category	Taxon	Zcn	Zcg	Zck	Zcu	References
Acritarchs and colonial organic-walled microfossils (continued)	*Volleyballia dehlerae*	–	Y	–	–	Nagy et al. 2009, Porter and Riedman 2016
	Unnamed Form A (Porter and Riedman 2016)	–	Y	–	–	Porter and Riedman 2016
	Unnamed Form B (Porter and Riedman 2016)	–	Y	–	–	Porter and Riedman 2016
	Unnamed Form C (Porter and Riedman 2016)	–	Y	–	–	Porter and Riedman 2016
	Unnamed Form D (Porter and Riedman 2016)	–	Y	Y	–	Porter and Riedman 2016
	Unnamed Form E (Porter and Riedman 2016)	–	Y	–	–	Porter and Riedman 2016
	Clusters of small cells, possibly *Eoentophysalis* sp. (cyanobacteria?)	–	Y	–	–	Vidal and Ford 1985
	Unspecified organic-walled microfossils	–	Y	Y	–	Bloeser et al. 1977; Vidal and Ford 1985; Horodyski 1993; Porter and Knoll 2000; Dehler et al. 2012, 2017
Filaments and sheaths	**Overall filaments and sheaths**	**–**	**Y**	**Y**	**–**	**–**
	Archaeotrichion spp.	–	–	Y	–	Horodyski 1993
	Rugosoopsis tenuis	–	Y	–	–	Porter and Riedman 2016
	Siphonophycus robustum	–	Y	Y	–	Porter and Riedman 2016
	Siphonophycus septatum	–	Y	Y	–	Porter and Riedman 2016
	Siphonophycus typicum	–	Y	Y	–	Porter and Riedman 2016
	Siphonophycus sp.	–	Y	Y	–	Horodyski 1993, Porter and Riedman 2016
	Taeniatum sp.	–	–	Y	Y	Vidal and Ford 1985, Nagy et al. 2009
	“Algal” filaments	–	Y	Y	–	Ford and Breed 1973a, Schopf et al. 1973, Bloeser et al. 1977, Ford 1990
	Bacterial filaments and filamentous bacteria	–	–	Y	–	Porter and Knoll 2000, Dehler et al. 2012
	Carbonaceous filaments	–	–	Y	–	Horodyski and Bloeser 1983
	Eukaryotic? filaments	–	–	Y	–	Porter and Knoll 2000, Dehler et al. 2012
	Filamentous sheaths, cyanobacteria?	–	Y	–	–	Vidal and Ford 1985

Appendix Table 4-A-2 (continued). Chuar Group fossil taxa. Zck = Kwagunt Formation, Zcg = Galeros Formation, and Zcn = Nankoweap Formation.

Category	Taxon	Zcn	Zcg	Zck	Zcu	References
Filaments and sheaths (continued)	Filamentous sheaths, septate?, possibly oscillatoriacean	–	–	Y	–	Vidal and Ford 1985
	Filamentous sheaths, similar to *Eomycetopsis*	–	–	Y	–	Schopf et al. 1973
	Nontubular filament fragments	–	–	Y	–	Horodyski 1993
	Unspecified filaments	–	–	Y	–	Dehler et al. 2017
Vase-shaped microfossils	**Overall vase-shaped microfossils**	**–**	**–**	**Y**	**–**	**–**
	Bombycion micron	–	–	Y	–	Porter et al. 2003
	Bonniea dacruchares	–	–	Y	–	Porter et al. 2003
	Bonniea pytinaia	–	–	Y	–	Porter et al. 2003
	Cycliocyrillium simplex	–	–	Y	–	Porter et al. 2003
	Cycliocyrillium torquata	–	–	Y	–	Porter et al. 2003
	Hemisphaeriella ornata	–	–	Y	–	Porter et al. 2003
	Limeta lageniformis	–	–	Y	–	Morais et al. 2019
	Melanocyrillium hexodiadema	–	–	Y	–	Bloeser 1985, Porter et al. 2003
	Melanocyrillium sp.	–	–	Y	–	Vidal and Ford 1985
	Melicerion poikilon	–	–	Y	–	Porter et al. 2003
	Obelix rootsii	–	–	Y	–	Morais et al. 2019
	Pakupaku kabin	–	–	Y	–	Morais et al. 2019
	Palaeoamphora urucumense	–	–	Y	–	Morais et al. 2019
	Palaeoarcella athanata	–	–	Y	–	Porter et al. 2003
	Trachycyrillium pudens	–	–	Y	–	Porter et al. 2003
	Trigonocyrillium fimbriatum	–	–	Y	–	Bloeser 1985, Porter et al. 2003
	Trigonocyrillium horodyskii	–	–	Y	–	Bloeser 1985, Porter et al. 2003
	Unspecified vase-shaped microfossils	–	–	Y	–	Bloeser et al. 1977; Porter and Knoll 2000; Dehler et al. 2012, 2017; Porter and Riedman 2016

Appendix Table 4-A-2 (continued). Chuar Group fossil taxa. Zck = Kwagunt Formation, Zcg = Galeros Formation, and Zcn = Nankoweap Formation.

Category	Taxon	Zcn	Zcg	Zck	Zcu	References
Miscellaneous fossils	**Overall miscellaneous fossils**	**–**	**Y**	**Y**	**Y**	**–**
	"Algae"	–	–	–	Y	GCM
	Organic detritus	–	Y	Y	–	Downie 1969
	Steranes	–	–	Y	–	Summons et al. 1988, Vogel et al. 2005, Dehler et al. 2012, Brocks et al. 2016
	Unspecified spheroids / unicells	–	Y	Y	–	Schopf et al. 1973, Bloeser et al. 1977, Vidal and Ford 1985, Ford 1990, Dehler et al. 2012
	Unspecified microfossils	–	Y	Y	–	Downie 1969, Ford 1990
Select pseudofossils and dubiofossils	"*Brooksella canyonensis*"	Y	–	–	–	Van Gundy 1937, 1951; Bassler 1941; Cloud 1968; Glaessner 1969; Kauffman and Steidtmann 1981; Kauffman and Fursich 1983; Ford 1990
	Various "fucoids" and other putative invertebrate trace fossils	Y	–	–	–	Ford 1990

Taxonomic Notes

Porter and Riedman (2016) should be considered the most authoritative account of GRCA's Precambrian organic-walled microfossils. A few taxa not cited in that report are included in the table here; a few taxa cited in Nagy et al. (2009) are exceptions. Some of the identifications in Nagy et al. (2009) were provisional, and because they were presented only in a list, they are difficult to evaluate (S. Porter, pers. comm., April 2019). It is likely they represent specimens later re-evaluated by Porter and Riedman (2016), so they were omitted from the table. An obvious overlap is the four forms of *Leiosphaeridia* differentiated by vesicle thickness in Nagy et al. (2009), which doubtless correspond to the four *Leiosphaeridia* species in Porter and Riedman (2016) (*L. crassa* and *L. jacutica* have thick vesicles and *L. minutissima* and *L. tenuissima* have thin vesicles). For reference, omitted taxa are:

- *Cerebrosphaera* sp. cf. *C. ananguae* (Zcg);
- *Eosynechococcus moorei* (Zcu);
- *Leiosphaeridia* sp. (very thin vesicle) (Zcg, Zck);
- *Leiosphaeridia* sp. (thin vesicle) (Zcg, Zck);
- *Leiosphaeridia* sp. (thick vesicle) (Zcg, Zck);
- *Leiosphaeridia* sp. (very thick vesicle) (Zcg, Zck);
- *Ostiana microcystis* (Zcu);
- *Pterospermopsimorpha insolita* (Zcg);
- *Trachysphaeridium levis* (Zcu);
- Unnamed Form B (Zcg, Zck)

Other taxonomic notes:

- *Cerebrosphaera buickii* of Nagy et al. (2009) = *Cerebrosphaera globosa*
- Clusters of micron-sized scales (Porter et al. 2013) = *Microlepidopalla mira*
- Cf. *Cymatiosphaeroides kullingii* of Vidal and Ford (1985) = possibly *Kildinosphaera verrucata* of Nagy et al. (2009) = *Lanulatisphaera laufeldii*
- *Collenia occidentale* = *Cryptozoon occidentale*; this also includes the so-called "Stromatopora" of some of Walcott's literature
- *Kildinosphaera lophostriata* = *Valeria lophostriata*
- *Kildinosphaera verrucata* of Vidal and Ford (1985) = *Vidalopalla* cf. *verrucata*
- *Melanocyrillium fimbriatum* = *Trigonocyrillium fimbriatum*
- *Melanocyrillium horodyskii* = *Trigonocyrillium horodyskii*
- *Leiosphaeridia* sp. A of Vidal and Ford (1985) and Nagy et al. (2009) = *Kaibabia gemmulella*
- *Satka colonialica* = *Squamosphaera colonialica*
- *Sphaerocongregus variabilis* = *Bavlinella faveolata*

- *Trachysphaeridium* sp. A of Vidal and Ford (1985), *Trachysphaeridium laminaritum* of Nagy et al. (2009) = *Culcitulisphaera revelata*
- *Trachysphaeridium laminaritum* of Vidal and Ford (1985) = *Lophosphaeridium laufeldi* of Nagy et al. (2009) = *Lanulatisphaera laufeldii*
- Species of *Trachysphaeridium* in general have been transferred to other genera (Porter and Riedman 2016), but some listed taxa are retained under *Trachysphaeridium* because it is not apparent where they should go
- Unnamed form A of Nagy et al. (2009) = *Volleyballia dehlerae*
- *Vandalosphaeridium walcottii = Galerosphaera walcottii*

Appendix 4-B. Precambrian Taxa Named From GRCA

Appendix Table 4-B-1. Precambrian taxa named from specimens found within GRCA.

Taxon	Citation	Age, Formation	Type Specimen	Notes
Cryptozoon? *occidentale*	Dawson 1897	Neoproterozoic, Kwagunt	USNM 33799 (USNM 60710 and 60711 per Rezak 1957)	Stromatolite
Chuaria circularis	Walcott 1899	Neoproterozoic, Kwagunt	Lectotype USNM 33800	Organic-walled microfossil
Kaibabia gemmulella	Porter and Riedman 2016	Neoproterozoic, Galeros	UCMP 36082a	Organic-walled microfossil
Microlepidopalla mira	Porter and Riedman 2016	Neoproterozoic, Galeros	UCMP 36104b	Organic-walled microfossil
Vandalosphaeridium walcottii	Vidal and Ford 1985	Neoproterozoic, Kwagunt	LO 5661	Organic-walled microfossil
Volleyballia dehlerae	Porter and Riedman 2016	Neoproterozoic, Galeros	UCMP 36080d	Organic-walled microfossil
Bombycion micron	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 62988	Vase-shaped microfossil
Bonniea dacruchares	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 64409	Vase-shaped microfossil
Bonniea pytinaia	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 64410	Vase-shaped microfossil
Cycliocyrillium simplex	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 64455	Vase-shaped microfossil
Cycliocyrillium torquata	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 64453	Vase-shaped microfossil
Hemisphaeriella ornata	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 62990	Vase-shaped microfossil
Melanocyrillium fimbriatum	Bloeser 1985	Neoproterozoic, Kwagunt	UCLA 58968 (now LACM?)	Vase-shaped microfossil
Melanocyrillium hexodiadema	Bloeser 1985	Neoproterozoic, Kwagunt	UCLA 58959 (now LACM?)	Vase-shaped microfossil
Melanocyrillium horodyskii	Bloeser 1985	Neoproterozoic, Kwagunt	UCLA 58976 (now LACM?)	Vase-shaped microfossil

Appendix Table 4-B-1 (continued). Precambrian taxa named from specimens found within GRCA.

Taxon	Citation	Age, Formation	Type Specimen	Notes
Melicerion poikilon	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 62990	Vase-shaped microfossil
Palaeoarcella athanata	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 62988	Vase-shaped microfossil
Trachycyrillium pudens	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 64413	Vase-shaped microfossil

Chapter 5. Paleozoic Invertebrate Paleontology of Grand Canyon National Park

By Linda Sue Lassiter[1], Justin S. Tweet[2], Frederick A. Sundberg[3], John R. Foster[4], and P. J. Bergman[5]

[1]Northern Arizona University
Department of Biological Sciences
Flagstaff, Arizona

[2]National Park Service
9149 79th Street S.
Cottage Grove, Minnesota 55016

[3]Museum of Northern Arizona
Research Associate
Flagstaff, Arizona

[4]Utah Field House of Natural History State Park Museum
Vernal, Utah

[5]Northern Arizona University
Flagstaff, Arizona

Introduction

As impressive as the Grand Canyon is to any observer from the rim, the river, or even from space, these cliffs and slopes are much more than an array of colors above the serpentine majesty of the Colorado River. The erosive forces of the Colorado River and feeder streams took millions of years to carve more than 290 million years of Paleozoic Era rocks. These exposures of Paleozoic Era sediments constitute 85% of the almost 5,000 km^2 (1,903 mi^2) of the Grand Canyon National Park (GRCA) and reveal important chronologic information on marine paleoecologies of the past. This expanse of both spatial and temporal coverage is unrivaled anywhere else on our planet. While many visitors stand on the rim and peer down into the abyss of the carved canyon depths, few realize that they are also staring at the history of life from almost 520 million years ago (Ma) where the Paleozoic rocks cover the great unconformity (Karlstrom et al. 2018) to 270 Ma at the top (Sorauf and Billingsley 1991). The Paleozoic rocks visible from the South Rim Visitors Center, are mostly from marine and some fluvial sediment deposits (Figure 5-1).

This report covers the most abundant group of fossils in GRCA, the skeletal remains of invertebrates, of which more than 99% are marine organisms. Invertebrate fossils include well-known groups such as arthropod trilobites, two-valved brachiopods and mollusk bivalves, corals, echinoderms, mollusk gastropods and cephalopods, sponges, microfossil foraminifera, and other taxonomic groups preserved because of their biomineralized tissues in calcitic, chitinophosphatic, silicic, and other durable body parts. There are more than 80 fossil invertebrate species named from GRCA.

The Paleozoic history of invertebrate life ranges from the Cambrian Period (541 Ma) and ends in the Permian Period (252 Ma), but the GRCA geologic strata cover approximately 520–270 My. The GRCA invertebrate fossil record starts in the Cambrian Tonto Group, then is interrupted for 70 My during the Ordovician, Silurian, and Early Devonian periods. This is a significant unconformity and gap in the Paleozoic fossil record at GRCA but should not be confused with the "Great Unconformity" (Elston et al. 1989). The record resumes in the Mississippian and continues with minor pauses into the middle Permian.



Figure 5-1. Grand Canyon view from South Rim Visitors Center with Paleozoic stratigraphic units marked within blue and orange bands. Blue bands delimit mostly marine and fluvial deposits that contain invertebrate body fossils; orange, terrestrial deposits with no invertebrate body fossils (LINDA S. LASSITER).

The Paleozoic faunal record of GRCA invertebrates offers glimpses of fossil communities important to the identification of fossil assemblages before and after major global extinction events. GRCA Cambrian fossils document assemblages after the late Cambrian (Stage 4) Sinsk Event of about 513 Ma (Zhuravlev and Wood 2018) in the Tonto Group. Other GRCA assemblages predate (Late Devonian Temple Butte Formation) and postdate (Early Mississippian Redwall Limestone) the Devonian Frasnian–Famennian global extinction event (Raup and Sepkoski 1982; Bond and Wignall 2005). Significant fossil assemblages are also recorded at GRCA before the Permian–Triassic mass extinction event (Kaibab Formation).

Even though the significance of the Paleozoic fossil record at GRCA is potentially as vast as the thousands of square kilometers within the park boundaries, a mere fraction of the park has been surveyed for invertebrate body fossils. Due to the rugged terrain of GRCA, most Paleozoic specimens inventoried are reported from near maintained trails, in the most-traveled inner corridor areas, and along the 277 river miles of the Colorado River. There is therefore an enormous untapped

reservoir of information at GRCA, especially pertaining to invertebrate body fossils of the Paleozoic Era.

Methods

More than 7,000 specimens (Appendix 5-A) and almost 270 species (Appendix 5-B) have been identified from the literature and museum records of the Grand Canyon National Park Museum (GRCA), Museum of Northern Arizona (MNA), and National Museum of Natural History (USNM). The GRCA museum has more than 5,800 fossil specimens interpreted as Paleozoic invertebrate body fossils, more than 4,600 of which were accessioned before 1980. The MNA has 214 invertebrate specimens known to be from GRCA localities. USNM online listings include more than 500 GRCA invertebrate body fossil specimens, with 42 specimens noted as holotypes. 80 species of Paleozoic invertebrates have been named from fossils found within GRCA (Appendix 5-C), and another 11 are based on fossils that may have been found in the park (Appendix 5-D). A number of other institutions reposit smaller collections of GRCA fossils (see for example Appendix D), and it was not possible to document them all to the same extent as the GRCA, MNA, and USNM collections. The literature acts as a partial proxy for these institutions; about 590 specimens have been identified from GRCA in the literature.

Localities were assessed as much as possible to verify that only those within park boundaries were included. It is inevitable that some reports have been omitted or overlooked, because of the long history of research, varying standards of locality reporting, and other factors. Conversely, some records that actually came from outside of GRCA may be included. Stratigraphy and taxonomy are also uncertain in some cases.

All taxonomic names have been updated to use currently accepted classifications and genus/species names. The original name in the literature or accession records was checked against several paleontology databases including the Global Biodiversity Information Facility (GBIF at gbif.org), the Paleobiology Database (PBDB at paleobiodb.org), and the World Foraminifera Database (marinespecies.org; Hayward et al. 2019). For additional details of the review process for this report, see the section on taxonomic uncertainty in Appendix 5-A.

Invertebrate Body Fossils

The Paleozoic history of Grand Canyon is recorded in the lineages of many phyla (Table 5-1), including the Arthropoda (trilobites, bradoriids, and insects), Brachiopoda (inarticulate and articulate brachiopods), Cnidaria (tabulate and rugosa corals), Echinodermata (eocrinoids, crinoids, sea stars, and sea urchins), Mollusca (bivalves, gastropods, cephalopods, and scaphopods), Porifera (sponges and sponge-like animals), Foraminifera (microfossils), and other groups (Problematica) of uncertain taxonomic affinity (*Chancelloria*, Hyolithida, *Margaretia*, *Scenella*, and *Tontoia*). The Paleozoic history of invertebrates at GRCA covers more than 180 million years, 13 named geologic formations, more than 7,000 collected fossils held at numerous repositories, and literature reports of more than 260 taxa.

Table 5-1. The number of Paleozoic invertebrate body fossils of GRCA by phylum as reported from the literature or as listed in the museum collections included in this report. Specimens in the collections that are not assigned to a phylum, or are composite blocks of undetermined specimens, comprise the undetermined category.

Phylum/Period	Cambrian	Devonian	Mississippian to Pennsylvanian	Permian	Total
Arthropoda	1419	2	10	45	1476
Brachiopoda	370	4	951	1062	2387
Bryozoa	0	0	96	113	209
Cnidaria	0	4	144	51	199
Echinodermata	9	0	39	139	187
Foraminifera	0	0	263	3	266
Mollusca	8	1	230	1260	1499
Porifera	7	0	0	40	47
Problematica	70	0	0	0	70
Undetermined	33	29	116	538	716
Total	**1916**	**40**	**1849**	**3251**	**7056**

Early Paleozoic faunas at GRCA are preserved in the Cambrian Tonto Group and are dominated by trilobite arthropods (Trilobita), but also contain phosphatic brachiopods (Linguliformea), calcitic brachiopods (Rhynchonelliformea), bradoriid arthropods (Bradoriida), eocrinoid echinoderms (Eocrinoidea), and the problematic chancelloriids (Chancelloriida) and hyoliths (Hyolithida). After the fossil record hiatus of the Ordovician to Early Devonian, GRCA fossil communities in the middle Paleozoic of the Temple Butte Formation are sparse and rarely identified beyond class, including Trilobita, Rhynchonellata, Anthozoa, Gastropoda, and potentially Stromatoporoidea. The Mississippian (Redwall and Surprise Canyon formations) invertebrates include a diverse assemblage of brachiopods, stenolaemate bryozoans (Bryozoa), echinoderms, rugose and tabulate corals (Anthozoa), and mollusks, primarily bivalves and gastropods but also cephalopods. Additionally, an extensive study of foraminifera has been done for these Carboniferous units (Skipp et al. 1966; Skipp 1969). The Pennsylvanian stratigraphic units (Supai Group formations of the Watahomigi, Manakacha, and Wescogame) are less fossiliferous due in part to the eolian and shallow water environments of deposition (Blakey 1990) but do have examples of lingulid, rhynchonellid, and strophomenid brachiopods, bivalves, gastropods, and crinoids. The Permian strata of GRCA includes the Supai Group Esplanade and Coconino Sandstone formations of primarily eolian, wind-driven origin (Blakey 1990) which lack macroscopic body fossils. The Hermit Formation is better known for flora but has a few noteworthy instances of arthropod insect body fragments and a possible eurypterid (Carpenter 1927, 1928; White 1929; Spamer 1984). With the return of shallow marine environments during the deposition of the Toroweap and Kaibab formations, GRCA invertebrate body fossils include brachiopods, bryozoans, corals, gastropods, bivalves, scaphopods, cephalopods, crinoids, echinoids, sponges, and a few proetid trilobites.

Stratigraphic Distribution of Fossils

The following sections document invertebrate body fossils found from the Paleozoic stratigraphic units of GRCA. Invertebrate body fossils have been confirmed from all Paleozoic formations at GRCA except for the Sixtymile Formation (recently found to be Cambrian in age and placed in the Tonto Group; Karlstrom et al. 2018, 2020), the Frenchman Mountain Dolostone, the Pakoon Limestone, and the Coconino Sandstone.

Cambrian Fauna

The early Paleozoic fossil record at GRCA is represented by the Cambrian Tonto Group, from which many invertebrate body fossils have been reported although Cambrian exposures cover only 13% of the park. The extensive Cambrian fauna of GRCA includes species from the groups Bradoriida, Brachiopoda, Echinodermata, Hyolithida, Porifera, and Trilobita. The Tonto Group includes five named formations (Karlstrom et al. 2020): the Sixtymile Formation, Tapeats Sandstone (Tapeats Formation), Bright Angel Shale (Bright Angel Formation), Muav Limestone (Muav Formation), and Frenchman Mountain Dolostone. No definite invertebrate body fossils have been reported from the Sixtymile Formation and Frenchman Mountain Dolostone at GRCA. Detailed geologic descriptions of the Tonto Group and other Paleozoic units of GRCA are discussed in Chapter 3 of this document. However, it is worth noting that the transition between the Tapeats Sandstone and Bright Angel Shale is difficult to distinguish, and for this report, the traditional interpretations and names of the stratigraphic units have been retained.

Tonto Group: Undifferentiated Geologic Unit Fauna

Species from the Tonto Group (Ctu, Cu) that are not assigned to one of the constituent geologic units include asaphid, corynexochid, olenellid, and ptychopariid trilobites, brachiopods of the Kutorginata, Lingulata, Paterinata, Rhynchonellata and Strophomenata, and hyoliths (Hyolitha). The GRCA brachiopods of the Cambrian are dominated by phosphatic types (Linguliformea) that first appeared globally during Series 2 (Stage 3) of the Cambrian (Kouchinsky et al. 2012).

Most arthropods from undifferentiated Cambrian rocks at GRCA are trilobites, represented by several orders. Corynexochids include *Anoria*, *Athabaskia kanabensis*, *Glossopleura* (*G. boccar*), *Kootenia* (*K. mckeei*), and *Ptarmigania.* Ptychopariids (Figure 5-2) include *Elrathia*, *Kochina angustata*, *Parehmania kwaguntensis*, *P. nitida*, *Proehmaniella hebes*, *Ptychoparia*, *Spencella* (*S. diligens*), and *Trachycheilus typicale* (Walcott 1890, 1916a; Schuchert 1918a; Noble 1922; Schenk and Wheeler 1942; Resser 1945; Foster 2011; Bonde et al. 2018). The asaphid *Glyphaspis kwaguntensis* (Resser 1945) and the olenellid *Olenoides* (*Olenellus*?) (Walcott 1890) have also been reported from undifferentiated Cambrian rocks at GRCA. Although there are specimens in the GRCA collections listed as agnostid trilobites, these are considered to be misidentified ptychopterids as agnostids are more typical of open ocean environments not found at GRCA. Additionally, there are fossils reported as ostracods by Walcott (1890) from the Cambrian Tonto Group, but these are presumably bradoriids.



Figure 5-2. Cambrian trilobite fragments of *Solenopleurella porcata* (USNM 108626; now *Spencella porcata* Rasetti 1963) from undifferentiated Tonto Group, possibly the Muav Limestone. Scale bar 1 cm (0.4 in) (NPS/ANNE MILLER).

Brachiopod species listed as from undifferentiated Tonto Group units include representatives from the inarticulate and phosphatic Linguliformea, and articulate calcitic Rhynchonelliformea (Williams et al. 1996). Brachiopods today still have these two kinds of biomineralization in their shells, a trait established during the early Cambrian (Zhuravlev and Wood 2018). Extant crinoids (Echinodermata), like brachiopods, may be called "Articulata" (Wright et al. 2017), adding confusion to the continued use of this term. Linguliforms (Lingulata) include *Acrothele*, several species of *Lingulella*, and *Lingulepis perattenuatus*. The paterinate linguliform species include *Iphidella pannula*, *Micromitra (Paterina) crenistria*, and *Iphidea ornatella.* Rhynchonelliforms include the kutorginid *Nisusia* (*N. kanabensis* and *N. obscura*), and protorthid *Protorthis* (Walcott 1890, 1912a, 1916a; Resser 1945).

Tonto Group: Tapeats Sandstone Fauna

The Tapeats Sandstone (Ctt) is mapped as covering about 88 km^2 (34 mi^2) or approximately 2% of the Paleozoic units exposed in GRCA (13% of the Tonto Group exposures). The characteristics of the depositional environment and multiple facies of the Tapeats Sandstone yield bioturbated surfaces better known for trace fossils than body fossils (Rose 2006). Thirteen invertebrate taxa are reported from the Tapeats Sandstone in various sources, although at least some of these are probably from the Bright Angel Shale instead. Arthropods include the redlichiid *Olenellus* (Resser 1945) and ptychopariid *Spencella*, and the bradoriids *Indianites* (*I. curta*, *I. impressa*) and *Walcottella*. The brachiopods are represented by five species of the lingulate genus *Lingula* (*L. chuarensis, L. euglypha, L. lineolata*, *L. spatulus*, and *L. zetus*), and two species of paterinate brachiopods, *Iphidella pannula* and *Paterina (Iphidea) crenistria.* Schuchert (1918a) reported an unidentified obolellid.

Tonto Group: Bright Angel Shale Fauna

The Bright Angel Shale (Ctba) produces proportionately more fossils per area in the Tonto Group and covers 5% of the GRCA geologic area (33% of the Tonto Group), about 266 km^2 (103 mi^2). This unit has locally abundant corynexochid and ptychopariid trilobites (Foster 2011) and chitinophosphatic brachiopods. As many as 59 species of fossil invertebrates have been named from

the Bright Angel Shale of GRCA (Appendix 5-C), although not all of them are currently considered valid. These 59 taxa include 23 bradoriids, 22 trilobites, 13 brachiopods, and the eocrinoid *Eocrinus multibrachiatus* (now classified in *Gogia*). The bradoriids were collected from only two localities within the park boundaries and include species from the genera *Dielymella*, *Indiana*, and *Walcottella* (Ulrich and Bassler 1931). The named trilobites from GRCA are in the genera *Acrocephalops*, *Albertella*, *Alokistocare* (*Amecephalus*), *Anoria*, *Ehmaniella*, *Elrathia*, *Glossopleura*, *Glyphaspis*, *Kootenia*, *Pachyaspis*, *Parehmania*, *Proehmaniella*, and *Spencia* (Walcott 1916a, 1916b; Resser 1945). The brachiopods named from GRCA are in the genera *Billingsella*, *Dictyonina*, *Iphidea*, *Lingulella*, *Nisusia*, and *Paterina* (Walcott 1897, 1898, 1902, 1905; Resser 1945).

Arthropoda

Trilobite specimens collected from GRCA are abundant but are often only fragments of the exoskeleton (Figure 5-3). GRCA fossil records capture some of the diversity of Cambrian trilobite assemblages (Schenk and Wheeler 1942; Resser 1945; Foster 2011). Trilobite genus and species taxonomy, and family-level assignments, are frequently revised. For example, ptychopariid trilobites of the genus *Amecephalus* (Figure 5-4A) are taxonomically listed as kochaspids, pending the new family description and assignments that are part of Sundberg et al. (in review).



Figure 5-3. Trilobite fragments including olenellid genal spines of unidentified species, and an olenellid pygidium boxed in red from the Cambrian Bright Angel Shale (GRCA 6598). Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).




Figure 5-4. The Cambrian trilobites (GRCA 9529) *Amecephalus althea* (A) and a dolichometopid (B) from the Bright Angel Shale. Scale bar 1 cm (0.4 in) (JOHN FOSTER).

Olenellid trilobites (Redlichiida: Olenellidae) have been found in the western portion of the Grand Canyon in the transition zone between the Tapeats Sandstone and Bright Angel Formation and within the Bright Angel Formation (Schenk and Wheeler 1942; Resser 1945). The olenellids are the earliest trilobites reported from GRCA in what is sometimes called the *Olenellus–Antagmus* horizon (McKee and Resser 1945).

tychopariid (Ptychopariida) trilobites differ from olenellids by possessing librigenae sutures, fewer thorax segments, and a distinct, but sometimes small pygidium (micropygous to subisopygous). Overall, ptychopariids are smaller than olenellids, particularly in the early Cambrian, but by the middle Cambrian ptychopariids such as dolichometopids (Figure 5-4B) could become quite large. A common ptychopariid of the Bright Angel Shale is *Amecephalus althea* (Figure 5-5B). Other ptychopariids in this unit are from the genera *Alokistocare* (*A. lepida*), *Antagmus* (*A. arizonaensis*), *Bolaspis* (*B. aemula*), *Ehmaniella*, *Elrathia* (*E. nitens*), *Elrathiella* (*E. decora*), *Kochina*, *Pachyaspis*, *Parehmania* (*P. kwaguntensis*, *P. tontoensis*), *Proehmaniella* (*P. basilica*), *Spencella* (*S. porcata*), and *Spencia* (*S. tontoensis*) (Walcott 1916a, 1916b; Schuchert 1918a; Noble 1922; Schenk and Wheeler 1942; McKee and Resser 1945; Resser 1945; Bonde et al. 2018).

Bright Angel Shale corynexochids are the most abundant trilobites of GRCA with more than 530 specimens in the GRCA museum collections (versus 221 specimens of ptychopariids). Corynexochid trilobites are represented at GRCA most commonly by the genera *Anoria* (*A. tontoensis* of Figure 5-5A), and *Glossopleura* (*G. boccar*, *G. mckeei*, *G. meriwitica*, *G. productus*) (Walcott 1916a; Schuchert 1918a; Noble 1922; Schenk and Wheeler 1942; Resser 1945). Other trilobite genera of this order from GRCA include *Albertella* (*A. schenki*), *Kootenia* (*K. schenki*, *K. simplex*), *Olenoides*, *Prozacanthoides*, *Ptarmigania*, and *Zacanthoides* (Schenk and Wheeler 1942; Resser 1945; Bonde et al. 2018).




Figure 5-5. Examples of common, well-preserved Bright Angel Shale ptychopariid trilobites. **A.** *Anoria tontoensis* (GRCA 11520) (NPS/REBECCA FOSTER). **B.** *Amecephalus althea* (GRCA 11612) (LINDA S. LASSITER). Scale bar 1 cm (0.4 in).

Another group of arthropods that are common in the Bright Angel Shale are the bradoriids. Bradoriids are extinct bivalved, phosphatic arthropods generally less than 20 mm (0.8 in) long and appear morphologically similar to the calcitic ostracods (Figure 5-6). Bradoriids can be quite common in some locations in the Bright Angel Shale. Recent phylogenetic analyses of all arthropods now place bradoriids as a basal member of the Euarthropoda, distinct from sister clades of the trilobites, chelicerates, and the derived clade of Mandibulata (Legg et al. 2013). However, some databases still place bradoriids in the class Ostracoda (GBIF) or resolved as a basal arthropod clade (PBDB).



Figure 5-6. Bradoriids from the Bright Angel Shale of GRCA. Scale bar 1 cm (0.4 in) (FREDERICK SUNDBERG).

Bradoriids have been described globally only from the Cambrian (Topper et al. 2011) and on the surface appear to have been highly endemic in eastern GRCA. Nearly two dozen bradoriid species have been named and described from the Bright Angel Shale of eastern GRCA by Ulrich and Bassler (1931) (Appendix 5-C). It would not be surprising if this figure is inflated by over-splitting. Fitting the GRCA bradoriids into the biogeography and distributions of other Cambrian faunal assemblages

might be useful as bradoriids became common worldwide before trilobites (Hou et al. 2010). GRCA bradoriids have been assigned to the genera *Bradoria, Dielasma*, *Indianites*, and *Walcottella* (Ulrich and Bassler 1931; Resser 1945); species are listed in Appendix Table 5-A-1 and 5-B-1.

Brachiopoda

Bright Angel Shale brachiopods from GRCA include phosphatic linguliform lingulids and paterinids, and examples of several orders of calcitic forms (Appendix Table 5-A-1 and 5-B-1). The similarity in size and chitinophosphatic composition make linguliform brachiopods easily confused with bradoriids in Cambrian stratigraphic units. Lingulid specimens are from *Acrotreta, Lingulella,* and *Lingula* (Frech 1893; Walcott 1898, 1902, 1912a, 1916a; Schuchert 1918a; Noble 1922; Resser 1945). Paterinids represented in the Bright Angel Shale include *Dictyonina, Iphidella, Micromitra,* and *Paterina* (Walcott 1897, 1912a, 1916a; Noble 1922; Resser 1945).

Calcitic-valved brachiopods of the Bright Angel Shale include obolellides, kutorginates (*Nisusia*), the rhynchonellid orthid *Wimanella,* an unidentified thecideid (*Diraphora?*), and the strophomenid *Billingsella* (Frech 1893; Schenk and Wheeler 1942; Resser 1945). Obolellides, such as *Obolella,* have oval, biconvex valves and a minute foramen opening at the tip of the pedicle valve. Kutorginates of the genus *Nisusia* have a strophic hinge, prominent, extended, curved beak over the interarea and can reach lengths of 1 to 2 cm (0.4 to 0.8 in). The only species of *Nisusia* from GRCA identified to a specific unit is *Nisusia noblei* from the Bright Angel Shale (Walcott 1924; Resser 1945). The orthid *Wimanella* is only about 4 mm (0.2 in) wide, can be poorly preserved and difficult to identify (Figure 5-7). Three specimens from the GRCA collections of the Bright Angel Shale have been incorrectly assigned to the genus *Composita,* which has no fossil record before the Devonian (Grinnell and Andrews 1964). Therefore, these specimens require reidentification; it is probable that they are orthids.



Figure 5-7. The phosphatic orthid brachiopod *Wimanella* (GRCA 8423) preserved mostly as outlines of the valves in a slab from the Bright Angel Shale. Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

The strophomenids *Billingsella* (Bright Angel Shale) and *Clitambonites* (Muav Limestone) are small brachiopods less than 2 cm (0.8 in) wide with strophic hinge lines and calcitic valves. *Billingsella* is

unique in possessing features that are associated with both strophomenids and orthids as a basal configuration, which could warrant placement in a distinct clade (Congreve et al. 2015).

Echinodermata

Eocrinoids (Eocrinoidea), a type of early stalked echinoderm, are reported from the Bright Angel Shale of GRCA (Kirk 1945; Robison 1965; Sprinkle 1973). *Eocrinus multibrachiatus*, named from a GRCA specimen in Kirk (1945), is now assigned to the genus *Gogia* (Appendix 5-C). *Gogia longidactylus* (GRCA 2641) is also known from Bright Angel Shale (Figure 5-8) and was illustrated in Sprinkle (1973). An undetermined cystoid (GRCA 2479) has been identified as a rhombiferid but has not been re-examined. An ascocystitid, *Eocystites*, was reported from this unit by Walcott (1916a) and Noble (1922), but the fossil was not in the collections surveyed for this report.



Figure 5-8. An eocrinoid, *Gogia longidactylus* (GRCA 2641), from the Bright Angel Shale. Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

Mollusca

Several Bright Angel Shale specimens in GRCA collections are identified to the molluscan classes Bivalvia (four specimens) and Gastropoda (three). Given the rarity of Cambrian mollusks in general, it is likely that these are misidentified.

Porifera

Three specimens (GRCA 2532, 2533, and 2537) are listed doubtfully as the archaeocyathid *Ethmophyllum chankensis*. However, archaeocyathids are not known later than the early Cambrian Series 3 (Lee et al. 2016) and other Bright Angel Shale specimens support a middle Cambrian age. If these fossils are confirmed as archaeocyaths, they would be among the youngest reported archaeocyathids in the world. Archaeocyaths were also reported in Resser (1945). There are a few other sponges listed but not identified at the GRCA museum.

Incertae Sedis

Several specimens of uncertain classification are reported from the Bright Angel Shale at GRCA. One (USNM 57660) is the holotype of *Tontoia kwaguntensis* (Figure 5-9), reported as a possible trilobite by Walcott (1912b). *Margaretia*, a stalked organism sometimes misspelled *Margaritia*, was described from the Burgess Shale as a fossil of algal or hemichordate affinities and is also reported from the Bright Angel Shale of GRCA (Schenk and Wheeler 1942). *Margaretia* specimens are listed in the GRCA collections with one specimen noted as having “thin fronds” or as possibly

dolichometopid trilobite fragments (GRCA 598, 605, 609, and 630). *Margaretia* is a genus used for algae, but the Burgess Shale specimens have recently interpreted as the tube enclosure of a hemichordate (Nanglu et al. 2016). This organism is of uncertain affinity and only listed in this invertebrate body fossil summary as a placeholder until taxonomic placement is resolved.



Figure 5-9. An illustration of *Tontoia kwaguntensis* (USNM 57660), which is described as a 25 mm (1 in) long animal of trilobite or other arthropod affinity (Walcott 1912b: Plate 24).

Chancelloria (several GRCA and MNA specimens) is a sponge-like animal of uncertain affinity (Elliott and Martin 1987), described as calcareous spicules of a sponge by Walcott (1920). It is interpreted as a sack-shaped animal with a spicular body (Elliott and Martin 1987). This animal is now considered to be related to *Wiwaxia* and halkieriids (Bengtson and Collins 2015) although it is *incertae sedis* at the phylum level. *Chancelloria* is associated with the class Coeloscleritophora (Elliott and Martin 1987). The spicules or sclerites from the Bright Angel Shale have a ring of five spines that are about 0.5 to 6.0 mm (0.02 to 0.2 in) in length with short central spines of 0.5 to 1 mm (0.02 to 0.04 in) length (Elliott and Martin 1987). The shapes of the hollow sclerites are interpreted as similar to, but distinct from, those of *Chancelloria pentacta*, not found at GRCA.

There are 38 specimens of hyoliths (Figure 5-10) in the collections at GRCA museum, all identified as *Hyolithes* sp. Hyoliths are an extinct group of animals with shells composed of aragonite. Their classification has long been debated, as *incertae sedis*, in their own phylum (Hyolitha), a class within Mollusca, or most recently as an ancestral lophophorate (Moysiuk and Caron 2017). Hyoliths from GRCA (Figure 5-10) are listed as in the genus *Hyolithes.* The specimens at GRCA superficially resemble the Burgess Shale and Spence Shale hyolith *Haplophrentis* (Babcock and Robison 1988) and may belong to that genus. Most of the time, these fossils are found disarticulated and the operculum may be mistaken for an inarticulate brachiopod.



Figure 5-10. Two hyoliths from the Bright Angel Shale (GRCA numbers uncertain). The hyolith on the right retains the operculum cap. Scale bar 1 cm (0.4 in) and U.S. quarter (diameter 24.26 mm or 0.955 in) (LINDA S. LASSITER).

Tonto Group: Muav Limestone Fauna

Outcrops of the Muav Limestone (Ctm) in GRCA represent the largest area of the Tonto Group (40%) and about 6% of the park (300 km^2 or 116 mi^2). However, the invertebrate body fossil assemblage is not as diverse as that found in the Bright Angel Shale. As with the other Tonto Group members, the Muav Limestone is trilobite-dominated with ptychopariid and corynexochid species. There are also several enigmatic fossils from the Muav Limestone including hyoliths, the single-shelled, mollusk *Helcionella*, and *Scenella hermitensis,* which is named from GRCA (Resser 1945). The hyoliths are discussed in more detail in the Bright Angel Shale section.

Seven fossil invertebrate taxa have been named from specimens found in the Muav Limestone of GRCA, with another 12 to 14 collected either from the Bright Angel Shale or Muav Limestone (Appendix 5-C). Those definitely from the Muav Limestone include *Scenella hermitensis*, the trilobite species *Bolaspis aemula*, *Glyphaspis tecta*, *Kootenia schenki*, *Spencella erosa*, and *S. porcata* (Resser 1945), and the brachiopod *Nisusia noblei* (Walcott 1924). Trilobites of the Muav Limestone may be attributed to similar assemblage zones as those in the Bright Angel Shale (Stage 5, Series 3), although this is still under current study.

Arthropoda

Identified trilobites of the Muav Limestone include the corynexochid genera *Anoria*, *Athabaskia*, *Dorypyge*, *Glossopleura*, *Kootenia*, *Neolenus,* and *Zacanthoides*; the asaphid *Glyphaspis*; and the ptychopariids *Alokistocare*, *Anomocarella*, *Bolaspis*, *Pachyaspis*, *Parehmania*, *Saukia,* and *Spencella* (Noble 1922; Stoyanow 1936; Schenk and Wheeler 1942; Resser 1945). Specimens identified to the species level include *Bolaspis aemula*, *Glossopleura meriwitica*, *Glyphaspis kwaguntensis*, *G. tecta*, *G. vulsa*, *Kootenia havasuensis, K. mckeei, K. schenki, K. simplex, Pachyaspis moorei, Parehmania kwaguntensis, Spencella porcata,* and *S. erosa* (Schenk and Wheeler 1942; Resser 1945). There are more than 60 unidentified or tentatively identified specimens of trilobites in the GRCA museum collections, including some identified tentatively as *Elrathiella*.

Brachiopoda

The strophomenid *Clitambonites* is found in the Muav Limestone (Noble 1922). The kutorginates are of the same diversity as in the Bright Angel Shale (*Nisusia*). Other Muav Limestone brachiopods are chintophosphatic forms in the genera *Dictyonina, Discina, Lingula, Lingulella (Lingulepis), Trematis,* and possibly *Acrotreta* (Walcott 1883, 1890; Noble 1922; Resser 1945). *Syntrophia* was identified by Noble (1922) but is typically found in Lower Ordovician to Silurian rocks which are missing from GRCA.

Echinodermata

Three undetermined cystoids (GRCA 2174, 11842, 11843) have been cataloged from the Muav Limestone in the GRCA museum collections but are doubtful in their identification and were excluded from the final listing (Appendix 5-A). Although James Sprinkle (UT-Austin) has interpreted one as surface impressions likely from a set of burrows, the other specimens have yet to be re-examined.

Mollusca

Helcionelloida is a class of snail-like mollusks that appeared in the Cambrian and went extinct in the Ordovician. *Helcionella* has been informally identified from the Muav Limestone (A. Palmer, USGS, written comm. to M. B. Ingham, 1963). The specimen from the Muav Limestone of GRCA is held at Lake Mead NRA (LAKE 2243), having been collected from a part of GRCA once within the recreation area.

Porifera

Resser (1945) indicated the presence of archaeocyathids in the Muav Limestone. However, these have most likely been misidentified.

Incertae Sedis

Hyolithes is reported from the Muav Limestone (Resser 1945) and in the collections at GRCA. Additionally, the Muav also has an unusual invertebrate fossil named *Scenella* (Resser 1945). First reported as a gastropod and classified in current databases (PBDB) as a monoplacophoran mollusk, *Scenella* is a simple, cap-shaped shell that resembles the modern limpet. *Scenella* from GRCA is generally 1 to 1.5 cm (0.4 to 0.6 in) long, oval in outline, and composed of either calcium phosphate or calcium carbonate. So far, it has only been reported from the Muav Limestone at GRCA (*Scenella hermitensis,* MNA 1640 and USNM 108568a, named in Resser 1945). Re-interpretations of similar shaped objects from other Cambrian localities have suggested that *Scenella* is an internal float device used by soft-bodied cnidarias (hydrozoans) (Yochelson and Cid 1984). *Chancelloria* may also be found in the Muav Limestone as mentioned by Schenk and Wheeler (1942) and Resser (1945).

Devonian Fauna

Temple Butte Formation Fauna

Two percent of the park has been marked as exposures of the Devonian Temple Butte Formation (Dtb) (95 km^2 or 37 mi^2), which is limestone or more commonly dolomitized (Rowland et al. 1995). This formation can be found as isolated channel-fills cut into the underlying Muav Limestone with many larger exposures of limestone and dolostone in the western GRCA (Beus 1990). The Temple

Butte Formation has probably not been sufficiently surveyed for fossils as reflected in the sparse number of specimens in the collections at the GRCA museum. Out of more than 7,000 Paleozoic invertebrate body fossils in GRCA collections, only about 40 are from the Temple Butte Formation and 30 of those are not even identified to phylum (Appendix Table 5-A-2). A few fossils, one brachiopod and one coral, are reported from locations of uncertain Devonian or Mississippian age from the literature (Beus 1990) that are believed to be from the Temple Butte Formation. Beus (1990) also noted the presence of possible stromatoporoids, but no further information on locality or current location is known. Otherwise, unidentified specimens of the Temple Butte Formation include a few rugose corals, gastropods, crinoids, and brachiopods (Beus 1990) but none are identified beyond genus level.

Arthropoda

There are unidentified trilobites associated with the Temple Butte Formation (GRCA 2668, 2669) that appear to be similar to *Elrathia*.

Brachiopoda

Spiriferids are the only named group of brachiopods reported from the Temple Butte Formation and are only identified as *Spirifer* sp. in the GRCA museum collections. Other unidentified brachiopods are mentioned in Walcott (1883) and Stoyanow (1936).

Cnidaria

Corals first appear in GRCA in the Temple Butte Formation. Only three specimens assigned to the order Stauriida have been identified from the literature (Walcott 1883; Noble 1922; Schenk and Wheeler 1942). One has been identified as *Streptelasma* sp. (Schenk and Wheeler 1942) but the location data is uncertain, and it may not have been within the current park boundaries.

Mollusca

One unidentified gastropod was mentioned by Walcott (1883) as being from unnamed Devonian rocks and is presumably from the Temple Butte Formation.

Mississippian Fauna

Mississippian units at GRCA include the Redwall Limestone and Surprise Canyon Formation (McKee and Gutschick 1969a; Beus 1990), which are thickest in western GRCA. These units cover deposition from the Early Mississippian (Kinderhookian, 360–355 Ma) in western GRCA to Late Mississippian (Chesterian) (McKee and Gutschick 1969b). These strata have had significant studies of the microfossil Foraminifera and macroscopic invertebrate body fossils.

Redwall Limestone Fauna

The Mississippian age Redwall Limestone (Mr) has exposures covering about 7% or 342 km^2 (132 mi^2) of the park but has produced about 20% of the specimens from the Paleozoic units of GRCA. There are four members present at GRCA, in ascending order the Whitmore Wash, Thunder Springs, Mooney Falls, and Horseshoe Mesa members (McKee 1963). The oldest member dates to the Early Mississippian (late Kinderhookian, equivalent to the middle Tournaisian), and the youngest, as Middle Mississippian (middle Meramecian, equivalent to the middle Visean) (McKee and Gutschick 1969b). The Redwall Limestone records two marine transgression-regression cycles, one in the

Whitmore Wash to Thunder Springs members, and one in the Mooney Falls to Horseshoe Mesa members (Beus 1989). These cycles created an intermittent shallow sea (Beus 2003) reflected in variations of the limestone that indicate changes in water depths and energy levels (McKee and Gutschick 1969c).

The Redwall Limestone faunal assemblages are dominated by rhynchonellid and strophomenid brachiopods. Besides the brachiopods, other invertebrate fossils include rugose and tabulate corals, bivalve and gastropod mollusks, and fenestrate and branching bryozoans. There are also a few trilobites, blastoids, and crinoids (chapters in McKee and Gutschick 1969a). A large study of foraminifera identified more than 360 specimens from the Redwall Limestone (Skipp et al. 1966; Skipp 1969).

Arthropoda

Trilobite presence is limited in the middle Paleozoic and this reduction in diversity and abundance is noted at GRCA with only a few species in the Redwall Limestone. Trilobites include *Aprathia* (Eodiscida) and the proetid trilobites *Breviphillipsia* and *Phillipsia peroccidens* (identified as *P. tuberculata* in GRCA collections; GRCA 9994) (McKee and Gutschick 1969d, 1969f; Cisne 1971; Brezinski 2017).

Brachiopoda

Most brachiopods of the Redwall Limestone are spiriferids (Rhynchonellata: Spiriferida) although by this time the orthotetids (Strophomenata) like *Derbyia* have also become common (Figure 5-11; Appendix Table 5-A-2 and 5-B-2). Carter et al. (2014) reported a number of brachiopod taxa from the Redwall Limestone of Arizona, based on specimens originally to have been described for McKee and Gutschick (1969a). Few of the studied specimens came from within GRCA itself, but it is likely that the taxa can also be found in the park.




Figure 5-11. An unidentified brachiopod with fine costae like the common *Derbyia* (Strophomenata: Orthotetida) on a fossiliferous slab from the Redwall Limestone (GRCA 11065). Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

Spiriferids represent more than 90% of Rhynchonellata specimens and the most frequently reported taxon is *Spirifer* sp. Rhynchonellata brachiopods besides the Spiriferida include representatives from the orders Athyridida, Orthida, Rhynchonellida, Spiriferinida, and Terebratulida (Frech 1893; Schuchert 1918b; Noble 1922; McKee and Gutschick 1969d; Carter et al. 2014; Bonde et al. 2018). Most reported taxa are only identified to the genus level (Appendix Table 5-A-2 and 5-B-2) (Noble 1922; McKee and Gutschick 1969d; Carter et al. 2014). This may be indicative of the poor preservation observed for most of the Redwall Limestone specimens, or relatively little study of the collections.

Of the 255 specimens of Strophomenata listed in the GRCA museum collections from the Redwall Limestone, 229 are identified as the orthid *Derbyia* sp. Aside from orthotetids, productids and strophomenids are also represented but in small numbers.

Bryozoa

Most bryozoans from the Redwall Limestone are fenestrate (Fenestrata and Fenestrida) or cystoporids (Cystoporida). Although there are several trepostome specimens at USNM reported from the Mooney Falls Member of the Redwall Limestone (Duncan 1969), none have been identified beyond order (Trepostomatida). Generally, bryozoans are fragmentary and molds with the best specimens are found in the oolitic limestone of the Mooney Falls Member (McKee and Gutschick 1969d). Encrusting, ramose, fenestrate, and branching bryozoan forms have been reported from GRCA, although most are not otherwise identified (Schuchert 1918b; McKee and Gutschick 1969d). Duncan (1969) did extensive work documenting the Redwall Limestone bryozoan assemblages and provides a helpful resource.

Fossil specimens of the Fenestrata include the genera *Fenestralia*, *Lyroporella*, *Neoreteporina*, and *Polyporella* (Duncan 1969; McKee and Gutschick 1969d). The Fenestrida from GRCA include *Fenestella* (*F. compressa, F. serratula*), *Penniretepora*, and *Polypora* (Schuchert 1918b; Duncan 1969; McKee and Gutschick 1969d). These are net-like, or lacy, zooaria with openings, or fenestrules, between the longitudinal sections of the overall colony form. The genera are distinguished by the shapes of the fenestrules between sections or by the position and numbers of the zooecium. For example, the genus *Polypora* (Fenestrida) has three to eight rows of autopore apertures on only one side (obverse) of the zoarium, and fenestrules are long and sometimes oval (Miller 1963). *Polypora* is helpful for stratigraphic correlation. It is rare in the Redwall but is also found in the Kaibab Formation. *Fenestralia* (Fenestrata) is found throughout the Redwall Limestone and has four zooecia per row that are divided by a keel into two rows on one side, and fenestrules are oval to rectangular (Burckle 1960). Most commonly, bryozoans are preserved so that the only visible diagnostic feature is the spacing and distribution of the fenestrules as the obverse side with zooecia not exposed (Duncan 1969). *Neoreteporina* is only reported from the Thunder Springs Member.

Cystoporids are found in the Redwall Limestone including *Cheilotrypa*, *Coscinotrypa*, *Dichotrypa*, and *Ramiporalia* (Duncan 1969; McKee and Gutschick 1969d). A distinctive middle Paleozoic bryozoan from GRCA is *Cystodictya*, which has lunaria in bifoliate colonies that may appear leaf-shaped when well-preserved.

Cnidaria

Redwall Limestone corals have been reviewed in detail in Sando (1969) and the GRCA fossils can be compared to specimens outside of GRCA (Sando and Bamber 1985). Tabulate corals are more diverse and abundant in the Redwall Limestone than other geologic units at GRCA, although many specimens are difficult to identify (Figure 5-12). There are specimens from the auloporids *Syringopora* (*S. aculeata*, S. *surcularia*), and *Cladochonus* (*Monilipora* of Noble 1922) (Schuchert 1918b; Noble 1922; Sando 1969; Sando and Bamber 1985; Bonde et al. 2018). *Syringopora* (Figure 5-13) has cylindrical tubes in irregular patterns with transverse connections at random locations. If the top is visible, these corals do not form the appearance of a chain but appear as masses of circular openings. Other tabulate forms include another Redwall Limestone species that may form small branches or look similar to *Syringopora* as a massive coral with conical tubes with thick, dividing walls and a classic "honeycomb" appearance on the surface. Favositid corals are represented in the GRCA collections (*Cladopora*).



Figure 5-12. A coral from the Redwall Limestone of GRCA (GRCA 4470) with poor preservation for clear diagnosis beyond phylum. Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

Rugose corals were more abundant than tabulates during the Mississippian and this is also reflected in GRCA fossil distributions. Solitary rugose corals are easily identified by their curved, ceratoid or horn-shape structure. In both individual and colonial rugosa, each polyp forms septa that radiate from the peripheral outer edge in towards the central axis. The rugose (Stauriida) coral *Vesiculophyllum incrassatum* is the most frequently reported coral from this unit (Easton and Gutschick 1953; McKee and Gutschick 1969d; Sando 1969; Bonde et al. 2018), but several other taxa are present in the Redwall of GRCA as well (Appendix Table 5-A-2 and 5-B-2).



Figure 5-13. Tabulate phaceloid coral (*Syringopora* sp.) from the Redwall Limestone of GRCA showing a massive form with individual corallites (GRCA 11055). Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

Echinodermata

Crinoid columnals are common but have limited diagnostic features for classification. In general, columnals are not usually diagnostic to genus or species, so it is usually not possible to classify isolated columnals. This helps to explain why nearly all crinoid specimens are only defined to the class level (Crinoidea). The blastoid *Pentremites* has been reported from the Redwall Limestone at GRCA (Macurda 1969); see the more detailed description in the Surprise Canyon Formation section.

Mollusca

Mollusks reported from the Redwall Limestone at GRCA include gastropods and a single cephalopod, *Rayonnoceras* (Breed 1969). A few specimens of bivalves have also been reported from the Redwall Limestone and are in GRCA collections, but were collected from localities outside of GRCA.

Many river guide texts refer to the nautiloid fossils found in abundance at Nautiloid Canyon in outcrops of the Whitmore Wash Member of the Redwall Limestone. At Nautiloid Canyon, the erosion surfaces create a false appearance of tentacles on the fossils (Billingsley and Breed 1976). They are attributed to *Rayonnoceras* sp., a longiconic orthocone (Breed 1969; Billingsley and Breed 1976). An undetermined coiled nautiloid was also briefly noted by Billingsley and Breed (1976).

Gastropods in the Redwall Limestone are often fragmented, worn, or may only be internal molds (steinkerns) (Figure 5-14), making species-level identification difficult (Yochelson 1969). Genera of the Euomphalina (Figure 5-14A) and Bellerophontida (Figure 5-14B) orders are the most common. Euomphalids at GRCA in the Redwall include the genera *Euomphalus*, *Platyceras*, and *Straparollus* (Noble 1922; Yochelson 1969). *Euomphalus* has a slight raised edge or shoulder or carinae on the whorls. *Platyceras* is a common gastropod of the middle Paleozoic with an asymmetrical spire and whorls that may not contact in later growth (evolute). *Straparollus* is similar to *Euomphalus* and distinguishing between these genera requires good preservation of the fossils. Bellerophonts include

specimens from *Bellerophon* (Noble 1922) and *Euphemites*, with the most common specimens at the GRCA museum being *E. subpapillosus*. *Bellerophon* is somewhat globular with unsculptured whorls. However, this genus has a distinctive band on the outer rim of the shell called a selenizone band. The aperture (opening) is flared, forming the widest part of the coiled shell. *Bellerophon* is about 2 cm (0.8 in) wide and found in Redwall Limestone and Kaibab Formation deposits (Figure 5-14).




Figure 5-14. Gastropods showing poor preservation common in fossils from the Redwall Limestone of the families Euomphalidae (GRCA 7641) **(A)** and Bellerophontidae (GRCA 7628) **(B)**. Ruler scale in millimeters (LINDA S. LASSITER).

Foraminifera

Fusulinids (Fusulinata), the "giant" calcareous foraminifera of the late Paleozoic, are well-represented in the Redwall Limestone at GRCA. Mississippian fusulinids were more commonly described from Russian literature before the studies conducted at GRCA (Skipp et al. 1966). Fusulinids of the order Endothyrida represent most of the diversity and are useful for biostratigraphy and zoning as with *Endothyra* found in the Horseshoe Mesa Member and *Tuberendothyra* in the Mooney Falls Member. The foram genera from GRCA include *Endothyra, Eoendothyranopsis, Septaglomospiranella,* and *Tournayella.* Foraminifera collected from GRCA and held at the USNM are listed in Table 5-2. There are a few specimens of the fusulinid *Schubertella* at the GRCA museum.

Table 5-2. Foraminifera from the Redwall Limestone of the Grand Canyon in USNM collections.

Class	Order	Species	#
Fusulinata	Archaediscida	*Pohlia henbesti*	18
	Archaediscida	*Septatournayella* cf. *S. pseudocamerata*	3
	Archaediscida	*Septatournayella?* cf. *"S. minuta"*	1
	Archaediscida	*Tournayella* aff. *"T. subangulata"*	6
	Archaediscida	*Tournayella discoidea*	18
	Earlandiida	*Earlandia* sp.	4

Table 5-2 (continued). Foraminifera from the Redwall Limestone of the Grand Canyon in USNM collections.

Class	Order	Species	#
Fusulinata (continued)	Earlandiida	*Paracaligella?* sp.	4
	Endothyrida	*Brunsiina?* sp.	1
	Endothyrida	*Chernyshinella?* aff. *"C. anteflexa"*	4
	Endothyrida	*Endothyra kleina*	1
	Endothyrida	*Endothyra* aff. *Spinothyra pauciseptata*	3
	Endothyrida	*Endothyra* cf. *E. incrassata*	2
	Endothyrida	*Endothyra gutschicki*	11
	Endothyrida	*Endothyra* sp.	12
	Endothyrida	*Endothyra tantula*	8
	Endothyrida	*Endothyra trachida*	15
	Endothyrida	*Endothyra?* aff. *"E. excellens"*	1
	Endothyrida	*Endothyra?* cf. *Plectogyra plectogyra*	4
	Endothyrida	*Eoendothyranopsis hinduensis*	4
	Endothyrida	*Eoendothyranopsis macra*	2
	Endothyrida	*Eoendothyranopsis scitula*	8
	Endothyrida	*Eoendothyranopsis* sp.	2
	Endothyrida	*Eoendothyranopsis spiroides*	15
	Endothyrida	*Eoendothyranopsis?* cf. *Banffella banffensis*	7
	Endothyrida	*Globoendothyra baileyi*	10
	Endothyrida	*Globoendothyra baileyi* subsp. *bridgensis*	4
	Endothyrida	*Globoendothyra baileyi* subsp. *poloumera*	6
	Endothyrida	*Granuliferelloides?* sp.	1
	Endothyrida	*Inflatoendothyra* aff. *I. nordvikensis*	1
	Endothyrida	*Inflatoendothyra eospiroides*	7
	Endothyrida	*Neoseptaglomospiranella dainae*	10
	Endothyrida	*Plectogyra turgida*	1
	Endothyrida	*Plectogyranopsis* aff. *P. eocompressa*	1
	Endothyrida	*Plectogyranopsis eocompressa*	8
	Endothyrida	*Quasiendothyra turbida*	3
	Endothyrida	*Septabrunsiina* (*Spinobrunsiina*) *parakrainica*	12
	Endothyrida	*Septabrunsiina mckeei*	12
	Endothyrida	*Septaglomospiranella chernoussovensis*	17
	Endothyrida	*Septaglomospiranella primaeva* subsp. *noda*	11
	Endothyrida	*Septaglomospiranella rossi*	2
	Endothyrida	*Skippella redwallensis*	12
	Endothyrida	*Spinobrunsiina torquida*	11
	Endothyrida	*Spinoendothyra bellicosta*	3
	Endothyrida	*Spinoendothyra spinosa*	13
	Endothyrida	*Spinoendothyra spinosa*	1

Table 5-2 (continued). Foraminifera from the Redwall Limestone of the Grand Canyon in USNM collections.

Class	Order	Species	#
Fusulinata (continued)	Endothyrida	*Spinoendothyra?* aff. *S. paraspinosa*	2
	Endothyrida	*Tuberendothyra* aff. *"Tuberendothyra tumulosa"*	2
	Endothyrida	*Tuberendothyra paratumula*	27
	Endothyrida	*Tuberendothyra safonovae*	8
	Endothyrida	*Tuberendothyra tuberculata*	22
	Lituolida	*Spiroplectamminoides?* cf. *Palaeospiroplectammina tchernyshinensis*	2
	Lituolida	*Spiroplectamminoides?* cf. *Spiroplectammina parva*	3
	Parathuramminida	*Calcisphaera* sp.	1
	Parathuramminida	*Tuberitina* sp.	5
Globothalamea	Lithuolida	*Glomospira?* sp.	1
Tubothalamea	Miliolida	*Cornuspira?* sp.	1

Surprise Canyon Formation Fauna

The Surprise Canyon Formation (Msc) covers only 20 km^2 (8 mi^2) as mapped but is a significant source of invertebrate body fossils (Beus 1999). The Surprise Canyon Formation is interpreted as a dendritic drainage system (Beus 2003). The formation is thickest in the western Grand Canyon, where the valleys eroded into the Redwall Limestone are deepest. The eastern GRCA exposures of the Surprise Canyon Formation are interpreted as fluvial with tidal influences only seen in the western GRCA deposits (Beus 1989). An alternate paleoenvironmental interpretation for the entire formation is a widespread shallow sea (Beus 2003).

Invertebrate body fossils have been reported from the marine-influenced middle and upper facies of the Surprise Canyon Formation (Figure 5-15) and include abundant and diverse brachiopods, bryozoans of ramose, fenestrate, and encrusting forms, echinoderms, corals, and a few bivalves and gastropods (Beus 1999; Billingsley and Beus 1999a). However, not all specimens collected and recorded in Beus (1999) from the Surprise Canyon Formation are from within GRCA or available for study from the museums.




Figure 5-15. A composite sample of unidentified brachiopods, crinoids and coral fragments from the Surprise Canyon Formation (GRCA 57636). Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

Arthropoda

Trilobite diversity continued to wane in the Surprise Canyon Formation, with only the proetid *Paladin* documented (M. Gordon, USGS, written comm. to E. D. McKee, 1979). Ostracods are also present (Billingsley and Beus 1999a).

Brachiopoda

Most of the brachiopods from the Surprise Canyon Formation belong to Rhynchonellata, with Strophomenata representing about one third of the brachiopod sample, but the museum collections are sparse (Appendix Table 5-A-2 and 5-B-2). Rhynchonellate genera are diverse with a few kinds of orthids, *Rhipidomella* and *Schizophoria* (Billingsley and McKee 1982), spiriferinids *Punctospirifer transversus* and *Reticulariina*, and terebratulids *Beecheria* and *Cranaena* (Billingsley and McKee 1982). The rhynchonellids are represented by *Cupularostrum*, *Leiorhynchoidea*, *Macropotamorhynchus*, *Rotaia neogenes*, and *Pugnoides* (Billingsley and McKee 1982). Notably absent when compared to the older Redwall Limestone assemblages are the formerly dominant *Derbyia* and *Spirifer*. The spiriferids of the Surprise Canyon Formation at GRCA include *Anthracospirifer*, *Brachythyris*, *Cleiothyridina*, *Composita*, and *Torynifer* (Billingsley and McKee 1982; Beus 1999). The Strophomenata are primarily productids (*Antiquatonia*, *Flexaria*, *Inflatia*, *Ovatia*) and the orthotetid *Orthotetes.*

Bryozoa

As is typical for a Carboniferous stratigraphic unit, the distinctive corkscrew-like bryozoan *Archimedes* is present (Beus 1999). *Archimedes* is only recorded from the Surprise Canyon Formation at GRCA. Currently, there are no formal reports of the other common bryozoan of this

period, *Polypora*, although it is reported from the Redwall Limestone (Duncan 1969; McKee and Gutschick 1969d).

Cnidaria

The corals include favositid tabulates and stauriid rugose corals. The tabulate coral *Michelinia* (Figure 5-16) has hexagonal patterns on the surface of the corallum with a raised center node in well-preserved specimens. Reported rugose corals include *Amplexus* and *Barytichisma* (Beus 1999).



Figure 5-16. Fragments of the coral *Michelinia* (Favositida) from the Surprise Canyon Formation (GRCA 7647). U.S. quarter for scale (diameter 24.26 mm or 0.955 in) (LINDA S. LASSITER).

Echinodermata

Few echinoderms are reported from Mississippian units at GRCA and almost all of the specimens identified beyond class are from the Surprise Canyon Formation. The only identified specimens of Asteroidea (sea stars) from GRCA come from this formation, with two specimens from GRCA (MNA N4010 and N4011) assigned to the order Uractinida (Uractinina in PBDB) (Beus 1999). Two additional unpublished blocks with asteroids are also in MNA collections (J. Tweet, pers. obs.). Blastoids (Blastoidea), represented by *Pentremites*, are uncommon. The only GRCA specimens in museum collections are from the Surprise Canyon Formation. Blastoids are extinct, stemmed echinoderms that have pentametric ambulacra distributed around the pear-shaped 1 to 2 cm (0.4 to 0.8 in) wide calyx or theca, thus explaining the name of *Pentremites.* The only Surprise Canyon crinoid attributed to a genus is *Cymbiocrinus*, although crinoid fragments are common (Beus 1999).

Mollusca

Few mollusks have been reported from the Surprise Canyon Formation, but the oldest definite bivalves at GRCA are found in this unit (Billingsley and McKee 1982). Bivalves include myalinids *Aviculopecten* and *Septimyalina* (M. Gordon, USGS, written comm. to E. D. McKee, 1979) and the trigoniid *Schizodus* (Billingsley and McKee 1982). The gastropods are typical of Mississippian assemblages, with *Bellerophon* (*Bellerophon*), *Euomphalus*, *Bellazona*, and *Straparollus* (M. Gordon, USGS, written comm. to E. D. McKee, 1979; Billingsley and McKee 1982).

Foraminifera

Only undetermined foraminifera have been reported from the Surprise Canyon Formation of GRCA (M. Gordon, USGS, written comm. to E. D. McKee, 1979). “Chesterian” foraminifera-producing sites on the Bright Angel Trail cited in McKee and Gutschick (1969b) are probably from the Horseshoe Mesa Member of the Redwall Limestone (Billingsley and Beus 1999b).

Pennsylvanian Fauna

Pennsylvanian geologic units at GRCA include three of the four formations of the Supai Group and represent different stages of several transgressive-regressive cycles, with the depositional setting fluctuating between continental eolian and shallow marine environments of a broad coastal plain (Blakey 2003). These units were deposited beginning in the latest Mississippian (Martin and Barrick 1999; Hodnett and Elliott 2018) and continuing through the Pennsylvanian. These strata have scattered areas of fossiliferous units with invertebrate body fossils in the carbonates most prominent in western GRCA (McKee 1982a). For convenience, the lower Permian Esplanade Sandstone of the Supai Group will also be considered here. It is a sparsely fossiliferous eolian sandstone, and invertebrates are only represented by microscopic bioclasts (fragments of fossils) (McKee 1982a). McKee (1982a) reported foraminifera, unspecified invertebrate fossils, and bioclasts of corals, bryozoans, bivalves or brachiopods, and echinoderms in thin-section from the Esplanade Sandstone. The bioclasts are possibly reworked materials from other units such as the Pakoon Limestone or may indicate that these samples were part of the Pakoon Limestone. Refinement of the presence of invertebrate body fossils in this unit would require a review of the slides but the repository for those specimens is unknown. The Pakoon Limestone, a fossiliferous marine limestone, intertongues to a limited extent with the lower Esplanade Limestone in western GRCA but has not been documented as producing fossils within the park and will not be considered further here. Outside of the park it has a fauna including corals, gastropod, brachiopods, fusulinids, and bryozoans (Spamer 1984).

Supai Group Fauna

The Supai Group (IPsu) has locally abundant marine fossils in the formations with limestone beds and covers about 14% of the park (700 km^2 or 270 mi^2). The rocks mostly date to the Pennsylvanian, and are divided into four formations, from oldest to youngest the Watahomigi Formation (IPswa), Manakacha Formation (IPsm), Wescogame Formation (IPswe), and Esplanade Sandstone (Pse) (McKee 1975, 1982; Blakey 1990). The Supai Group lies unconformably on the Surprise Canyon Formation indicating a short hiatus in deposition (Beus 1989). Of more than 500 specimens and about 80 genera identified from the Supai Group formations, 66 genera are listed from the Watahomigi Formation. Fossils in the Watahomigi Formation suggest low energy conditions (McKee 1982a) but possibly too energetic or with too much sand and silt for extensive coral growth (Gordon 1982), which explains the dominance of brachiopods in the assemblages. Fossils are sparse in the overlying Manakacha, Wescogame, and Esplanade formations at GRCA, only containing a few foraminifera, unspecified marine fossils, and bioclasts of unspecific bryozoans, brachiopods, bivalves, pelmatozoan (stalked) echinoderms, and foraminifera (Gordon 1982; McKee 1982a). Larger but still not notably diverse faunas are known for the Manakacha and Wescogame formations from areas just outside of GRCA in the western Grand Canyon area, where the marine influence was more pronounced.

The most abundant invertebrate specimens from the Supai Group within GRCA reflect a shallow marine, nearshore environment of brachiopods and bivalve mollusks (Figure 5-17). Less abundant taxa include gastropods (*Euomphalus*), disarticulated corals and crinoids. There are many unidentified specimens in the GRCA museum collections including some tentatively identified as ostracods.



Figure 5-17. Examples of brachiopods and bivalves from the Supai Group (GRCA 11326) including prominent examples of *Anthracospirifer tanoensis* (center and right), with the bivalves *Promytilus* and *Permophorus* reported as present as well. U.S. quarter for scale (diameter 24.26 mm or 0.955 in) (LINDA S. LASSITER).

Arthropoda

The arthropod community is primarily a scattered number of proetid trilobites (Gordon 1982; McKee 1982a) of the genus *Paladin* at GRCA. This genus is a common Mississippian to early Permian trilobite with an oval shape and isopygous form (Cisne 1967). The reduction of trilobite diversity at GRCA is reflected in global patterns and may also be indicative of the paleoenvironment of the time. Ostracods are recognized in bioclasts from the Manakacha Formation within GRCA (McKee 1982a).

Brachiopoda

Brachiopods from the Supai Group include species from the classes Lingulata, Rhynchonellata, and Strophomenata (Appendix Table 5-A-3 and 5-B-3) (Gordon 1982; McKee 1982a, 1982b). The lingulids *Orbiculoidea* (Figure 5-18A) and *Lingula* are not as common as other brachiopods and include the species *Orbiculoidea meekana* (McKee 1982a). The cone-like shape of orbiculoids supports interpretations of high-energy environments where collected (Gordon 1982; Spamer 1984).

The most common Supai Group brachiopods from GRCA include the spiriferid *Composita*, and, as was true in the Redwall Limestone, the ubiquitous strophomenate *Derbyia* (Figure 5-18B). There are also a few productids, but they are not as dominant as in the Permian Period. Spiriferids are classified based upon their internal lophophores with the typical, wide, hinge line and tapered points seen in

Spirifer and *Anthracospirifer* (Figure 5-17). *Composita*, however, looks more like a terebratulid in external morphology and is abundant in the Watahomigi Formation (McKee 1975).




Figure 5-18. Supai Group brachiopods with reasonable preservation (LINDA S. LASSITER). **A.** *Orbiculoidea* (Lingulida) (GRCA 1126) showing concentric growth lines. **B.** *Derbyia* (GRCA 20577) with fine costae. Scale bar 1 cm (0.4 in).

Bryozoa

A small bryozoan fauna has been reported from the Watahomigi Formation (Gordon 1982; McKee 1982a). Encrusting bryozoans are reported on the surface of brachiopods (Gordon 1982) and assist interpretations of paleoenvironments accordingly. Higher in the Supai Group, poor environmental conditions for bryozoans has resulted in only bioclasts recording their presence at GRCA for the Supai Group, primarily in the Wescogame Formation (McKee 1982a). As McKee (1982a) points out, the sandy beds created an "unfavorable" environment.

Cnidaria

Conulariids have been reported from the Watahomigi Formation (Gordon 1982; McKee 1982a). Conulariids have a unique herringbone surface pattern and four sides. The morphological features have led to many different interpretations for phylogenetic placement of conulariids. Most recently, conulariids are considered to be ancestral cnidarians related to sea jellies of the class Scyphozoa (Van Iten et al. 2006) or medusozoans (Staurozoa) with a stalk (Marques and Collins 2004).

Tabulate and rugose corals are uncommon. Undetermined corals and the favositid *Michelinia* have been reported from the Watahomigi Formation (McKee 1982a, 1982b), and a few other taxa have been found near but outside of GRCA (McKee 1982a) and so may eventually be found within the park. These fossils are typical of quieter environments. Above the Watahomigi, the only cnidarian remains reported from the Supai Group within GRCA are coral bioclasts in the Esplanade Sandstone (McKee 1982a).

Echinodermata

There are literature references that indicate echinoid spines and crinoid columnals are found in the Watahomigi Formation (Gordon 1982; McKee 1982a). Several GRCA museum specimens may be echinoderms but are not in the collections surveyed in this report.

Mollusca

The Watahomigi Formation is dominated by myalinid bivalves, including *Aviculopecten* (*A. gravidus*), *Myalina* (*M. cuneiformis*, *M. perattenuata*), and *Septimyalina* (see Appendix Table 5-A-3 and 5-B-3). Outside of GRCA, there are reports of cardiids of *Oriocrassatella*, *Wilkingia terminalis*, and *Permophorus* in the Wescogame Formation (Gordon 1982; McKee 1982a).

The only other class of mollusks known from the Supai Group of GRCA is Gastropoda, but cephalopods and rostroconchs (extinct bivalve-like mollusks which had single folded shells) have been found near the park (Gordon 1982). All of the Supai gastropods reported from within GRCA are from the Watahomigi Formation, but large bellerophontid gastropods are known outside of the park in the Manakacha and Wescogame formations (McKee 1982a). The Watahomigi gastropods include bellerophontids and euomphalinids (*Euomphalus* and *Straparollus*) (Gordon 1982; McKee 1982a).

Foraminifera

Foraminifera or foraminiferal bioclasts are known from all of the formations of the Supai Group at GRCA, with the most taxa reported from the Manakacha Group (McKee 1982a, 1982b). McKee (1975, 1982a) used fusulinids to determine the age of the Manakacha Formation (Atokan, Middle Pennsylvanian) and Wescogame Formation (Virgilian, Late Pennsylvanian).

Permian Fauna

Permian invertebrate body fossils are known from the Esplanade Sandstone (mentioned above), the Hermit Formation, the Toroweap Formation, and the Kaibab Formation within GRCA. The Hermit Formation is dated to the early Permian (Cisuralian epoch) and represents low-energy, fluvial deposition across a broad coastal plain (Blakey 2003), in a semi-arid climate with long hot and dry seasons (White 1929). The overlying eolian Coconino Sandstone (Cisuralian) interrupts the invertebrate body fossil history of the Permian Period at GRCA with a lack of body fossils but does have invertebrate trace fossils (Chapter 8). Invertebrate body fossils are once again reported in the Toroweap Formation (McKee 1938). The Toroweap records another transgression-regression cycle, with deposition varying from tidal flat, evaporites, eolian, and shallow-marine environments in a semi-arid to arid climate (Rawson and Turner 1974; Turner 1990). The Kaibab Formation forms the cap of the Grand Canyon rim at GRCA and most of the surface exposures, and was formed by a shallow sea (Noble 1914; McKee 1938).

Hermit Formation Fauna

The Hermit Formation (Ph), which covers 8% of the surface area of the park (397 km^2 or 153 mi^2), is significant because it contains the only specimens of pre-Quaternary insects from GRCA (Spamer 1984). Although sometimes called the Hermit Shale, most of the formation is sandstone, siltstone, and mudstone in a reddish-brown slope-forming unit (Blakey 2003). Invertebrate body fossils are

uncommon and often poorly preserved due to the terrestrial and arid conditions of preservation in a fluvial to deltaic environment, and the unfavorable grain size (Spamer 1984).

Arthropoda

Two fossil insect wings from the Hermit Formation of GRCA have been described as species of *Tupus*: *T. gilmorei* (Carpenter 1927), *T. whitei* (Figure 5-19) and *T. permianus* (Carpenter 1928), originally under the obsolete spelling "*Typus*". Both species are meganeurids, members of an extinct clade resembling modern dragonflies (Odonata). The finds are significant for GRCA and for dating the Hermit Formation to the early Permian (Spamer 1984).



Figure 5-19. Hermit Formation insect wing of *Tupus whitei* (GRCA 3090). Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

Other fossils reported as Insecta are a partial wing of an odonate (now lost) and a forewing of an unnamed blattoid or cockroach (USNM 71712, collected by David White in 1927) (Spamer 1984). The only reported eurypterid, *Hastimima*, from GRCA is from this unit, but details are limited as the fossil is only an external mold (White 1929) with insufficient detail to be included in recent Eurypterida phylogenetic analyses (Lamsdell et al. 2010).

Toroweap Formation Fauna

The Toroweap Formation (Pt) was included in the Kaibab Formation before McKee (1938) distinguished it as a separate formation. The Toroweap is dated to the late early Permian (approximately 273 Ma) (McKee and Breed 1969; Mathis and Bowman 2007) and is divided into three members, from oldest to youngest the Seligman, Brady Canyon, and Woods Ranch members. The Toroweap represents about 4% or 189 km^2 (73 mi^2) of exposures at the park. In the Grand Canyon, western faunal assemblages are typical of open but shallow marine environments with corals, bryozoans, brachiopods, and crinoids. The eastern areas are interpreted as a restricted-marine environment with a molluscan assemblage of bivalves, gastropods, a few nautiloid cephalopods, and scaphopods (Rawson and Turner 1974). The Brady Canyon Member is the most fossiliferous, with bryozoans, brachiopods, bivalves, nautiloids, gastropods, scaphopods, ostracods, crinoids, and sea urchins (McKee 1938). The Seligman Member has poorly preserved mollusks (McKee 1938), and the top, thin beds of the Woods Ranch Member have abundant specimens of the smooth-surfaced clam *Schizodus*, leading McKee (1938) to refer to the unit as the *Schizodus* limestone. The three

members formed during a transgression-regression cycle, with the Seligman Member as the early part of the transgression and tidal flat, the Brady Canyon Member as the maximum transgression of low-energy, shallow marine deposits, and the Woods Ranch Member as the regressive phase of an evaporitic marine shelf (Rawson and Turner 1974).

Many fossil specimens from the Toroweap Formation of GRCA are not identified to the genus level, as with the assemblage of bivalves shown in Figure 5-20, although some brachiopods, bivalves, gastropods, scaphopods and others have been (McKee 1938; Spamer 1984; Sorauf and Billingsley 1991). There are bryozoans and sponges identified only to phylum in MNA and GRCA museum collections. Some mollusks and ostracods were not identified fully in McKee (1938). Spamer (1992) summarized assemblages with sponges, corals, bryozoans, brachiopods, bivalves, nautiloid cephalopods, gastropods, scaphopods, trilobites, and echinoids from across the Grand Canyon area. There are also fossils from the GRCA area reported from undifferentiated Toroweap Formation/Kaibab Formation rocks in the museum collections. These include the bivalve *Wilkingia wyomingensis*, the gastropod *Glyptospira cristulata*, spiriferid *Squamularia*, orthotetid *Meekella occidentalis*, productid *Rugatia paraindica*, and a trepostomatid bryozoan of the genus *Rhombotrypella*.



Figure 5-20. Toroweap Formation slab with unidentified bivalves (GRCA 15324). Scale bar is 1 cm (0.4 in) (LINDA S. LASSITER).

Arthropoda

Ostracods are uncommon at GRCA, having only been reported from the Toroweap Formation (McKee 1938), Surprise Canyon Formation (Billingsley and Beus 1999a), and as bioclasts from the Manakacha Formation (McKee 1982a). The relative absence of ostracods from Devonian and later units at GRCA is puzzling. Leperditicopids are common in Devonian stratigraphic units elsewhere, as are palaeocopids from the Devonian through the Permian.

Brachiopoda

Brachiopods identified to genus or species from the Toroweap of GRCA are the productid *Peniculauris bassi* and the common spiriferid *Composita* (Schuchert 1918b; McKee 1938). A species of brachiopod (*Productus ivesi*, now *Peniculauris ivesi*) was named from a specimen found in the Toroweap Formation or Kaibab Formation at a location possibly within the park (Newberry 1861) (Appendix 5-D).

Bryozoa

Unidentified fenestrate forms are recorded in composite blocks collected from the Toroweap Formation of GRCA (Figure 5-21). McKee (1938) identified multiple beds with bryozoans in Toroweap limestone beds, but the fossils are poorly preserved.




Figure 5-21. Example of impressions of fenestrate-type bryozoan colonies (GRCA 11373) from the Toroweap Formation. Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

Echinodermata

The echinoid *Archaeocidaris* (sometimes reported under the suppressed name *Echinocrinus*) has been reported or collected from the Toroweap Formation (Newberry 1861; McKee 1938; Sorauf and Billingsley 1991; GRCA Museum records). Many specimens are cataloged at GRCA. Three echinoid species, *Archaeocidaris gracilis*, *A. longispinus*, and *A. ornatus* (now *A. coloradensis*), were possibly named from the Toroweap or Kaibab formations of the park (Appendix 5-D). Echinoids, or sea urchins, are preserved as disarticulated spines. Crinoids are noted in Toroweap Formation limestone lenses at GRCA (McKee 1938) but are mostly columnal fragments and otherwise not identifiable to genus or species.

Mollusca

The Toroweap Formation has a widespread bed in the upper Woods Ranch Member with abundant fossils of the bivalve *Schizodus* in limestone lenses (McKee 1938). The Toroweap bivalve community includes the pholadomyid *Allorisma*, the ostreid *Pteria*, the adapedont *Edmondia*, the protobranch *Solemya*, and the trigoniid *Schizodus* (McKee 1938). There are also some scallops

(*Deltopecten*) and the asymmetrically valved *Nuculana* (including *Leda* and *Nucula* of McKee 1938). Gastropods are represented by several genera not defined to species (McKee 1938). McKee (1938) mentions the cephalopod *Domatoceras*.

There are scaphopods in the Toroweap Formation, but unlike the Kaibab forms, the specimens have not been identified beyond genus. Scaphopods in the museum collections at GRCA from the Toroweap are labeled as *Dentalium* and show longitudinal ridges (costate). Scaphopods are known informally as tusk shells due to their elongate, conical shells, which have openings at both ends. Scaphopods from the museum collections are fragments about 1 to 2 cm (0.4 to 0.8 in) in length and a few millimeters in diameter.

Kaibab Formation Fauna

The Kaibab Formation (Pk), also known as the Kaibab Limestone, covers more than 1,057 km^2 (408 mi^2) or more than 21% of the surface of the GRCA park and provides evidence of Permian life from a vast, shallow sea. The Kaibab Formation has proven difficult to date precisely but appears to include both the late early Permian (McKee and Breed 1969) and early middle Permian (Thompson 1995; Blakey and Middleton 2012). This predates the Permian–Triassic boundary, when a major extinction event wiped out about 95% of marine life (Sepkoski 1997). Within GRCA, Kaibab Formation fossils document marine life before the extinction event. Invertebrate body fossils reported in the literature from the Kaibab Formation of GRCA include sponges, rugose corals, conulariids, bryozoans, brachiopods, bivalves, nautiloids, gastropods, scaphopods, trilobites, crinoids, and echinoids (McKee 1935, 1938; Griffin 1966; Cisne 1971).

The two members of the Kaibab Formation are the Fossil Mountain Member (formerly the gamma and beta units) and the younger Harrisburg Member (formerly alpha unit) (McKee 1938; Blakey and Knepp 1989). The Fossil Mountain Member is a cliff-forming unit of limestone, dolomite, and sandy limestone, with some fossiliferous chert nodules. The overlying Harrisburg Member is a slope-former with some ledges and is a mixed unit of limestone, dolomite, siltstone, sandstone, gypsum, and chert. There are differences in the faunal assemblages between the Fossil Mountain Member and the overlying Harrisburg Member. The Fossil Mountain Member includes sponges, rugose corals, conulariids, and echinoids, all of which are missing in the Harrisburg Member. The Fossil Mountain Member is more fossiliferous in western GRCA than eastern GRCA, corresponding to the predominance of limestone instead of dolomite (Beus and Billingsley 1989). The Harrisburg Member also has a gypsiferous facies with brachiopods in western GRCA, but non-gypsiferous facies are dominated by mollusks in eastern GRCA. The Harrisburg Member nautiloids have not been reported from the Fossil Mountain Member assemblages (McKee 1938; Miller and Unklesbay 1942; Miller and Youngquist 1949).

At least nine fossil invertebrates have been named from specimens found in the Kaibab Formation of GRCA, including four brachiopods, three bryozoans, a conulariid, and a trilobite (Appendix 5-C). In addition, five Kaibab brachiopod species (*Avonia dorsoconcava*, *Chonetes kaibabensis*, *Marginifera meridionalis*, *Productus bassi*, and *Productus paraindicus*; all now under different genera) have been named from specimens possibly found within GRCA and three sea urchins (*Archaeocidaris gracilis*,

A. longispinus, and *A. ornatus*) may have been named from the Toroweap or Kaibab formations of GRCA (Appendix 5-D).

Arthropoda

Arthropods are only represented by trilobites at GRCA in the Kaibab Formation. These proetids include the genera *Ameura*, *Ditomopyge*, *Paladin*, and the species *Delaria macclintocki*, *D. sevilloidia*, *D. snowi*, *Ditomopyge scitulus*, and *Novoameura mckeei* (McKee 1938; Cisne 1971). *Delaria macclintocki* was named from GRCA (Cisne 1971).

Brachiopoda

Brachiopods are diverse and abundant in the Kaibab Formation, with a number of taxa represented at GRCA (Appendix Table 5-A-4 and 5-B-4). The most common brachiopods from the Kaibab Formation are the productids (Figure 5-22), which are abundant at many Permian localities. The common *Peniculauris bassi* was also known formerly as *Productus bassi* or *Dictyoclostus bassi*, and many GRCA museum specimens still carry the original name in the catalog. *Avonia subhorrida newberryi* (McKee 1938), *Chonetes quadratus* (King 1931), *Composita arizonica* (McKee 1938), and *Derbyia arizonensis* (McKee 1941) are brachiopods named from GRCA's Kaibab Formation (Appendix 5-C).



Figure 5-22. Kaibab Limestone productids from GRCA (LINDA S. LASSITER). **A.** *Peniculauris bassi* (GRCA 21372). **B.** A drawer full of productid brachiopods (GRCA 11055). Scale bar 1 cm (0.4 in).

The diverse brachiopod orders of the Kaibab Formation include Athyridida (McKee 1938), Rhynchonellida (McKee 1938; Sorauf and Billingsley 1991), Spiriferida (McKee 1938; Sorauf and Billingsley 1991), Spiriferinida (McKee 1938), Terebratulida (McKee 1938), Orthida (Condra and Elias 1944; McKee 1938), Orthotetida (McKee 1938; Sorauf and Billingsley 1991), and Productida (McKee 1938; Sorauf and Billingsley 1991; Thayer 2009). GRCA museum collections have hundreds of Kaibab Formation productids of the genera *Chonetes*, "*Productus*", and *Rugatia* and the equally common spiriferid genera *Composita* and *Squamularia.* The once-dominant strophomenid orthotetid *Derbyia* (*D. arizonensis*, *D. multistriata*, *D. nasuta*) is still common but not present in the quantities seen in the earlier stratigraphic units of the Carboniferous. In a section described by McKee (1938) along the Hermit Trail, assemblages of *Avonia*, *Chonetes*, "*Productus*", *Rugatia*, *Composita*, *Squamularia*, and *Derbyia* are visible with bivalves, crinoids, and bryozoans.

Bryozoa

Three bryozoan species, *Bicorbis arizonica*, originally *Bicorbula* (Condra and Elias 1945a), *Bascomella subsphaerica* (Condra and Elias 1944), and *Girtypora maculata* (McKinney 1983), have been named from the GRCA Kaibab Formation (Appendix 5-C). Fenestrate bryozoans (Figure 5-23) with characteristic fan shapes and other forms can be seen in many GRCA Kaibab exposures but are only identifiable to species by destructive analysis of the zooecia. Identified fenestrate bryozoans from the Kaibab include the genera *Bicorbis*, *Chasmatopora*, *Fenestella*, and *Septopora*. The most common genus is *Septopora*, which has oval fenestrules with three to four zooecia per row on the obverse side. The reverse side is smooth with some pores. *Fistulipora* are distinctive in that the zooecium has a raised, horse-shoe shaped area on one side of the aperture called a lunarium. Another common cystoporidan of the Kaibab Formation is *Meekopora*, which lacks the lunarium, although discerning these features is difficult without preparation and magnification of the specimen.



Figure 5-23. An unidentified fenestrate bryozoan from the Kaibab Formation. Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

Several genera of cryptostome (Cryptostomida) bryozoans have been reported from the Kaibab Formation of GRCA. *Rhombopora* forms cylindrical zooaria that may or may not be branched and have acanthopore ridges in polygonal raised areas around the zooecium. Other cryptostomid bryozoans include *Girtypora*, *Rhabdomeson*, and *Streblotrypa* (McKinney 1983).

Trepostome bryozoans form ramose, or erect, branching zooaria. The most common trepostome is *Stenopora*, but there are also specimens from the genera *Stenodiscus* and *Bascomella*. The bryozoan *Bascomella subsphaerica* (Condra and Elias 1944) is named from GRCA (Appendix 5-C).

Cnidaria

The only true coral reported from the Kaibab Formation is the solitary rugosan genus *Lophophyllum* (McKee 1938; Sorauf and Billingsley 1991).

Conulariids have been reported from the Kaibab Formation. McKee (1935) named the Kaibab species *Conularia kaibabensis* (Figure 5-24) for a GRCA specimen (Appendix 5-C). This conulariid is also sometimes referred to as *Paraconularia kaibabensis* (Sinclair 1948; McKee and Breed 1969, in which it is described as a coral, perhaps because of the then-recent placement of conulariids within Cnidaria). The specimen was first thought to be a fish, as described in Spamer (1984).



Figure 5-24. Conulariid *Conularia kaibabensis* holotype from the Kaibab Formation (USNM 102289). Scale bar 1 cm (0.4 in) (NPS/ANNE MILLER).

Echinodermata

Echinoids, or sea urchins, are often only preserved by spines that are about 2 cm (0.8 in) long (McKee 1938; Sorauf and Billingsley 1991) and at GRCA are found primarily in the Kaibab Formation. Crinoid columnals are found in many Kaibab Formation rocks along the rim trail (Figure 5-25). Multiple specimens identified as the sea urchin genera *Archaeocidaris* (superseded name "*Echinocrinus*" of some references) are reported from the Kaibab Formation. Three species of echinoids potentially from the Kaibab Formation of GRCA were described in Newberry (1861) as *Archaeocidaris gracilis*, *A. longispinus*, and *A. ornatus*. An additional undescribed specimen (GRCA 20263) may also be *Archaeocidaris*.



Figure 5-25. Crinoid stems and individual columnals found in the Kaibab Formation (GRCA 11468). Ruler includes mm tick marks and cm boxes (LINDA S. LASSITER).

Mollusca

Most bivalve specimens from GRCA are from the Permian and some became quite large (Figure 5-26). This is a typical population distribution as bivalves grew in diversity and numbers during the late Paleozoic (Clapham and Bottjer 2007). The trigoniid bivalve *Schizodus* and the myalinid *Aviculopecten* are common in the Kaibab Formation, with about 170 specimens of the former in GRCA collections. *Aviculopecten* (Figure 5-27) can be distinguished by the coarse costae which *Schizodus* lacks. Another myalinid from the Kaibab Formation at GRCA is the genus *Myalina.* Along the Hermit Trail, fossil assemblages often include representatives of the bivalves *Schizodus*, *Aviculopecten*, and *Solemya* (=obsolete *Solenomya*) (McKee 1938).

The Kaibab bivalve community consists of scallop-like pectinids of the genera *Acanthopecten coloradoensis*, *Deltopecten* (*D. caneyanus*), *Pecten*, *Pseudomonotis*, and *Pterinopecten.* Other bivalves are the nuculanids *Nucula* (*N. levatiforme*), *Nuculana*, *Yoldiella lucida*, the pholadomyid *Allorisma* (*A. capax*), the ostreids *Bakewellia parva* and *Pteria*, and the modiomorphid *Modiomorpha* (Shimer 1919; McKee 1938). *Nuculana* valves have an asymmetrical, swept appearance and are easily recognized. The cardiid bivalves are *Astartella* (*A. gurleyi*, *A. subquadrata*), *Permophorus* (*P. albequus*), and *Wilkingia wyomingensis* (Shimer 1919; McKee 1938).

There are gastropods from GRCA of the orders Bellerophontida, Cycloneritida, Euomphalina, Murchisoniina, and Pleurotomariida (Shimer 1919; McKee 1938). The most common gastropods of the GRCA museum collections from the Kaibab Formation are bellerophontids (*Bellerophon*, *Bucanopsis*, *Euphemites*), but other taxa are also present (Appendix Table 5-A-4 and 5-B-4).

Specimens are better preserved in the Kaibab Formation but may still be difficult to identify as they are often preserved as steinkerns which preserve no external details.



Figure 5-26. A large bivalve of the Kaibab Formation, *Allorisma capax* (GRCA 13501). U.S. quarter for scale (diameter 24.26 mm or 0.955 in) (LINDA S. LASSITER).



Figure 5-27. A common myalinid bivalve, *Aviculopecten* (GRCA 3621), from the Kaibab Formation. Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

Scaphopods also occur in the Kaibab Formation of GRCA. Shimer (1919) and McKee (1938) reported *Prodentalium canna* (Figure 5-28). *P. canna* is distinguished by a straight shape to the shell and ribbing which is not obvious, but concentric growth rings may be. Permian scaphopods are also

sometimes referred to under the names *Dentalium* and *Plagioglypta*, and many specimens are still labeled under either name at the GRCA museum. Scaphopods from the Kaibab Formation are small fragments often no wider than a millimeter and no longer than 1 cm (0.4 in).



Figure 5-28. A specimen of the scaphopod *Prodentalium canna* (GRCA 14373) showing the conical shape and small size. It was collected from the Kaibab Limestone outside of the park boundaries. U.S. quarter for scale (diameter 24.26 mm or 0.955 in) (LINDA S. LASSITER).

Coiled Permian cephalopods are rare at GRCA and the lack of ammonoids was noted by McKee (1938). For the Kaibab Formation, there are reports and museum records of the genera *Domatoceras*, *Metacoceras*, and *Stearoceras* (formerly *Titanoceras*) (Miller and Unklesbay 1942; Miller and Youngquist 1949). The orthoconic nautiloid *Orthoceras* was reported from the Kaibab Formation in McKee (1938), but this is a wastebasket identification for a longiconic form. Several specimens of possible nautiloids are also mentioned (McKee 1938; Miller and Unklesbay 1942; Miller and Youngquist 1949).

Porifera

One sponge species is identified from the Kaibab Formation of GRCA, the lithistid demosponge *Actinocoelia maeandrina* (GRCA 3592) (Griffin 1966; Thayer 2009). This sponge is frequently preserved in silica nodules and is relatively common. It is considered an index fossil for age-comparable stratigraphic units to the Kaibab Formation (Finks et al. 1961). Many other Kaibab sponge specimens at the GRCA museum have not been assigned to a genus (Figure 5-29). Six demosponges at MNA are from the Kaibab Formation at GRCA, one of which has been attributed to *Actinocoelia* (MNA 7248), the other five only identified to phylum.



Figure 5-29. An unidentified sponge from the Kaibab Formation (GRCA 3590). Scale bar 1 cm (0.4 in) (LINDA S. LASSITER).

Summary of the Paleozoic Invertebrate Body Fossils

More than 6,500 specimens were reviewed from catalog listings from museum collections (GRCA, MNA, USNM), and another 592 were identified from the literature. Only 269 species from GRCA are documented as being from a known and verified species. The Grand Canyon Museum (GRCA) has more than 5,800 specimens identified as invertebrate body fossils from the Paleozoic. More than 5,400 of those specimens were collected before 1980 and some of the localities may now be from outside of the current park boundaries. The Smithsonian (USNM) paleobiology collections have more than 500 specimens from the Grand Canyon and the Museum of Northern Arizona (MNA) has more than 240 specimens. Although a determined effort was made to check locations against the current park boundaries, there is much more work to be done to refine and correct the summary of invertebrate body fossils from GRCA. The numbers of specimens per species or entry from literature reports only documents that the kind of organism was reported. Within the text of the literature are more accurate numbers of specimens, in most cases. There are more specimens, more museums, and probably more records that need review and incorporation into this survey. The GRCA compilation of Paleozoic invertebrate body fossils and documentation of this report is only a starting point.

Some discrepancies in these findings require further investigation. For example, the record of archaeocyathid sponges in the Tonto Group, if confirmed, could indicate a refugia of this group otherwise assumed to have gone extinct during the Sinsk extinctions (Zhuravlev and Wood 2018). There are enigmatic fossils from the Bright Angel Shale that may be resolved by comparison with records from Cambrian lagerstätten localities outside of GRCA. The GRCA fossil record is remarkably limited for cephalopods, ostracods, and rostroconchs. The limited occurrence of key bryozoans *Archimedes* and *Polypora* is odd. There are other puzzles that may be solved using the GRCA resources of exposed Paleozoic units and specimens in the collections; these are but a few.

Conclusions

The Grand Canyon records life on our planet immediately before or after several of Earth's major extinction events. Globally, the Sinsk extinction event of the Stage 4 early Cambrian (513 Ma) (Zhuravlev and Wood 2018) preceded the rocks of the Tonto Group (510–505 Ma). The depositional

record of marine fossils at GRCA resumes only after the Ordovician radiation and during the final convergence of the paleocontinents of Laurentia, Baltica, and Avalonia into Laurussia. The GRCA Devonian fossil record during the Frasnian stage immediately precedes the Hangenberg extinction event of the Frasnian–Famennian (372 Ma) (Caplan and Bustin 1999). The major extinctions of the middle and end-Permian took place after the deposition of the Kaibab Formation. The invertebrate body fossils reported here are only those collated from three museums and the literature. Admittedly, there are other fossils in inventories at other museums and private collections that are not yet identified. There are many undiscovered clues about how life evolved and changed in the marine environments during the Paleozoic as captured in these rocks. As more work is done, the final story will be epic, just like the views from the top, and even from space.

Acknowledgments

The compilation of this report required the help of several curators including Kim Besom and Colleen Hyde of the Grand Canyon National Park Museum who provided access to the extensive collections of Paleozoic invertebrate body fossils. Kim was instrumental in building a basic starting point of the inventory by pulling records of all specimens that could be classified as invertebrate body fossils (more than 8,000 records). Additionally, Janet Gillette, Museum of Northern Arizona, provided a current listing of GRCA sourced specimens at MNA, even during a time when the museum faced the threat of a nearby forest fire (aptly named the Museum Fire). Anne Miller provided input for the project including photographs and information from the Smithsonian Institute on invertebrate body fossil holotypes. Diana Boudreau and Klara Widrig, Geoscientists in the Park, assisted with a short survey of common fossils exposed on the South Rim properties and identified several potential survey sites for future studies. Ted McClure assisted with obtaining several needed publications and references. Mark Nebel provided essential information to estimate geologic provenance and areas within the GRCA park boundaries. Anne Miller, Diana Boudreau, and Vincent Santucci also assisted with the early revisions and drafting of this chapter. Two anonymous reviewers also provided very helpful final edit recommendations and comments.

Literature Cited

Babcock, L. E., and R. A. Robison. 1988. Taxonomy and paleobiology of some middle Cambrian *Scenella* (Cnidaria) and hyolithids (Mollusca) from western North America. Paleontological Contributions, The University of Kansas 121.

Bengtson, S., and D. Collins. 2015. Chancelloriids of the Cambrian Burgess Shale. Palaeontologia Electronica 18.1.6A:1–67. Available at: https://palaeo-electronica.org/content/2015/1031-chancelloriids (accessed January 2020).

Beus, S. S. 1987. Geology along the South Kaibab Trail, eastern Grand Canyon, Arizona. Pages 371–378 *in* S. S. Beus, editor. Rocky Mountain section of the Geological Society of America. Geological Society of America, Boulder, Colorado. Centennial field guide 2.

Beus, S. S. 1989. Devonian and Mississippian geology of Arizona. Pages 287–311 *in* J. P. Jenney and S. J. Reynolds, editors. Geologic evolution of Arizona. Arizona Geological Society Digest 17.

Beus, S. S. 1990. Temple Butte Formation. Pages 107–117 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology. Oxford University Press, New York, New York.

Beus, S. S. 1999. Megafossil paleontology. Pages 69–96 *in* G. H. Billingsley and S. S. Beus, editors. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

Beus, S. S. 2003. Redwall Limestone and Surprise Canyon Formation. Pages 115–135 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Beus, S. S., and G. H. Billingsley. 1989. Paleozoic strata of the Grand Canyon, Arizona. Pages 122–127 *in* D. P. Elston, G. H. Billingsley, and R. A. Young, editors. Geology of Grand Canyon, northern Arizona (with Colorado River guides): Lees Ferry to Pierce Ferry, Arizona.

Billingsley, G. H., and S. S. Beus. 1985. The Surprise Canyon Formation, an Upper Mississippian and Lower Pennsylvanian (?) rock unit in the Grand Canyon, Arizona. U.S. Geological Survey, Washington, D.C. Bulletin 1605:A27–A33.

Billingsley, G. H., and S. S. Beus, editors. 1999a. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

Billingsley, G. H., and S. S. Beus. 1999b. Overview of the Surprise Canyon Formation. Pages 1–8 *in* G. H. Billingsley and S. S. Beus, editors. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

Billingsley, G. H., and W. J. Breed. 1976. Mississippian nautiloids of the Grand Canyon. Plateau 48:67–70.

Billingsley, H., and E. D. McKee. 1982. Pre-Supai buried valleys. Pages 137–153 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

Blakey, R. C. 1990. Stratigraphy and geologic history of Pennsylvanian and Permian rocks, Mogollon Rim region, central Arizona and vicinity. Geological Society of America Bulletin 102:1189–1217.

Blakey, R. C. 2003. Supai Group and Hermit Formation. Pages 136–162 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Blakey, R. C., and R. Knepp. 1989. Pennsylvanian and Permian geology of Arizona. Pages 313–347 *in* J. P. Jenney and S. J. Reynolds, editors. Geologic evolution of Arizona. Arizona Geological Society Digest 17.

Blakey, R. C., and L. T. Middleton. 2012. Geologic history and paleogeography of Paleozoic and early Mesozoic sedimentary rocks, eastern Grand Canyon, Arizona. Pages 81–92 *in* J. M. Timmons and K. E. Karlstrom, editors. Grand Canyon geology; two billion years of Earth's history. Geological Society of America, Boulder, Colorado. Special Paper 489.

Bond, D., and P. B. Wignall. 2005. Chapter 9. Evidence for Late Devonian (Kellwasser) anoxic events in the Great Basin, western United States. Developments in Palaeontology and Stratigraphy 20:225–262.

Bonde, A., V. L. Santucci, J. S. Tweet, E. Eichenberg, and B. Moore. 2018. Lake Mead National Recreation Area: Paleontological resources inventory. Natural Resource Report NPS/LAKE/NRR—2018/1618. National Park Service, Fort Collins, Colorado.

Breed, W. J. 1969. The discovery of orthocone nautiloids in the Redwall Limestone—Marble Canyon, Arizona. Page 134 *in* D. L. Bass, editor. Geology and natural history of the Grand Canyon region. Four Corners Geological Society, Flagstaff, Arizona. Field Conference 5.

Brezinski, D. K. 2017. Trilobites from the Redwall Limestone (Mississippian) of Arizona. Annals of Carnegie Museum 84:165–172.

Burckle, L. H. 1960. Some Mississippian fenestrate Bryozoa from central Utah. Journal of Paleontology 34:1077–1098.

Caplan, M. L., and R. M. Bustin. 1999. Devonian–Carboniferous Hangenberg mass extinction event, widespread organic-rich mudrock and anoxia: Causes and consequences. Palaeogeography, Palaeoclimatology, Palaeoecology 148:187–207.

Carpenter, F. M. 1927. A fossil insect from the lower Permian of the Grand Canyon. Proceedings of the United States National Museum 71(23).

Carpenter, F. M. 1928. A new protodonatan from the Grand Canyon. Psyche 35:186–189.

Carter, J. L., D. K. Brezinski, A. D. Kollar, and J. T. Dutro, Jr. 2014. Brachiopoda taxonomy and biostratigraphy of the Redwall Limestone (Lower Mississippian) of Arizona. Annals of Carnegie Museum 82(3):257–290.

Cisne, J. L. 1967. Two new Mississippian trilobites of the genus *Paladin*. Journal of Paleontology 41(5):1267–1273.

Cisne, J. L. 1971. Paleoecology of trilobites of the Kaibab limestone (Permian) in Arizona, Utah, and Nevada. Journal of Paleontology 45(3):525–533.

Clapham, M. E., and D. J. Bottjer. 2007. Permian marine paleoecology and its implications for large-scale decoupling of brachiopod and bivalve abundance and diversity during the Lopingian (Late Permian). Palaeogeography, Palaeoclimatology, Palaeoecology 249:283–301.

Condra, G. E., and M. K. Elias. 1944. Carboniferous and Permian ctenostomatous Bryozoa. Geological Society of America Bulletin 55:517–568.

Condra, G. E., and M. K. Elias. 1945a. *Bicorbula*, a new Permian bryozoan, probably a bryozoan-algal consortium. Journal of Paleontology 19(2):116–125.

Condra, G. E., and M. K. Elias. 1945b. *Bicorbis arizonica* Condra and Elias, new name for *B. arizonica*. Journal of Paleontology 19:411.

Congreve, C. R., A. Z. Krug, and M. E. Patzkowsky. 2015. Phylogenetic revision of the Strophomenida, a diverse and ecologically important Palaeozoic brachiopod order. Palaeontology 58:743–758.

Cooper, G. A., and R. E. Grant. 1975. Permian brachiopods of west Texas, III. Smithsonian Contributions to Paleobiology 19.

Duncan, H. 1969. Bryozoans. Pages 345–433 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Easton, W. H., and R. C. Gutschick. 1953. Corals from the Redwall Limestone (Mississippian) of Arizona. Bulletin, Southern California Academy of Sciences 52(1):1–27.

Elliott, D. K., and D. L. Martin. 1987. *Chancelloria*, an enigmatic fossil from the Bright Angel Shale (Cambrian) of Grand Canyon, Arizona. Journal of the Arizona-Nevada Academy of Science 21(2):67–72.

Elston, D. P., G. H. Billingsley, and R. A. Young, editors. 1989. Geology of Grand Canyon, northern Arizona (with Colorado River guides): Lees Ferry to Pierce Ferry, Arizona. Hanshaw, P. M. Field trips for the 28th international geological congress. American Geophysical Union, Washington, D.C.

Finks, R. M., E. L. Yochelson, and R. P. Sheldon. 1961. Stratigraphic implications of a Permian Sponge occurrence in the Park City Formation of Western Wyoming. Journal of Paleontology 35:564–568.

Foster, J. R. 2011. Trilobites and other fauna from two quarries in the Bright Angel Shale (middle Cambrian, Series 3; Delamaran), Grand Canyon National Park, Arizona. Pages 99–120 *in* J. S. Hollingsworth, F. A. Sundberg, and J. R. Foster, editors. Cambrian stratigraphy and paleontology of northern Arizona and southern Nevada. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 67.

Frech, F. 1893. Section in Congress Canyon opposite Point Sublime. Pages 476–481 *in* Compte rendu. International Geologic Congress [Congres geologique international], 5th session. Government Printing Office, Washington, D.C.

Gordon, M., Jr. 1973/01/03. OER-72-5. USGS internal memo to E. D. McKee (E&R file).

Gordon, M., Jr. 1979/01/23. O&G-78-10. USGS internal memo to E. D. McKee (E&R file).

Gordon, M., Jr. 1982. Biostratigraphy of the Watahomigi Formation. Pages 113–135 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

Grant, R. E. 1963/10/16. O-60-55. USGS internal memo to M. B. Ingham (E&R file).

Griffin, L. R. 1966. *Actinocoelia maeandrina* Finks, from the Kaibab Limestone of northern Arizona. Brigham Young University Geology Studies 13:105–108.

Grinnell, R. S., and G. W. Andrews. 1964. Morphologic studies of the brachiopod genus *Composita*. Journal of Paleontology 38:227–248.

Hayward, B. W., F. Le Coze, D. Vachard, and O. Gross. 2019. World Foraminifera Database. Accessed at http://www.marinespecies.org/foraminifera on 2020-01-03. doi:10.14284/305

Hodnett, J. P. M., and D. K. Elliott. 2018. Carboniferous chondrichthyan assemblages from the Surprise Canyon and Watahomigi formations (latest Mississippian–Early Pennsylvanian) of the western Grand Canyon, northern Arizona. Journal of Paleontology 92, Memoir 77.

Hou, X., M. Williams, D. J. Siveter, D. J. Siveter, R. J. Aldridge, and R. S. Sansom. 2010. Soft-part anatomy of the Early Cambrian bivalved arthropods *Kunyangella* and *Kunmingella*: Significance for the phylogenetic relationships of Bradoriida. Proceedings of the Royal Society B: Biological Sciences 277:1835–1841.

Karlstrom, K., J. Hagadorn, G. Gehrels, W. Matthews, M. Schmitz, L. Madronich, J. Mulder, M. Pecha, D. Giesler, and L. Crossey. 2018. Cambrian Sauk transgression in the Grand Canyon region redefined by detrital zircons. Nature Geoscience 11:438–443.

Karlstrom, K. E., M. T. Mohr, M. D. Schmitz, F. A. Sundberg, S. M. Rowland, R. Blakey, J. R. Foster, L. J. Crossey, C. M. Dehler, and J. W. Hagadorn. 2020. Redefining the Tonto Group of Grand Canyon and recalibrating the Cambrian timescale. Geology in press. doi: https://doi.org/10.1130/G46755.1.

King, R. E. 1931. The geology of the Glass Mountains. Part II, Faunal summary and correlation of the Permian formations with description of Brachiopoda. Texas University Bulletin 3042.

Kirk, E. 1945. *Eocrinus* (*E. multibrachiatus*). Pages 185–187 *in* E. D. McKee and C. E. Resser. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

Kouchinsky, A., S. Bengtson, B. Runnegar, C. Skovsted, M. Steiner, and M. Vendrasco. 2012. Chronology of early Cambrian biomineralization. Geological Magazine 149:221–251.

Lamsdell, J. C., S. J. Braddy, and O. E. Tetlie. 2010. The systematics and phylogeny of the Stylonurina (Arthropoda: Chelicerata: Eurypterida). Journal of Systematic Palaeontology 8:49–61.

Lee, J.-H., J. Hong, S.-J. Choh, D.-J. Lee, J. Woo, and R. Riding. 2016. Early recovery of sponge framework reefs after Cambrian archaeocyath extinction: Zhangxia Formation (early Cambrian Series 3), Shandong, North China. Palaeogeography, Palaeoclimatology, Palaeoecology 457:269–276.

Legg, D. A., M. D. Sutton, and G. D. Edgecombe. 2013. Arthropod fossil data increase congruence of morphological and molecular phylogenies. Nature Communications 4:2485.

Macurda, D. B., Jr. 1969. Blastoids. Pages 457–473 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Marques, A. C., and A. G. Collins. 2004. Cladistic analysis of Medusozoa and cnidarian evolution. Invertebrate Biology 123:23–42.

Martin, H., and J. E. Barrick. 1999. Conodont biostratigraphy. Pages 97–116 *in* G. H. Billingsley and S. S. Beus, editors. 1999. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

Mathis, A., and C. Bowman. 2007. Telling time at Grand Canyon National Park. Park Science 24(2):78–83.

McKee, E. D. 1935. A Conularia from the Permian of Arizona. Journal of Paleontology 9:427–429.

McKee, E. D. 1938. The environment and history of the Toroweap and Kaibab formations of northern Arizona and southern Utah. Carnegie Institution of Washington Publication 492.

McKee, E. D. 1941. *Derbya arizonensis*, new name for *D. regularis* McKee. Journal of Paleontology 15(1):91.

McKee, E. D. 1945. Stratigraphy and ecology of the Grand Canyon Cambrian. Pages 5–168 *in* E. D. McKee and C. E. Resser. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

McKee, E. D. 1963. Nomenclature for lithologic subdivisions of the Mississippian Redwall Limestone, Arizona: Article 65. Pages C21-C22 *in* Geological Survey Research 1963. U.S. Geological Survey, Washington, D.C. Professional Paper 475-C.

McKee, E. D. 1969. Paleozoic rocks of Grand Canyon. Pages 78–90 *in* D. L. Baars, F. G. Bowman, Jr., and C. W. Spencer, editors. Geology and natural history of the Grand Canyon region. Four Corners Geological Society, Durango, Colorado. Field conference guidebook 5.

McKee, E. D. 1975. The Supai Group: Subdivision and nomenclature. U.S. Geological Survey, Washington, D.C. Bulletin 1395-J.

McKee, E. D. 1982a. Distribution and age of flora and fauna. Pages 75–112 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D. 1982b. Measured stratigraphic sections. Pages 389–496 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D. 1982c. Environment of deposition of sandstone bodies. Pages 245–262 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C.

McKee, E. D., and W. J. Breed. 1969. The Toroweap Formation and Kaibab Limestone. Pages 12–26 *in* W. K. Summers and F. E. Kottlowski, editors. The San Andres Limestone, a reservoir for oil and water in New Mexico. New Mexico Geological Survey, Socorro, New Mexico. Special Publication 3.

McKee, E. D., and R. C. Gutschick. 1969a. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969b. Sequence of sediments and unconformities. Pages 13–95 in E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969c. Interpretation of environments. Pages 553–570 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969d. The Redwall faunas. Pages 125–172 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969e. Appendix: Descriptions of stratigraphic sections of the Redwall Limestone. Pages 613–692 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969f. Miscellaneous fossil groups - algae and stromatolites, holothurians, trilobites, ostracodes, and fish. Pages 545–552 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and C. E. Resser. 1945. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

McKinney, F. K. 1983. Ectoprocta (Bryozoa) from the Permian Kaibab Formation, Grand Canyon National Park, Arizona. Fieldiana: Geology (New Series) 13(1341).

Miller, A. K., and A. G. Unklesbay. 1942. Permian nautiloids from western United States. Journal of Paleontology 16(6):719–738.

Miller, A. K., and W. Youngquist. 1949. American Permian nautiloids. Geological Society of America, New York, New York. Memoir 41.

Miller, T. G. 1963. The bryozoan genus *Polypora* M'Coy. Palaeontology 6:166–171.

Moysiuk, J., M. R. Smith, and J.-B. Caron. 2017. Hyoliths are Palaeozoic lophophorates. Nature 541:394–397.

Nanglu, K., J.-B. Caron, S. Conway Morris, and C. B. Cameron. 2016. Cambrian suspension-feeding tubicolous hemichordates. BMC Biology 14.

Newberry, J. S. 1861. Geological report. *In* Ives, J. C., Report upon the Colorado River of the West, explored in 1857 and 1858. U.S. 36th Congress, 1st Session, House Executive Document 90, Serial 1058, Part 3.

Newell, N. D. 1937. Late Paleozoic pelecypods: Pectinacea. Kansas Geological Survey, Lawrence, Kansas. Bulletin 10(1).

Noble, L. F. 1914. The Shinumo quadrangle, Grand Canyon district, Arizona. U.S. Geological Survey, Washington, D.C. Bulletin 549.

Noble, L. F. 1922. A section of the Paleozoic formations of the Grand Canyon at Bass Trail. Pages 23–73 *in* Shorter contributions to general geology, 1922. U.S. Geological Survey, Washington, D.C. Professional Paper 131-B.

Palmer, A. R. 1963. O-60-55. USGS internal memo to M. B. Ingham (E&R file).

Raup, D. M., and J. J. Sepkoski. 1982. Mass extinctions in the marine fossil record. Science 215:1501–1503.

Rasetti, F. 1963. Middle Cambrian ptychoparioid trilobites from the conglomerates of Quebec. Journal of Paleontology 37:575–594.

Rawson, R. R., and C. E. Turner. 1974. The Toroweap Formation: A new look. Pages 155–190 *in* T. N. V. Karlstrom, G. A. Swann, and R. L. Eastwood, editors. Geology of northern Arizona, with notes on archaeology and paleoclimate. Geological Society of America, Rocky Mountain Section, Flagstaff, Arizona. Guidebook 27, Part 1, regional studies.

Resser, C. E. 1935. Nomenclature of some Cambrian trilobites. Smithsonian Miscellaneous Collections 93(5):1–46.

Resser, C. E. 1945. Cambrian fossils of the Grand Canyon. Pages 169–220 *in* E. D. McKee and C. E. Resser. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

Robison, R. A. 1965. Middle Cambrian eocrinoids from western North America. Journal of Paleontology 39:355–364.

Rose, E. C. 2006. Nonmarine aspects of the Cambrian Tonto Group of the Grand Canyon, USA, and broader implications. Palaeoworld 15:223–241.

Rowland, S. M., G. D. Osborn, and D. J. Graber. 1995. Lower Paleozoic stratigraphy of Fern Glen Canyon, central Grand Canyon, Arizona. Journal of the Arizona-Nevada Academy of Science 28(1-2):1–11.

Sando, W. J. 1963/10/14. O-60-55. USGS internal memo to M. B. Ingham (E&R file).

Sando, W. J. 1969. Corals. Pages 257–343 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Sando, W. J., and E. W. Bamber. 1985. Coral zonation of the Mississippian System in the Western Interior Province of North America. U.S. Geological Survey, Washington, D.C. Professional Paper 1334.

Schenk, E. T., and H. E. Wheeler. 1942. Cambrian sequence in western Grand Canyon, Arizona. Journal of Geology 50(7):822–899.

Schuchert, C. 1918a. The Cambrian of the Grand Canyon of Arizona. American Journal of Science (4^{th} series) 45(269):362–369.

Schuchert, C. 1918b. On the Carboniferous of the Grand Canyon of Arizona. American Journal of Science (4^{th} series) 45(269):347–361.

Sepkoski, J. J. 1997. Biodiversity: Past, Present, and Future. Journal of Paleontology 71:533–539.

Shimer, H. W. 1919. Permo-Triassic of northwestern Arizona. Geological Society of America Bulletin 30(4):471–498.

Skipp, B. 1969. Foraminifera. Pages 173–255 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Skipp, B., L. D. Halcomb, and R. C. Gutschick. 1966. Tournayellinae, calcareous foraminifera, in Mississippian rocks of North America; with translations from the original Russian of descriptions of several key genera and species by Ivan Mittin and Betty Skipp. Cushman Foundation for Foraminiferal Research Special Publication 9.

Sorauf, J. E., and G. H. Billingsley. 1991. Members of the Toroweap and Kaibab formations, Lower Permian, northern Arizona and southwestern Utah. The Mountain Geologist 28(1):9–24.

Spamer, E. E. 1984. Paleontology in the Grand Canyon of Arizona: 125 years of lessons and enigmas from the late Precambrian to the present. The Mosasaur 2:45-128.

Spamer, E. E. 1992. The Grand Canyon fossil record: A source book in paleontology of the Grand Canyon and vicinity, northwestern Arizona and southeastern Nevada. Geological Society of America, Boulder, Colorado. Microform Publication 24.

Sprinkle, J. 1973. Morphology and evolution of blastozoan echinoderms. Harvard University Museum of Comparative Zoology Special Publication.

Stoyanow, A. A. 1936. Correlation of Arizona Paleozoic formations. Geological Society of America Bulletin 37:459–540.

Strother, P. K., and J. H. Beck. 2000. Spore-like microfossils from Middle Cambrian strata: expanding the meaning of the term cryptospore. Pages 413–424 *in* M. M. Harley, C. M. Morton, and S. Blackmore, editors. Pollen and spores: morphology and biology. Royal Botanic Gardens, Kew, United Kingdom.

Sundberg, F. A., K. E. Karlstrom, G. Geyer, J. R. Foster, M. D. Schmitz, J. W. Hagadorn, M. T. Mohr, and L. J. Crossey. In review. A mid-Cambrian conundrum: Testing the chronology of a major faunal turnover. Geology.

Thayer, D. 2009. An introduction to Grand Canyon fossils. Grand Canyon Association, Grand Canyon, Arizona.

Thompson, K. L. 1995. Paleoecology and biostratigraphy of the Fossil Mountain Member, Kaibab Formation in northwestern Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Topper, T. P., C. B. Skovsted, G. A. Brock, and J. R. Paterson. 2011. The oldest bivalved arthropods from the early Cambrian of east Gondwana: Systematics, biostratigraphy and biogeography. Gondwana Research 19:310–326.

Turner, C. E. 1990. Toroweap Formation. Pages 203–223 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology. Oxford University Press, New York, New York.

Ulrich, E. O., and R. S. Bassler. 1931. Cambrian bivalved Crustacea of the order Conchostraca. U.S. National Museum Proceedings 78 (article 4).

Van Iten, H. V., J. M. Leme, M. G. Simões, A. C. Marques, and A. G. Collins. 2006. Reassessment of the phylogenetic position of conulariids (Ediacaran–Triassic) within the subphylum Medusozoa (Phylum Cnidaria). Journal of Systematic Palaeontology 4:109–118.

Walcott, C. D. 1883. Pre-Carboniferous strata in the Grand Cañon of the Colorado, Arizona. American Journal of Science (3rd series) 26:437–442.

Walcott, C. D. 1890. Description of a new genus and species of inarticulate brachiopod from the Trenton limestone. Proceedings of the United States National Museum 12:365–366.

Walcott, C. D. 1897. Cambrian Brachiopoda; genera *Iphidea* and *Yorkia*; with descriptions of new species of each, and of the genus Acrothele. U.S. National Museum Proceedings 19:707–718.

Walcott, C. D. 1898. Cambrian Brachiopoda: *Obolus* and *Lingulella*, with description of new species. U.S. National Museum Proceedings 21:385–420.

Walcott, C. D. 1902. Cambrian Brachiopoda: *Acrotreta*; *Linnarssonella*; *Obolus*; with descriptions of new species. U.S. National Museum Proceedings 25:577–612.

Walcott, C. D. 1905. Cambrian Brachiopoda; with descriptions of new genera and species. U.S. National Museum Proceedings 28:227–337.

Walcott, C. D. 1908. Cambrian geology and paleontology, No. 3: Cambrian Brachiopoda: Descriptions of new genera and species. Smithsonian Miscellaneous Collections 53:53–165.

Walcott, C. D. 1912a. Cambrian Brachiopoda. U.S. Geological Survey, Washington, D.C. Monograph 51.

Walcott, C. D. 1912b. Cambrian geology and paleontology II. Middle Cambrian Branchiopoda, Malacostraca, Trilobita, and Merostomata. Smithsonian Miscellaneous Collections 57(6):145–228.

Walcott, C. D. 1916a. Cambrian geology and paleontology III. Cambrian trilobites. Smithsonian Miscellaneous Collections 64(3):157–258.

Walcott, C. D. 1916b. Cambrian geology and paleontology II. Cambrian trilobites. Smithsonian Miscellaneous Collections 64(5):303–570.

Walcott, C. D. 1920. Cambrian geology and paleontology IV:6—Middle Cambrian Spongiae. Smithsonian Miscellaneous Collections 67:261–364.

Walcott, C. D. 1924. Cambrian geology and paleontology IV. No. 9. Cambrian and Ozarkian Brachiopoda; Ozarkian Cephalopoda and Notostraca. Smithsonian Miscellaneous Collections 67(9):477–554.

White, C. D. 1929. Flora of the Hermit shale, Grand Canyon, Arizona. Carnegie Institution of Washington Publication 405.

Williams, A., S. J. Carlson, H. C. Brunton, L. E. Holmer, and L. E. Popov. 1996. A supra-ordinal classification of the Brachiopoda. Proceedings of the Royal Society B: Biological Sciences 351:1171–1193.

Wright, D. F., W. I. Ausich, S. R. Cole, M. E. Peter, and E. C. Rhenberg. 2017. Phylogenetic taxonomy and classification of the Crinoidea (Echinodermata). Journal of Paleontology 91:829–846.

Yochelson, E. L. 1969. Gastropods and pelecypods. Pages 439–451 in E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Yochelson, E. L., and D. G. Cid. 1984. Reevaluation of the systematic position of *Scenella*. Lethaia 17:331–340.

Zhuravlev, A. Y., and R. A. Wood. 2018. The two phases of the Cambrian Explosion. Scientific Reports 8:1–10.

Appendix 5-A. Grand Canyon (GRCA) Paleozoic Invertebrate Specimens List

This appendix lists identified specimens of GRCA fossils from Paleozoic geologic units of GRCA in tables. The information was assembled from four sources: three museums and the literature. The three museums surveyed for this report are the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), and the Smithsonian Museum (USNM). Some entries listed in museum catalogs may be specimens reported from the literature, creating some duplication. While reviewing listings, species names were reviewed for consistency and current usage; the methods are listed in the following section on taxonomic uncertainty. Further work to resolve duplicates or underreported numbers of specimens is needed and recognized. To aid navigation and readability, the full list has been split into four shorter tables: Appendix Table 5-A-1, Cambrian; Appendix Table 5-A-2, Devonian and Mississippian; Appendix Table 5-A-3, Supai Group; and Appendix Table 5-A-4, post-Supai Permian. Additional electronic updates and details of this appendix will be published in a digital repository (DRYAD) at a later date.

Appendix Table 5-A-1. Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctu	Other	Arthropoda	Ostracoda	Leperditicopida	Undetermined	1
	Other	Arthropoda	Trilobita	Asaphida	*Glyphaspis kwaguntensis*	1
	Other	Arthropoda	Trilobita	Corynexochida	*Anoria* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Athabaskia kanabensis*	1
	Other	Arthropoda	Trilobita	Corynexochida	*Glossopleura boccar*	1
	Other	Arthropoda	Trilobita	Corynexochida	*Glossopleura* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Kootenia mckeei*	1
	Other	Arthropoda	Trilobita	Corynexochida	*Kootenia* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Olenoides* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Ptarmigania* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Elrathiella?* aff. *E. insueta*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Kochina angustata*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Parehmania kwaguntensis*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Parehmania nitida*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Proehmaniella hebes*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Ptychoparia* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Spencella diligens*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Spencella* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Trachycheilus typicale*	1
	Other	Brachiopoda	Kutorginata	Kutorginida	*Nisusia kanabensis*	1
	USNM	Brachiopoda	Kutorginata	Kutorginida	*Nisusia* sp.	1
	Other	Brachiopoda	Lingulata	Lingulida	*Acrothele* sp.	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella kanabensis*	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella* sp.	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella winona* aff. *L. winona* subsp. *convexa*	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctu (cont.)	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella zetus*	1
	USNM	Brachiopoda	Lingulata	Lingulida	*Lingulepis perattenuatus*	1
	Other	Brachiopoda	Paterinata	Paterinida	*Iphidea ornatella*	1
	USNM	Brachiopoda	Paterinata	Paterinida	*Iphidella pannula*	3
	Other	Brachiopoda	Paterinata	Paterinida	*Micromitra (Paterina) crenistria*	1
	Other	Brachiopoda	Rhynchonellata	Protorthida	*Protorthis* sp.	1
	Other	Problematica	Hyolitha	Hyolithida	*Hyolithes* sp.	1
Ctt	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Indianites* cf. *I. faba*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Indianites curta*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Indianites impressa*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella* sp.	1
	MNA	Arthropoda	Trilobita	Ptychopariida	*Spencella* sp.	1
	Other	Arthropoda	Trilobita	Redlichiida	*Olenellus* sp.	1
	USNM	Brachiopoda	Lingulata	Lingulida	*Lingulella chuarensis*	21
	USNM	Brachiopoda	Lingulata	Lingulida	*Lingulella euglypha*	2
	USNM	Brachiopoda	Lingulata	Lingulida	*Lingulella lineolatus*	5
	USNM	Brachiopoda	Lingulata	Lingulida	*Lingulella spatulus*	3
	USNM	Brachiopoda	Lingulata	Lingulida	*Lingulella zetus*	4
	Other	Brachiopoda	Obolellata	Obolellida	Undetermined	1
	USNM	Brachiopoda	Paterinata	Paterinida	*Iphidella pannula*	1
	USNM	Brachiopoda	Paterinata	Paterinida	*Paterina (Iphidea)? crenistria*	7
Ctba	Other	Arthropoda	Bradoriida clade	Bradoriida	*Bradoria tontoensis*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Bradoria tontoensis*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Indianites* aff. *I. faba?*	1
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Indianites curtus*	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctba (cont.)	Other	Arthropoda	Bradoriida clade	Bradoriida	*Indianites faba*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Indianites faba*	5
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Indianites faba* subsp. *intermedia*	1
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Indianites impressus*	1
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Indianites intermedius*	1
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella apicalis*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella apicalis*	5
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella breviuscula*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella breviuscula*	2
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella concentrica*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella concentrica*	3
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella leperditoides*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella leperditoides*	1
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella limatula*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella limatula*	3
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella longula*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella longula*	1
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella nitida*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella nitida*	1
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella oblonga*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella oblonga*	2
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella obsoleta*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella obsoleta*	1
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella pulchella*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella pulchella*	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctba (cont.)	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella scitula*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella scitula*	1
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella subtruncata*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella subtruncata*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella subtruncata*	2
	Other	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella ventrosa*	1
	USNM	Arthropoda	Bradoriida clade	Bradoriida	*Walcottella ventrosa*	1
	Other	Arthropoda	Bradoriida clade	Indianidae	*Dielymella appressa*	1
	USNM	Arthropoda	Bradoriida clade	Indianidae	*Dielymella appressa*	3
	Other	Arthropoda	Bradoriida clade	Indianidae	*Dielymella dorsalis*	1
	USNM	Arthropoda	Bradoriida clade	Indianidae	*Dielymella dorsalis*	1
	Other	Arthropoda	Bradoriida clade	Indianidae	*Dielymella nasuta*	1
	USNM	Arthropoda	Bradoriida clade	Indianidae	*Dielymella nasuta*	3
	Other	Arthropoda	Bradoriida clade	Indianidae	*Dielymella recticardinalis*	1
	USNM	Arthropoda	Bradoriida clade	Indianidae	*Dielymella recticardinalis*	5
	Other	Arthropoda	Bradoriida clade	Indianidae	*Dielymella recticardinalis* aff. *D. recticardinalis* subsp. *angustata*	1
	USNM	Arthropoda	Bradoriida clade	Indianidae	*Dielymella recticardinalis* aff. *D. recticardinalis* subsp. *angustata*	1
	GRCA	Arthropoda	Trilobita	Asaphida	*Glyphaspis* sp.	1
	Other	Arthropoda	Trilobita	Asaphida	*Glyphaspis* sp.	1
	Other	Arthropoda	Trilobita	Asaphida	*Glyphaspis vulsa*	1
	GRCA	Arthropoda	Trilobita	Composite	Composite of *Alokistocare* sp. and *Glossopleura mckeei*	1
	GRCA	Arthropoda	Trilobita	Composite	Composite of *Amecephalus althea* and *Glossoplerua producta*	3

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctba (cont.)	GRCA	Arthropoda	Trilobita	Composite	Composite of *Anoria tontoensis* and *Amecephalus althea*	2
	Other	Arthropoda	Trilobita	Corynexochida	*Albertella schenki*	1
	Other	Arthropoda	Trilobita	Corynexochida	*Albertella* sp.	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Anoria* sp.	29
	MNA	Arthropoda	Trilobita	Corynexochida	*Anoria* sp.	3
	Other	Arthropoda	Trilobita	Corynexochida	*Anoria* sp.	2
	GRCA	Arthropoda	Trilobita	Corynexochida	*Anoria tontoensis*	149
	MNA	Arthropoda	Trilobita	Corynexochida	*Anoria tontoensis*	2
	Other	Arthropoda	Trilobita	Corynexochida	*Anoria tontoensis*	1
	Other	Arthropoda	Trilobita	Corynexochida	*Athabaskia* sp.	2
	Other	Arthropoda	Trilobita	Corynexochida	*Glossopleura* aff. *"G. walcotti"*	1
	Other	Arthropoda	Trilobita	Corynexochida	*Glossopleura boccar*	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Glossopleura mckeei*	134
	MNA	Arthropoda	Trilobita	Corynexochida	*Glossopleura mckeei*	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Glossopleura meriwitica*	27
	GRCA	Arthropoda	Trilobita	Corynexochida	*Glossopleura productus*	48
	GRCA	Arthropoda	Trilobita	Corynexochida	*Glossopleura* sp.	55
	Other	Arthropoda	Trilobita	Corynexochida	*Glossopleura* sp.	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Glossopleura* sp. or *Anoria* sp.	24
	GRCA	Arthropoda	Trilobita	Corynexochida	*Glossopleura* sp. or *Clavaspidella* sp.	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Kootenia* aff. *"K. simplia"*	3
	GRCA	Arthropoda	Trilobita	Corynexochida	*Kootenia schenki*	2
	Other	Arthropoda	Trilobita	Corynexochida	*Kootenia simplex*	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Kootenia* sp.	13
	Other	Arthropoda	Trilobita	Corynexochida	*Kootenia* sp.	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctba (cont.)	Other	Arthropoda	Trilobita	Corynexochida	*Olenoides* sp.	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Prozacanthoides* sp.	2
	Other	Arthropoda	Trilobita	Corynexochida	*Ptarmigania* sp.	1
	GRCA	Arthropoda	Trilobita	Corynexochida	Undetermined	2
	Other	Arthropoda	Trilobita	Corynexochida	*Zacanthoides* cf. *Z. walapai*	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Zacanthoides* sp.	42
	Other	Arthropoda	Trilobita	Corynexochida	*Zacanthoides* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Acrocephalops?* cf. *A. arizonaensis*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Alokistocare lepida*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Alokistocare* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Alokistocare* sp. or *Ehmaniella* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Alokistocare?* sp.	1
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Amecephalus althea*	76
	MNA	Arthropoda	Trilobita	Ptychopariida	*Amecephalus althea*	6
	Other	Arthropoda	Trilobita	Ptychopariida	*Amecephalus althea*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Amecephalus* cf. *A. althea*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Amecephalus* cf. *A. packi*	1
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Amecephalus* sp.	86
	MNA	Arthropoda	Trilobita	Ptychopariida	*Amecephalus* sp.	1
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Antagmus arizonaensis*	2
	Other	Arthropoda	Trilobita	Ptychopariida	*Antagmus arizonaensis*	1
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Antagmus* sp.	3
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Bolaspis aemula*	6
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Ehmaniella* aff. "*E. arizonaensis*"	21
	Other	Arthropoda	Trilobita	Ptychopariida	*Ehmaniella* aff. "*E. arizonaensis*"	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctba (cont.)	Other	Arthropoda	Trilobita	Ptychopariida	*Ehmaniella* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Elrathia nitens*	1
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Elrathia* sp.	24
	Other	Arthropoda	Trilobita	Ptychopariida	*Elrathia* sp.	2
	MNA	Arthropoda	Trilobita	Ptychopariida	*Elrathiella?* aff. *E. insueta*	1
	MNA	Arthropoda	Trilobita	Ptychopariida	*Kochina* aff. "*K. adunca*"	1
	MNA	Arthropoda	Trilobita	Ptychopariida	*Pachyaspis* aff. *"P. arenosa"*	1
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Pachyaspis* cf. *P. fonticola*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Pachyaspis fonticola*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Pachyaspis moorei*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Pachyaspis* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Parehmania kwaguntensis*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Parehmania* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Parehmania tontoensis*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Proehmaniella basilica*	1
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Spencella porcata*	2
	MNA	Arthropoda	Trilobita	Ptychopariida	*Spencia tontoensis*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Spencia tontoensis*	1
	GRCA	Arthropoda	Trilobita	Redlichiida	*Olenellus* sp.	113
	Other	Arthropoda	Trilobita	Redlichiida	*Olenellus* sp.	1
	Other	Arthropoda	Trilobita	Redlichiida	Undetermined	1
	GRCA	Arthropoda	Trilobita	Undetermined	Undetermined	258
	MNA	Arthropoda	Trilobita	Undetermined	Undetermined	3
	Other	Arthropoda	Trilobita	Undetermined	Undetermined	1
	Other	Brachiopoda	Kutorginata	Kutorginida	*Nisusia* aff. *N. noblei*	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctba (cont.)	GRCA	Brachiopoda	Kutorginata	Kutorginida	*Nisusia* aff. *N. obscura*	1
	Other	Brachiopoda	Kutorginata	Kutorginida	*Nisusia* aff. *N. obscura*	1
	GRCA	Brachiopoda	Kutorginata	Kutorginida	*Nisusia noblei*	1
	GRCA	Brachiopoda	Kutorginata	Kutorginida	*Nisusia* sp.	22
	Other	Brachiopoda	Kutorginata	Kutorginida	*Nisusia* sp.	1
	Other	Brachiopoda	Kutorginata	Kutorginida	*Nisusia?* sp.	1
	GRCA	Brachiopoda	Lingulata	Lingulida	*Acrotreta* sp.	6
	GRCA	Brachiopoda	Lingulata	Lingulida	*Lingula* sp.	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingula* sp.	2
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella acutangula*	1
	MNA	Brachiopoda	Lingulata	Lingulida	*Lingulella* aff. *“L. noblei”*	1
	GRCA	Brachiopoda	Lingulata	Lingulida	*Lingulella chuarensis*	98
	MNA	Brachiopoda	Lingulata	Lingulida	*Lingulella chuarensis*	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella chuarensis*	1
	MNA	Brachiopoda	Lingulata	Lingulida	*Lingulella euglypha*	5
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella euglypha*	1
	MNA	Brachiopoda	Lingulata	Lingulida	*Lingulella lineolata*	5
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella lineolata*	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella mckeei*	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella* sp.	1
	GRCA	Brachiopoda	Lingulata	Lingulida	*Lingulella spatula*	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella spatula*	1
	MNA	Brachiopoda	Lingulata	Lingulida	*Lingulella spatula*	5
	MNA	Brachiopoda	Lingulata	Lingulida	*Lingulella themis*	2
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella zetus*	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctba (cont.)	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella?* aff. *O. themis*	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella?* cf. *L. monticula*	1
	GRCA	Brachiopoda	Lingulata	Lingulida	*Lingulella?* sp.	4
	MNA	Brachiopoda	Lingulata	Lingulida	*Lingulella?* sp.	2
	GRCA	Brachiopoda	Lingulata	Lingulida	*Obolus* sp.	21
	Other	Brachiopoda	Obolellata	Obolellida	*Obolella aff. O. polita*	1
	MNA	Brachiopoda	Obolellata	Obolellida	*Obolella* sp.	1
	Other	Brachiopoda	Obolellata	Obolellida	*Obolella* sp.	2
	Other	Brachiopoda	Paterinata	Paterinida	*Dictyonina arizonaensis*	1
	MNA	Brachiopoda	Paterinata	Paterinida	*Iphidella arizonaensis*	1
	GRCA	Brachiopoda	Paterinata	Paterinida	*Iphidella pannula* aff. *I. pannula* subsp. *aladensis*	2
	GRCA	Brachiopoda	Paterinata	Paterinida	*Iphidella* sp.	4
	Other	Brachiopoda	Paterinata	Paterinida	*Micromitra (Paterina) crenistria*	1
	Other	Brachiopoda	Paterinata	Paterinida	*Micromitra (Paterina) superba*	1
	Other	Brachiopoda	Paterinata	Paterinida	*Micromitra pealei*	1
	MNA	Brachiopoda	Paterinata	Paterinida	*Micromitra superba*	4
	GRCA	Brachiopoda	Paterinata	Paterinida	*Paterina?* sp.	3
	Other	Brachiopoda	Paterinata	Paterinida	*Paterina?* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Orthida	*Wimanella* sp.	4
	Other	Brachiopoda	Rhynchonellata	Thecideida	*Diraphora?* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Undetermined	Undetermined	3
	GRCA	Brachiopoda	Strophomenata	Billingsellida	*Billingsella* sp.	13
	GRCA	Brachiopoda	Undetermined	Undetermined	Undetermined	51
	MNA	Brachiopoda	Undetermined	Undetermined	Undetermined	5
	Other	Brachiopoda	Undetermined	Undetermined	Undetermined	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctba (cont.)	GRCA	Composite	Composite	Composite	Composite of *Billingsella highlandensis* and *Glossopleura* or *Anoria*	2
	GRCA	Composite	Composite	Composite	Composite of *Billingsella highlandensis* and *Glossopleura* sp.	2
	MNA	Composite	Composite	Composite	Composite of *Hyolithes* and *Anoria tontoensis* and *Amecephalus althea*	1
	GRCA	Echinodermata	Cystoidea?	Undetermined	Undetermined	1
	Other	Echinodermata	Eocrinoidea	Ascocystitida	*Eocystites* sp.	1
	GRCA	Echinodermata	Eocrinoidea	Gogiida	*Gogia longidactylus*	1
	Other	Echinodermata	Eocrinoidea	Gogiida	*Gogia multibrachiatus*	1
	USNM	Echinodermata	Eocrinoidea	Gogiida	*Gogia multibrachiatus*	1
	Other	Echinodermata	Eocrinoidea	Gogiida	*Gogia?* aff. *G. longidactylus*	1
	GRCA	Echinodermata	Eocrinoidea	Gogiida	Undetermined	1
	GRCA	Echinodermata	Not assigned	Rhombifera	Undetermined	1
	GRCA	Echinodermata	Undetermined	Undetermined	Undetermined	1
	GRCA	Mollusca	Bivalvia	Undetermined	Undetermined	4
	GRCA	Mollusca	Gastropoda	Undetermined	Undetermined	3
	GRCA	Porifera	Archaeocyatha	Ajacicyathida	*Ethmophyllum chankensis*	3
	Other	Porifera	Archaeocyatha	Undetermined	Undetermined	1
	GRCA	Porifera	Undetermined	Undetermined	Undetermined	2
	GRCA	Problematica	Coeloscleritophora	Chancelloriida	*Chancelloria* cf. *C. eros*	3
	MNA	Problematica	Coeloscleritophora	Chancelloriida	*Chancelloria* sp.	7
	Other	Problematica	Hyolitha	Hyolithida	*Hyolithes* sp.	3
	GRCA	Problematica	Hyolitha	Hyolithida	*Hyolithes?* sp.	38
	MNA	Problematica	Hyolitha	Hyolithida	*Hyolithes?* sp.	3
	Other	Problematica	Undetermined	Undetermined	*Margaretia?* sp.	2

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctba (cont.)	Other	Problematica	Undetermined	Undetermined	*Tontoia kwaguntensis*	1
	USNM	Problematica	Undetermined	Undetermined	*Tontoia kwaguntensis*	1
	GRCA	Undetermined	Undetermined	Undetermined	Undetermined	27
Ctm	GRCA	Arthropoda	Trilobita	Asaphida	*Glyphaspis kwaguntensis*	1
	Other	Arthropoda	Trilobita	Asaphida	*Glyphaspis kwaguntensis*	1
	GRCA	Arthropoda	Trilobita	Asaphida	*Glyphaspis* sp.	4
	Other	Arthropoda	Trilobita	Asaphida	*Glyphaspis* sp.	1
	GRCA	Arthropoda	Trilobita	Asaphida	*Glyphaspis tecta*	1
	Other	Arthropoda	Trilobita	Asaphida	*Glyphaspis tecta*	1
	GRCA	Arthropoda	Trilobita	Asaphida	*Glyphaspis vulsa*	1
	Other	Arthropoda	Trilobita	Asaphida	*Glyphaspis?* sp.	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Anoria* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Anoria* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Athabaskia* sp.	2
	Other	Arthropoda	Trilobita	Corynexochida	*Dorypyge* sp.	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Glossopleura meriwitica*	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Glossopleura* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Glossopleura* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Glossopleura* sp. or *Anoria* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Kootenia havasuensis*	1
	Other	Arthropoda	Trilobita	Corynexochida	*Kootenia mckeei*	1
	Other	Arthropoda	Trilobita	Corynexochida	*Kootenia schenki*	1
	Other	Arthropoda	Trilobita	Corynexochida	*Kootenia simplex*	1
	GRCA	Arthropoda	Trilobita	Corynexochida	*Kootenia* sp.	3
	Other	Arthropoda	Trilobita	Corynexochida	*Kootenia* sp.	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctm (cont.)	Other	Arthropoda	Trilobita	Corynexochida	*Kootenia?* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Neolenus* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Pagodia?* sp.	1
	Other	Arthropoda	Trilobita	Corynexochida	*Zacanthoides* sp.	1
	Other	Arthropoda	Trilobita	Proetida	*Bathyurus?* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Alokistocare* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Anomocarella* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Bolaspis aemula*	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Bolaspis*? sp.	1
	USNM	Arthropoda	Trilobita	Ptychopariida	*Elrathiella?* aff. *E. insueta*	3
	Other	Arthropoda	Trilobita	Ptychopariida	*Pachyaspis moorei*	1
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Parehmania kwaguntensis*	3
	Other	Arthropoda	Trilobita	Ptychopariida	*Ptychoparia* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Ptychoparia?* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Saukia* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Spencella erosa*	1
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Spencella porcata*	10
	Other	Arthropoda	Trilobita	Ptychopariida	*Spencella porcata*	1
	GRCA	Arthropoda	Trilobita	Ptychopariida	*Spencella* sp.	1
	MNA	Arthropoda	Trilobita	Ptychopariida	*Spencella* sp.	1
	Other	Arthropoda	Trilobita	Ptychopariida	*Spencella* sp.	1
	GRCA	Arthropoda	Trilobita	Undetermined	Undetermined	68
	MNA	Arthropoda	Trilobita	Undetermined	Undetermined	1
	Other	Arthropoda	Trilobita	Undetermined	Undetermined	1
	Other	Brachiopoda	Kutorginata	Kutorginida	*Nisusia noblei*	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctm (cont.)	Other	Brachiopoda	Kutorginata	Kutorginida	*Nisusia noblei*	1
	Other	Brachiopoda	Kutorginata	Kutorginida	*Nisusia* sp.	1
	Other	Brachiopoda	Kutorginata	Kutorginida	*Nisusia?* sp.	1
	Other	Brachiopoda	Lingulata	Lingulida	*Acrotreta?* sp.	1
	Other	Brachiopoda	Lingulata	Lingulida	*Discina* sp.	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingula* sp.	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulella (Lingulepis)* sp.	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingulepis prima*	1
	Other	Brachiopoda	Lingulata	Lingulida	*Trematis* sp.	1
	Other	Brachiopoda	Paterinata	Paterinida	*Dictyonina* sp.	1
	USNM	Brachiopoda	Rhynchonellata	Kutorginida	*Nisusia noblei*	3
	Other	Brachiopoda	Rhynchonellata	Orthida	*Finkelnburgia* sp.	1
	Other	Brachiopoda	Rhynchonellata	Pentamerida	*Syntrophia* sp.	1
	Other	Brachiopoda	Strophomenata	Billingsellida	*Clitambonites* sp.	1
	Other	Brachiopoda	Undetermined	Undetermined	Undetermined	1
	MNA	Composite	Composite	Composite	Composite unidentified	1
	Other	Mollusca	Helcionelloida	Helcionelliformes	*Helcionella* sp.	1
	Other	Porifera	Archaeocyatha	Undetermined	Undetermined	1
	Other	Problematica	Coeloscleritophora	Chancelloriida	*Chancelloria* cf. *C. eros*	1
	Other	Problematica	Coeloscleritophora	Chancelloriida	*Chancelloria?* sp.	1
	Other	Problematica	Hyolitha	Hyolithida	*Hyolithes* aff. *H. primordialis*	1
	Other	Problematica	Hyolitha	Hyolithida	*Hyolithes* sp.	1
	GRCA	Problematica	Hyolitha	Hyolithida	*Hyolithes?* sp.	4
	Other	Problematica	Hyolitha	Hyolithida	*Hyolithes?* sp.	1
	MNA	Problematica	Monoplacophora?	Cyrtonellida?	*Scenella hermitensis*	1

Appendix Table 5-A-1 (continued). Specimens of invertebrate body fossils from GRCA Cambrian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ctm	Other	Problematica	Monoplacophora?	Cyrtonellida?	*Scenella hermitensis*	1
Cu	USNM	Brachiopoda	Kutorginata	Kutorginida	*Nisusia obscura*	1

Appendix Table 5-A-2. Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Dtb	GRCA	Arthropoda	Trilobita	Undetermined	Undetermined	2
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer* sp.	2
	MNA	Brachiopoda	Undetermined	Undetermined	Undetermined	1
	Other	Brachiopoda	Undetermined	Undetermined	Undetermined	1
	Other	Cnidaria	Anthozoa	Stauriida	*Streptelasma?* sp.	1
	Other	Cnidaria	Anthozoa	Stauriida	Undetermined	1
	Other	Cnidaria	Anthozoa	Stauriida	Undetermined	1
	Other	Cnidaria	Anthozoa	Undetermined	Undetermined	1
	Other	Mollusca	Gastropoda	Undetermined	Undetermined	1
	GRCA	Undetermined	Undetermined	Undetermined	Undetermined	28
	Other	Undetermined	Undetermined	Undetermined	Undetermined	1
Mr	Other	Arthropoda	Trilobita	Eodiscida	*Aprathia* sp.	1
	Other	Arthropoda	Trilobita	Proetida	*Breviphillipsia* sp.	1
	GRCA	Arthropoda	Trilobita	Proetida	*Phillipsia tuberculata*	1
	GRCA	Arthropoda	Trilobita	Undetermined	Undetermined	1
	USNM	Brachiopoda	Rhynchonellata	Athyridida	*Cleiothyridina* sp.	1
	Other	Brachiopoda	Rhynchonellata	Orthida	*Rhipidomella* sp.	1

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Mr (cont.)	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Camarotoechia* aff. "*C. nutallica"*	2
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	*Camarotoechia* sp.	1
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	*Camarotoechia?* sp.	2
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Pugnax osagensis*	1
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Pugnax* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Pugnoides* sp.	5
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	*Pugnoides* sp.	1
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	Undetermined	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita?* sp.	1
	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Mirifusella* cf. *M. fortunata*	2
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Neospirifer striatus*	1
	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Prospira* sp.	2
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Punctospirifer kentuckyensis*	4
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer* aff. *S. incertus*	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer centronatus*	2
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer centronatus*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer* cf. *Elivina occidentalis*	1
	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer redwallensis*	4
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer* sp.	325
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer* sp.	3
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer* sp.	1
	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Torynifer* cf. *T. setiger*	2
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Unispirifer minnewankensis*	1

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Mr (cont.)	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Unispirifer minnewankensis*	3
	Other	Brachiopoda	Rhynchonellata	Spiriferinida	*Syringothyris?* sp.	1
	USNM	Brachiopoda	Rhynchonellata	Terebratulida	*Dielasma* sp.	1
	Other	Brachiopoda	Rhynchonellata	Terebratulida	Undetermined	1
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Derbyia* sp.	229
	Other	Brachiopoda	Strophomenata	Orthotetida	*Floweria chemungensis*	1
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Meekella* sp.	1
	MNA	Brachiopoda	Strophomenata	Orthotetida	*Orthotetes* sp.	1
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Streptorhynchus* sp.	5
	GRCA	Brachiopoda	Strophomenata	Productida	*Chonetes* sp.	7
	GRCA	Brachiopoda	Strophomenata	Productida	*Productus* sp.	4
	USNM	Brachiopoda	Strophomenata	Productida	*Tomiproductus gallatinensis*	2
	GRCA	Brachiopoda	Strophomenata	Productida	Undetermined	2
	Other	Brachiopoda	Strophomenata	Productida	Undetermined	1
	Other	Brachiopoda	Strophomenata	Strophomenida	*Leptagonia* sp.	1
	Other	Brachiopoda	Strophomenata	Undetermined	Undetermined	1
	GRCA	Brachiopoda	Undetermined	Undetermined	Undetermined	12
	MNA	Brachiopoda	Undetermined	Undetermined	Undetermined	1
	Other	Brachiopoda	Undetermined	Undetermined	Undetermined	1
	USNM	Brachiopoda	Undetermined	Undetermined	Undetermined	2
	Other	Bryozoa	Stenolaemata	Cryptostomida	Undetermined	1
	USNM	Bryozoa	Stenolaemata	Cystoporida	*Cheilotrypa* sp.	2
	USNM	Bryozoa	Stenolaemata	Cystoporida	*Coscinotrypa* sp.	1
	Other	Bryozoa	Stenolaemata	Cystoporida	*Cystodictya* sp.	1
	Other	Bryozoa	Stenolaemata	Cystoporida	*Dichotrypa* sp.	2

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Mr (cont.)	USNM	Bryozoa	Stenolaemata	Cystoporida	*Dichotrypa* sp.	6
	USNM	Bryozoa	Stenolaemata	Cystoporida	*Dichotrypa?* sp.	3
	USNM	Bryozoa	Stenolaemata	Cystoporida	*Ramiporalia* sp.	2
	GRCA	Bryozoa	Stenolaemata	Fenestrata	*Fenestralia* sp.	1
	USNM	Bryozoa	Stenolaemata	Fenestrata	*Fenestralia* sp.	2
	Other	Bryozoa	Stenolaemata	Fenestrata	*Fenestralia?* sp.	1
	USNM	Bryozoa	Stenolaemata	Fenestrata	*Lyroporella* sp.	8
	USNM	Bryozoa	Stenolaemata	Fenestrata	*Lyroporella?* sp.	1
	USNM	Bryozoa	Stenolaemata	Fenestrata	*Neoreteporina?* sp.	1
	USNM	Bryozoa	Stenolaemata	Fenestrata	*Polyporella* sp.	3
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* aff. *F. albida*	1
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* aff. *F. cingulata*	2
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* aff. *F. compressa*	4
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* aff. *F. serratula*	4
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* aff. *F. subflexuosa*	2
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* cf. *F. cingulata*	1
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* cf. *F. compressa*	2
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* cf. *F. serratula*	1
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Fenestella compressa*	5
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Fenestella serratula*	1
	GRCA	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* sp.	2
	Other	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* sp.	8
	Other	Bryozoa	Stenolaemata	Fenestrida	*Penniretepora* sp.	1
	Other	Bryozoa	Stenolaemata	Fenestrida	*Polypora* sp.	1
	USNM	Bryozoa	Stenolaemata	Fenestrida	*Polypora* sp.	1

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Mr (cont.)	Other	Bryozoa	Stenolaemata	Fenestrida	*Polypora*? sp.	1
	Other	Bryozoa	Stenolaemata	Fenestrida	Undetermined	1
	USNM	Bryozoa	Stenolaemata	Trepostomatida	Undetermined	5
	GRCA	Bryozoa	Undetermined	Undetermined	Undetermined	2
	MNA	Bryozoa	Undetermined	Undetermined	Undetermined	2
	Other	Bryozoa	Undetermined	Undetermined	Undetermined	1
	Other	Cnidaria	Anthozoa	Auloporida	*Cladochonus?* sp.	1
	Other	Cnidaria	Anthozoa	Auloporida	*Syringopora aculeata*	1
	Other	Cnidaria	Anthozoa	Auloporida	*Syringopora* aff. *S. surcularia*	1
	Other	Cnidaria	Anthozoa	Auloporida	*Syringopora* cf. *S. surcularia*	1
	GRCA	Cnidaria	Anthozoa	Auloporida	*Syringopora* sp.	1
	MNA	Cnidaria	Anthozoa	Auloporida	*Syringopora* sp.	2
	Other	Cnidaria	Anthozoa	Auloporida	*Syringopora* sp.	1
	Other	Cnidaria	Anthozoa	Auloporida	*Syringopora surcularia*	1
	GRCA	Cnidaria	Anthozoa	Auloporida	*Syringopora surcularia*	13
	USNM	Cnidaria	Anthozoa	Auloporida	*Syringopora surcularia*	1
	GRCA	Cnidaria	Anthozoa	Favositida	*Cladopora* sp.	1
	Other	Cnidaria	Anthozoa	Stauriida	*Amplexizaphrentis* sp.	1
	Other	Cnidaria	Anthozoa	Stauriida	*Clisiophyllum* sp.	1
	Other	Cnidaria	Anthozoa	Stauriida	*Diphyphyllum?* cf. *Lithostrotion* sp.	1
	Other	Cnidaria	Anthozoa	Stauriida	*Diphyphyllum?* sp.	1
	USNM	Cnidaria	Anthozoa	Stauriida	*Dorlodotia* cf. *D. arizelum*	1
	Other	Cnidaria	Anthozoa	Stauriida	*Dorlodotia* cf. *D. inconstans*	1
	GRCA	Cnidaria	Anthozoa	Stauriida	*Dorlodotia inconstans*	3
	MNA	Cnidaria	Anthozoa	Stauriida	*Dorlodotia inconstans*	1

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Mr (cont.)	USNM	Cnidaria	Anthozoa	Stauriida	*Dorlodotia inconstans*	2
	Other	Cnidaria	Anthozoa	Stauriida	*Dorlodotia inconstans*	1
	Other	Cnidaria	Anthozoa	Stauriida	*Dorlodotia* sp.	1
	USNM	Cnidaria	Anthozoa	Stauriida	*Ekvasophyllum* cf. *E. inclinatum*	1
	Other	Cnidaria	Anthozoa	Stauriida	*Homalophyllites paucicinctus*	1
	Other	Cnidaria	Anthozoa	Stauriida	*Homalophyllites subcrassus*	1
	USNM	Cnidaria	Anthozoa	Stauriida	*Homalophyllites subcrassus*	2
	GRCA	Cnidaria	Anthozoa	Stauriida	*Lithostrotion* (*Diphyphyllum*)*? inconstans*	3
	Other	Cnidaria	Anthozoa	Stauriida	*Menophyllum excavatum*	1
	Other	Cnidaria	Anthozoa	Stauriida	*Sychnoelasma* sp.	1
	GRCA	Cnidaria	Anthozoa	Stauriida	*Triplophyllites* (*Triplophyllites*) *persimilia*	2
	GRCA	Cnidaria	Anthozoa	Stauriida	*Triplophyllites* sp.	2
	Other	Cnidaria	Anthozoa	Stauriida	Undetermined	1
	GRCA	Cnidaria	Anthozoa	Stauriida	*Vesiculophyllum incrassatum*	20
	Other	Cnidaria	Anthozoa	Stauriida	*Vesiculophyllum incrassatum*	1
	USNM	Cnidaria	Anthozoa	Stauriida	*Vesiculophyllum incrassatum*	2
	Other	Cnidaria	Anthozoa	Stauriida	*Vesiculophyllum* sp.	1
	Other	Cnidaria	Anthozoa	Stauriida	*Zaphrentites persimilis*	1
	Other	Cnidaria	Anthozoa	Undetermined	Undetermined	1
	GRCA	Cnidaria	Undetermined	Undetermined	Undetermined	51
	Other	Echinodermata	Blastoidea	Spiraculata	*Pentremites* sp.	1
	GRCA	Echinodermata	Crinoidea	Undetermined	Undetermined	13
	Other	Echinodermata	Crinoidea	Undetermined	Undetermined	1
	Other	Echinodermata	Crinoidea	Undetermined	Undetermined	1
	USNM	Foraminifera	Fusulinata	Archaediscida	*Pohlia henbesti*	18

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Mr (cont.)	Other	Foraminifera	Fusulinata	Archaediscida	*Pohlia henbesti*	2
	Other	Foraminifera	Fusulinata	Archaediscida	*Septatournayella*? sp.	1
	Other	Foraminifera	Fusulinata	Archaediscida	*Tournayella* sp.	1
	Other	Foraminifera	Fusulinata	Archaediscida	*Tournayella?* sp.	1
	Other	Foraminifera	Fusulinata	Earlandiida	*Earlandia* sp.	1
	USNM	Foraminifera	Fusulinata	Earlandiida	*Earlandia* sp.	4
	Other	Foraminifera	Fusulinata	Earlandiida	*Paracaligella?* sp.	1
	USNM	Foraminifera	Fusulinata	Earlandiida	*Paracaligella?* sp.	4
	Other	Foraminifera	Fusulinata	Earlandiida	*Paramillerella?* sp.	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra* aff. *E. gutschicki*	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra* aff. *E. tantala*	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra* aff. *E. trachida*	2
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra kleina*	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Endothyra kleina*	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra* sp.	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra* sp.	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Endothyra* sp.	2
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra tantula*	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Endothyra tantula*	8
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra trachida*	2
	USNM	Foraminifera	Fusulinata	Endothyrida	*Endothyra trachida*	15
	USNM	Foraminifera	Fusulinata	Endothyrida	*Endothyra?* aff. *"E. excellens"*	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Eoendothyranopsis* aff. *E. spiroides*	2
	USNM	Foraminifera	Fusulinata	Endothyrida	*Eoendothyranopsis* sp.	2
	Other	Foraminifera	Fusulinata	Endothyrida	*Eoendothyranopsis spiroides*	2

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Mr (cont.)	USNM	Foraminifera	Fusulinata	Endothyrida	*Eoendothyranopsis spiroides*	15
	Other	Foraminifera	Fusulinata	Endothyrida	*Globoendothyra baileyi*	2
	USNM	Foraminifera	Fusulinata	Endothyrida	*Globoendothyra baileyi*	10
	Other	Foraminifera	Fusulinata	Endothyrida	*Globoendothyra baileyi* subsp. *poloumera*	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Globoendothyra baileyi* subsp. *poloumera*	6
	Other	Foraminifera	Fusulinata	Endothyrida	*Inflatoendothyra eospiroides*	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Inflatoendothyra eospiroides*	7
	Other	Foraminifera	Fusulinata	Endothyrida	*Septabrunsiina* (*Spinobrunsiina*) *parakrainica*	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Septabrunsiina* (*Spinobrunsiina*) *parakrainica*	12
	Other	Foraminifera	Fusulinata	Endothyrida	*Septabrunsiina* sp.	2
	Other	Foraminifera	Fusulinata	Endothyrida	*Septaglomospiranella chernoussovensis*	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Septaglomospiranella chernoussovensis*	17
	Other	Foraminifera	Fusulinata	Endothyrida	*Septaglomospiranella rossi*	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Septaglomospiranella rossi*	2
	Other	Foraminifera	Fusulinata	Endothyrida	*Septaglomospiranella* sp.	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Spinobrunsiina* aff. *S. torquida*	2
	USNM	Foraminifera	Fusulinata	Endothyrida	*Spinobrunsiina torquida*	11
	Other	Foraminifera	Fusulinata	Endothyrida	*Spinoendothyra* aff. *S. spinosa*	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Spinoendothyra spinosa*	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Spinoendothyra spinosa*	13
	Other	Foraminifera	Fusulinata	Endothyrida	*Tuberendothyra paratumula*	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Tuberendothyra paratumula*	27
	Other	Foraminifera	Fusulinata	Endothyrida	*Tuberendothyra* sp.	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Tuberendothyra* sp.*?*	1

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Mr (cont.)	Other	Foraminifera	Fusulinata	Endothyrida	*Tuberendothyra tuberculata*	1
	USNM	Foraminifera	Fusulinata	Endothyrida	*Tuberendothyra tuberculata*	22
	Other	Foraminifera	Fusulinata	Endothyrida	*Spinobrunsiina torquida*	1
	GRCA	Foraminifera	Fusulinata	Fusulinida	*Schubertella* sp.	4
	Other	Foraminifera	Fusulinata	Lituolida	*Spiroplectamminoides?* cf. *Spiroplectammina parva*	1
	USNM	Foraminifera	Fusulinata	Lituolida	*Spiroplectamminoides?* cf. *Spiroplectammina parva*	3
	Other	Foraminifera	Fusulinata	Parathuramminida	*Calcisphaera* sp.	1
	USNM	Foraminifera	Fusulinata	Parathuramminida	*Calcisphaera* sp.	1
	Other	Mollusca	Cephalopoda	Actinocerida	*Rayonnoceras* sp.	1
	GRCA	Mollusca	Gastropoda	Bellerophontida	*Bellerophon* sp.	2
	Other	Mollusca	Gastropoda	Bellerophontida	*Bellerophon* sp.	1
	GRCA	Mollusca	Gastropoda	Bellerophontida	*Euphemites subpapillosus*	17
	Other	Mollusca	Gastropoda	Bellerophontida	*Euphemites?* sp.	1
	Other	Mollusca	Gastropoda	Bellerophontida	Undetermined	1
	GRCA	Mollusca	Gastropoda	Euomphalina	*Euomphalus* sp.	3
	Other	Mollusca	Gastropoda	Euomphalina	*Euomphalus?* sp.	1
	GRCA	Mollusca	Gastropoda	Euomphalina	*Straparollus* sp.	15
	GRCA	Mollusca	Gastropoda	Euomphalina	Undetermined	7
	Other	Mollusca	Gastropoda	Euomphalina	Undetermined	1
	GRCA	Mollusca	Gastropoda	Murchisoniina	*Loxonema* sp.	1
	Other	Mollusca	Gastropoda	Murchisoniina	Undetermined	1
	Other	Mollusca	Gastropoda	Pleurotomariida	*Euconospira* aff. *"E. montezuma"*	1
	Other	Mollusca	Gastropoda	Undetermined	Undetermined	1
	GRCA	Undetermined	Undetermined	Undetermined	Undetermined	60

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Mr (cont.)	Other	Undetermined	Undetermined	Undetermined	Undetermined	1
Msc	Other	Arthropoda	Trilobita	Proetida	*Paladin* cf. *P. chesterensis*	1
	Other	Arthropoda	Trilobita	Proetida	*Paladin* sp.	1
	Other	Brachiopoda	Rhynchonellata	Athyrida	*Cleiothyridina* sp.	1
	Other	Brachiopoda	Rhynchonellata	Athyrida	*Eumetria* sp.	1
	Other	Brachiopoda	Rhynchonellata	Orthida	*Rhipidomella nevadensis*	1
	GRCA	Brachiopoda	Rhynchonellata	Orthida	*Rhipidomella* sp.	1
	Other	Brachiopoda	Rhynchonellata	Orthida	*Schizophoria* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Cupularostrum* aff. *“C. purduie”*	1
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	*Leiorhynchoidea carbonifera*	1
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Leiorhynchoidea* sp.	2
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	*Macropotamorhynchus* cf. *M. purduei*	1
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Macropotamorhynchus* sp.	1
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	*Pugnoides* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Rotaia neogenes*	1
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	*Rotaia neogenes*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer* aff. *A. curvilateralis*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer bifurcatus*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer* cf. *A. curvilateralis*	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer curvilateralis*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer curvilateralis*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Brachythyris (Spirifer) subcardiiformis*	1
	MNA	Brachiopoda	Rhynchonellata	Spiriferida	*Cleiothyridina orbicularis*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita gibbosa*	1

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Msc (cont.)	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita laevis*	1
	MNA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita laevis*	2
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita laevis*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita ovata*	1
	MNA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita subquadrata*	2
	MNA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita subquadrata*	2
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita subquadrata*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Torynifer* aff. *T. setiger*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Torynifer* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	Undetermined	2
	MNA	Brachiopoda	Rhynchonellata	Spiriferinida	*Punctospirifer transversus*	2
	GRCA	Brachiopoda	Rhynchonellata	Spiriferinida	*Reticulariina* sp.	1
	Other	Brachiopoda	Rhynchonellata	Terebratulida	*Beecheria* cf. B. *arkansanum*	1
	Other	Brachiopoda	Rhynchonellata	Terebratulida	*Beecheria* sp.	1
	Other	Brachiopoda	Rhynchonellata	Terebratulida	*Cranaena* sp.	1
	MNA	Brachiopoda	Strophomenata	Orthotetida	*Orthotetes* sp.	1
	Other	Brachiopoda	Strophomenata	Orthotetida	*Orthotetes* sp.	1
	Other	Brachiopoda	Strophomenata	Orthotetida	Undetermined	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Antiquatonia* sp.	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Flexaria* sp.	3
	Other	Brachiopoda	Strophomenata	Productida	*Flexaria* sp.	1
	Other	Brachiopoda	Strophomenata	Productida	*Inflatia* aff. *I. clydensis*	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Inflatia* sp.	2

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Msc (cont.)	Other	Brachiopoda	Strophomenata	Productida	*Inflatia* sp.	1
	Other	Brachiopoda	Strophomenata	Productida	*Inflatia* sp. or *Sandia* sp.	1
	Other	Brachiopoda	Strophomenata	Productida	*Inflatia?* sp.	1
	Other	Brachiopoda	Strophomenata	Productida	*Ovatia* sp.	2
	Other	Brachiopoda	Strophomenata	Productida	Undetermined	2
	Other	Brachiopoda	Undetermined	Undetermined	Undetermined	2
	MNA	Bryozoa	Stenolaemata	Fenestrida	*Archimedes* sp.	1
	Other	Bryozoa	Stenolaemata	Fenestrida	*Archimedes* sp.	1
	Other	Bryozoa	Undetermined	Undetermined	Undetermined	4
	GRCA	Cnidaria	Anthozoa	Favositida	*Michelinia* sp.	2
	Other	Cnidaria	Anthozoa	Favositida	*Michelinia* sp.	1
	GRCA	Cnidaria	Anthozoa	Stauriida	*Amplexus* aff. *"Barytchisma" zaphrentiformis*	1
	Other	Cnidaria	Anthozoa	Stauriida	*Amplexus* sp.	1
	Other	Cnidaria	Anthozoa	Stauriida	*Barytichisma* sp.	1
	Other	Cnidaria	Anthozoa	Stauriida	Undetermined	1
	Other	Cnidaria	Anthozoa	Undetermined	Undetermined	1
	MNA	Echinodermata	Asteroidea	Uractinida	Undetermined	4
	GRCA	Echinodermata	Blastoidea	Spiraculata	*Pentremites* sp.	2
	MNA	Echinodermata	Blastoidea	Spiraculata	*Pentremites* sp.	1
	Other	Echinodermata	Blastoidea	Undetermined	Undetermined	1
	MNA	Echinodermata	Crinoidea	Ampelocrinida	*Cymbiocrinus* sp.	1
	Other	Echinodermata	Crinoidea	Ampelocrinida	*Cymbiocrinus* sp.	1
	Other	Echinodermata	Crinoidea	Undetermined	Undetermined	1
	Other	Echinodermata	Echinoidea	Undetermined	Undetermined	2
	Other	Echinodermata	Undetermined	Undetermined	Undetermined	1

Appendix Table 5-A-2 (continued). Specimens of invertebrate body fossils from GRCA Devonian and Mississippian units as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Msc (cont.)	Other	Echinodermata	Undetermined	Undetermined	Undetermined	1
	Other	Foraminifera	Undetermined	Undetermined	Undetermined	1
	Other	Mollusca	Bivalvia	Myalinida	*Aviculopecten* sp.	1
	Other	Mollusca	Bivalvia	Myalinida	*Septimyalina* sp.	1
	Other	Mollusca	Bivalvia	Myalinida	*Septimyalina*? sp.	1
	Other	Mollusca	Bivalvia	Pectinida	Undetermined	1
	Other	Mollusca	Bivalvia	Pholadomyida	*Edmondia* sp.	1
	Other	Mollusca	Bivalvia	Trigoniida	*Schizodus* sp.	1
	GRCA	Mollusca	Bivalvia	Undetermined	Undetermined	1
	Other	Mollusca	Bivalvia	Undetermined	Undetermined	1
	MNA	Mollusca	Gastropoda	Archaeogastropoda	Undetermined	1
	Other	Mollusca	Gastropoda	Bellerophontida	*Bellerophon* (*Bellerophon*) sp.	1
	Other	Mollusca	Gastropoda	Bellerophontida	*Bellerophon* spp.	1
	Other	Mollusca	Gastropoda	Eotomarioidea	*Glabrocingulum* sp.	1
	Other	Mollusca	Gastropoda	Euomphalina	*Euomphalus* sp.	1
	Other	Mollusca	Gastropoda	Euomphalina	*Straparollus?* sp.	1
	MNA	Mollusca	Gastropoda	Murchisoniina	*Bellazona* sp.	1
	Other	Mollusca	Gastropoda	Murchisoniina	*Bellazona* sp.	1
	Other	Mollusca	Gastropoda	Murchisoniina	*Loxonema* sp.	1
	Other	Mollusca	Undetermined	Undetermined	Undetermined	1
	GRCA	Undetermined	Undetermined	Undetermined	Undetermined	2
	Other	Undetermined	Undetermined	Undetermined	Undetermined	1
	Other	Undetermined	Undetermined	Undetermined	Undetermined	1

Appendix Table 5-A-3. Specimens of invertebrate body fossils from the Supai Group of GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
IPsu	GRCA	Brachiopoda	Lingulata	Lingulida	*Orbiculoidea* sp.	28
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	94
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer* sp.	25
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Derbyia* sp.	41
	GRCA	Brachiopoda	Strophomenata	Productida	*Linoproductus* sp.	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Productus* sp.	6
	GRCA	Cnidaria	Undetermined	Undetermined	Undetermined	3
	GRCA	Echinodermata	Crinoidea	Undetermined	Undetermined	5
	GRCA	Mollusca	Bivalvia	Myalinida	*Aviculopecten* sp.	2
	GRCA	Mollusca	Bivalvia	Myalinida	*Myalina* aff. *M. swallovi*	6
	GRCA	Mollusca	Bivalvia	Myalinida	*Myalina cuneiformis*	2
	GRCA	Mollusca	Bivalvia	Myalinida	*Myalina perattenuata*	6
	GRCA	Mollusca	Bivalvia	Myalinida	*Myalina* sp.	73
	GRCA	Mollusca	Bivalvia	Mytilida	*Modiolus* sp.	4
	GRCA	Mollusca	Bivalvia	Ostreida	*Cornellites* sp.	2
	GRCA	Mollusca	Bivalvia	Undetermined	Undetermined	30
	GRCA	Mollusca	Gastropoda	Euomphalina	*Euomphalus?* sp.	4
	MNA	Mollusca	Gastropoda	Undetermined	Undetermined	1
	GRCA	Undetermined	Undetermined	Undetermined	Undetermined	50
IPwa	Other	Arthropoda	Trilobita	Proetida	*Paladin* sp.	1
	Other	Arthropoda	Trilobita	Proetida	*Paladin?* aff. "*Librogenae*"	1
	Other	Arthropoda	Trilobita	Proetida	*Paladin?* sp.	1
	Other	Brachiopoda	Lingulata	Lingulida	*Lingula* sp.	2
	Other	Brachiopoda	Lingulata	Lingulida	*Orbiculoidea* aff. *O. meekana*	1
	Other	Brachiopoda	Lingulata	Lingulida	*Orbiculoidea meekana*	1

Appendix Table 5-A-3 (continued). Specimens of invertebrate body fossils from the Supai Group of GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
IPswa (cont.)	USNM	Brachiopoda	Lingulata	Lingulida	*Orbiculoidea meekana*	1
	Other	Brachiopoda	Lingulata	Lingulida	*Orbiculoidea* sp.	1
	Other	Brachiopoda	Rhynchonellata	Orthida	*Schizophoria* aff. S. *altirostris*	1
	USNM	Brachiopoda	Rhynchonellata	Orthida	*Schizophoria* aff. S. *altirostris*	2
	Other	Brachiopoda	Rhynchonellata	Orthida	*Schizophoria altirostris*	2
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	*Cupularostrum?* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer newberryi*	1
	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer newberryi*	2
	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer occiduus*	4
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer tanoensis*	1
	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Anthracospirifer tanoensis*	2
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita ovata*	1
	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Composita ovata*	3
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	1
	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita subtilita*	1
	USNM	Brachiopoda	Rhynchonellata	Spiriferida	*Composita subtilita*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Spiriferina?* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferinida	*Punctospirifer transversus*	1
	USNM	Brachiopoda	Rhynchonellata	Spiriferinida	*Punctospirifer transversus*	3
	Other	Brachiopoda	Rhynchonellata	Spiriferinida	*Reticulariina gonionota*	2
	Other	Brachiopoda	Strophomenata	Orthotetida	*Derbyia* aff. *D. robusta*	1
	Other	Brachiopoda	Strophomenata	Orthotetida	*Derbyia* sp.	1
	USNM	Brachiopoda	Strophomenata	Orthotetida	*Derbyia* sp.?	1

Appendix Table 5-A-3 (continued). Specimens of invertebrate body fossils from the Supai Group of GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
IPswa (cont.)	Other	Brachiopoda	Strophomenata	Orthotetida	*Orthotetes* sp.	1
	USNM	Brachiopoda	Strophomenata	Orthotetida	*Orthotetes* sp.	1
	Other	Brachiopoda	Strophomenata	Productida	Undetermined	1
	USNM	Brachiopoda	Undetermined	Undetermined	Undetermined	1
	Other	Bryozoa	Stenolaemata	Cystoporata	Undetermined	1
	Other	Bryozoa	Stenolaemata	Cystoporida	*Cystiodictya?* sp.	1
	Other	Bryozoa	Stenolaemata	Cystoporida	Undetermined	1
	Other	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* sp.	1
	Other	Bryozoa	Stenolaemata	Trepostomatida	Undetermined	1
	Other	Bryozoa	Undetermined	Undetermined	Undetermined	1
	Other	Cnidaria	Anthozoa	Favositida	*Michelinia* sp.	1
	Other	Cnidaria	Anthozoa	Undetermined	Undetermined	1
	Other	Cnidaria	Staurozoa?	Conulatae	Undetermined	1
	Other	Echinodermata	Crinoidea	Undetermined	Undetermined	1
	Other	Echinodermata	Undetermined	Undetermined	Undetermined	1
	Other	Foraminifera	Fusulinata	Fusulinida	*Fusulinella* sp.	1
	Other	Foraminifera	Fusulinata	Fusulinida	*Pseudostaffella* sp.	1
	Other	Mollusca	Bivalvia	Cardiida	*Oriocrassatella* sp.	1
	Other	Mollusca	Bivalvia	Cardiida	*Permophorus* sp.	1
	USNM	Mollusca	Bivalvia	Cardiida	*Permophorus* sp.	1
	Other	Mollusca	Bivalvia	Myalinida	*Aviculopecten* aff. *A. gravidus*	1
	Other	Mollusca	Bivalvia	Myalinida	*Aviculopecten gravidus*	1
	USNM	Mollusca	Bivalvia	Myalinida	*Aviculopecten gravidus*	1
	Other	Mollusca	Bivalvia	Myalinida	*Aviculopecten* sp.	1
	Other	Mollusca	Bivalvia	Myalinida	*Aviculopecten* sp.	1

Appendix Table 5-A-3 (continued). Specimens of invertebrate body fossils from the Supai Group of GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
IPswa (cont.)	USNM	Mollusca	Bivalvia	Myalinida	*Aviculopecten* sp.	1
	USNM	Mollusca	Bivalvia	Myalinida	*Myalina (Myalina)* sp.	1
	Other	Mollusca	Bivalvia	Myalinida	*Myalina* sp.	2
	Other	Mollusca	Bivalvia	Myalinida	*Septimyalina* sp.	1
	USNM	Mollusca	Bivalvia	Myalinida	*Septimyalina* sp.	2
	Other	Mollusca	Bivalvia	Myalinida	Undetermined	1
	Other	Mollusca	Bivalvia	Ostreida	*Leptodesma* sp.	1
	Other	Mollusca	Bivalvia	Trigoniida	*Schizodus* sp.	1
	USNM	Mollusca	Bivalvia	Trigoniida	*Schizodus* sp.	2
	Other	Mollusca	Bivalvia	Unionida	*Promytilus* sp.	1
	USNM	Mollusca	Bivalvia	Unionida	*Promytilus* sp.	1
	Other	Mollusca	Gastropoda	Bellerophontida	Undetermined	1
	Other	Mollusca	Gastropoda	Euomphalina	*Euomphalus* sp.	1
	Other	Mollusca	Gastropoda	Euomphalina	*Straparollus* sp.	1
	Other	Mollusca	Gastropoda	Pleurotomariida	Undetermined	1
	Other	Mollusca	Gastropoda	Undetermined	Undetermined	1
	Other	Mollusca	Gastropoda	Undetermined	Undetermined	1
IPsm	Other	Arthropoda	Ostracoda	Undetermined	Undetermined	1
	Other	Echinodermata	Undetermined	Undetermined	Undetermined	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra media*	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra* sp.	1
	Other	Foraminifera	Fusulinata	Endothyrida	*Endothyra teres*	1
	Other	Foraminifera	Fusulinata	Fusulinida	*Eoschubertella* sp.	1
	Other	Foraminifera	Fusulinata	Fusulinida	*Fusulinella* sp.	1
	Other	Foraminifera	Fusulinata	Fusulinida	*Pseudostaffella* sp.	1

Appendix Table 5-A-3 (continued). Specimens of invertebrate body fossils from the Supai Group of GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
IPsm (cont.)	Other	Foraminifera	Fusulinata	Fusulinida	*Schubertella* sp.	2
	Other	Foraminifera	Fusulinata	Fusulinida	Undetermined	1
	Other	Foraminifera	Undetermined	Undetermined	Undetermined	1
	Other	Undetermined	Undetermined	Undetermined	Calcisphere bioclasts	1
IPswe	Other	Bryozoa	Undetermined	Undetermined	Undetermined	1
	Other	Foraminifera	Fusulinata	Fusulinida	Undetermined	1
	Other	Foraminifera	Undetermined	Undetermined	Undetermined	1
Pse	Other	Bryozoa	Undetermined	Undetermined	Undetermined	1
	Other	Cnidaria	Undetermined	Undetermined	Undetermined	1
	Other	Echinodermata	Undetermined	Undetermined	Undetermined	1
	Other	Foraminifera	Fusulinata	Fusulinida	*Schubertella* sp.	2
	Other	Foraminifera	Undetermined	Undetermined	Undetermined	1

Appendix Table 5-A-4. Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ph	Other	Arthropoda	Subphylum Chelicerata	Eurypterida	*Hastimima?* sp.	1
	USNM	Arthropoda	Subphylum Chelicerata	Eurypterida	*Hastimima?* sp.	2
	USNM	Arthropoda	Insecta	Blattodea	Undetermined	1
	USNM	Arthropoda	Insecta	Meganisoptera	*Tupus gilmorei*	1
	USNM	Arthropoda	Insecta	Meganisoptera	*Tupus whitei*	1

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Ph (cont.)	Other	Arthropoda	Insecta	Odonata	Undetermined	1
	GRCA	Undetermined	Undetermined	Undetermined	Undetermined	10
	USNM	Undetermined	Undetermined	Undetermined	Undetermined	1
Pt	Other	Arthropoda	Ostracoda	Undetermined	Undetermined	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	25
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Peniculauris bassi*	7
	MNA	Brachiopoda	Strophomenata	Productida	*Peniculauris ivesi*	1
	Other	Brachiopoda	Strophomenata	Productida	*Peniculauris ivesi*	1
	MNA	Bryozoa	Undetermined	Undetermined	Undetermined	1
	Other	Bryozoa	Undetermined	Undetermined	Undetermined	1
	GRCA	Echinodermata	Crinoidea	Undetermined	Undetermined	2
	MNA	Echinodermata	Crinoidea	Undetermined	Undetermined	1
	Other	Echinodermata	Crinoidea	Undetermined	Undetermined	1
	GRCA	Echinodermata	Echinoidea	Not assigned	*Archaeocidaris* sp.	1
	MNA	Echinodermata	Echinoidea	Not assigned	*Archaeocidaris* sp.	3
	Other	Echinodermata	Echinoidea	Not assigned	*Archaeocidaris* sp.	1
	GRCA	Echinodermata	Echinoidea	Not assigned	*Archaeocidaris* sp. (*Echinocrinus* sp.)	24
	MNA	Echinodermata	Echinoidea	Undetermined	Undetermined	1
	Other	Mollusca	Bivalvia	Adapedonta	*Edmondia* sp.	1
	Other	Mollusca	Bivalvia	Cardiida	*Permophorus* sp.	1
	GRCA	Mollusca	Bivalvia	Nuculanida	*Nuculana* sp.	3
	Other	Mollusca	Bivalvia	Nuculanida	*Nuculana?* sp.	1
	Other	Mollusca	Bivalvia	Ostreida	*Pteria* sp.	1
	GRCA	Mollusca	Bivalvia	Pectinida	*Deltopecten* sp.	1

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pt (cont.)	MNA	Mollusca	Bivalvia	Pectinida	*Pecten* sp.	1
	GRCA	Mollusca	Bivalvia	Pholadomyida	*Allorisma* sp.	1
	MNA	Mollusca	Bivalvia	Pholadomyida	*Allorisma* sp.	1
	Other	Mollusca	Bivalvia	Pholadomyida	*Allorisma* sp.	1
	Other	Mollusca	Bivalvia	Protobranchia	*Solemya* sp.	1
	GRCA	Mollusca	Bivalvia	Trigoniida	*Schizodus* sp.	5
	MNA	Mollusca	Bivalvia	Trigoniida	*Schizodus* sp.	1
	Other	Mollusca	Bivalvia	Trigoniida	*Schizodus* sp.	1
	GRCA	Mollusca	Bivalvia	Undetermined	Undetermined	1
	Other	Mollusca	Bivalvia	Undetermined	Undetermined	1
	Other	Mollusca	Cephalopoda	Nautilida	*Domatoceras?* sp.	1
	Other	Mollusca	Gastropoda	Bellerophontida	*Bellerophon* sp.	2
	GRCA	Mollusca	Gastropoda	Bellerophontida	*Euphemites* sp.	2
	Other	Mollusca	Gastropoda	Bellerophontida	*Euphemites* sp.	1
	Other	Mollusca	Gastropoda	Cycloneritida	*Naticopsis* sp.	1
	GRCA	Mollusca	Gastropoda	Euomphalina	*Euomphalus* sp.	3
	Other	Mollusca	Gastropoda	Euomphalina	*Euomphalus* sp.	1
	Other	Mollusca	Gastropoda	Murchisoniina	*Aclisina* sp.	1
	Other	Mollusca	Gastropoda	Murchisoniina	*Goniospira* sp.	1
	Other	Mollusca	Gastropoda	Undetermined	Undetermined	1
	GRCA	Mollusca	Scaphopoda	Dentaliida	*Dentalium* sp.	2
	Other	Mollusca	Undetermined	Undetermined	Undetermined	1
	GRCA	Porifera	Undetermined	Undetermined	Undetermined	1
	GRCA	Undetermined	Undetermined	Undetermined	Undetermined	23

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
	GRCA	Arthropoda	Trilobita	Proetida	*Ameura* sp.	3
	GRCA	Arthropoda	Trilobita	Proetida	*Anisopyge (=Novoameura) mckeei*	1
	GRCA	Arthropoda	Trilobita	Proetida	*Delaria macclintocki*	1
	Other	Arthropoda	Trilobita	Proetida	*Delaria sevilloidia*	1
	GRCA	Arthropoda	Trilobita	Proetida	*Delaria snowi*	2
	Other	Arthropoda	Trilobita	Proetida	*Delaria snowi*	1
	GRCA	Arthropoda	Trilobita	Proetida	*Ditomopyge scitulus*	1
	GRCA	Arthropoda	Trilobita	Proetida	*Ditomopyge* sp.	1
	Other	Arthropoda	Trilobita	Proetida	*Ditomopyge* sp.	1
	Other	Arthropoda	Trilobita	Proetida	*Novoameura mckeei*	1
	GRCA	Arthropoda	Trilobita	Undetermined	Undetermined	23
	Other	Arthropoda	Trilobita	Undetermined	Undetermined	1
Pk	GRCA	Brachiopoda	Lingulata	Lingulida	*Orbiculoidea* sp.	5
	GRCA	Brachiopoda	Rhynchonellata	Athyridida	*Hustedia* aff. *H. meekana*	2
	Other	Brachiopoda	Rhynchonellata	Athyridida	*Hustedia* aff. *H. meekana*	1
	Other	Brachiopoda	Rhynchonellata	Athyridida	*Hustedia* sp.	1
	Other	Brachiopoda	Rhynchonellata	Orthida	*Rhipidomella hessensis*	1
	Other	Brachiopoda	Rhynchonellata	Orthida	*Rhipidomella transversa*	1
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	*Phrenophoria pinguis*	1
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Pugnax osagensis*	6
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Pugnax* sp.	3
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Pugnoides* sp.	21
	Other	Brachiopoda	Rhynchonellata	Rhynchonellida	*Pugnoides* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Rhynchonella* sp.	6
	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Stenoscisma hueconianum*	1

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	GRCA	Brachiopoda	Rhynchonellata	Rhynchonellida	*Stenoscisma* sp.	9
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita arizonica*	11
	MNA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita arizonica*	2
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita arizonica*	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	62
	MNA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	2
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Composita subtilita*	3
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita subtilita*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Composita?* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Neophricadothyris* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Phricodothyris guadalupensis*	2
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Phricodothyris guadalupensis*	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Spirifer* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Spiriferina* sp.	7
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Spiriferina?* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Squamularia* sp.	105
	MNA	Brachiopoda	Rhynchonellata	Spiriferida	*Squamularia* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferida	*Squamularia* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferida	*Timaniella pseudocameratus*	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferinida	*Punctospirifer* sp.	1
	Other	Brachiopoda	Rhynchonellata	Spiriferinida	*Punctospirifer?* sp.	1
	GRCA	Brachiopoda	Rhynchonellata	Spiriferinida	*Spiriferellina hilli*	1
	MNA	Brachiopoda	Rhynchonellata	Spiriferinida	*Spiriferellina hilli*	1
	Other	Brachiopoda	Rhynchonellata	Spiriferinida	*Spiriferellina hilli*	1

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	MNA	Brachiopoda	Rhynchonellata	Terebratulida	*Dielasma bovidens*	1
	GRCA	Brachiopoda	Rhynchonellata	Terebratulida	*Dielasma phosphoriense*	1
	Other	Brachiopoda	Rhynchonellata	Terebratulida	*Dielasma phosphoriense*	1
	GRCA	Brachiopoda	Rhynchonellata	Terebratulida	*Dielasma* sp.	6
	Other	Brachiopoda	Strophomenata	Orthotetida	*Derbyia arizonensis*	1
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Derbyia arizonensis*	4
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Derbyia multistriata*	1
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Derbyia nasuta*	1
	Other	Brachiopoda	Strophomenata	Orthotetida	*Derbyia nasuta*	1
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Derbyia* sp.	22
	MNA	Brachiopoda	Strophomenata	Orthotetida	*Derbyia sp.*	1
	Other	Brachiopoda	Strophomenata	Orthotetida	*Derbyia* sp.	1
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Meekella occidentalis*	5
	USNM	Brachiopoda	Strophomenata	Orthotetida	*Meekella occidentalis*	1
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Meekella pyramidalis*	14
	MNA	Brachiopoda	Strophomenata	Orthotetida	*Meekella pyramidalis*	2
	Other	Brachiopoda	Strophomenata	Orthotetida	*Meekella pyramidalis*	1
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Meekella* sp.	41
	MNA	Brachiopoda	Strophomenata	Orthotetida	*Meekella sp.*	1
	Other	Brachiopoda	Strophomenata	Orthotetida	*Meekella* sp.	1
	GRCA	Brachiopoda	Strophomenata	Orthotetida	*Orthotetes* sp.	16
	MNA	Brachiopoda	Strophomenata	Productida	*Antiquatonia* sp.	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Avonia* sp.	57
	MNA	Brachiopoda	Strophomenata	Productida	*Avonia* sp.	3
	Other	Brachiopoda	Strophomenata	Productida	*Avonia* sp.	1

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	Other	Brachiopoda	Strophomenata	Productida	*Bathymyonia nevadensis*	1
	MNA	Brachiopoda	Strophomenata	Productida	*Bathymyonia nevadensis*	2
	GRCA	Brachiopoda	Strophomenata	Productida	*Chonetes* aff. "*C. hillianus*"	12
	GRCA	Brachiopoda	Strophomenata	Productida	*Chonetes permianus*	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Chonetes* sp.	141
	MNA	Brachiopoda	Strophomenata	Productida	*Chonetes* sp.	1
	Other	Brachiopoda	Strophomenata	Productida	*Chonetes* sp.	2
	GRCA	Brachiopoda	Strophomenata	Productida	*Dictyoclostus semireticulatus*	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Dictyoclostus* sp.	1
	Other	Brachiopoda	Strophomenata	Productida	*Dyoros* (*Tetragonetes) tetragonus*	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Dyoros kaibabensis*	1
	Other	Brachiopoda	Strophomenata	Productida	*Dyoros kaibabensis*	1
	Other	Brachiopoda	Strophomenata	Productida	*Dyoros subliratus*	1
	MNA	Brachiopoda	Strophomenata	Productida	*Dyoros subliratus*	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Echinauris dorsoconcava*	1
	Other	Brachiopoda	Strophomenata	Productida	*Echinauris dorsoconcava*	1
	Other	Brachiopoda	Strophomenata	Productida	*Echinauris newberryi*	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Echinauris newberryi*	1
	Other	Brachiopoda	Strophomenata	Productida	*Echinauris* sp.	1
	Other	Brachiopoda	Strophomenata	Productida	*Kozlowskia* sp.	1
	MNA	Brachiopoda	Strophomenata	Productida	*Kutorginella meridionalis*	5
	Other	Brachiopoda	Strophomenata	Productida	*Liosotella popei*	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Liosotella popei*	3
	MNA	Brachiopoda	Strophomenata	Productida	*Marginifera* sp.	1
	Other	Brachiopoda	Strophomenata	Productida	*Marginifera* sp.	1

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	GRCA	Brachiopoda	Strophomenata	Productida	*Peniculauris bassi*	66
	MNA	Brachiopoda	Strophomenata	Productida	*Peniculauris bassi*	2
	MNA	Brachiopoda	Strophomenata	Productida	*Peniculauris bassi*	4
	Other	Brachiopoda	Strophomenata	Productida	*Peniculauris bassi*	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Peniculauris ivesi*	43
	Other	Brachiopoda	Strophomenata	Productida	*Peniculauris ivesi*	1
	USNM	Brachiopoda	Strophomenata	Productida	*Peniculauris ivesi*	3
	MNA	Brachiopoda	Strophomenata	Productida	*Peniculauris* sp.	1
	Other	Brachiopoda	Strophomenata	Productida	*Peniculauris* sp.	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Productus* aff. *"P. iiginac"*	2
	GRCA	Brachiopoda	Strophomenata	Productida	*Productus* sp.	87
	GRCA	Brachiopoda	Strophomenata	Productida	*Rugatia occidentalis*	133
	Other	Brachiopoda	Strophomenata	Productida	*Rugatia occidentalis*	1
	USNM	Brachiopoda	Strophomenata	Productida	*Rugatia occidentalis*	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Rugatia paraindica*	2
	MNA	Brachiopoda	Strophomenata	Productida	*Rugatia paraindica*	2
	Other	Brachiopoda	Strophomenata	Productida	*Rugatia paraindica*	1
	USNM	Brachiopoda	Strophomenata	Productida	*Rugatia paraindica*	3
	Other	Brachiopoda	Strophomenata	Productida	*Rugatia* sp.	1
	GRCA	Brachiopoda	Strophomenata	Productida	Undetermined	1
	GRCA	Brachiopoda	Strophomenata	Productida	*Waagenoconcha irginae*	9
	GRCA	Brachiopoda	Strophomenata	Productida	*Waagenoconcha* sp.	5
	Other	Brachiopoda	Strophomenata	Productida	*Waagenoconcha* sp.	1
	GRCA	Brachiopoda	Undetermined	Undetermined	Undetermined	9
	MNA	Brachiopoda	Undetermined	Undetermined	Undetermined	1

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	MNA	Brachiopoda	Undetermined	Undetermined	Undetermined	1
	MNA	Brachiopoda	Undetermined	Undetermined	Undetermined	1
	MNA	Brachiopoda	Undetermined	Undetermined	Undetermined	2
	Other	Brachiopoda	Undetermined	Undetermined	Undetermined	1
	Other	Bryozoa	Stenolaemata	Cryptostomida	*Girtypora maculata*	1
	Other	Bryozoa	Stenolaemata	Cryptostomida	*Rhabdomeson* sp.	1
	GRCA	Bryozoa	Stenolaemata	Cryptostomida	*Rhombopora lepidodendroides*	3
	Other	Bryozoa	Stenolaemata	Cryptostomida	*Rhombopora lepidodendroides*	1
	Other	Bryozoa	Stenolaemata	Cryptostomida	*Streblotrypa sp.*	1
	GRCA	Bryozoa	Stenolaemata	Cystoporida	*Dichotrypa* sp.	1
	MNA	Bryozoa	Stenolaemata	Cystoporida	*Fistulipora* aff. *“F. arizonensis”*	1
	GRCA	Bryozoa	Stenolaemata	Cystoporida	*Fistulipora* sp.	3
	MNA	Bryozoa	Stenolaemata	Cystoporida	*Fistulipora* sp.	2
	Other	Bryozoa	Stenolaemata	Cystoporida	*Fistulipora* sp.	2
	MNA	Bryozoa	Stenolaemata	Cystoporida	*Meekopora* aff. “*M. robusta”*	13
	MNA	Bryozoa	Stenolaemata	Cystoporida	*Meekopora* aff*. M. tenuis*	1
	Other	Bryozoa	Stenolaemata	Cystoporida	*Meekopora parilis*	1
	GRCA	Bryozoa	Stenolaemata	Cystoporida	*Meekopora* sp.	2
	MNA	Bryozoa	Stenolaemata	Cystoporida	*Meekopora* sp.	3
	Other	Bryozoa	Stenolaemata	Cystoporida	*Meekopora* sp.	1
	MNA	Bryozoa	Stenolaemata	Cystoporida	*Meekopora* sp. aff*. M. tenuis*	1
	Other	Bryozoa	Stenolaemata	Fenestrida	*Bicorbis arizonica*	1
	GRCA	Bryozoa	Stenolaemata	Fenestrida	*Chasmatopora* sp.	2
	GRCA	Bryozoa	Stenolaemata	Fenestrida	*Fenestella* sp.	1
	GRCA	Bryozoa	Stenolaemata	Fenestrida	*Polypora* aff*. P. spinulifera*	5

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	GRCA	Bryozoa	Stenolaemata	Fenestrida	*Polypora* sp.	2
	GRCA	Bryozoa	Stenolaemata	Fenestrida	*Polypora spinulifera*	4
	Other	Bryozoa	Stenolaemata	Fenestrida	*Polypora spinulifera*	1
	GRCA	Bryozoa	Stenolaemata	Fenestrida	*Septopora biserialis*	3
	Other	Bryozoa	Stenolaemata	Fenestrida	*Septopora biserialis*	1
	GRCA	Bryozoa	Stenolaemata	Fenestrida	*Septopora* sp.	9
	Other	Bryozoa	Stenolaemata	Fenestrida	*Septopora* sp.	1
	Other	Bryozoa	Stenolaemata	Fenestrida	Undetermined	1
	MNA	Bryozoa	Stenolaemata	Trepostomatida	*Bascomella subsphaerica*	3
	Other	Bryozoa	Stenolaemata	Trepostomatida	*Bascomella subsphaerica*	1
	Other	Bryozoa	Stenolaemata	Trepostomatida	*Stenodiscus* sp.	1
	MNA	Bryozoa	Stenolaemata	Trepostomatida	*Stenopora* aff. *"S. diagonalis"*	4
	MNA	Bryozoa	Stenolaemata	Trepostomatida	*Stenopora* aff. "*S. kaibabensis"*	9
	MNA	Bryozoa	Stenolaemata	Trepostomatida	*Stenopora* aff. *"S. polygona"*	5
	MNA	Bryozoa	Stenolaemata	Trepostomatida	*Stenopora intercalaris*	1
	GRCA	Bryozoa	Stenolaemata	Trepostomatida	*Stenopora* sp.	6
	Other	Bryozoa	Stenolaemata	Trepostomatida	*Stenopora* sp.	2
	Other	Bryozoa	Undetermined	Fenestrida	Undetermined	1
	MNA	Bryozoa	Undetermined	Undetermined	*Timanodyctia* sp.	1
	MNA	Bryozoa	Undetermined	Undetermined	Undetermined	5
	Other	Bryozoa	Undetermined	Undetermined	Undetermined	1
	GRCA	Cnidaria	Anthozoa	Stauriida	*Lophophyllum* sp.	12
	Other	Cnidaria	Anthozoa	Stauriida	*Lophophyllum* sp.	2
	Other	Cnidaria	Anthozoa	Stauriida	Undetermined	1
	USNM	Cnidaria	Staurozoa?	Conulatae	*Conularia kaibabensis*	1

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	GRCA	Cnidaria	Undetermined	Undetermined	Undetermined	32
	MNA	Cnidaria	Undetermined	Undetermined	Undetermined	2
	GRCA	Echinodermata	Crinoidea	Undetermined	Undetermined	79
	MNA	Echinodermata	Crinoidea	Undetermined	Undetermined	3
	Other	Echinodermata	Crinoidea	Undetermined	Undetermined	1
	GRCA	Echinodermata	Echinoidea	Not assigned	"*Echinocrinus*?" aff. *"E. coloradensis"*	1
	MNA	Echinodermata	Echinoidea	Not assigned	*Archaeocidaris* sp.	1
	Other	Echinodermata	Echinoidea	Not assigned	*Archaeocidaris* sp.	1
	GRCA	Echinodermata	Echinoidea	Not assigned	*Archaeocidaris* sp. (including *Echinocrinus* sp.)	10
	GRCA	Echinodermata	Undetermined	Undetermined	Undetermined	8
	Other	Mollusca	Bivalvia	Cardiida	*Astartella gurleyi*	1
	GRCA	Mollusca	Bivalvia	Cardiida	*Astartella* sp.	12
	GRCA	Mollusca	Bivalvia	Cardiida	*Astartella subquadrata*	10
	MNA	Mollusca	Bivalvia	Cardiida	*Astartella subquadrata*	1
	GRCA	Mollusca	Bivalvia	Cardiida	*Permophorus albequus*	4
	MNA	Mollusca	Bivalvia	Cardiida	*Permophorus albequus*	1
	GRCA	Mollusca	Bivalvia	Cardiida	*Permophorus* sp.	12
	Other	Mollusca	Bivalvia	Cardiida	*Permophorus* sp.	1
	USNM	Mollusca	Bivalvia	Cardiida	*Wilkingia wyomingensis*	1
	GRCA	Mollusca	Bivalvia	Modiomorphida	*Modiomorpha* sp.	2
	GRCA	Mollusca	Bivalvia	Myalinida	*Aviculopecten* sp.	23
	Other	Mollusca	Bivalvia	Myalinida	*Aviculopecten* sp.	3
	MNA	Mollusca	Bivalvia	Myalinida	*Aviculopecten* sp.	2
	MNA	Mollusca	Bivalvia	Myalinida	*Aviculopecten?* (*Pseudomonotis?*) sp.	1
	MNA	Mollusca	Bivalvia	Myalinida	*Myalina* (*Myalina*) sp.	2

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	GRCA	Mollusca	Bivalvia	Nuculanida	*Nucula levatiforme*	1
	GRCA	Mollusca	Bivalvia	Nuculanida	*Nuculana* sp.	1
	Other	Mollusca	Bivalvia	Nuculanida	*Nuculana?* sp.	1
	GRCA	Mollusca	Bivalvia	Nuculanida	*Yoldiella lucida*	6
	Other	Mollusca	Bivalvia	Ostreida	*Bakewellia parva*	1
	GRCA	Mollusca	Bivalvia	Ostreida	*Pteria* sp.	3
	Other	Mollusca	Bivalvia	Ostreida	*Pteria* sp.	1
	Other	Mollusca	Bivalvia	Pectinida	*Acanthopecten coloradoensis*	1
	GRCA	Mollusca	Bivalvia	Pectinida	*Acanthopecten coloradoensis*	5
	GRCA	Mollusca	Bivalvia	Pectinida	*Deltopecten caneyanus*	17
	GRCA	Mollusca	Bivalvia	Pectinida	*Deltopecten* sp.	11
	GRCA	Mollusca	Bivalvia	Pectinida	*Pecten* sp.	9
	MNA	Mollusca	Bivalvia	Pectinida	*Pseudomonotis* sp.	3
	GRCA	Mollusca	Bivalvia	Pectinida	*Pseudomonotis* sp.	1
	GRCA	Mollusca	Bivalvia	Pectinida	*Pterinopecten* sp.	1
	MNA	Mollusca	Bivalvia	Pectinida	Undetermined	1
	GRCA	Mollusca	Bivalvia	Pholadomyida	*Allorisma capax*	12
	GRCA	Mollusca	Bivalvia	Pholadomyida	*Allorisma* sp.	6
	Other	Mollusca	Bivalvia	Protobranchia	*Solemya* sp.	1
	GRCA	Mollusca	Bivalvia	Protobranchia	*Solemya trapezoides*	32
	GRCA	Mollusca	Bivalvia	Trigoniida	*Kaibabella curvilenata*	1
	GRCA	Mollusca	Bivalvia	Trigoniida	*Schizodus canalis*	1
	GRCA	Mollusca	Bivalvia	Trigoniida	*Schizodus* sp.	176
	Other	Mollusca	Bivalvia	Trigoniida	*Schizodus* sp.	1
	GRCA	Mollusca	Bivalvia	Undetermined	Undetermined	447

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	MNA	Mollusca	Bivalvia	Undetermined	Undetermined	4
	Other	Mollusca	Bivalvia	Undetermined	Undetermined	1
	Other	Mollusca	Cephalopoda	Nautilida	*Domatoceras bradyi*	1
	Other	Mollusca	Cephalopoda	Nautilida	*Domatoceras simplex*	1
	GRCA	Mollusca	Cephalopoda	Nautilida	*Domatoceras* sp.	2
	GRCA	Mollusca	Cephalopoda	Nautilida	*Metacoceras* sp.	1
	GRCA	Mollusca	Cephalopoda	Nautilida	*Stearoceras rotundatum*	5
	Other	Mollusca	Cephalopoda	Nautilida	*Stearoceras rotundatum*	1
	GRCA	Mollusca	Cephalopoda	Nautilida	*Stearoceras sanandreasense*	8
	Other	Mollusca	Cephalopoda	Nautilida	*Stearoceras sanandreasense*	1
	GRCA	Mollusca	Cephalopoda	Nautilida	*Stearoceras* sp.	13
	GRCA	Mollusca	Cephalopoda	Orthocerida	*Orthoceras* sp.	1
	Other	Mollusca	Cephalopoda	Orthocerida	*Orthoceras* sp.	1
	GRCA	Mollusca	Cephalopoda	Undetermined	Undetermined	1
	GRCA	Mollusca	Gastropoda	Bellerophontida	*Bellerophon deflectus*	24
	GRCA	Mollusca	Gastropoda	Bellerophontida	*Bellerophon majusculus*	1
	Other	Mollusca	Gastropoda	Bellerophontida	*Bellerophon majusculus*	1
	GRCA	Mollusca	Gastropoda	Bellerophontida	*Bellerophon* sp.	90
	GRCA	Mollusca	Gastropoda	Bellerophontida	*Bucanopsis* sp.	40
	GRCA	Mollusca	Gastropoda	Bellerophontida	*Euphemites aequisulcatus*	1
	Other	Mollusca	Gastropoda	Bellerophontida	*Euphemites* cf. *E. carbonarius*	1
	GRCA	Mollusca	Gastropoda	Bellerophontida	*Euphemites* sp.	36
	Other	Mollusca	Gastropoda	Bellerophontida	*Euphemites* sp.	1
	GRCA	Mollusca	Gastropoda	Bellerophontida	*Warthia* sp.	1
	GRCA	Mollusca	Gastropoda	Cycloneritida	*Naticopsis kaibabensis*	3

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	GRCA	Mollusca	Gastropoda	Cycloneritida	*Naticopsis* sp.	2
	Other	Mollusca	Gastropoda	Cycloneritida	*Naticopsis* sp.	1
	GRCA	Mollusca	Gastropoda	Euomphalina	*Euomphalus kaibabensis*	2
	MNA	Mollusca	Gastropoda	Euomphalina	*Euomphalus kaibabensis*	1
	GRCA	Mollusca	Gastropoda	Euomphalina	*Euomphalus* sp.	25
	MNA	Mollusca	Gastropoda	Euomphalina	*Euomphalus* sp.	1
	Other	Mollusca	Gastropoda	Euomphalina	*Euomphalus* sp.	1
	GRCA	Mollusca	Gastropoda	Euomphalina	*Glyptospira cristulata*	6
	GRCA	Mollusca	Gastropoda	Murchisoniina	*Ananias franciscanus*	6
	GRCA	Mollusca	Gastropoda	Murchisoniina	*Ananias franciscanus*	14
	GRCA	Mollusca	Gastropoda	Murchisoniina	*Goniospira* sp.	1
	Other	Mollusca	Gastropoda	Murchisoniina	*Goniospira* sp.	1
	GRCA	Mollusca	Gastropoda	Not assigned	*Orthonema* sp.	1
	GRCA	Mollusca	Gastropoda	Not assigned	*Orthonema striatonodosum*	4
	GRCA	Mollusca	Gastropoda	Pleurotomariida	*Murchisonia geminocarinata*	2
	Other	Mollusca	Gastropoda	Pleurotomariida	*Murchisonia?* cf. *M. terebra*	1
	GRCA	Mollusca	Gastropoda	Pleurotomariida	*Pleurotomaria* sp.	8
	Other	Mollusca	Gastropoda	Pleurotomariida	*Pleurotomaria* sp.	1
	GRCA	Mollusca	Gastropoda	Undetermined	Undetermined	1
	MNA	Mollusca	Gastropoda	Undetermined	Undetermined	1
	Other	Mollusca	Gastropoda	Undetermined	Undetermined	1
	GRCA	Mollusca	Scaphopoda	Dentaliida	*Dentalium* sp.	22
	GRCA	Mollusca	Scaphopoda	Dentaliida	*Plagioglypta* sp.	12
	GRCA	Mollusca	Scaphopoda	Dentaliida	*Prodentalium canna*	10
	Other	Mollusca	Scaphopoda	Dentaliida	*Prodentalium canna*	1

Appendix Table 5-A-4 (continued). Specimens of invertebrate body fossils from post-Supai Permian units at GRCA as reported from the Grand Canyon Museum (GRCA), Museum of Northern Arizona (MNA), Smithsonian Institution (USNM), or otherwise listed in the literature (Other). Specimens reported are listed with the stratigraphic unit as reported by the sources, if available, although some reports may have been based on superseded stratigraphic interpretations as described in the methods section.

Unit	Source	Phylum	Class	Order	Species	#
Pk (cont.)	GRCA	Mollusca	Scaphopoda	Undetermined	Undetermined	28
	Other	Mollusca	Undetermined	Undetermined	Undetermined	1
	GRCA	Porifera	Demospongea	Lithistida	*Actinocoelia maeandrina*	1
	Other	Porifera	Demospongea	Lithistida	*Actinocoelia maeandrina*	1
	MNA	Porifera	Demospongea	Lithistida	*Actinocoelia* sp.	1
	Other	Porifera	Demospongea	Lithistida	*Actinocoelia* sp.	1
	GRCA	Porifera	Undetermined	Undetermined	Undetermined	29
	MNA	Porifera	Undetermined	Undetermined	Undetermined	5
	Other	Porifera	Undetermined	Undetermined	Undetermined	1
	GRCA	Undetermined	Undetermined	Undetermined	Undetermined	477
Pu	MNA	Brachiopoda	Rhynchonellata	Spiriferida	*Squamularia* sp.	1
	USNM	Brachiopoda	Strophomenata	Orthotetida	*Meekella occidentalis*	1
	USNM	Brachiopoda	Strophomenata	Productida	*Rugatia paraindica*	1
	USNM	Bryozoa	Stenolaemata	Trepostomatida	*Rhombotrypella* sp.	1
	USNM	Mollusca	Bivalvia	Cardiida	*Wilkingia wyomingensis*	2
	USNM	Mollusca	Gastropoda	Euophalina	*Glyptospira cristulata*	8
	GRCA	Undetermined	Undetermined	Undetermined	Undetermined	27

Taxonomic Uncertainty

Taxonomic uncertainty is a common issue for museum curators and researchers. Out of more than 7,000 entries reviewed for this appendix, only about 5,500 had valid genus and/or species names that could be verified in an online, searchable database of paleontologic specimens or in the global names index (gni.globalnames.org). Most of the specimens had names that could not be verified in the databases, indicating that GRCA specimens are difficult to verify.

For this report, the following uncertainty notations were used after Bengtson (1988) and are as follows. When a specimen was originally labeled with "aff." or "cf." and the species name is not currently valid, the inventory interpretation for the genus and/or species will include the questionable name in quotes ("). The name was also revised to include the full genus name in lieu of using an initial abbreviation for clarity for all species names, including those following aff. or cf. If the original entry was noted as n. sp., this was interpreted as a new species to be defined at a future time if the name could not be confirmed by any other means. This notation was difficult to interpret if the specimen noted with n. sp. had not been formally described or listed with a reference for the source of the notation. For example, a specimen listed in an inventory as *Composita* n. sp. could be considered equivalent to sp.? or simply sp. with the second notation preferred. If a museum catalog entry had n. sp. for a museum specimen, the specimen name herein is referred to by the valid genus name followed by "sp." to indicate the uncertain interpretation of this specimen to species level. Taxonomists use "?" as a prefix or suffix identifier for a taxon name that appears similar to a genus known to the person making the entry. However, for this inventory, the question mark is appended consistently at the end of the genus or species name for ease in spreadsheet sorting filters and consistency. Taxonomists of the past were not subject to the interpretations of software protocols that use the question mark as an individual character placeholder. For example, *Composita* sp. for a museum specimen would indicate that the specimen matched the criteria for the genus, but further diagnosis to species level was not made by the author or collector. *Composita*? sp. uses a valid genus name but may not be the best genus match for this specimen. *Composita* n. sp. was found as the listed species name for many museum catalog entries but is confusing. This notation (n. sp.) would mean that the specimen does not meet the diagnosis criteria for any known species of the genus according to the definitions available at the time the specimen was cataloged. However, unless a reference was linked to the entry that actually described a new species (n. sp.), the entry was changed to sp.? for this inventory. Otherwise, the lack of a published description associated with a new species would make the name a *nomem nudum*.

Conforming to Bengtson (1988), the use of sp. is sufficient for this report of known specimens and their species interpretations. A question mark after the genus name would therefore be similar to some uses of cf. but indicates a tentative identification to that genus and is preferred. Generally, *Composita* sp. should be interpreted herein as a specimen probably of the *Composita* genus, but further identification has not been made with certainty. The specimen may be able to be refined to a species name, require a new description as n. sp., or may be too difficult to ascertain due to the preservation of the fossil. The use of cf. and aff. is also difficult to interpret as cf. means to "compare to" and aff. may mean "compare with", but both were retained as the original source applied if a species was provided for comparison.

Institutional Abbreviations—**GRCA**, Grand Canyon National Park Museum, Grand Canyon, Arizona; **MNA**, Museum of Northern Arizona, Flagstaff, Arizona; **Other**, from literature sources; **USNM**, National Museum of Natural History Paleobiology Collections, Washington, D.C.

Geologic Unit Abbreviations—**Cambrian: Ctu**, Tonto Group undifferentiated; **Ctt**, Tonto Group, Tapeats Sandstone; **Ctba**, Tonto Group, Bright Angel Shale; **Ctm**, Tonto Group, Muav Limestone; **Cu**, Cambrian undifferentiated; **Devonian: Dtb**, Temple Butte Formation; **Mississippian: Mr**, Redwall Limestone; **Msc**, Surprise Canyon Formation; **Pennsylvanian: IPsu**, Supai Group undifferentiated (most records are probably from the Watahomigi Formation); **IPswa**, Supai Group: Watahomigi Formation; **IPsm**, Supai Group: Manakacha Formation; **IPswe**, Supai Group: Wescogame Formation; **Permian: Pse**, Supai Group: Esplanade Sandstone; **Ph**, Hermit Formation; **Pt**, Toroweap Formation; **Pk**, Kaibab Formation; **Pu**, Permian undifferentiated.

Literature Cited

Bengtson, P. 1988. Open nomenclature. Palaeontology 31:223–227.

Taxonomic Notes

Because taxonomic updates have been applied, a list of changes is appended to make it simpler to translate from historic usage to what is presented here. This is by no means exhaustive. It is broken into four sections, for the four stratigraphic blocks (Cambrian, Devonian–Mississippian, Supai Group, and post-Supai Permian). The usage favored in the tables is on the right. These lists are applicable to Appendix 5-B as well as 5-A.

Cambrian

- *Alokistocare althea = Amecephalus althea*
- *Billingsella obscura = Nisusia obscura*
- *Clavaspidella kanabensis = Athabaskia kanabensis*
- *Clavaspidella enucleata* possibly = *Clavaspidella* sp. =*Athabaskia*
- *Dolichometopus productus* (in part) = *Dolichometopus tontoensis* (in part) = *Glossopleura mckeei = Glossopleura boccar*
- *Dolichometopus tontoensis* (in part) = *Anoria tontoensis*
- *Ehmaniella basilica = Proehmaniella basilica*
- *Ehmaniella hebes = Proehmaniella hebes*
- *Eocrinus = Gogia*
- *Finkelnburgia noblei = Nisusia noblei*
- *Glossopleura mckeei = Glossopleura boccar*
- *Indiana = Indianites*
- *Iphidea crenistria = Micromitra* (*Paterina*) *crenistria*
- *Iphidea superba = Micromitra* (*Paterina*) *superba*

- *Kootenia* n. sp. of Schenk and Wheeler (1942) = *Kootenia schenki*
- *Lingulella perattenuata* ("*attenuata*" of Resser 1945) = *Lingulella spatula*
- *Micromitra* (*Iphidella*) *pannula* (in part) = *Dictyonina arizonaensis*
- *Nisusia*? (*Jamesella*) *kanabensis = Nisusia kanabensis*
- *Obolus* (*Lingulella*) *chuarensis = Obolus* (*Westonia*) *chuarensis = Lingulella chuarensis*
- *Obolus* (*Lingulella*) *euglyphus = Obolus* (*Westonia*) *euglyphus = Lingulella euglypha*
- *Obolus* (*Lingulella*) *lineolatus = Lingulella lineolata*
- *Obolus* (*Lingulella*) *spatulus = Lingulella attenuata* (in part) = *Lingulella* (*Lingulepis*) *spatula = Lingulepis spatula = Lingulella spatula*
- *Obolus* (*Lingulella*) *zetus = Lingulella zetus*
- *Obolus* (*Westonia*) *themis = Lingulella themis*
- *Orthisina = Clitambonites*
- *Solenopleurella diligens = Spencella diligens*
- *Solenopleurella erosa = Spencella erosa*
- *Solenopleurella porcata = Spencella porcata*
- *Solenopleurella* n. sp. of Schenk and Wheeler (1941) = *Solenopleurella porcata = Spencella porcata*

Devonian–Mississippian

- *Caninophyllum incrassatum = Vesiculophyllum incrassatum*
- *Endothyra baileyi = Globoendothyra baileyi*
- *Endothyra baileyi poloumera = Globoendothyra baileyi poloumera*
- *Endothyra eospiroides = Inflatoendothyra eospiroides*
- *Endothyra spinosa = Spinoendothyra spinosa*
- *Endothyra torquida = Spinobrunsiina torquida*
- *Eomillerella spiroides = Eoendothyranopsis spiroides*
- *Lithostrotion* (*Diphyphyllum*) *inconstans = Dorlodotia inconstans*
- *Monilipora*? sp. = *Cladochonus* sp.
- *Phillipsia sampsoni = Aprathia* sp.
- *Phillipsia tuberculata = Phillipsia peroccidens*
- *Schuchertella chemungensis = Floweria chemungensis*
- *Septaglomospiranella primaeva = Septaglomospiranella chernoussovensis*
- *Septatournayella henbesti = Pohlia henbesti*

- *Spirifer striatus = Neospirifer striatus*
- *Straparollus* (*Euomphalus*) = *Euomphalus*
- *Triplophyllites* (*Homalophyllites*) *paucicinctus = Homalophyllites paucicinctus*

Supai Group

- *Anthracospirifer curvilateralis* subsp. *tanoensis = Anthracospirifer tanoensis*
- *Derbyia* may also be spelled *Derbya* in some references
- *Straparollus* (*Euomphalus*) = *Euomphalus*

Post-Supai Permian

- *Anisopyge mckeei = Novoameura mckeei*
- *Avonia dorsoconcava = Echinauris dorsoconcava*
- *Avonia subhorrida newberryi = Echinauris newberryi*
- *Bicorbula arizonica = Bicorbis arizonica*
- *Chonetes hillanus = Dyoros hillanus*
- *Chonetes kaibabensis = Quadrochonetes kaibabensis = Dyoros kaibabensis*
- *Chonetes* (*Lissochonetes*) *subliratus = Dyoros subliratus*
- *Chonetes quadratus = Dyoros tetragonus*
- *Delaria macclintocki = Delaria sevilloidia*
- *Dentalium canna = Plagioglypta canna = Prodentalium canna*
- *Derbyia* may also be spelled *Derbya* in some references
- *Derbyia regularis = Derbyia arizonensis*
- *Echinocrinus = Archaeocidaris* (suppressed for *Archaeocidaris*)
- *Euphemus = Euphemites*
- *Griffithides scitulus = Ditomoyge scitulus*
- *Marginifera meridionalis = Kozlowskia meridionalis = Kutorginella meridionalis*
- *Marginifera popei = Liosotella popei*
- *Leda* sp. and *Nucula* sp. of McKee 1938 = *Nuculana*? sp.
- *Pleurophorus = Permophorus*
- *Productus irginae = Waagenoconcha irginae*
- *Productus subhorridus = Avonia subhorrida = Echinauris subhorrida*
- *Productus* (*Dictyoclostus*) *bassi = Peniculauris bassi*
- *Productus* (*Dictyoclostus*) *ivesi = Dictyoclostus ivesi = Peniculauris ivesi*
- *Productus* (*Dictyoclostus*) *occidentalis = Dictyoclostus occidentalis = Rugatia occidentalis*

- *Productus* (*Dictyoclostus*) *paraindicus = Rugatia paraindicus*
- *Productus montpelierensis = Waagenoconcha montpelieriensis = Bathymyonia nevadensis*
- *Pugnoides pinguis = Wellerella pinguis = Phrenophoria pinguis*
- *Retzia meekana = Hustedia meekana*
- *Squamularia guadalupensis = Phricodothyris guadalupensis*
- *Spiriferina hilli = Spiriferellina hilli*
- *Straparollus* (*Euomphalus*) *= Euomphalus*
- *Titanoceras rotundatum = Stearoceras rotundatum*
- *Titanoceras sanandreasense = Stearoceras sanandreasense*
- *Typus gilmorei = Tupus gilmorei*

Typus whitei = Tupus whitei

Appendix 5-B. Paleozoic Invertebrate Body Fossils From GRCA From the Literature

This appendix lists Paleozoic invertebrates which have been reported from the literature from GRCA localities with their stratigraphic occurrences and references (see the main "Literature Cited" of this chapter for full citations). The same taxonomic caveats discussed in Appendix 5-A apply here. As with the tables in Appendix 5-B, the tables have been split into four to improve navigation and readability: Appendix Table 5-B-1, Cambrian; Appendix Table 5-B-2, Devonian and Mississippian; Appendix Table 5-B-3, Supai Group; and Appendix Table 5-B-4, post-Supai Permian.

Appendix Table 5-B-1. Cambrian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Ctu	Arthropoda	Ostracoda	Undetermined	Walcott 1890
	Arthropoda	Trilobita	*Anoria* sp.	Walcott 1890
	Arthropoda	Trilobita	*Athabaskia kanabensis*	Resser 1945
	Arthropoda	Trilobita	*Elrathiella?* aff. *E. insueta*	Resser 1945
	Arthropoda	Trilobita	*Glossopleura boccar*	Palmer E&R 1963/10/17, Bonde et al. 2018
	Arthropoda	Trilobita	*Glossopleura* sp.	Palmer E&R 1963/10/17, Bonde et al. 2018
	Arthropoda	Trilobita	*Glyphaspis kwaguntensis*	Resser 1945
	Arthropoda	Trilobita	*Kochina angustata*	Resser 1945
	Arthropoda	Trilobita	*Kootenia mckeei*	Resser 1945
	Arthropoda	Trilobita	*Kootenia* sp.	Resser 1945
	Arthropoda	Trilobita	*Olenoides* sp.	Walcott 1890
	Arthropoda	Trilobita	*Parehmania kwaguntensis*	Bonde et al. 2018
	Arthropoda	Trilobita	*Parehmania nitida*	Resser 1945
	Arthropoda	Trilobita	*Proehmaniella hebes*	Resser 1945
	Arthropoda	Trilobita	*Ptarmigania* sp.	Resser 1945
	Arthropoda	Trilobita	*Spencella diligens*	Resser 1945
	Arthropoda	Trilobita	*Spencella* sp.	Resser 1945
	Arthropoda	Trilobita	*Trachycheilus typicale*	Resser 1945
	Brachiopoda	Kutorginata	*Nisusia kanabensis*	Walcott 1908, 1912a, Resser 1945
	Brachiopoda	Lingulata	*Acrothele* sp.	Walcott 1890
	Brachiopoda	Lingulata	*Lingulella* aff. *L. winona* subsp. *convexa*	Walcott 1912a
	Brachiopoda	Lingulata	*Lingulella kanabensis*	Resser 1945
	Brachiopoda	Lingulata	*Lingulella* sp.	Walcott 1890, Resser 1945
	Brachiopoda	Lingulata	*Lingulella zetus*	Walcott 1898, 1912a, 1916a, Resser 1945
	Brachiopoda	Paterinata	*Iphidea ornatella*	Walcott 1890
	Brachiopoda	Paterinata	*Micromitra (Paterina) crenistria*	Walcott 1897, 1912a, 1916a, Resser 1945
	Brachiopoda	Rhynchonellata	*Protorthis* sp.	Walcott 1912a
	Problematica	Hyolitha	*Hyolithes* sp.	Walcott 1890, Resser 1945
Ctt	Arthropoda	Trilobita	*Olenellus* sp.	Resser 1945
	Brachiopoda	Obolellata	Undetermined	Schuchert 1918a
Ctba	Arthropoda	Bradoriida clade	*Bradoria tontoensis*	Ulrich & Bassler 1931, Resser 1945

Appendix Table 5-B-1 (continued). Cambrian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Ctba (cont.)	Arthropoda	Bradoriida clade	*Dielymella* aff. *D. recticardinalis* subsp. *angustata*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Dielymella appressa*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Dielymella dorsalis*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Dielymella nasuta*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Dielymella recticardinalis*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Indianites curtus*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Indianites faba*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Indianites impressus*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Indianites intermedius*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella apicalis*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella breviuscula*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella concentrica*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella leperditoides*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella limatula*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella longula*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella nitida*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella oblonga*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella obsoleta*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella pulchella*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella scitula*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella subtruncata*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Bradoriida clade	*Walcottella ventrosa*	Ulrich & Bassler 1931, Resser 1945
	Arthropoda	Trilobita	*Acrocephalops?* cf. *A. arizonaensis*	Resser 1945
	Arthropoda	Trilobita	*Albertella schenki*	Resser 1945
	Arthropoda	Trilobita	*Albertella* sp.	Schenk & Wheeler 1942; Bonde et al. 2018
	Arthropoda	Trilobita	*Alokistocare lepida*	Schenk & Wheeler 1942
	Arthropoda	Trilobita	*Alokistocare* sp.	Schenk & Wheeler 1942, Resser 1945
	Arthropoda	Trilobita	*Alokistocare* sp. or *Ehmaniella* sp.	Bonde et al. 2018
	Arthropoda	Trilobita	*Alokistocare?* sp.	Schenk & Wheeler 1942
	Arthropoda	Trilobita	*Amecephalus althea*	Walcott 1916a, 1916b, Schuchert 1918a, Noble 1922, Schenk & Wheeler 1942, Resser 1945, Foster 2011

Appendix Table 5-B-1 (continued). Cambrian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Ctba (cont.)	Arthropoda	Trilobita	*Amecephalus* cf. *A. althea*	Schenk & Wheeler 1942
	Arthropoda	Trilobita	*Amecephalus* cf. *A. packi*	Foster 2011
	Arthropoda	Trilobita	*Anoria* sp.	Schenk & Wheeler 1942, Resser 1945
	Arthropoda	Trilobita	*Anoria tontoensis*	Walcott 1916a, Schuchert 1918a, Noble 1922, Resser 1945, Foster 2011
	Arthropoda	Trilobita	*Antagmus arizonaensis*	Resser 1945, Bonde et al. 2018
	Arthropoda	Trilobita	*Athabaskia* sp.	Resser 1945
	Arthropoda	Trilobita	*Ehmaniella* aff. "*E. arizonaensis*"	Resser 1945
	Arthropoda	Trilobita	*Ehmaniella* sp.	Bonde et al. 2018
	Arthropoda	Trilobita	*Elrathia nitens*	Resser 1945
	Arthropoda	Trilobita	*Elrathia* sp.	Schenk & Wheeler 1942, Resser 1945
	Arthropoda	Trilobita	*Glossopleura* aff. *"G. walcotti"*	Foster 2011
	Arthropoda	Trilobita	*Glossopleura boccar*	Walcott 1916a, Schuchert 1918a, Noble 1922, Resser 1935, 1945, Schenk & Wheeler 1942, Palmer E&R 1963/10/17, Foster 2011, Bonde et al. 2018
	Arthropoda	Trilobita	*Glossopleura* sp.	Resser 1945, Elliott & Martin 1987
	Arthropoda	Trilobita	*Glyphaspis* sp.	Schenk & Wheeler 1942, Resser 1945
	Arthropoda	Trilobita	*Glyphaspis vulsa*	Resser 1945
	Arthropoda	Trilobita	*Kootenia simplex*	Resser 1945
	Arthropoda	Trilobita	*Kootenia* sp.	Foster 2011
	Arthropoda	Trilobita	*Olenellus* sp.	Schenk & Wheeler 1942, Resser 1945
	Arthropoda	Trilobita	*Olenoides* sp.	Schenk & Wheeler 1942, Resser 1945
	Arthropoda	Trilobita	*Pachyaspis fonticola*	Resser 1945
	Arthropoda	Trilobita	*Pachyaspis moorei*	Resser 1945, Bonde et al. 2018
	Arthropoda	Trilobita	*Pachyaspis* sp.	Resser 1945
	Arthropoda	Trilobita	*Parehmania kwaguntensis*	Bonde et al. 2018
	Arthropoda	Trilobita	*Parehmania* sp.	Resser 1945
	Arthropoda	Trilobita	*Parehmania tontoensis*	Resser 1945
	Arthropoda	Trilobita	*Proehmaniella basilica*	Bonde et al. 2018
	Arthropoda	Trilobita	*Ptarmigania* sp.	Resser 1945
	Arthropoda	Trilobita	*Spencia tontoensis*	Resser 1945
	Arthropoda	Trilobita	Undetermined	Palmer E&R 1963/10/17
	Arthropoda	Trilobita	Undetermined (Olenellidae)	Bonde et al. 2018

Appendix Table 5-B-1 (continued). Cambrian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Ctba (cont.)	Arthropoda	Trilobita	*Zacanthoides* cf. *Z. walapai*	Resser 1945
	Arthropoda	Trilobita	*Zacanthoides* sp.	Resser 1945
	Brachiopoda	Kutorginata	*Nisusia obscura*	Walcott 1905, 1912a, 1916a, Resser 1945
	Brachiopoda	Kutorginata	*Nisusia* sp.	Schenk & Wheeler 1942
	Brachiopoda	Kutorginata	*Nisusia?* sp.	Resser 1945
	Brachiopoda	Lingulata	*Lingula* sp.	Strother & Beck 2000
	Brachiopoda	Lingulata	*Lingulella acutangulus*	Walcott 1912a, Noble 1922
	Brachiopoda	Lingulata	*Lingulella chuarensis*	Walcott 1898, 1912a, 1916a, Schuchert 1918a, Noble 1922, Resser 1945, Foster 2011
	Brachiopoda	Lingulata	*Lingulella euglypha*	Walcott 1898, 1912a, 1916a, Resser 1945
	Brachiopoda	Lingulata	*Lingulella lineolata*	Walcott 1898, 1912a, 1916a, Noble 1922, Resser 1945
	Brachiopoda	Lingulata	*Lingulella mckeei*	Resser 1945, Foster 2011, Bonde et al. 2018
	Brachiopoda	Lingulata	*Lingulella* sp.	Walcott 1890, Resser 1945
	Brachiopoda	Lingulata	*Lingulella spatula*	Walcott 1902, 1912a, Noble 1922, Resser 1945
	Brachiopoda	Lingulata	*Lingulella zetus*	Walcott 1898, 1912a, 1916a, Resser 1945
	Brachiopoda	Lingulata	*Lingulella?* aff. *L. themis*	Walcott 1905, 1912a, Noble 1922, Resser 1945
	Brachiopoda	Lingulata	*Lingulella?* cf. *L. monticula*	Frech 1893
	Brachiopoda	Obolellata	*Obolella* aff. *O. polita*	Frech 1893
	Brachiopoda	Obolellata	*Obolella* sp.	Frech 1893
	Brachiopoda	Paterinata	*Dictyonina arizonaensis*	Walcott 1912a, Noble 1922, Resser 1945, Bonde et al. 2018
	Brachiopoda	Paterinata	*Micromitra (Paterina) crenistria*	Walcott 1897, 1912a, 1916a, Resser 1945
	Brachiopoda	Paterinata	*Micromitra (Paterina) superba*	Walcott 1897, 1912a, 1916a, Noble 1922, Resser 1945
	Brachiopoda	Paterinata	*Micromitra pealei*	Walcott 1912a, 1916a
	Brachiopoda	Paterinata	*Paterina?* sp.	Resser 1945
	Brachiopoda	Rhynchonellata	*Diraphora?* sp.	Resser 1945
	Brachiopoda	Undetermined	Undetermined	McKee 1945
	Echinodermata	Eocrinoidea	*Eocystites* sp.	Walcott 1916a, Noble 1922
	Echinodermata	Eocrinoidea	*Gogia multibrachiatus*	Resser 1945, Foster 2011
	Echinodermata	Eocrinoidea	*Gogia?* aff. *G. longidactylus*	Foster 2011
	Porifera	Archaeocyatha	Undetermined	Resser 1945
	Problematica	Coeloscleritophora	*Chancelloria* sp.	Elliott & Martin 1987

Appendix Table 5-B-1 (continued). Cambrian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Ctba (cont.)	Problematica	Hyolitha	*Hyolithes* sp.	Walcott 1916a, Noble 1922, Schenk & Wheeler 1942, Resser 1945, Foster 2011, Bonde et al. 2018
	Problematica	Undetermined	*Margaretia?* sp.	Schenk & Wheeler 1942
	Problematica	Undetermined	*Tontoia kwaguntensis*	Walcott 1912b, Resser 1945
Ctm	Arthropoda	Trilobita	*Alokistocare* sp.	Schenk & Wheeler 1942, Resser 1945
	Arthropoda	Trilobita	*Anomocarella* sp.	Noble 1922
	Arthropoda	Trilobita	*Anoria* sp.	Schenk & Wheeler 1942, Resser 1945
	Arthropoda	Trilobita	*Athabaskia* sp.	Resser 1945
	Arthropoda	Trilobita	*Bathyurus?* sp.	Noble 1922
	Arthropoda	Trilobita	*Bolaspis aemula*	Resser 1945
	Arthropoda	Trilobita	*Bolaspis*? sp.	Resser 1945
	Arthropoda	Trilobita	*Dorypyge* sp.	Stoyanow 1936
	Arthropoda	Trilobita	*Glossopleura* sp.	Schenk & Wheeler 1942, Resser 1945, Palmer E&R 1963/10/17, Bonde et al. 2018
	Arthropoda	Trilobita	*Glossopleura* sp. or *Anoria* sp.	Resser 1945
	Arthropoda	Trilobita	*Glyphaspis* sp.	Schenk & Wheeler 1942, Resser 1945
	Arthropoda	Trilobita	*Glyphaspis tecta*	Resser 1945
	Arthropoda	Trilobita	*Glyphaspis?* sp.	Resser 1945
	Arthropoda	Trilobita	*Kootenia havasuensis*	Resser 1945
	Arthropoda	Trilobita	*Kootenia mckeei*	Resser 1945
	Arthropoda	Trilobita	*Kootenia schenki*	Schenk & Wheeler 1942, Resser 1945, Palmer E&R 1963/10/17, Bonde et al. 2018
	Arthropoda	Trilobita	*Kootenia simplex*	Resser 1945
	Arthropoda	Trilobita	*Kootenia* sp.	Schenk & Wheeler 1942, Resser 1945, Palmer E&R 1963/10/17
	Arthropoda	Trilobita	*Kootenia?* sp.	Resser 1945
	Arthropoda	Trilobita	*Neolenus* sp.	Noble 1922
	Arthropoda	Trilobita	*Pachyaspis moorei*	Resser 1945, Bonde et al. 2018
	Arthropoda	Trilobita	*Pagodia?* sp.	Noble 1922
	Arthropoda	Trilobita	*Ptychoparia* sp.	Walcott 1883, 1890, Noble 1922
	Arthropoda	Trilobita	*Ptychoparia?* sp.	Noble 1922
	Arthropoda	Trilobita	*Saukia* sp.	Noble 1922

Appendix Table 5-B-1 (continued). Cambrian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Ctm (cont.)	Arthropoda	Trilobita	*Spencella erosa*	Resser 1945
	Arthropoda	Trilobita	*Spencella porcata*	Schenk & Wheeler 1942, Resser 1945, Palmer E&R 1963/10/17, Bonde et al. 2018
	Arthropoda	Trilobita	*Spencella* sp.	Resser 1945
	Arthropoda	Trilobita	Undetermined	Stoyanow 1936, Palmer E&R 1963/10/17
	Arthropoda	Trilobita	*Zacanthoides* sp.	Resser 1945
	Brachiopoda	Kutorginata	*Nisusia* aff. *N. noblei*	Resser 1945
	Brachiopoda	Kutorginata	*Nisusia noblei*	Walcott 1924, Resser 1945
	Brachiopoda	Kutorginata	*Nisusia* sp.	Palmer E&R 1963/10/17, Bonde et al. 2018
	Brachiopoda	Kutorginata	*Nisusia?* sp.	Palmer E&R 1963/10/17
	Brachiopoda	Lingulata	*Acrotreta?* sp.	Noble 1922
	Brachiopoda	Lingulata	*Discina* sp.	Noble 1922
	Brachiopoda	Lingulata	*Lingula* sp.	Noble 1922
	Brachiopoda	Lingulata	*Lingulella (Lingulepis)* sp.	Walcott 1883, 1890, Noble 1922
	Brachiopoda	Lingulata	*Lingulepis prima*	Noble 1922
	Brachiopoda	Lingulata	*Trematis* sp.	Noble 1922
	Brachiopoda	Paterinata	*Dictyonina* sp.	Resser 1945
	Brachiopoda	Rhynchonellata	*Finkelnburgia* sp.	Noble 1922
	Brachiopoda	Rhynchonellata	*Syntrophia* sp.	Noble 1922
	Brachiopoda	Strophomenata	*Clitambonites* sp.	Noble 1922
	Brachiopoda	Undetermined	Undetermined	McKee 1945
	Mollusca	Helcionelloida	*Helcionella* sp.	Palmer E&R 1963/10/17, Bonde et al. 2018
	Porifera	Archaeocyatha	Undetermined	Resser 1945
	Problematica	Coeloscleritophora	*Chancelloria* cf. *C. eros*	Schenk & Wheeler 1942
	Problematica	Coeloscleritophora	*Chancelloria?* sp.	Resser 1945
	Problematica	Hyolitha	*Hyolithes* aff. *H. primordialis*	Noble 1922
	Problematica	Hyolitha	*Hyolithes* sp.	Noble 1922, Schenk & Wheeler 1942, Resser 1945
	Problematica	Hyolitha	*Hyolithes?* sp.	Resser 1945
	Problematica	Monoplacophora?	*Scenella hermitensis*	Resser 1945

Appendix Table 5-B-2. Devonian and Mississippian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Dtb	Brachiopoda	Undetermined	Undetermined	Walcott 1883, Stoyanow 1936
	Cnidaria	Anthozoa	*Streptelasma?* sp.	Schenk & Wheeler 1942
	Cnidaria	Anthozoa	Undetermined (“cup corals”)	Noble 1922
	Cnidaria	Anthozoa	Undetermined (“cyathophylloids”)	Walcott 1883
	Mollusca	Gastropoda	Undetermined	Walcott 1883
	Undetermined	Undetermined	Undetermined	McKee 1969, McKee & Gutschick 1969b, 1969e
Mr	Arthropoda	Trilobita	*Aprathia* sp.	McKee & Gutschick 1969d, 1969f, Brezinski 2017
	Arthropoda	Trilobita	*Breviphillipsia* sp.	Cisne 1971
	Brachiopoda	Rhynchonellata	*Camarotoechia* sp.	Noble 1922
	Brachiopoda	Rhynchonellata	*Camarotoechia?* sp.	Noble 1922
	Brachiopoda	Rhynchonellata	*Composita?* sp.	Noble 1922
	Brachiopoda	Rhynchonellata	*Dielasma* sp.	Carter et al. 2014
	Brachiopoda	Rhynchonellata	*Mirifusella* cf. *M. fortunata*	Carter et al. 2014
	Brachiopoda	Rhynchonellata	*Neospirifer striatus*	Frech 1893
	Brachiopoda	Rhynchonellata	*Prospira* sp.	Carter et al. 2014
	Brachiopoda	Rhynchonellata	*Pugnoides* sp.	Billingsley & McKee 1982
	Brachiopoda	Rhynchonellata	*Rhipidomella* sp.	Noble 1922
	Brachiopoda	Rhynchonellata	*Spirifer* aff. *S. incertus*	Noble 1922
	Brachiopoda	Rhynchonellata	*Spirifer centronatus*	Schuchert 1918b, Noble 1922
	Brachiopoda	Rhynchonellata	*Spirifer* cf. *Elivina occidentalis*	Grant E&R 1963/10/16, Bonde et al. 2018
	Brachiopoda	Rhynchonellata	*Spirifer redwallensis*	Carter et al. 2014
	Brachiopoda	Rhynchonellata	*Spirifer* sp.	Schuchert 1918b, Grant E&R 1963/10/16, McKee & Gutschick 1969d
	Brachiopoda	Rhynchonellata	*Spirifer* sp.	Grant E&R 1963/10/16
	Brachiopoda	Rhynchonellata	*Syringothyris?* sp.	Noble 1922, Grant E&R 1963/10/16, Bonde et al. 2018
	Brachiopoda	Rhynchonellata	*Unispirifer minnewankensis*	Carter et al. 2014
	Brachiopoda	Rhynchonellata	Undetermined	McKee & Gutschick 1969b, 1969d
	Brachiopoda	Strophomenata	*Floweria chemungensis*	Noble 1922

Appendix Table 5-B-2 (continued). Devonian and Mississippian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Mr (cont.)	Brachiopoda	Strophomenata	*Leptagonia* sp.	McKee & Gutschick 1969b
	Brachiopoda	Strophomenata	*Tomiproductus gallatinensis*	Carter et al. 2014
	Brachiopoda	Strophomenata	Undetermined	McKee & Gutschick 1969b, 1969d, 1969e
	Brachiopoda	Undetermined	Undetermined	McKee & Gutschick 1969b, 1969d
	Bryozoa	Stenolaemata	*Cystodictya* sp.	McKee & Gutschick 1969d
	Bryozoa	Stenolaemata	*Dichotrypa* sp.	Duncan 1969, McKee & Gutschick 1969d
	Bryozoa	Stenolaemata	*Fenestella* sp.	McKee & Gutschick 1969d
	Bryozoa	Stenolaemata	*Fenestralia?* sp.	McKee & Gutschick 1969d
	Bryozoa	Stenolaemata	*Penniretepora* sp.	McKee & Gutschick 1969d
	Bryozoa	Stenolaemata	*Polypora* sp.	Schuchert 1918b, McKee & Gutschick 1969d
	Bryozoa	Stenolaemata	*Polypora*? sp.	McKee & Gutschick 1969d
	Bryozoa	Stenolaemata	Undetermined	McKee & Gutschick 1969d
	Bryozoa	Undetermined	Undetermined	McKee & Gutschick 1969e
	Cnidaria	Anthozoa	*Amplexizaphrentis* sp.	Sando & Bamber 1985
	Cnidaria	Anthozoa	*Cladochonus?* sp.	Noble 1922
	Cnidaria	Anthozoa	*Clisiophyllum* sp.	Schuchert 1918b
	Cnidaria	Anthozoa	*Diphyphyllum?* cf. *Lithostrotion* sp.	Noble 1922
	Cnidaria	Anthozoa	*Diphyphyllum?* sp.	Noble 1922
	Cnidaria	Anthozoa	*Dorlodotia* cf. *D. inconstans*	Easton & Gutschick 1953
	Cnidaria	Anthozoa	*Dorlodotia inconstans*	Easton & Gutschick 1953
	Cnidaria	Anthozoa	*Dorlodotia* sp.	Sando & Bamber 1985
	Cnidaria	Anthozoa	*Homalophyllites paucicinctus*	Easton & Gutschick 1953
	Cnidaria	Anthozoa	*Homalophyllites subcrassus*	McKee & Gutschick 1969d, Sando 1969
	Cnidaria	Anthozoa	*Menophyllum excavatum*	Schuchert 1918b
	Cnidaria	Anthozoa	*Sychnoelasma* sp.	Sando & Bamber 1985
	Cnidaria	Anthozoa	*Syringopora aculeata*	McKee & Gutschick 1969d, Sando 1969
	Cnidaria	Anthozoa	*Syringopora* aff. *S. surcularia*	Sando E&R 1963/10/14

Appendix Table 5-B-2 (continued). Devonian and Mississippian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Mr (cont.)	Cnidaria	Anthozoa	*Syringopora* cf. *S. surcularia*	Schuchert 1918b
	Cnidaria	Anthozoa	*Syringopora* sp.	McKee & Gutschick 1969d, Sando & Bamber 1985
	Cnidaria	Anthozoa	*Syringopora surcularia*	Sando E&R 1963/10/14, Bonde et al. 2018
	Cnidaria	Anthozoa	Undetermined	McKee & Gutschick 1969d, 1969e
	Cnidaria	Anthozoa	*Vesiculophyllum incrassatum*	Easton & Gutschick 1953, Sando E&R 1963/10/14, 1969, McKee & Gutschick 1969d, Bonde et al. 2018
	Cnidaria	Anthozoa	*Vesiculophyllum* sp.	Sando & Bamber 1985
	Cnidaria	Anthozoa	*Zaphrentites persimilis*	Sando E&R 1963/10/14, Bonde et al. 2018
	Echinodermata	Blastoidea	*Pentremites* sp.	McKee & Gutschick 1969d, Macurda 1969
	Echinodermata	Crinoidea	Undetermined	Frech 1893, Schuchert 1918b, Noble 1922, McKee & Gutschick 1969b, 1969d, 1969e, Beus 1987
	Foraminifera	Fusulinata	*Calcisphaera* sp.	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Earlandia* sp.	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Endothyra* aff. *E. gutschicki*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Endothyra* aff. *E. tantala*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Endothyra* aff. *E. trachida*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Endothyra kleina*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Endothyra* sp.	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Endothyra tantula*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Endothyra trachida*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Eoendothyranopsis* aff. *E. spiroides*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Eoendothyranopsis spiroides*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Globoendothyra baileyi*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Globoendothyra baileyi* subsp. *poloumera*	McKee & Gutschick 1969d, Skipp 1969
	Foraminifera	Fusulinata	*Inflatoendothyra eospiroides*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Paracaligella?* sp.	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Paramillerella?* sp.	McKee & Gutschick 1969d

Appendix Table 5-B-2 (continued). Devonian and Mississippian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Mr (cont.)	Foraminifera	Fusulinata	*Pohlia henbesti*	Skipp et al. 1966, McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Septabrunsiina* (*Spinobrunsiina*) *parakrainica*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Septabrunsiina* sp.	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Septaglomospiranella chernoussovensis*	Skipp et al. 1966, McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Septaglomospiranella rossi*	McKee & Gutschick 1969d, Skipp 1969
	Foraminifera	Fusulinata	*Septaglomospiranella* sp.	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Septatournayella*? sp.	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Spinobrunsiina* aff. *S. torquida*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Spinobrunsiina torquida*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Spinoendothyra* aff. *S. spinosa*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Spinoendothyra spinosa*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Spiroplectamminoides?* cf. *Spiroplectammina parva*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Tournayella* sp.	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Tournayella?* sp.	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Tuberendothyra paratumula*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Tuberendothyra* sp.	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Tuberendothyra* sp.*?*	McKee & Gutschick 1969d
	Foraminifera	Fusulinata	*Tuberendothyra tuberculata*	McKee & Gutschick 1969d
	Mollusca	Cephalopoda	*Rayonnoceras* sp.	Breed 1969
	Mollusca	Gastropoda	*Bellerophon* sp.	Noble 1922
	Mollusca	Gastropoda	*Euconospira* aff. *"E. montezuma"*	Bonde et al. 2018
	Mollusca	Gastropoda	*Euomphalus?* sp.	Noble 1922
	Mollusca	Gastropoda	*Euphemites?* sp.	Bonde et al. 2018
	Mollusca	Gastropoda	Undetermined	McKee & Gutschick 1969d, Yochelson 1969
	Undetermined	Undetermined	Undetermined	McKee 1969, McKee & Gutschick 1969b, 1969e

Appendix Table 5-B-2 (continued). Devonian and Mississippian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
	Arthropoda	Trilobita	*Paladin* cf. *P. chesterensis*	Beus 1999
	Arthropoda	Trilobita	*Paladin* sp.	Gordon E&R 1979/1/23, Beus 1999
	Brachiopoda	Rhynchonellata	*Anthracospirifer* aff. *A. curvilateralis*	Billingsley & McKee 1982
	Brachiopoda	Rhynchonellata	*Anthracospirifer bifurcatus*	Gordon E&R 1973/1/3
	Brachiopoda	Rhynchonellata	*Anthracospirifer* cf. *A. curvilateralis*	Gordon E&R 1979/1/23
	Brachiopoda	Rhynchonellata	*Anthracospirifer curvilateralis*	Beus 1999
	Brachiopoda	Rhynchonellata	*Anthracospirifer* sp.	Gordon E&R 1979/1/23, Billingsley & McKee 1982
	Brachiopoda	Rhynchonellata	*Beecheria* cf. *B. arkansanum*	Beus 1999
	Brachiopoda	Rhynchonellata	*Beecheria* sp.	Billingsley & McKee 1982
	Brachiopoda	Rhynchonellata	*Brachythyris (Spirifer) subcardiiformis*	Gordon E&R 1973/1/3
	Brachiopoda	Rhynchonellata	*Cleiothyridina* sp.	Beus 1999
	Brachiopoda	Rhynchonellata	*Composita gibbosa*	Billingsley & McKee 1982
	Brachiopoda	Rhynchonellata	*Composita laevis*	Beus 1999
Msc	Brachiopoda	Rhynchonellata	*Composita ovata*	Billingsley & McKee 1982
	Brachiopoda	Rhynchonellata	*Composita* sp.	Gordon E&R 1979/1/23
	Brachiopoda	Rhynchonellata	*Composita subquadrata*	Beus 1999
	Brachiopoda	Rhynchonellata	*Cranaena* sp.	Gordon E&R 1979/1/23
	Brachiopoda	Rhynchonellata	*Eumetria* sp.	Beus 1999
	Brachiopoda	Rhynchonellata	*Flexaria* sp.	Beus 1999
	Brachiopoda	Rhynchonellata	*Leiorhynchoidea carbonifera*	Beus 1999
	Brachiopoda	Rhynchonellata	*Leiorhynchoidea* sp.	Gordon E&R 1979/1/23
	Brachiopoda	Rhynchonellata	*Macropotamorhynchus* cf. *M. purduei*	Beus 1999
	Brachiopoda	Rhynchonellata	*Pugnoides* sp.	Billingsley & McKee 1982
	Brachiopoda	Rhynchonellata	*Rhipidomella nevadensis*	Beus 1999
	Brachiopoda	Rhynchonellata	*Rotaia neogenes*	Beus 1999
	Brachiopoda	Rhynchonellata	*Schizophoria* sp.	Billingsley & McKee 1982
	Brachiopoda	Rhynchonellata	*Torynifer* aff. *T. setiger*	Billingsley & McKee 1982

Appendix Table 5-B-2 (continued). Devonian and Mississippian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Msc (cont.)	Brachiopoda	Rhynchonellata	*Torynifer* sp.	Beus 1999
	Brachiopoda	Rhynchonellata	Undetermined	Billingsley & Beus 1999a, Beus 1999, Bonde et al. 2018
	Brachiopoda	Strophomenata	*Inflatia* aff. *I. clydensis*	Gordon E&R 1979/1/23, Billingsley & McKee 1982
	Brachiopoda	Strophomenata	*Inflatia* sp.	Gordon E&R 1979/1/23
	Brachiopoda	Strophomenata	*Inflatia* sp. or *Sandia* sp.	Billingsley & McKee 1982
	Brachiopoda	Strophomenata	*Inflatia?* sp.	Gordon E&R 1973/1/3
	Brachiopoda	Strophomenata	*Orthotetes* sp.	Beus 1999
	Brachiopoda	Strophomenata	*Ovatia* sp.	Gordon E&R 1979/1/23, Billingsley & McKee 1982
	Brachiopoda	Strophomenata	Undetermined	Billingsley & McKee 1982, Beus 1999, Billingsley & Beus 1999a
	Brachiopoda	Undetermined	Undetermined	Billingsley & McKee 1982, Billingsley & Beus 1985, 1999a, Hodnett & Elliott 2018
	Bryozoa	Fenestrida	*Archimedes* sp.	Beus 1999
	Bryozoa	Undetermined	Undetermined	Billingsley & Beus 1985, 1999a, Beus 1999, Hodnett & Elliott 2018
	Cnidaria	Anthozoa	*Amplexus* sp.	Beus 1999
	Cnidaria	Anthozoa	*Barytichisma* sp.	Beus 1999
	Cnidaria	Anthozoa	*Michelinia* sp.	Billingsley & McKee 1982
	Cnidaria	Anthozoa	Undetermined	Billingsley & Beus 1985, Hodnett & Elliott 2018
	Cnidaria	Anthozoa	Undetermined (Rugosa)	Billingsley & Beus 1999a
	Echinodermata	Asteroidea	Undetermined	Beus 1999
	Echinodermata	Blastoidea	*Pentremites* sp.	Beus 1999
	Echinodermata	Crinoidea	*Cymbiocrinus* sp.	Beus 1999
	Echinodermata	Crinoidea	Undetermined	Beus 1999, Billingsley and Beus 1999a, Hodnett & Elliott 2018
	Echinodermata	Echinoidea	Undetermined	Billingsley & Beus 1999a
	Echinodermata	Undetermined	Undetermined	Billingsley & Beus 1985, 1999a
	Echinodermata	Undetermined	Undetermined (Pelmatozoa)	Billingsley & McKee 1982

Appendix Table 5-B-2 (continued). Devonian and Mississippian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Msc (cont.)	Foraminifera	Undetermined	Undetermined	Gordon E&R 1979/1/23
	Mollusca	Bivalvia	*Aviculopecten* sp.	Gordon E&R 1979/1/23
	Mollusca	Bivalvia	*Edmondia* sp.	Beus 1999
	Mollusca	Bivalvia	*Schizodus* sp.	Billingsley & McKee 1982
	Mollusca	Bivalvia	*Septimyalina* sp.	Beus 1999
	Mollusca	Bivalvia	*Septimyalina*? sp.	Gordon E&R 1979/1/23
	Mollusca	Bivalvia	Undetermined	Billingsley & McKee 1982
	Mollusca	Bivalvia	Undetermined	Billingsley & McKee 1982
	Mollusca	Gastropoda	*Bellazona* sp.	Beus 1999
	Mollusca	Gastropoda	*Bellerophon* (*Bellerophon*) sp.	Gordon E&R 1979/1/23, Billingsley & McKee 1982
	Mollusca	Gastropoda	*Bellerophon* sp.	Beus 1999
	Mollusca	Gastropoda	*Euomphalus* sp.	Gordon E&R 1979/1/23
	Mollusca	Gastropoda	*Glabrocingulum* sp.	Beus 1999
	Mollusca	Gastropoda	*Loxonema* sp.	Beus 1999
	Mollusca	Gastropoda	*Straparollus?* sp.	Billingsley & McKee 1982
	Mollusca	Undetermined	Undetermined	Billingsley & Beus 1985
	Undetermined	Undetermined	Undetermined	Billingsley & Beus 1985, Beus 1995

Appendix Table 5-B-3. Supai Group invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
IPswa	Arthropoda	Trilobita	*Paladin* sp.	Gordon 1982, McKee 1982a
	Arthropoda	Trilobita	*Paladin?* aff. librogenae	Gordon 1982, McKee 1982a
	Arthropoda	Trilobita	*Paladin?* sp.	Gordon 1982
	Brachiopoda	Lingulata	*Lingula* sp.	Gordon 1982, McKee 1982a, 1982b
	Brachiopoda	Lingulata	*Orbiculoidea* aff. *O. meekana*	Gordon 1982
	Brachiopoda	Lingulata	*Orbiculoidea meekana*	Gordon 1982, McKee 1982a
	Brachiopoda	Lingulata	*Orbiculoidea* sp.	McKee 1982b

Appendix Table 5-B-3 (continued). Supai Group invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
IPswa (cont.)	Brachiopoda	Rhynchonellata	*Anthracospirifer newberryi*	Gordon 1982, McKee 1982a
	Brachiopoda	Rhynchonellata	*Anthracospirifer tanoensis*	Gordon 1982, McKee 1982a
	Brachiopoda	Rhynchonellata	*Composita ovata*	Gordon 1982, McKee 1982a
	Brachiopoda	Rhynchonellata	*Composita* sp.	Gordon 1982, McKee 1982a, 1982b
	Brachiopoda	Rhynchonellata	*Composita subtilita*	Gordon 1982, McKee 1982a
	Brachiopoda	Rhynchonellata	*Cupularostrum?* sp.	Gordon 1982
	Brachiopoda	Rhynchonellata	*Punctospirifer transversus*	Gordon 1982, McKee 1982a
	Brachiopoda	Rhynchonellata	*Reticulariina gonionota*	Gordon 1982, McKee 1982a
	Brachiopoda	Rhynchonellata	*Schizophoria* aff. S. *altirostris*	Gordon 1982
	Brachiopoda	Rhynchonellata	*Schizophoria altirostris*	Gordon 1982, McKee 1982a
	Brachiopoda	Rhynchonellata	*Spirifer* sp.	McKee 1982b
	Brachiopoda	Rhynchonellata	*Spiriferina?* sp.	McKee 1982b
	Brachiopoda	Strophomenata	*Derbyia* aff. *D. robusta*	Gordon 1982
	Brachiopoda	Strophomenata	*Derbyia* sp.	McKee 1982a, 1982b
	Brachiopoda	Strophomenata	*Orthotetes* sp.	Gordon 1982, McKee 1982a
	Brachiopoda	Strophomenata	Undetermined	McKee 1982a, 1982b
	Bryozoa	Stenolaemata	*Cystiodictya?* sp.	Gordon 1982, McKee 1982a
	Bryozoa	Stenolaemata	*Fenestella* sp.	Gordon 1982, McKee 1982a
	Bryozoa	Stenolaemata	Undetermined	Gordon 1982, McKee 1982a
	Bryozoa	Undetermined	Undetermined	McKee 1982b
	Cnidaria	Anthozoa	*Michelinia* sp.	Gordon 1982, McKee 1982a
	Cnidaria	Anthozoa	Undetermined	McKee 1982b
	Cnidaria	Staurozoa	Undetermined (Conularia)	Gordon 1982, McKee 1982a
	Echinodermata	Crinoidea	Undetermined	McKee 1982b, Billingsley & Beus 1999a
	Echinodermata	Undetermined	Undetermined	McKee 1982a
	Foraminifera	Fusulinata	*Fusulinella* sp.	McKee 1982a, 1982b
	Foraminifera	Fusulinata	*Pseudostaffella* sp.	McKee 1982a, 1982b
	Mollusca	Bivalvia	*Aviculopecten* aff. *A. gravidus*	Gordon 1982
	Mollusca	Bivalvia	*Aviculopecten gravidus*	Gordon 1982, McKee 1982a
	Mollusca	Bivalvia	*Aviculopecten* sp.	Gordon 1982, McKee 1982a
	Mollusca	Bivalvia	*Leptodesma* sp.	Gordon 1982, McKee 1982a
	Mollusca	Bivalvia	*Myalina* sp.	Gordon 1982, McKee 1982a

Appendix Table 5-B-3 (continued). Supai Group invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
IPswa (cont.)	Mollusca	Bivalvia	*Oriocrassatella* sp.	Gordon 1982
	Mollusca	Bivalvia	*Permophorus* sp.	McKee 1938
	Mollusca	Bivalvia	*Promytilus* sp.	Gordon 1982, McKee 1982a
	Mollusca	Bivalvia	*Schizodus* sp.	Gordon 1982
	Mollusca	Bivalvia	Undetermined	Gordon 1982, McKee 1982a
	Mollusca	Gastropoda	*Euomphalus* sp.	Gordon 1982
	Mollusca	Gastropoda	*Straparollus* sp.	McKee 1982a
	Mollusca	Gastropoda	Undetermined	Gordon 1982, McKee 1982a, 1982b, Billingsley & Beus 1999a
IPsm	Arthropoda	Ostracoda	Undetermined	McKee 1982a
	Bryozoa	Undetermined	Undetermined	McKee 1982b
	Echinodermata	Undetermined	Undetermined	McKee 1982a
	Foraminifera	Fusulinata	*Endothyra media*	McKee 1982a
	Foraminifera	Fusulinata	*Endothyra* sp.	McKee 1982c
	Foraminifera	Fusulinata	*Endothyra teres*	McKee 1982a
	Foraminifera	Fusulinata	*Eoschubertella* sp.	McKee 1982a, 1982b
	Foraminifera	Fusulinata	*Fusulinella* sp.	McKee 1982a, 1982b
	Foraminifera	Fusulinata	*Pseudostaffella* sp.	McKee 1982a, 1982b
	Foraminifera	Fusulinata	*Schubertella* sp.	McKee 1982a, 1982b
	Foraminifera	Fusulinata	Undetermined	McKee 1982b, 1982c
	Foraminifera	Undetermined	Undetermined	McKee 1982a
	Undetermined	Undetermined	Calcisphere bioclasts	McKee 1982a
IPswe	Bryozoa	Undetermined	Undetermined	McKee 1982b
	Echinodermata	Undetermined	Undetermined	Gordon 1982
	Foraminifera	Fusulinata	Undetermined	McKee 1982b, 1982c
	Foraminifera	Undetermined	Undetermined	McKee 1982a
Pse	Bryozoa	Undetermined	Undetermined	McKee 1982b
	Cnidaria	Undetermined	Undetermined	McKee 1982a
	Echinodermata	Undetermined	Undetermined	McKee 1982a
	Foraminifera	Fusulinata	*Schubertella* sp.	McKee 1982a
	Foraminifera	Undetermined	Undetermined	McKee 1982a

Appendix Table 5-B-4. Post-Supai Group Permian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Ph	Arthropoda	Subphylum Chelicerata	*Hastimima?* sp.	White 1929
	Arthropoda	Insecta	*Tupus gilmorei*	Carpenter 1927, White 1929
	Arthropoda	Insecta	*Tupus whitei*	Carpenter 1928, White 1929
	Arthropoda	Insecta	Undetermined (Blattodea)	Carpenter 1928, Spamer 1984
	Arthropoda	Insecta	Undetermined (Odonata)	Carpenter 1928
Pt	Arthropoda	Ostracoda	Undetermined	McKee 1938
	Brachiopoda	Rhynchonellata	*Composita* sp.	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Strophomenata	*Peniculauris ivesi*	McKee 1938
	Bryozoa	Undetermined	Undetermined	McKee 1938, Sorauf & Billingsley 1991
	Echinodermata	Crinoidea	Undetermined	McKee 1938, Sorauf & Billingsley 1991
	Echinodermata	Echinoidea	*Archaeocidaris* sp.	McKee 1938, Sorauf & Billingsley 1991
	Mollusca	Bivalvia	*Allorisma* sp.	McKee 1938
	Mollusca	Bivalvia	*Edmondia* sp.	McKee 1938
	Mollusca	Bivalvia	*Nuculana?* sp.	McKee 1938
	Mollusca	Bivalvia	*Permophorus* sp.	McKee 1938
	Mollusca	Bivalvia	*Pteria* sp.	McKee 1938
	Mollusca	Bivalvia	*Schizodus* sp.	McKee 1938
	Mollusca	Bivalvia	*Solemya* sp.	McKee 1938
	Mollusca	Bivalvia	Undetermined	McKee 1938
	Mollusca	Cephalopoda	*Domatoceras?* sp.	McKee 1938
	Mollusca	Gastropoda	*Aclisina* sp.	McKee 1938
	Mollusca	Gastropoda	*Bellerophon* sp.	McKee 1938
	Mollusca	Gastropoda	*Euomphalus* sp.	McKee 1938
	Mollusca	Gastropoda	*Euphemites* sp.	McKee 1938
	Mollusca	Gastropoda	*Goniospira* sp.	McKee 1938
	Mollusca	Gastropoda	*Naticopsis* sp.	McKee 1938
	Mollusca	Gastropoda	Undetermined	McKee 1938

Appendix Table 5-B-4 (continued). Post-Supai Group Permian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Pt (cont.)	Mollusca	Undetermined	Undetermined	McKee 1938
Pk	Arthropoda	Trilobita	*Delaria sevilloidia*	Cisne 1971
	Arthropoda	Trilobita	*Delaria snowi*	Cisne 1971
	Arthropoda	Trilobita	*Ditomopyge* sp.	McKee 1938, Sorauf & Billingsley 1991, Thayer 2009
	Arthropoda	Trilobita	*Novoameura mckeei*	Cisne 1971
	Arthropoda	Trilobita	Undetermined	Billingsley & Beus 1999a
	Brachiopoda	Rhynchonellata	*Composita arizonica*	McKee 1938
	Brachiopoda	Rhynchonellata	*Composita* sp.	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Rhynchonellata	*Composita subtilita*	McKee 1938, Gordon 1982
	Brachiopoda	Rhynchonellata	*Composita?* sp.	McKee 1938
	Brachiopoda	Rhynchonellata	*Dielasma phosphoriense*	McKee 1938
	Brachiopoda	Rhynchonellata	*Hustedia* aff. *H. meekana*	McKee 1938
	Brachiopoda	Rhynchonellata	*Hustedia* sp.	Thompson 1995
	Brachiopoda	Rhynchonellata	*Neophricadothyris* sp.	Thompson 1995
	Brachiopoda	Rhynchonellata	*Phrenophoria pinguis*	McKee 1938
	Brachiopoda	Rhynchonellata	*Phricodothyris guadalupensis*	McKee 1938
	Brachiopoda	Rhynchonellata	*Pugnoides* sp.	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Rhynchonellata	*Punctospirifer?* sp.	McKee 1938
	Brachiopoda	Rhynchonellata	*Rhipidomella hessensis*	Condra & Elias 1944
	Brachiopoda	Rhynchonellata	*Rhipidomella transversa*	McKee 1938
	Brachiopoda	Rhynchonellata	*Spiriferellina hilli*	McKee 1938
	Brachiopoda	Rhynchonellata	*Spiriferina?* sp.	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Rhynchonellata	*Squamularia* sp.	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Strophomenata	*Avonia* sp.	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Strophomenata	*Bathymyonia nevadensis*	McKee 1938
	Brachiopoda	Strophomenata	*Chonetes* sp.	McKee 1938

Appendix Table 5-B-4 (continued). Post-Supai Group Permian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Pk (cont.)	Brachiopoda	Strophomenata	*Derbyia arizonensis*	McKee 1938, 1941
	Brachiopoda	Strophomenata	*Derbyia nasuta*	McKee 1938
	Brachiopoda	Strophomenata	*Derbyia* sp.	McKee 1938, Sorauf & Billingsley 1991, Thayer 2009
	Brachiopoda	Strophomenata	*Dyoros* (*Tetragonetes*) *tetragonus*	King 1931, Cooper and Grant 1975
	Brachiopoda	Strophomenata	*Dyoros kaibabensis*	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Strophomenata	*Dyoros subliratus*	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Strophomenata	*Echinauris dorsoconcava*	McKee 1938
	Brachiopoda	Strophomenata	*Echinauris newberryi*	McKee 1938
	Brachiopoda	Strophomenata	*Echinauris* sp.	Thompson 1995
	Brachiopoda	Strophomenata	*Kozlowskia* sp.	Thompson 1995
	Brachiopoda	Strophomenata	*Kutorginella meridionalis*	McKee 1938
	Brachiopoda	Strophomenata	*Liosotella popei*	McKee 1938
	Brachiopoda	Strophomenata	*Marginifera* sp.	McKee 1938
	Brachiopoda	Strophomenata	*Meekella pyramidalis*	Schuchert 1918b, McKee 1938
	Brachiopoda	Strophomenata	*Meekella* sp.	McKee 1938, Sorauf & Billingsley 1991, Thayer 2009
	Brachiopoda	Strophomenata	*Peniculauris bassi*	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Strophomenata	*Peniculauris ivesi*	Schuchert 1918b, McKee 1938
	Brachiopoda	Strophomenata	*Peniculauris* sp.	Thayer 2009
	Brachiopoda	Strophomenata	*Rugatia occidentalis*	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Strophomenata	*Rugatia paraindica*	McKee 1938
	Brachiopoda	Strophomenata	*Rugatia* sp.	Thompson 1995
	Brachiopoda	Strophomenata	*Waagenoconcha* sp.	McKee 1938, Sorauf & Billingsley 1991
	Brachiopoda	Undetermined	Undetermined	McKee 1938
	Bryozoa	Stenolaemata	*Bascomella subsphaerica*	Condra & Elias 1944
	Bryozoa	Stenolaemata	*Bicorbis arizonica*	Condra & Elias 1945a, 1945b, McKinney 1983
	Bryozoa	Stenolaemata	*Fistulipora* sp.	McKee 1938, McKinney 1983
	Bryozoa	Stenolaemata	*Girtypora maculata*	McKinney 1983

Appendix Table 5-B-4 (continued). Post-Supai Group Permian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Pk (cont.)	Bryozoa	Stenolaemata	*Meekopora parilis*	McKinney 1983
	Bryozoa	Stenolaemata	*Meekopora* sp.	McKee 1938
	Bryozoa	Stenolaemata	*Polypora spinulifera*	McKee 1938
	Bryozoa	Stenolaemata	*Rhabdomeson* sp.	McKinney 1983
	Bryozoa	Stenolaemata	*Rhombopora lepidodendroides*	McKee 1938
	Bryozoa	Stenolaemata	*Septopora biserialis*	McKee 1938
	Bryozoa	Stenolaemata	*Septopora* sp.	McKee 1938
	Bryozoa	Stenolaemata	*Stenodiscus* sp.	McKinney 1983
	Bryozoa	Stenolaemata	*Stenopora* sp.	McKee 1938
	Bryozoa	Stenolaemata	*Streblotrypa sp.*	McKinney 1983
	Bryozoa	Stenolaemata	Undetermined	McKinney 1983
	Bryozoa	Undetermined	Undetermined	McKee 1938, McKinney 1983, Sorauf & Billingsley 1991, Thompson 1995
	Cnidaria	Anthozoa	*Lophophyllum* sp.	McKee 1938, Sorauf & Billingsley 1991
	Cnidaria	Anthozoa	Undetermined	McKee 1938
	Cnidaria	Staurozoa?	*Conularia kaibabensis*	McKee 1935, Sinclair 1948, Spamer 1984
	Echinodermata	Crinoidea	Undetermined	Shimer 1919, McKee 1938, Sorauf & Billingsley 1991, Thayer 2009
	Echinodermata	Echinoidea	*Archaeocidaris* sp.	McKee 1938, Sorauf & Billingsley 1991
	Mollusca	Bivalvia	*Acanthopecten coloradoensis*	Newell 1937
	Mollusca	Bivalvia	*Astartella gurleyi*	Shimer 1919, McKee 1938
	Mollusca	Bivalvia	*Aviculopecten* sp.	McKee 1938, Sorauf & Billingsley 1991
	Mollusca	Bivalvia	*Bakewellia parva*	Shimer 1919, McKee 1938
	Mollusca	Bivalvia	*Nuculana?* sp.	McKee 1938
	Mollusca	Bivalvia	*Permophorus* sp.	McKee 1938
	Mollusca	Bivalvia	*Pteria* sp.	McKee 1938
	Mollusca	Bivalvia	*Schizodus* sp.	McKee 1938
	Mollusca	Bivalvia	*Solemya* sp.	McKee 1938

Appendix Table 5-B-4 (continued). Post-Supai Group Permian invertebrate taxa reported from GRCA localities in the literature. For abbreviations see text.

Unit	Phylum	Class	Species	Reference
Pk (cont.)	Mollusca	Bivalvia	Undetermined	McKee 1938
	Mollusca	Cephalopoda	*Domatoceras bradyi*	Miller & Unklesbay 1942, Miller & Youngquist 1949
	Mollusca	Cephalopoda	*Domatoceras simplex*	McKee 1938
	Mollusca	Cephalopoda	*Orthoceras* sp.	McKee 1938
	Mollusca	Cephalopoda	*Stearoceras rotundatum*	Miller & Unklesbay 1942, Miller & Youngquist 1949
	Mollusca	Cephalopoda	*Stearoceras sanandreasense*	Miller & Unklesbay 1942, Miller & Youngquist 1949
	Mollusca	Gastropoda	*Bellerophon majusculus*	McKee 1938
	Mollusca	Gastropoda	*Euomphalus* sp.	McKee 1938
	Mollusca	Gastropoda	*Euphemites* cf. *E. carbonarius*	Shimer 1919, McKee 1938
	Mollusca	Gastropoda	*Euphemites* sp.	McKee 1938
	Mollusca	Gastropoda	*Goniospira* sp.	McKee 1938
	Mollusca	Gastropoda	*Murchisonia?* cf. *M. terebra*	Shimer 1919, McKee 1938
	Mollusca	Gastropoda	*Naticopsis* sp.	McKee 1938
	Mollusca	Gastropoda	*Pleurotomaria* sp.	McKee 1938
	Mollusca	Gastropoda	Undetermined	McKee 1938
	Mollusca	Scaphopoda	*Prodentalium canna*	Shimer 1919, McKee 1938
	Mollusca	Undetermined	Undetermined	McKee 1938
	Porifera	Demospongiae	*Actinocoelia maeandrina*	Griffin 1966
	Porifera	Demospongiae	*Actinocoelia* sp.	Thayer 2009
	Porifera	Undetermined	Undetermined	Schuchert 1918b, McKee 1938, Sorauf & Billingsley 1991

Appendix 5-C. Paleozoic Invertebrate Body Fossil Taxa Named From GRCA

Befitting a park with such a long history of paleontological investigations and diversity of fossiliferous rocks, GRCA has been the source for numerous type specimens. At least 80 species of fossil invertebrates have been named from specimens discovered within 2019 GRCA boundaries. They are listed in Appendix Table 5-C-1 in alphabetical order, using the original species names. An additional 11 invertebrate species have been named from specimens possibly derived from GRCA locations (see the following Appendix 5-D). The great majority of these taxa were collected and named during the National Park era (1919–present), although not all of the discovery localities were within GRCA at the time of discovery due to park boundary changes. For example, some type specimens recovered from what is now far western GRCA were in Lake Mead National Recreation Area (LAKE) during the 1930s. Pre-1919 taxa are primarily Walcott's Cambrian invertebrates and a few species from the very early era of exploration. Land once part of southern GRCA was transferred to the Havasupai Indian Reservation in 1975. Although there are some fossil sites described in the pre-1975 literature as within GRCA that are now outside the park boundaries, no type specimen locations were affected by the change in ownership.

Appendix Table 5-C-1. Fossil invertebrate taxa named from specimens collected from GRCA. Species are listed with stratigraphic unit, taxonomic group, original name, source citation, specimen accession identifiers, and notes.

Unit	Phylum	Original Name	Citation	Type Specimen	Notes
Ctba	Arthropoda	*Acrocephalops? arizonaensis*	Resser 1945	USNM 108624	Trilobite
	Arthropoda	*Albertella schenki*	Resser 1945	USNM 108583	Trilobite
	Arthropoda	*Alokistocare althea*	Walcott 1916a	Lectotype USNM 61574	Trilobite
	Arthropoda	*Dielymella appressa*	Ulrich & Bassler 1931	USNM 56506	Bradoriid
	Arthropoda	*Dielymella dorsalis*	Ulrich & Bassler 1931	USNM 56505	Bradoriid
	Arthropoda	*Dielymella nasuta*	Ulrich & Bassler 1931	Cotypes USNM 56508, 56509	Bradoriid
	Arthropoda	*Dielymella recticardinalis*	Ulrich & Bassler 1931	USNM 56510	Bradoriid
	Arthropoda	*Dielymella recticardinalis angustata*	Ulrich & Bassler 1931	USNM 56511	Bradoriid
	Arthropoda	*Dolichometopus tontoensis*	Walcott 1916b	Lectotype USNM 62685	Trilobite, *Anoria tontoensis*
	Arthropoda	*Ehmaniella arizonaensis*	Resser 1945	USNM 108603a	Trilobite
	Arthropoda	*Elrathia nitens*	Resser 1945	USNM 108625	Trilobite
	Arthropoda	*Glossopleura mckeei*	Resser 1935	USNM 62714	Trilobite
	Arthropoda	*Glyphaspis vulsa*	Resser 1945	USNM 108595a	Trilobite
	Arthropoda	*Indiana curta*	Ulrich & Bassler 1931	USNM 56466	Bradoriid
	Arthropoda	*Indiana impressa*	Ulrich & Bassler 1931	USNM 56463	Bradoriid
	Arthropoda	*Kootenia simplex*	Resser 1945	USNM 108591a	Trilobite
	Arthropoda	*Pachyaspis fonticola*	Resser 1945	USNM 108608	Trilobite
	Arthropoda	*Parehmania tontoensis*	Resser 1945	USNM 108614	Trilobite
	Arthropoda	*Spencia tontoensis*	Resser 1945	USNM 108611a	Trilobite
	Arthropoda	*Walcottella breviuscula*	Ulrich & Bassler 1931	USNM 56481	Bradoriid
	Arthropoda	*Walcottella leperditoides*	Ulrich & Bassler 1931	USNM 56484	Bradoriid
	Arthropoda	*Walcottella limatula*	Ulrich & Bassler 1931	USNM 56488	Bradoriid
	Arthropoda	*Walcottella longula*	Ulrich & Bassler 1931	USNM 56491	Bradoriid
	Arthropoda	*Walcottella nitida*	Ulrich & Bassler 1931	USNM 56485	Bradoriid
	Arthropoda	*Walcottella oblonga*	Ulrich & Bassler 1931	USNM 56486	Bradoriid
	Arthropoda	*Walcottella obsoleta*	Ulrich & Bassler 1931	USNM 56487	Bradoriid

Appendix Table 5-C-1 (continued). Fossil invertebrate taxa named from specimens collected from GRCA. Species are listed with stratigraphic unit, taxonomic group, original name, source citation, specimen accession identifiers, and notes.

Unit	Phylum	Original Name	Citation	Type Specimen	Notes
Ctba	Arthropoda	*Walcottella pulchella*	Ulrich & Bassler 1931	USNM 56483	Bradoriid
	Arthropoda	*Walcottella scitula*	Ulrich & Bassler 1931	USNM 56482	Bradoriid
	Arthropoda	*Walcottella subtruncata*	Ulrich & Bassler 1931	USNM 56490	Bradoriid
	Arthropoda	*Walcottella ventrosa*	Ulrich & Bassler 1931	USNM 56489	Bradoriid
	Brachiopoda	*Billingsella obscura*	Walcott 1905	USNM 52258a	*Nisusia obscura*
	Brachiopoda	*Iphidea crenistria*	Walcott 1897	USNM 26431a	*Micromitra* (*Paterina*) *crenistria*
	Brachiopoda	*Lingulella mckeei*	Resser 1945	USNM 108561a	Brachiopod
	Brachiopoda	*Obolus* (*Lingulella*) *euglyphus*	Walcott 1898	USNM 27316a	*Lingulella euglypha*
	Brachiopoda	*Obolus* (*Lingulella*) *spatulus*	Walcott 1902	USNM 35290a	*Lingulella spatula*
	Brachiopoda	*Obolus* (*Lingulella*) *zetus*	Walcott 1898	USNM 27347b	*Lingulella zetus*
	Brachiopoda	*Obolus* (*Westonia*) *themis*	Walcott 1905	USNM 51732a	*Lingulella themis*
	Echinodermata	*Eocrinus multibrachiatus*	Kirk (in Resser 1945)	USNM 108556a	*Gogia multibrachiatus*
Ctba or Ctm	Arthropoda	*Clavaspidella kanabensis*	Resser 1945	USNM 108578a	Trilobite
	Arthropoda	*Ehmaniella hebes*	Resser 1945	USNM 108612a	Trilobite
	Arthropoda	*Elrathiella? insueta*	Resser 1945	USNM 108621a	Trilobite
	Arthropoda	*Glyphaspis kwanguntensis*	Resser 1945	USNM 108618a	Trilobite
	Arthropoda	*Kochina? angustata*	Resser 1945	USNM 108610	Trilobite
	Arthropoda	*Kootenia mckeei*	Resser 1945	USNM 108588a	Trilobite
	Arthropoda	*Parehmania kwaguntensis*	Resser 1945	USNM 108620a	Trilobite
	Arthropoda	*Parehmania nitida*	Resser 1945	USNM 108613a	Trilobite
	Arthropoda	*Solenopleurella diligens*	Resser 1945	USNM 108627a	Trilobite
	Arthropoda	*Trachycheilus typicale*	Resser 1945	USNM 108619	Trilobite
	Brachiopoda	*Lingulella kanabensis*	Resser 1945	USNM 33829a	Brachiopod
	Brachiopoda	*Nisusia* (*Jamesella*) *kanabensis*	Walcott 1908	USNM 52300	Brachiopod
Ctba?	Arthropoda	*Indiana faba*	Ulrich & Bassler 1931	USNM 56458	Bradoriid
	Arthropoda	*Indiana faba intermedia*	Ulrich & Bassler 1931	USNM 56462	Bradoriid

Appendix Table 5-C-1 (continued). Fossil invertebrate taxa named from specimens collected from GRCA. Species are listed with stratigraphic unit, taxonomic group, original name, source citation, specimen accession identifiers, and notes.

Unit	Phylum	Original Name	Citation	Type Specimen	Notes
Ctba? (cont.)	Arthropoda	*Walcottella apicalis*	Ulrich & Bassler 1931	Cotypes USNM 56477, 56478	Bradoriid
	Arthropoda	*Walcottella concentrica*	Ulrich & Bassler 1931	Cotypes USNM 56479, 56480	Bradoriid
	Brachiopoda	*Dictyonina arizonaensis*	Resser 1945	USNM 108557	Brachiopod
	Brachiopoda	*Iphidea superba*	Walcott 1897	USNM 26429a	*Micromitra* (*Paterina*) *superba*
	Brachiopoda	*Obolus* (*Lingulella*) *chuarensis*	Walcott 1898	USNM 57020a	*Lingulella chuarensis*
	Brachiopoda	*Obolus* (*Lingulella*) *lineolatus*	Walcott 1898	USNM 27325a	*Lingulella lineolata*
	Problematica	*Tontoia kwaguntensis*	Walcott 1912b	USNM 57660	*Incertae sedis*
Ctm	Arthropoda	*Bolaspis aemula*	Resser 1945	USNM 108602a	Trilobite
	Arthropoda	*Glyphaspis tecta*	Resser 1945	USNM 108596a	Trilobite
	Arthropoda	*Kootenia schenki*	Resser 1945	USNM 108586a	Trilobite
	Arthropoda	*Solenopleurella erosa*	Resser 1945	USNM 108616a	Trilobite
	Arthropoda	*Solenopleurella porcata*	Resser 1945	USNM 108626a	Trilobite
	Brachiopoda	*Finkelnburgia noblei*	Walcott 1924	USNM 69750–69752	*Nisusia noblei*
	Problematica	*Scenella hermitensis*	Resser 1945	USNM 108568a	*Incertae sedis*
Ctu	Arthropoda	*Bradoria tontoensis*	Ulrich & Bassler 1931	USNM 81377	Bradoriid
Mr	Foraminifera	*Endothyra baileyi poloumera*	Skipp 1969	USNM 641727	*Globoendothyra baileyi poloumera*
	Foraminifera	*Septaglomospiranella rossi*	Skipp 1969	USNM 641578	–
Ph	Arthropoda	*Typus gilmorei*	Carpenter 1927	USNM 71279	Insect, *Tupus gilmorei*
	Arthropoda	*Typus whitei*	Carpenter 1928	USNM 71713	Insect, *Tupus whitei*
Pk	Arthropoda	*Delaria macclintocki*	Cisne 1971	GCNPM 3949	Trilobite
	Brachiopoda	*Avonia subhorrida newberryi*	McKee 1938	USNM 102301	*Echinauris newberryi*
	Brachiopoda	*Chonetes quadratus*	King 1931	YPM 10830a	*Dyoros* (*Tetragonetes*) *tetragonus*
	Brachiopoda	*Composita arizonica*	McKee 1938	Syntypes USNM 102303, 102304, 102305	Brachiopod

Appendix Table 5-C-1 (continued). Fossil invertebrate taxa named from specimens collected from GRCA. Species are listed with stratigraphic unit, taxonomic group, original name, source citation, specimen accession identifiers, and notes.

Unit	Phylum	Original Name	Citation	Type Specimen	Notes
Pk (cont.)	Brachiopoda	*Derbyia regularis*	McKee 1938	Syntypes USNM 102290, 102291	*Derbyia arizonensis*
	Bryozoa	*Bascomella subsphaerica*	Condra & Elias 1944	NGS 449	–
	Bryozoa	*Bicorbula arizonica*	Condra & Elias 1945a	NGS 264	*Bicorbis arizonica*
	Bryozoa	*Girtypora maculata*	McKinney 1983	FMNH PE 24301	–
	Cnidaria	*Conularia kaibabensis*	McKee 1935	USNM 102289 (was GCNPM FK211)	Conulariid, also *Paraconularia kaibabensis*

Appendix 5-D. Paleozoic Invertebrate Body Fossil Taxa Potentially Named From GRCA

A small number of taxa have been named from fossils discovered somewhere in the Grand Canyon area, but with insufficient provenance information to determine the exact location. Some, all, or none of the type specimens may have come from GRCA. Names are listed as first defined. Some names may be revised in other sources. However, not all changes are reflected in the inventory at this time.

Appendix Table 5-D-1. Fossil invertebrate taxa named from specimens possibly found within GRCA. Species are listed with stratigraphic unit, taxonomic group, original name, source citation, specimen accession identifiers, and notes.

Unit	Phylum	Original Name	Citation	Type Specimen	Notes
Ctm	Arthropoda	*Kootenia havasuensis*	Resser 1945	USNM 108601a	Trilobite
	Arthropoda	*Pachyaspis moorei*	Resser 1945	USNM 108606a	Trilobite
Pt	Brachiopoda	*Productus ivesi*	Newberry 1861	Syntypes USNM 5356a–c	*Peniculauris ivesi*
Pk	Brachiopoda	*Avonia dorsoconcava*	McKee 1938	GCNPM 9993	*Echinauris dorsoconcava*
	Brachiopoda	*Chonetes kaibabensis*	McKee 1938	Syntypes USNM 102292, 102293, 102294	*Dyoros kaibabensis*
	Brachiopoda	*Marginifera meridionalis*	McKee 1938	Syntype USNM 102302	*Kutorginella meridionalis*
	Brachiopoda	*Productus (Dictyoclostus) bassi*	McKee 1938	Lectotype USNM 102295a	*Peniculauris bassi*
	Brachiopoda	*Productus (Dictyoclostus) paraindicus*	McKee 1938	Lectotype USNM 102297	*Rugatia paraindica*
Pt or Pk	Echinodermata	*Archaeocidaris gracilis*	Newberry 1861	Syntypes CU 604, USNM 5412	–
	Echinodermata	*Archaeocidaris longispinus*	Newberry 1861	CU 6419G	–
	Echinodermata	*Archaeocidaris ornatus*	Newberry 1861	Syntypes CU 6000G	*Archaeocidaris coloradensis*

Chapter 6. Paleozoic Vertebrate Paleontology of Grand Canyon National Park: Research History, Resources, And Potential

By John-Paul Michael Hodnett[1] and David Kenneth Elliott[2]

[1]Maryland-National Capital Park and Planning Commission
Department of Parks and Recreation, Archaeology Program
8204 McClure Road
Upper Marlboro, Maryland 20772

[2]Northern Arizona University
School of Earth and Sustainability, Geology Division
Flagstaff, Arizona 86011

Introduction

The Grand Canyon is an awe-inspiring place and is a physical testament of deep time. More than a 150 years' worth of geologic research has yielded a rich paleontological record, documenting approximately a half billion years of life on this planet. Much of the Grand Canyon is composed of Paleozoic rocks, with the oldest going back to the Cambrian. The fossil record of these Paleozoic rocks is dominated by a rich invertebrate heritage. The vertebrate record, however, is typically portrayed as unusually sparse. The Paleozoic vertebrate record at the Grand Canyon is best known for the varied trace fossils of tetrapods from the Pennsylvanian–early Permian Supai Group and the Permian Coconino Sandstone, where multiple different ichnotaxa have been described (e. g., McKee 1982; Spamer 1984; Elliott and Blakey 2005; Hunt et al. 2005). However, no tetrapod body fossils have been identified from these beds. Much of the research and literature on the body fossil record of vertebrates from the Grand Canyon has been focused on Pleistocene vertebrates from cave sites along the length of the canyon (e. g., Lucas and Morgan 2005; Mead 2005).

Nevertheless, Paleozoic vertebrate body fossils have been reported from the Grand Canyon. Walcott (1880) made the first report of vertebrate fossils from the Devonian Temple Butte Limestone during the initial surveys made by the United States Geological Survey (USGS). McKee (1938) reported and described the first Permian vertebrate body fossil, a chondrichthyan tooth plate, from the Kaibab Formation. In 1944, the chondrichthyan *Megactenopetalus kaibabanus* was described from a large tooth collected from the North Rim at Point Sublime, and this is the first designated holotype of a Paleozoic vertebrate fossil from the Grand Canyon (David 1944). Other reports of vertebrate body fossils are from the Mississippian Redwall Limestone (McKee and Gutschick 1969) and the Pennsylvanian–Permian Supai Group (McKee 1982). Reviews of these vertebrate body fossils from the Paleozoic of the Grand Canyon have been made in the past (Spamer 1984; Elliott and Blakey 2005; Hunt et al. 2005). Recently, a rich assemblage of chondrichthyans was described from the Upper Mississippian Surprise Canyon Formation from the western Grand Canyon (Hodnett and Elliott 2018).

The purpose of the report presented here is to both collate previous documentation of Paleozoic vertebrate body fossils from the Grand Canyon and present the current physical record of specimens that have been collected from Grand Canyon National Park (GRCA). We have identified the presence of Paleozoic body fossils from seven geologic horizons within the Grand Canyon, which includes records outside of GRCA boundaries. We have included these records as part of our review as a "total record" approach to understanding the taxonomic richness of these seven geologic horizons. This additional information should assist in fossil resource planning, management, and priority assessment for further Paleozoic vertebrate research at GRCA.

Institutional Abbreviations—**GRCA**, Grand Canyon National Park Museum, Grand Canyon, Arizona; **MNA**, Museum of Northern Arizona, Flagstaff, Arizona; **USNM**, National Museum of Natural History Paleobiology Collections, Washington, D.C.

Devonian

Temple Butte Limestone

The Temple Butte Limestone is the only Devonian formation exposed at GRCA, and it also extends into southeast Nevada. The Temple Butte Limestone consists of thin, discontinuous lenses no more than 30 m (100 ft) thick deposited in paleo-valleys cut into the underlying Muav Limestone within the eastern Grand Canyon (Beus 2003a). The central and western sections of the Temple Butte Limestone are more continuous (Beus 2003a). The Temple Butte Limestone consists of a westward thickening wedge of interbedded dolomite, sandy dolomite, sandstones, mudstones, and limestones, ranging from purple, reddish-purple, and dark to light gray (Beus 2003a). The eastern paleo-valleys of the Temple Butte strata consist of interbedded mudstones, sandstones, dolomite, and conglomerates (Beus 2003a).

The upper and lower contacts of the Temple Butte Limestone represent major unconformities with the underlying middle Cambrian Muav Limestone and the overlying Lower Mississippian Redwall Limestone (Beus 2003a). The model for deposition of the Temple Butte Limestone is similar to that of the younger Surprise Canyon Formation (Beus 2003a). The Temple Butte Limestone is a contemporaneous equivalent to the Martin Formation found in central and southern Arizona, which is known for a diverse vertebrate assemblage (Elliott and Blakey 2005). Both the Temple Butte Limestone and the Martin Formation are considered to be early Late Devonian (Frasnian) in age (Elliott and Blakey 2005).

History

The first report of a vertebrate fossil from the Grand Canyon was by Charles Doolittle Walcott (1880), who reported: "The purple sandstones deposited in the hollows of the Silurian limestones are characterized by the presence of Placoganoid fishes of a Devonian type," which he identified during a field excursion in 1879. Schuchert (1918) communicated with L. Noble on a series of scales and plates of fish collected from the Temple Butte Limestone in 1916. These fish fossils were identified by J. W. Gidley of the Smithsonian Institution as the plates of the antiarch placoderm fish *Bothriolepis nitidens* and the scales of the sarcopterygian *Holoptychius* (Schuchert 1918; Noble 1922). Noble (1922) made no report of *Holoptychius* scales in the summary he published from

Gidley's analysis. Denison (1951) accepted the occurrence of *Holoptychius* and revised the Temple Butte placoderm to *Bothriolepis coloradensis* based on similar outer dermal plate ornamentation morphology.

GRCA Resources

At present there are only two specimens identified in the National Museum of Natural History Paleobiology collections that come from the Temple Butte Limestone of the Grand Canyon (Figure 6-1). USNM PAL 328639 consists of part and counterpart of the distal part of a pectoral appendage (Figures 6-1A and 1B) and fragments and impressions of indeterminate plates of *Bothriolepis* cf. *B. coloradensis* that were collected by C. D. Walcott on January 17, 1883 as part of the United States Geological Survey study. This specimen is here considered to be the first discovered and oldest fossil vertebrate to be catalogued from the Grand Canyon. In addition, USNM PAL 328643 (Figures 6-1C–E) is a collection of approximately 30+ elements consisting of pectoral appendage, dorsal and ventral shield, and cephalic plates of *Bothriolepis coloradensis* collected by L. F. Noble in April 1916 and later mentioned by Denison (1951). At present, no specimens of *Holoptychius* from the Temple Butte Limestone have been identified in a museum collection.

Potential

Notes by Noble (1922) suggest that vertebrate remains from the Temple Butte are common but fragmentary approximately 1 m (3 ft) above the base of the Temple Butte Limestone and its contact with the Muav Limestone, in an impure dolomitic limestone. Noble (1922) speculated that other deposits within the Temple Butte Limestone could similarly be rich in vertebrate material.




Figure 6-1. Antiarch placoderm, *Bothriolepis,* fossils from the Temple Butte Limestone, Grand Canyon (JOHN-PAUL HODNETT). **A–B.** USNM PAL 328639, *Bothriolepis* cf. *B. coloradensis*, the proximal part of a pectoral appendage collected by C. D. Walcott. **C–E.** USNM PAL 328643, *Bothriolepis coloradensis*, a selection of plates collected by L. F. Noble. Scale equals 1 cm (0.4 in).

Mississippian

Redwall Limestone

The Redwall Limestone is a cliff-forming unit of early to middle Mississippian age sedimentary rocks (Beus 2003b). It consists primarily of light-olive-gray to light gray, thin- to thick bedded cherty limestones, interbedded crystalline dolomites, and limestone layers with chert lenses (Beus 2003b). The Redwall Limestone is divided into four lithologic units starting with the Whitmore Wash Member at the base, and ascending up to the Thunder Springs Member, Mooney Falls Member, and Horseshoe Mesa Member, respectively (McKee and Gutschick 1969). The Redwall Limestone was formed through two marine transgression and regression events from a seaway to the west (McKee and Gutschick 1969). The Whitmore Wash Member represents the first transgression event, and the Thunder Springs Member represents the first regression event, followed by the second transgression event of the Mooney Falls Member, and the final regression of the Horseshoe Mesa Member (McKee and Gutschick 1969). The Redwall Limestone has an upper unconformable contact with the Surprise Canyon Formation, which represents a time of uplift and karstic paleo-valleys cutting into the Redwall Limestone (Beus 2003). The Redwall Limestone has a rich invertebrate record. Regional variation of fossil faunas are noted within it, particularly of the corals and foraminifers (McKee and Gutschick 1969).

History

McKee and Gutschick (1969) reported a number of durophagous holocephalan chondrichthyan tooth plates and a spine from multiple localities within the Redwall Limestone in northern and central Arizona. The fossils were submitted to D. H. Dunkle, then at the National Museum of Natural History, for identification. Dunkle identified from the Mooney Falls Member the euchondrocephalan *Orodus major* and the holocephalans *Psephodus* sp., *Deltodus* sp., *Helodus* sp., and *Psammodus* sp., as well as a few unidentified fish teeth (McKee and Gutschick 1969). A single dorsal spine of *Physonemus* was also identified (McKee and Gutschick 1969). An indeterminate fish tooth is reported from the Horseshoe Member at Bright Angel Trail (McKee and Gutschick 1969).

D. H. Dunkle noted that much of the Redwall fish fauna was similar to taxa found in the early Mississippian Burlington Limestone of Iowa (St. John and Worthen 1875; McKee and Gutschick 1969). Elliott and Blakey (2005) noted that the Redwall Limestone is approximately contemporaneous with the Escabrosa Limestone in southern Arizona, which has a rich but poorly studied fish assemblage (Gass 1963). Outside of the Grand Canyon, a dolostone nodule from the Redwall preserved the external mold and some bony debris of a large gyracanthid acanthodian pectoral spine (USNM 409810, Babcock and Feldmann 1986). Babcock and Feldmann (1986) identified this specimen as *Oracanthus* sp., and it was collected from the top of a mesa between Rock and Blye canyons south of Peach Springs Arizona. This is the first record of a gyracanthid fish from the state of Arizona.

GRCA Resources

Unfortunately, none of the fish taxa reported in McKee and Gutschick (1969) is represented by specimens in a museum collection with the exception of GRCA 20037 (Figure 6-2A), a tooth of *Helodus* sp. from Kaibab Trail, “Member C [Mooney Falls Member] of Redwall Limestone.” This

specimen was collected by E. D. McKee on March 4, 1959. An uncataloged specimen collected by George Billingsley from the Mooney Falls Member of the Redwall Limestone at Surprise Canyon, was given to us (DKE) for identification (Figure 6-2B). The specimen consists of a partial tooth whorl of two teeth that are similar to *Helodus* (*Psephodus*) *didymus* of the Mississippian (Visean) of Ireland (Stahl 1999). This specimen is currently under study by the authors.




Figure 6-2. Chondrichthyan fossils from the Mooney Fall Member, Redwall Limestone (JOHN-PAUL HODNETT). **A.** GRCA 20037, *Helodus* sp. from Kaibab Trail. **B.** Uncatalogued specimen, cf. *Helodus* (*Psephodus*) *didymus*, from Surprise Canyon. Scale equals 1 cm (0.4 in).

Potential

There seems to be a high potential for more vertebrate materials to be collected from the Redwall Limestone at GRCA.

Surprise Canyon Formation

The Surprise Canyon Formation is the name given to a series of channel fills and karstic cave deposits of Late Mississippian (Serpukhovian) age in Grand Canyon (Billingsley and Beus 1985). These represent a considerable hiatus between the Redwall Limestone and the overlying Supai Group, and were originally recognized by McKee and Gutschick (1969), who gave several examples and descriptions of these deposits, although at that time considering them to be part of the basal Supai Group. The channels were originally described by Billingsley (1978) and later interpreted as paleo-valleys by Billingsley and McKee (1982). The formation is nowhere continuous and consists of isolated, lens-shaped outcrops scattered over Grand Canyon and Marble Canyon to the east, representing deposits in deltaic and tidal channels that drained into an estuary to the east. Outcrops are generally up to 45 m (150 ft) thick in central Grand Canyon close to the presumed headwaters of

the paleochannels but reach 122 m (400 ft) in the west in proximity to the estuary (Billingsley and Beus 1985). Description of the extensive fossil invertebrate assemblages was carried out by Beus (1985, 1986, 1999), who showed that the lower beds are fluvial, whereas the upper part of the succession is marine.

The Surprise Canyon Formation was originally divided into a lower unit consisting of fluvial clastics and an upper marine unit composed of siltstones and limestones (Billingsley and Beus 1985). Subsequent studies indicated the presence of three units: a lower fluvial chert pebble conglomerate interbedded with coarse- to fine grained red-brown sandstone and siltstone mainly of terrestrial origin; a middle marine unit of grey-yellow or reddish-brown, coarsely crystalline, thin-bedded limestone separated from the lower unit by an erosional unconformity; and an upper marine unit of reddish-brown, calcareous siltstone, with minor limestone. Some of the lower fluvial rocks in western Grand Canyon include interbedded limestone and shale suggesting brief marine incursions into paleo-river drainages (Billingsley and Beus 1999).

The sequence in the paleo-valley fills is interpreted as representing deposition in channels developed in the Redwall Limestone. These were shallow deltaic and tidal drainage channels formed during the westward retreat of the sea in which the Redwall Limestone had accumulated. Development of karst and entrenchment of the channels probably occurred in the early Serpukhovian (Billingsley and Beus 1999) and led to the development of a network of drainages. Highlands to the east became a source area for detrital material that was incorporated into deposits of the Surprise Canyon Formation. In the late Chesterian/Serpukhovian a period of subsidence allowed marine waters to gradually flood the eroded valleys forming local estuaries (Billingsley and Beus 1999). As the sea transgressed the estuaries also moved eastwards, their deposits forming the marine Middle and Upper members of the Surprise Canyon Formation. A minor unconformity between the Surprise Canyon Formation and the overlying Watahomigi Formation (the basal formation of the Supai Group) suggests a regional interval of uplift and erosion (Billingsley and Beus 1999).

History

McKee (1982) reported “a shark dentition” from the middle unit of the then-undefined Surprise Canyon Formation, which was below the contact of the Watahomigi Formation. This shark dentition was not identified. Billingsley and Beus (1999) reported the occurrence of vertebrate remains from the lower basal conglomerate member and from the Middle and Upper limestone members. These large vertebrate fossils (macro-teeth and spines) were sent to Richard Lund of Adelphi University for identification and later deposited at the National Museum of Natural History (R. Lund, pers comm. 2016). Martin (1992) and Martin and Barrick (1999) recorded the occurrence of micro-vertebrate remains from conodont residues from multiple localities in the Lower and Middle members of the Surprise Canyon Formation, but they did not identify these vertebrate taxa.

Hodnett and Elliott (2018) reviewed both the macro and micro chondrichthyan teeth and spines from the Surprise Canyon Formation and identified 31 taxa from the Lower, Middle, and Upper members at seven localities within the Grand Canyon, some from GRCA and others from Hualapai Nation land. Of the 31 taxa from the Surprise Canyon Formation, four new taxa were named: *Novaculodus billingsleyi*, *Microklomax carrieae*, *Cooleyella platera*, and *Amaradontus santuccii*. The

chondrichthyans of the Lower Member include the taxa *Microklomax carrieae, Cooleyella platera, Amaradontus santuccii*, *Heteropetalus* sp., orodontid indet., *Deltodus* cf. *D. angularis*, *Deltodus* cf. *D. cingulatus*, *Amelacanthus* sp., and *Acondylacanthus* sp. The chondrichthyans of the Middle Member include the taxa *Thrinacodus gracia, Bransonella nebraskensis*, stethacanthid indet., falcatid indet. 1, falcatid indet. 2, *Denaea williamsi*, cf. "*Ctenacanthus*" *costellatus*, ctenacanthid indet., *Clairina* sp., *Microklomax carrieae*, *Novaculodus billingsleyi*, cf. *Mesodmodus* sp., *Hamiltonichthys* sp., *Cooleyella fordi*, *Cooleyella platera*, *Amaradontus santuccii*, *Srianta* cf. *S. srianta*, eugenodontid indet., petalodontid indet., *Cochliodus* cf. *C. contortus*, and *Deltodus* sp. The chondrichthyans from the Upper Member include the taxa *Cladodus* cf. *C. marginatus*, *Srianta* cf. *S. srianta*, *Helodus*? sp., and *Deltodus* cf. *D. angularis*.

Aside from the chondrichthyans, there were also a number of isolated teeth, scales, and bone of actinopterygian fishes from the Lower and Middle member samples collected by Martin (1992). From Blue Mountain Canyon small tetrapods are represented by a few jaws with closely spaced teeth, which are the first record of Paleozoic tetrapods from the Grand Canyon. The actinopterygians and tetrapods are currently under study by the authors of this report.

GRCA Resources

The collection of vertebrates described from the Surprise Canyon Formation is housed in the National Museum of Natural History Paleobiology collections and the Museum of Northern Arizona Geology Department. It should be noted that the type specimens of *Novaculodus billingsleyi*, *Microklomax carrieae*, *Cooleyella platera*, and *Amaradontus santuccii* were collected from Hualapai Nation land in the Grand Canyon. The following list contains all the material collected from GRCA:

USNM: *Cladodus* cf. *C. marginatus*, USNM PAL 412169 (Figure 6-3A); *Helodus*? sp., USNM PAL 603799; *Cochliodus* cf. *C. contortus* (Figure 6-3C), USNM PAL 412147; *Deltodus* cf. *D. angularis*, USNM PAL 412170, 603798 (Figure 6-3B), 412168; *Deltodus* cf. *D. cingulatus; Deltodus* sp., USNM PAL 412173, 412174. *Amelacanthus* sp., USNM PAL 412150; Acondylacanthus sp., USNM PAL 412149. Specimens not used in Hodnett and Elliott (2018), indeterminate cladodont-grade tooth, USNM PAL 603801; indeterminate bone, USNM PAL 412172, 412171.

MNA: *Microklomax carrieae*, MNA V11301 (Figure 6-3G); *Cooleyella platera*, MNA V11308, V11309 (Figure 6-3F); *Amaradontus santuccii*, MNA V11314, V11315 (Figure 6-3E); *Heteropetalus* sp., MNA V11320 (Figure 6-3D), V 11321; orodontid indet. MNA V11322.




Figure 6-3. Chondrichthyans from the Surprise Canyon Formation collected from Grand Canyon National Park (JOHN-PAUL HODNETT). **A.** *Cladodus* cf. *C. marginatus*, USNM PAL 412169. **B.** *Deltodus* cf. *D. angularis*, USNM PAL 603798. **C.** *Cochliodus* cf. *C. contortus*, USNM PAL 412147. **D.** *Heteropetalus* sp., MNA V11320. **E.** *Amaradontus santuccii*, MNA V11315. **F.** *Cooleyella platera*, MNA V11309. **G.** *Microklomax carrieae*, MNA V11301. Scale for **A–C** equals 1 cm (0.4 in), scale for **D–G** equals 500 µm (0.02 in).

Potential

It is evident from the extensive chondrichthyan fauna (Hodnett and Elliott 2018) and the number of undescribed micro-remains of bony fish and tetrapods (Hodnett and Elliott pers. obs.) that the Surprise Canyon Formation has a relatively rich assemblage of vertebrates. However, we still have a relatively poor understanding of the species richness between the three members of the Surprise Canyon Formation. It is recommended that more extensive fieldwork in the Surprise Canyon Formation be conducted, particularly in the canyons north of the Colorado River, to collect additional specimens for study.

Pennsylvanian

Supai Group

The Supai Group is a slope-forming unit composed of Pennsylvanian to early Permian red bed deposits found throughout the Grand Canyon. The Supai Group is made up of four geologic

formations, which are, in ascending order, the Watahomigi Formation, Manakacha Formation, Wescogame Formation, and Esplanade Sandstone/Pakoon Limestone. Presently, only the Pennsylvanian Watahomigi and Wescogame formations and the early Permian Pakoon Limestone have records of vertebrates.

Watahomigi Formation

The Watahomigi Formation is the lowest formation in the Supai Group. This geologic unit began as an eastwards transgression during the Late Mississippian (Surprise Canyon Formation) (McKee 1982). This marine transgression event spread over the relatively flat Redwall Limestone and the filled Surprise Canyon Formation paleo-valleys (McKee 1982). The Watahomigi Formation averages 91 m (300 ft) thick in the western portion of the Grand Canyon, 61 m (200 ft) thick in the central area, and about 30 m (100 ft) thick in the eastern sections of the canyon (McKee 1982). The lower section of the Watahomigi is made up of thin gravel sheets, followed by layers of mud, silt, and eventually carbonate deposits (McKee 1982).

The initial age determination of the Redwall and Watahomigi formations was based on calcareous foraminiferans (Skipp 1969) and corals (Sando 1969). Based on these studies the youngest Redwall strata were thought to be Osagean or early Meramecian and the lowest Watahomigi beds were thought to be middle Pennsylvanian. This meant that the erosional unconformity between them represented the Late Mississippian and the early Pennsylvanian. The subsequent discovery of an erosional remnant of Redwall containing late Meramecian or early Chesterian foraminifera and corals (Skipp 1969; Sando 1969) reduced the gap between the Redwall and the Watahomigi formations to the Chesterian and part of the Morrowan. This gap was further narrowed by the determination of a mid-Morrowan (Bashkirian) age for the lowermost units of the Watahomigi Formation based on brachiopods (Gordon 1982).

History

McKee (1982) noted the occurrence of tooth plates of the holocephalan *Deltodus* and an indeterminate fish spine from the Grand Canyon. Hodnett and Elliott (2018) described a tooth plate of *Deltodus* sp. Martin (1992) and Martin and Barrick (1999) noted chondrichthyan remains from their conodont residues. Hodnett and Elliott (2018) reviewed this material, which contained a new xenacanth shark taxon *Hokomata parva* (Figure 6-4A–C). Not included in the study by Hodnett and Elliott (2018) were a few chondrichthyan dermal denticles and the jaw of a small actinopterygian fish (Figure 6-4D–E).




Figure 6-4. Fish fossils from the Watahomigi Formation (JOHN-PAUL HODNETT). **A–C.** Holotype (MNA V11264) of the xenacanth shark *Hokomata parva*: **A.** Labial view; **B.** Oral view; **C.** Aboral view. **D–E.** Uncatalogued actinopterygian fish jaw: **D.** Medial view; **E.** Lateral view. **F.** *Deltodus* sp., USNM PAL 412145. **A–E**, scale equals 500 µm (0.02 in); **F**, scale equals 1 cm (0.4 in).

GRCA Resources

The *Deltodus* teeth mentioned by McKee have not been located in any museum collection. The *Deltodus* sp. tooth described by Hodnett and Elliott (2018) is in the National Museum of Natural History Paleobiology Collections (USNM PAL 412145) (Figure 6-4F). The microvertebrate fossils from Martin (1992), Martin and Barrick (1999), and Hodnett and Elliott (2018) come from the Hualapai Reservation and are held in the collections of the Museum of Northern Arizona Geology Department along with the type specimen to *Hokomata parva* (MNA V11264). The other microvertebrate fossils are currently under study by the authors.

Potential

The Watahomigi Formation has a high potential for other vertebrate remains to be discovered, in the form of both macro- and micro-remains.

Wescogame Formation

The Wescogame Formation represents another large marine transgression event after an unconformity with the underlying Manakacha Formation (McKee 1982). The lower beds of the Wescogame Formation are marked by numerous channel beds made up of gravels of siltstone, limestone and conglomerate. The upper beds consist of carbonates, marine clays and sandstones (McKee 1982). The age of the Wescogame is considered to be Virgilian based on the invertebrate assemblages (McKee 1982).

History

McKee (1982) reported the tooth plates of *Deltodus* sp., identified by D. H. Dunkle, from Grand Canyon and the adjacent Lake Mead National Recreation Area.

GRCA Resources

The *Deltodus* tooth plates mentioned in McKee (1982) have not been located in any museum collection.

Potential

There is a good potential for additional vertebrate fossils to be collected from this area.

Permian

Pakoon Limestone

The Pakoon Limestone is the western lateral marine equivalent to the Esplanade Sandstone in the Grand Canyon and consists of thin- to thick-bedded dolomitic limestones, which inter-tongue with the Esplanade Sandstone (McKee 1982). These beds mark the last member of the deposition of the Supai Group, which is followed by the Hermit Formation. The Pakoon Limestone extends from the western Grand Canyon and north into southwestern Utah and is Wolfcampian (early Permian) in age (McKee 1982). An abundance of marine fossils, which include corals, pelmatozoans, and bivalves, are known from the western margins of the Pakoon Limestone, whereas the Esplanade Sandstone only has a fragmentary invertebrate fossil record (McKee 1982).

History

McKee (1982) recorded the presence of two chondrichthyan tooth types from the Pakoon Limestone, *Cladodus* sp. and *Deltodus* sp., from the westernmost end of the Grand Canyon. Both taxa were identified by David Dunkle. The *Cladodus* sp. identification is most likely another cladodont shark taxon as *Cladodus* historically served as a "waste bin" for Paleozoic cladodont-grade chondrichthyan teeth. *Cladodus sensu strictu* is a genus of ctenacanth shark found only in the Mississippian (Duffin and Ginter 2006). Hunt et al. (2005) erroneously referred these specimens from the Pakoon Formation to the younger Kaibab Formation.

GRCA Resources

At present, no specimens of *Cladodus* or *Deltodus* from the Pakoon Limestone of the Grand Canyon have been transferred to any museum collection and are considered here to be missing.

Potential

The fossiliferous nature of the Pakoon Formation with its high abundance of marine invertebrates (McKee 1982) and the record of at least two possible chondrichthyan taxa indicates a strong potential for more vertebrate material to be recovered where the Pakoon Formation is exposed.

Kaibab Formation

The marine limestones, dolomitic limestones, and sandstones of the Kaibab Formation are extensive throughout northern Arizona, southern Utah, east-central Nevada and southeast California. At the Grand Canyon, the Kaibab Formation forms the upper rim, which many park visitors stand upon while looking out over the canyon. The age of the Kaibab Formation is late early Permian (Leonardian/Kungurian) to early middle Permian (Guadalupian/Roadian) based on conodont analysis and the invertebrate assemblages (Thompson 1995; Hopkins and Thompson 2003). The depositional environment of the Kaibab represents a broad continental shelf exceeding approximately 125 km

(200 mi) in width (Hopkins and Thompson 2003). Frequent periods of sea level changes led to major lateral shifts of supratidal, subtidal, and shallow-marine environments that led to complex interlayering of different types of carbonate and clastic sediments (Hopkins and Thompson 2003). McKee (1938) proposed the division of the Kaibab Formation into three members, from oldest to youngest, the gamma, beta, and alpha. Sorauf (1962) combined the beta and gamma members into a single unit, the Fossil Mountain Member, and renamed the alpha member as the Harrisburg Member.

Previously, vertebrate fossils have been considered rare within the Kaibab Formation (David 1944). However, vertebrate fossils from the Kaibab are now known from multiple locations within GRCA and southward towards the Flagstaff metropolitan area (Hodnett et al. 2011a, 2011b, 2012; Elliott and Hodnett 2013). Between GRCA collections and those from the Flagstaff localities the Kaibab Formation has one of the most extensive marine vertebrate assemblages for the global Permian, with an estimated 40+ chondrichthyan taxa and a large number of indeterminate osteichthyans including actinopterygians and coelacanths (Hodnett et al. 2011a, 2011b, 2012; Elliott and Hodnett 2013). This high number of taxa comes from both macro and micro vertebrate remains.

History

The earliest account of a vertebrate fossil from the Kaibab of the Grand Canyon came from an anonymous note in 1926 that reported fish remains consisting of “imprint of ribs and backbone” coming from the Kaibab Formation and collected by A. L. Brown from the North Rim. Spamer (1984) later relayed a personal communication with E. D. McKee that this fish specimen was in fact the scyphozoan *Conularia kaibabensis*. McKee (1938) reported the first definitive fish remains from the Grand Canyon. He examined three teeth in the USGS paleontology collection that had been collected by C. D. Walcott in 1882, which he indicated came from the Alpha Member (now Harrisburg Member) of the Kaibab Formation. McKee gave no taxonomic identification for the three teeth Walcott collected. McKee collected and figured (1938: Plate 15, Figure 4) a mandibular tooth plate of *Deltodus mercurii* (Figure 6-6B) from an undisclosed locality in the Beta Member (Fossil Mountain Member) at the Grand Canyon.

Hussakoff (1943) described a number of fish teeth and a fin spine from a site known locally as the “Bottomless Pits,” northwest of Walnut Canyon National Monument and is on the east side of Flagstaff, Arizona. The taxa identified by Hussakoff (1943) include the teeth of the chondrichthyans *Deltodus mercurii*, *Psephodus* sp., and *Janassa* sp., a euselachian shark spine fragment referred to “*Hybodus* sp.,” and isolated actinopterygian teeth. A year later, David (1944) described the type specimen of *Megactenopetalus kaibabanus*, which consisted of an upper symphysial tooth of a large species of petalodontimorph chondrichthyan from the North Rim of GRCA (Figure 6-5A–B). Additional specimens of *Megactenopetalus* have been collected from the Kaibab Formation near Sedona, Arizona, and juvenile teeth have recently been identified from Flagstaff, Arizona (Ossian 1976; Hodnett et al. 2011b). Outside of the Kaibab Formation, *Megactenopetalus* has been found in the approximately contemporaneous Permian sediments of the Concha Limestone in southeastern Arizona (Hansen 1978) and New Mexico (Ossian 1978; Hunt and Lucas 2005), the Glass Mountain region of Texas (Ossian 1976; Hunt and Lucas 2005), China (Young 1950), and Iran (Golshani and Janvier 1974).




Figure 6-5. Chondrichthyan fossils collected from the Kaibab Formation on the North Rim of Grand Canyon National Park (JOHN-PAUL HODNETT). **A–B.** MNA V. 1367, holotype of *Megactenopetalus kaibabanus:* **A.** Labial view; **B.** Orolingual view. **C–D**, MNA Pl. 562, *Psephodus* sp.: **C.** Oral view; **D.** Distal view. Scale equals 1 cm (0.4 in).

Gass (1963) briefly described a large tooth plate he referred to *Psephodus* sp. that was also collected on the North Rim of GRCA. The genus *Psephodus* is known primarily from the early Carboniferous (Mississippian) of Europe and North America, but there are records of Permian *Psephodus* from Pakistan and Russia (Stahl 1999). The GRCA *Psephodus* would be the first Permian record of this taxon in North America. Johnson and Zidek (1981) reported on the dental palatal plates of platysomid fishes from the GRCA Bright Angel Trail (Figure 6-6C) and a locality within the Kaibab National Forest. Additional Kaibab platysomid tooth plates and isolated teeth were collected within the Flagstaff vicinity and in central Arizona (Johnson and Zidek 1981). Recently the mold of a large platysomid tooth plate was found in situ in a block of Kaibab Formation along the Hermit Trail by S.G. Lucas (New Mexico Museum of Natural History and Science, Albuquerque) in 2018 but not collected. Platysomid fishes are deep-bodied actinopterygians and are known from the Pennsylvanian to the Permian of Europe and North America (Johnson and Zidek 1981; Zidek 1992; Mickle and Bader 2009).




Figure 6-6. Kaibab vertebrate fossils in GRCA collections (JOHN-PAUL HODNETT). **A.** GRCA 10030, euselachii indeterminate dorsal fin spine. **B.** GRCA 9368, *Deltodus mercurii* mandibular tooth plate. **C.** Platysomid indeterminate fish tooth plate. **D.** GRCA 10326, indeterminate actinopterygian cleithrum. Scale equals 1 cm (0.4 in).

Thompson (1995) briefly described a number of vertebrate micro-remains from the Fossil Mountain type section within GRCA. Vertebrate fossils include the teeth of an indeterminate hybodont (identified as *Polyacrodus* cf. *P. witchitaensis* in Hodnett et al. 2011a) and the early neoselachian *Cooleyella peculiaris*, a taxon related to *Cooleyella platera* from the Upper Mississippian Surprise Canyon Formation (Hodnett and Elliott 2018). Thompson also reported a tooth of *Symmorium* cf. *S. reniforme*, a symmoriid chondrichthyan from the Upper Pennsylvanian of the Black Shales (Ginter et al. 2010). We reviewed these specimens and they appear to be worn teeth of *Heslerodus divergens*, a taxon known from the Kaibab Formation near Flagstaff (Hodnett et al. 2012) and a new record of a jalododont chondrichthyan. Thompson (1995) also identified a number of isolated dermal and buccal membrane denticles of chondrichthyians such as *Cooperella striatulata, Sturgenonella quinqueloba, Moreyella typicalis, Kirkella typicalis*, and a *Petrodus*-like taxon. These denticles are often associated with multiple different family groups of chondrichthyans including hybodonts, petalodonts, symoriids, and holocephalans. A number of isolated indeterminate actinopterygian teeth

have also been identified from the Fossil Mountain locality (Thompson 1995). Hodnett et al. (2011a, 2011b, 2012, 2013) and Elliott and Hodnett (2013) reported on a number of new chondrichthyan taxa from the Kaibab Formation. These include at least seven taxa of ctenacanthiform sharks, a xencanthimorph, seven hybodontiform sharks, and six petalodontimorph chondrichthyans; most of which were collected from the vicinity of Flagstaff, Arizona. Stable and radiogenic isotope analyses of the Flagstaff Kaibab chondrichthyan teeth showed depleted δ^{18}OP values giving a signature of extreme salinity and warm waters for the Kaibab assemblage (Fisher et al. 2014). However, it was determined that the Kaibab tooth samples used had gone through a diagenetic alteration that skewed the results, making accurate interpretation not possible (Fisher et al. 2014).

GRCA Resources

At present, the teeth collected by C. D. Walcott in 1882 reported in McKee (1938) have not been relocated in the USGS Collections.

MNA: The two teeth collected on the North Rim of GRCA are housed at the Museum of Northern Arizona, Flagstaff. The type of *Megactenopetalus kaibabanus* (MNA V. 1367) is on permanent display in the Geology Hall of the museum gallery, while the *Psephodus* sp. (MNA Pl. 562) tooth plate is in the paleontology collections (Figures 6-5C and 5D). The vertebrate samples reported in Thompson (1995) are all housed at the Museum of Northern Arizona. The platysomid tooth plates reported in Johnson and Zidek (1981) are a cast (MNA PL. 556) and a tooth plate (MNA PL. 43).

GRCA: Specimens (excluding casts) collected from the Kaibab at Grand Canyon National Park include: GRCA 9368, *Deltodus mercurii* mandibular tooth plate (from McKee 1938: 166, Plate 15, Figure 14) GRCA; GRCA 10030, euselachian indeterminate dorsal fin spine; GRCA 10229 (Figure 6-6A), platysomid tooth plate (may be the same specimen listed as a cast in Johnson and Zidek 1981 under MNA PL. 556); GRCA 10326, indeterminate actinopterygian cleithrum (Figure 6-6D).

Potential

Based on previous reports and new work conducted at fossil sites in the Flagstaff vicinity, there is an extremely high potential for the Kaibab exposed at GRCA to yield a significant amount of vertebrate fossils.

Conclusions

As demonstrated above, the Grand Canyon has a surprisingly rich Paleozoic body fossil record, although the majority of these records are of the teeth and dermal spines of chondrichthyan fishes. This is not surprising as much of the geologic history of the Paleozoic at the Grand Canyon is the record of multiple marine transgression and regression events. Unfortunately, even though the literature lists multiple records of vertebrate body fossil remains, many of those records are not supported by physical specimens in museum collections for verification or further research. As such, it our opinion that a need to return to these sites to look for and collect additional vertebrate material is warranted.

The Devonian through Permian rocks of Grand Canyon National Park have a high potential to reveal a further wealth of information to understand the dynamics of a constantly changing physical world

and its impact on the organisms that lived during those times. In terms of an advance in our knowledge of paleoichthyology, the geographical position and the amount of time covered at Grand Canyon could give us a large dataset of how fishes evolved during the Paleozoic, as there are few localities in the American southwest that expose such an extensive geological sequence of time for marine and near-marine sedimentary rocks. There is good potential for the occurrence of early tetrapods in the Devonian Temple Butte Formation, as well as collecting additional tetrapod remains from the Upper Mississippian Surprise Canyon Formation.

Acknowledgments

We would like to thank Colleen Hyde of the Grand Canyon Museum for her assistance in locating and providing photographs of specimens under her care. We thank Janet and David Gillette for their assistance with viewing and documenting fossils at the Museum of Northern Arizona and their continued support of our research. We also thank Amanda Millhouse for her assistance in facilitating access to the Paleobiology Collections at the National Museum of Natural History. Special thanks to Vincent Santucci and Justin Tweet of the National Park Service Paleontology Program (Geologic Resource Division) for assistance with specimen records and references. Justin Tweet, Anne Miller, Diana Boudreau, Klara Widrig, Tim Connors, and Vincent Santucci made comments and suggestions that improved this manuscript. We thank Spencer Lucas and Matt Friedman for reviewing the manuscript.

Literature Cited

Anonymous. 1926. Fossil fish. Grand Canyon Nature Notes 1(7):4.

Babcock, L. E., and R. E. Feldmann. 1986. Devonian and Mississippian conulariids of North America. Part B: *Parconularia*, *Reticularconularia*, new genus and organisms rejected from Conulariida. Annals of Carnegie Museum 55(16):411–479.

Beus, S. S. 1985. Late Mississippian fossil communities in the Surprise Canyon Formation, an estuarine sequence in Grand Canyon, Arizona. Abstracts of the Symposium on South Western Geology and Paleontology, Museum of Northern Arizona, Flagstaff: 1.

Beus, S. S. 1986. Biostratigraphy of the Surprise Canyon Formation (Latest Mississippian and Earliest Pennsylvanian) in western Grand Canyon. Abstracts with Programs - Geological Society of America 18(5):341.

Beus, S. S. 1999. Megafossil paleontology. Pages 69–96 *in* G. H. Billingsley and S. S. Beus, editors. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

Beus, S. S. 2003a. Temple Butte Formation. Pages 107–114 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Beus, S. S. 2003b. Redwall Limestone and Surprise Canyon Formation. Pages 115–134 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Billingsley, G. H. 1978. A synopsis of the stratigraphy in the western Grand Canyon, Arizona. Museum of Northern Arizona Press Research Paper 16:7–9.

Billingsley, G. H., and S. S. Beus. 1985. The Surprise Canyon Formation: An Upper Mississippian and Lower Pennsylvanian(?) rock unit in the Grand Canyon, Arizona. Pages A27–A33 *in* Stratigraphic notes, 1984. U.S. Geological Survey, Washington, D.C. Bulletin 1605-A.

Billingsley, H., and E. D. McKee. 1982. Pre-Supai buried valleys. Pages 137–153 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

Denison, R. H. 1951. Late Devonian fresh-water fishes from the western United States. Fieldiana: Geology 11(5):221–261.

Duffin, C. J., and M. Ginter. 2006. Comments on the selachian genus *Cladodus* Agassiz, 1843. Journal of Vertebrate Paleontology 26(2):253–266.

Elliott, D. K., and R. C. Blakey. 2005. The Pre-Permian vertebrate record in Arizona. New Mexico Museum of Natural History and Science Bulletin 29:1–8

Elliott, D. K, and J.-P. M. Hodnett. 2013. A new species of *Bransonella* (Chondrichthyes, Xenacanthimorpha, Bransonelliformes) from the Middle Permian Kaibab Formation of Northern Arizona. Journal of Paleontology 87(6):1136–1142.

Fisher, J., J. W. Schneider, J.-P. M. Hodnett, D. K. Elliott, G. D. Johnson, S. Voight, M. M. Joachimski, M. Tichomirowa, and J. Gotze. 2014. Stable and radiogenic isotope analysis on shark teeth from the Early to the Middle Permian (Sakmarian-Roadian) of the southwestern USA. Historical Biology 26(6):710–727.

Gass. H. L. 1963. A review of the Paleozoic fish of Arizona. Thesis. University of Arizona, Tucson, Arizona.

Gordon, M., Jr. 1982. Biostratigraphy of the Watahomigi Formation. Pages 113–135 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

Hansen, M. C. 1978. A presumed lower dentition and spine of a Permian petalodontiform chondrichthyan, *Megactenopetalus kaibabanus.* Journal of Paleontology 52(1):55–60.

Hodnett, J.-P. M., and D. K. Elliott. 2018. Carboniferous chondrichthyan assemblages from the Surprise Canyon and Watahomigi formations (latest Mississippian–Early Pennsylvanian) of the western Grand Canyon, northern Arizona. Journal of Paleontology 92, Memoir 77.

Hodnett, J.-P. M., D. K. Elliott, and T. J. Olson. 2011a. The hybodontiform sharks (Chondrichthyes) from the marine Permian (Leonardian) Kaibab Formation of northern Arizona. Ichthyolith Issues, Special Publication 12:23–24.

Hodnett, J.-P. M., D. K. Elliott, and T. J. Olson. 2011b. The Petalodontiformes (Chondrichthyes; Euchondrocephali) from the marine Permian (Leonardian/Guadeloupian) Kaibab Formation, Northern Arizona. Journal of Vertebrate Paleontology 31(supplement 1):126.

Hodnett, J.-P. M., D. K. Elliott, T. J. Olson, and J. H. Wittke. 2012. Ctenacanthiform sharks from the Permian Kaibab Formation, northern Arizona. Historical Biology 24(4):381–395.

Hodnett, J.-P. M., D. K. Elliott, and T. J. Olson. 2013. A new basal hybodont (Chondrichthyes, Hybodontiformes) from the Middle Permian (Roadian) Kaibab Formation, of Northern Arizona. New Mexico Museum of Natural History and Science 60:103–108

Hopkins, R. L., and K. L. Thompson. 2003. Kaibab Formation. Pages 196–211 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Hunt, A. P., and S. G. Lucas. 2005. The chondrichthyan *Megactenopetalus kaibabanus* from the Early-?Middle Permian of southern New Mexico and adjacent area of Texas. New Mexico Museum of Natural History and Science Bulletin 30:117–118.

Hunt, A. P., S. G. Lucas, V. L. Santucci, and D. K. Elliott. 2005. Permian vertebrates of Arizona. New Mexico Museum of Natural History and Science Bulletin 29:10–15.

Hussakoff, L. 1943. Permian fishes from the Kaibab Formation. Geological Society of America Bulletin 54(12):1834.

Johnson, G. D., and J. Zidek. 1981. Late Paleozoic phyllodont tooth plates. Journal of Paleontology 55(3):524–536.

Lucas, S. G., and G. S. Morgan. 2005. Pleistocene mammals of Arizona: An overview. New Mexico Museum of Natural History and Science Bulletin 29:152–157.

Martin, H., and J. E. Barrick. 1999. Conodont biostratigraphy. Pages 97–116 *in* G. H. Billingsley and S. S. Beus, editors. 1999. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

McKee, E. D. 1938. The environment and history of the Toroweap and Kaibab formations of northern Arizona and southern Utah. Carnegie Institution of Washington Publication 492.

McKee, E. D. 1982. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D., and R. C. Gutschick. 1969. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Mead, J. I. 2005. Late Pleistocene (Rancholabrean) amphibians and reptiles of Arizona. New Mexico Museum of Natural History and Science Bulletin 29:136–151.

Mickel, K. E., and K. Bader. 2009. A new platysomid from the Upper Carboniferous of Kansas (USA) and remarks on the systematics of deep-bodied lower actinopterygians. Acta Zoologica 90(supplement 1):211–219

Noble, L. F. 1922. A section of the Paleozoic formations of the Grand Canyon at Bass Trail. Pages 23–73 *in* Shorter contributions to general geology, 1922. U.S. Geological Survey, Washington, D.C. Professional Paper 131-B.

Ossian, C. R. 1976. Redescription of *Megactenopetalus kaibabanus* David 1944 (Chondrichthyes: Petalodontidae) with comments on its geographic and stratigraphic distribution. Journal of Paleontology 50(3):392–397.

Sando, W. J. 1969. Corals. Pages 257–343 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Schuchert, C. 1918. On the Carboniferous of the Grand Canyon of Arizona. American Journal of Science (4^{th} series) 45:347–361.

Skipp, B. 1969. Foraminifera. Pages 173–255 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Sorauf, J. E. 1962. Structural geology and stratigraphy of the Whitmore area, Mohave County, Arizona. Dissertation. University of Kansas, Lawrence, Kansas.

Spamer, E. E. 1984. Paleontology in the Grand Canyon of Arizona: 125 years of lessons and enigmas from the Late Precambrian to the present. The Mosasaur 2:45–128.

Stahl, B. J. 1999. Chondrichthyes III, Holocephali. Volume 4 of Handbook of Paleoichthyology, H.-P. Schultze, editor. Verlag Dr. Friedrich Pfeil, Munich, Germany.

Thompson, K. L. 1995. Paleoecology and biostratigraphy of the Fossil Mountain Member, Kaibab Formation, in northwestern Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Walcott, C. D. 1880. The Permian and other Paleozoic groups of the Kanab Valley, Arizona. The American Journal of Science (3^{rd} series) 20:221–225.

Zidek, J. 1992. Late Pennsylvanian Chondrichthyes, Acanthodii, and deep-bodied Actinopterygii from the Kinney Quarry, Manzanita Mountains, New Mexico. New Mexico Bureau of Mines and Minerals Resources Bulletin 138:145–182.

Chapter 7. Paleozoic Paleobotany of Grand Canyon National Park

By Cassi Knight[1]

[1]Paleoworks Consulting
2045 NW 29th Ave
Portland, Oregon 97210

Introduction

Grand Canyon National Park (GRCA) is certainly one of the most awe-inspiring natural sculptures on Earth. From the rich black Vishnu Schist lining the depths of the canyon, to the shining white cliffs of the Kaibab Formation perched on the canyon rim, the exposed rocks display approximately one-third of all of Earth's history in a well-exposed geologic section. Each of the rock layers that form the intricacies of Grand Canyon's walls represents a different environment that existed in this place as the Colorado Plateau changed over hundreds of millions of years. Many of the sediments deposited in these different environments preserve fossils that provide an immense wealth of information about the evolution of life on earth.

Grand Canyon exposes rocks that were formed starting in the Precambrian through the middle Permian in the Paleozoic and are ~1800–270 million years (Ma) old. The most visible rocks in the canyon—the flat lying layers that dominate many vistas from the rim—are the Paleozoic portion of Grand Canyon's stratigraphy. During the early and middle Paleozoic, the Colorado Plateau was largely inundated by shallow seas and the marine sediments deposited during this time (~545–335 Ma) reflect these conditions. Local sea level started falling during the Mississippian, and the resulting deposits of the Surprise Canyon Formation, Supai Group, Hermit Formation, and Coconino Sandstone represent a range from nearshore to estuarine to terrestrial environments. The fossil plants reported from Grand Canyon National Park are found in the Surprise Canyon Formation, members of the Supai Group, and the Hermit Formation.

The Hermit Formation preserves the most diverse and well-preserved flora from Grand Canyon, with 42 different species reported, and 25 type specimens known from the associated collections (Appendix 7-A). There are seven identified plant taxa reported from the Supai Group and 12 from the Surprise Canyon Formation, many of which are fragmentary and more poorly preserved. Whereas plant fossils are not overly abundant in Grand Canyon, they are certainly significant because they add to the limited record of Mississippian, Pennsylvanian, and Permian age paleofloras from the western margin of the Pangean supercontinent, a region subject to considerably different climatic conditions than contemporaneous areas in the more central Euramerican regions.

History of Research

The first paleobotanical remains from the Grand Canyon were discovered by Charles Schuchert in 1915 in the Hermit Formation along the Hermit Trail in the Waldron Basin at a locality called Red Top. Schuchert sent a small collection of these specimens to U.S. Geological Survey (USGS) paleobotanist David White for identification. White noted *Walchia*, "*Gigantopteris"* (likely *Supaia*),

Sphenophyllum, and *Callipteris* in this initial sampling, and these findings were published in 1918 (Schuchert 1918; White 1929b). A few more plant fossils were found and reported to White by Levi F. Noble and Charles Gilmore between White's 1918 publication and his Carnegie Institution funded collecting trips during the summers of 1926–1928. White's systematic collecting during these field trips yielded the specimens that he would publish in his 1929 monograph on the Hermit Formation paleoflora. Most of his plant collection came from a quarry he made along the Yaki Trail (now called the South Kaibab Trail) and is marked today by the "fossil fern exhibit" constructed by the Civilian Conservation Corps (CCC) in 1937 just to the west of the trail on Cedar Ridge (Figure 7-1).



Figure 7-1. A. Paleobotanist David White (left) and John C. Merriam working in the Cedar Ridge fossil quarry during field work in the summer of 1927 (NPS). **B.** Fossil Fern Exhibit constructed by the Civilian Conservation Corps (CCC) at the site of White's quarry on Cedar Ridge, South Kaibab Trail (NPS).

David White also spent time searching the Supai Group for plant fossils, particularly during the summer of 1928, and published a mention of his findings in his 1929 Hermit Formation monograph (White 1929a). White reported the conifer *Walchia* and algal trace fossil *Rivularites* from multiple horizons in the lower and middle Supai, and an assemblage of *Walchia*, *Taeniopteris, Neuropteris*, (*incertae sedis*), *Cordaites* (conifer), and *Calamites* (horsetails) from 7.6 m (25 ft) above the base of the Watahomigi Formation near the South Kaibab Trail (McKee 1982). Plant fossils were next reported from the Supai Group by Edwin D. McKee in a large USGS report, having been found during field work supporting this publication. From many measured sections throughout the Supai, occurrences of *Walchia*, stems, ferns, *Rivularites*, stromatolites, miscellaneous algae, and bioclasts are noted (McKee 1982).

The Surprise Canyon Formation is the most recently recognized sediment package in GRCA. USGS geologist George Billingsley first noticed the large dark red-brown channel fills sitting on top of the Redwall Limestone during mapping reconnaissance flights in the 1970s (Beus 1986). Between 1975 and 1983, Billingsley and Beus measured sections for many of the larger Surprise Canyon outcrops and observed a fairly consistent assemblage of plant fossils in the terrestrial deposits at the base of these channels. They repeatedly reported tree impressions, wood fragments, and leaves, and spores attributable to ferns and lycopods like *Lepidodendron* (Beus 1986). A brief description of this flora was published in Billingsley and McKee (1982), with USGS paleobotanist Serge Mamay identifying

two ferns and five sphenophytes. Other unidentified plant fragments and 22 spore and pollen taxa were described in this report as well, but none of the plant or palynological specimens were figured (Tidwell et al. 1992). Beus later collected 40 additional plant specimens, and these contributed to a closer review of the Surprise Canyon flora by Tidwell et al. (1992) who identified a total of 12 plant taxa.

Stratigraphic Distribution of Fossils

The Paleozoic rocks of Grand Canyon include, from oldest to youngest, the Tapeats Sandstone, Bright Angel Shale, Muav Limestone, Temple Butte Formation, Redwall Limestone, Surprise Canyon Formation, the Supai Group, Hermit Formation, Coconino Sandstone, Toroweap Formation, and Kaibab Formation (McKee 1982). Some portions of the Surprise Canyon Formation, the Supai Group, and all of the Hermit Formation were deposited under terrestrial and/or estuarine conditions and account for the Paleozoic floras reported from GRCA. The Surprise Canyon Formation represents ancient coastal plain and estuarine environments and records a sparse assemblage of fossil plants. The Supai Group preserves sporadic plants in some of its members, in the sandy and silty terrestrial deposits left by intervals of low sea level. The Hermit Formation was deposited in a seasonally dry (Cecil 2003; Gastaldo and Demko 2011) lowland environment, an interpretation based on the composition of the paleoflora, and preserves the most robust fossil plant assemblage in the canyon.

Plant fossils from the Hermit Formation and the Supai Group are mostly known from eastern portions of GRCA, especially from outcrops near the South Kaibab, Bright Angel, and Hermit Trails. This is somewhat unsurprising, given the difficulty and exposure of the general terrain—prospecting for new fossil sites can be rather challenging. Conversely, the Surprise Canyon flora is mostly reported from sites in western GRCA, where the isolated channels of this formation are largest. Most of the access to these sections was helicopter supported through a USGS project, so the fossil sites reflect a larger spatial distribution and are largely inaccessible by foot.

Surprise Canyon Formation

The Surprise Canyon Formation is latest Mississippian in age (~325 Ma) and is exposed throughout Grand Canyon as isolated lenses filling old erosional valleys, caves, and karst collapse structures in the top of the Redwall Limestone (Billingsley and Beus 1985; Beus 1986). The conglomerate and mudstone beds that make up the channel fills were originally considered to be part of the Watahomigi Formation, the basal unit of the Supai Group. The rocks that form the Surprise Canyon Formation were only identified as a unique sediment package in 1979 by George Billingsley (Billingsley 1979), and formally described by Billingsley and Beus (1985). The Surprise Canyon Formation occurs throughout Grand Canyon and in parts of Marble Canyon, and is rather difficult to identify because of its discontinuous nature and because its larger, more visible outcrops occur in remote areas of western Grand Canyon (Tidwell et al. 1992).

In the western Grand Canyon, the Surprise Canyon Formation occurs as thicker sections in western Grand Canyon with channels typically 45 to 60 m (150 to 200 ft) wide, and as thick as 120 m (400 ft) (Tidwell 1992). It is divided into three major units: 1) terrestrial conglomerate and sandstone; 2) marine skeletal limestone; and 3) marine siltstone and silty to sandy limestone (Tidwell 1992). Most

channels, however, only contain the basal terrestrial and the upper fine-grained clastic and carbonate rocks. Plant fossils are found only in the sandstone, siltstone, and shale beds of the lower unit, particularly in western GRCA, and include tree impressions, wood fragments, and plant compressions (Billingsley and Beus 1985).

Identified taxa include the horsetails *Calamites cistiiformis*, *Calamites* sp., the lycopods *Lepidodendron aculeatum*, *L. mannabachanese*, *L. volkmannianum*, *Lepidostrobus* sp. cf. *L. ornatus*, *Lepidostrobophyllum* sp., *Stigmaria ficoides*, the ferns *Pecopteris* cf. *P. aspera*, the pteridosperm seed *Wardia* sp., and *Cyperites* sp. and *Knorria* sp. (*incertae sedis*) (Beus 1990, 1995; Tidwell et al. 1992; Hodnett and Elliot 2018). There are also poorly preserved plants that Billingsley and McKee (1982) reported, based on personal communication with Mamay, as two ferns (one described as "a very delicate, small-pinnuled sphenopterid or pecopterid"), five sphenopsids ("the sphenopsid fragments are parts of the fruiting organs of a phyllothecoid plant"), and other unidentifiable plant fragments. A brief mention of stromatolites at the top of the Surprise Canyon sequence at the Quartermaster Canyon section was made by Beus (1990). While this flora is neither well-preserved, abundant, or diverse, it is significant because Mississippian and early Permian age floras are rare in the western US, and this is the first flora of this age reported from Arizona (Tidwell et al. 1992).

In addition to the megaflora, an assemblage of 22 pollen and spore taxa was recovered from the Surprise Canyon Formation (see list below) and suggests that this plant community was more diverse than its megafossil record implies (Billingsley and McKee 1982; Tidwell et al. 1992). The palynological samples were collected from the basal terrestrial sand and conglomerate unit at the Granite Park section locality (Billingsley and McKee 1982).

Per Billingsley and McKee (1982), pollen and spore taxa known from the Surprise Canyon Formation include the following:

- *Anapiculatisporites concinnus*
- *Anaplanisporites globulus*
- *Auroraspora solisortus*
- *Calamospora* sp.
- *Convolutispora florida*
- *Convolutispora* sp.
- Dictyotriletes cf. *D. clatriformis*
- *Granulatisporites granulatus*
- *Granulatisporites* sp.
- *Hadrohercos stereo*
- *Knoxisporites triradiatus*
- *Lycospora* spp.
- *Punctatisporites* cf. *P. heterofiliferous*

- *Punctatisporites* cf. *P. nitidus*
- *Punctatisporites pseudolevatus*
- *Punctatisporites solidus*
- *Punctatisporites validus*
- *Punctatisporites* spp.
- *Reticulatisporites* sp.
- *Schulzospora rara*
- *Triquitrites* sp.
- Monosaccates indet.
- Unassigned (? Algal)

Lepidodendrales

Several fossils from the Surprise Canyon flora represent different organ or tissue types of arborescent lycopsid, an extinct plant group commonly known as scale trees or giant clubmosses. The trunk impressions that preserve different leaf cushion patterns belong to three species of the genus *Lepidodendron*, the spore producing lycopsid cones are assigned to *Lepidostrobus* sp. *cf. L. ornatus*, isolated sporophyll impressions to *Lepidostrobophyllum* sp., and the root structures to *Stigmaria ficoides* (Tidwell et al. 1992). Based on the leaf cushion patterns on the trunks' exterior, three different types of lycopsid bark were identified: *Lepidodendron aculeatum*, *L. mannabachanese*, and *L. volkmannianum* (Tidwell et al. 1992).

Filicales and Pteridosperms

Fragmentary remains of the marattialean tree fern frond *Pecopteris* sp. were reported by Tidwell et al. (1992), who mentioned it closely resembles *P. aspera*, but there is not sufficient evidence to conclusively make this identification. Seeds that have been recovered from limestone beds were reported as *Wardia* sp.—platyspermic winged seeds: bilaterally flattened, spherical bodies with small wing-like projections along their lateral flanks. Based on other floras from North America, *Wardia* sp. is associated with the compound, pinnate (fern-like) foliage of the pteridosperm *Aneimites* (Tidwell et al. 1992).

Incertae Sedis

Several impressions of what appear to be the subsurface bark layer of lycopsid trunks are tentatively reported from the Surprise Canyon flora, and have been listed as *Knorria* sp. (Tidwell et al. 1992). These fossils show the pattern of leaf vasculature where it connects from the lycopsid trunk to its leaves, and the best preserved specimen shows helically arranged leaf bases, but lacking features allowing more detailed classification. Arborescent lycopsids belong to the still-extant order Isoetales, commonly known as the quillworts, one of three extant orders in the class Lycopsida, the earliest still extant group of vascular plants (Taylor et al. 2009). Additionally, isolated, fragmentary leaves of *Cyperites* sp. are poorly preserved in the Surprise Canyon Formation and their taxonomic

relationships to particular lycopsid stem taxa is indeterminate (most of the stem forms produced very similar, grass-like leaves that are difficult to differentiate in compression preservation).

Supai Group

The Supai Group spans the Early Pennsylvanian to the middle early Permian (315–285 Ma), and crops out prominently as a red, stair-stepped slope throughout the Grand Canyon (McKee 1982). The erosion resistant Esplanade Sandstone at its top creates the iconic Esplanade Platform in the western reaches of GRCA. Sediments of the Supai Group were deposited during the late Paleozoic Ice Age. The sediment packages range from terrestrial to fully marine deposits (with intermediate environments also represented), reflecting a time of great sea-level fluctuation caused by global glacial-interglacial cycles (Fielding et al. 2008; Montañez and Poulsen 2013). The Supai Group records four separate marine transgressions, and each distinct event corresponds to one of the four formations in the Supai Group: the Watahomigi Formation, Manakacha Formation, Wescogame Formation, and the Esplanade Sandstone (McKee 1982). With each transgression, the sea level rose progressively higher and flooded increasingly large portions of Arizona and the San Juan Basin (New Mexico, Utah, and Colorado) (McKee 1982).

Each of the four members of the Supai Group have some portion of terrestrial deposits that contain plant fossils belonging to the horsetails, seed ferns, conifers, and plant remains of undetermined affiliation (White 1929a; McKee 1982). These terrestrial sediments largely consist of siltstones and mudstones, and somewhat commonly preserve mud crack and raindrop surfaces (White 1929a; McKee 1982). From the marine limestones of the Supai, fossils identified as algae or algal structures, and several types of bioclasts have also been reported (McKee 1982).

Watahomigi Formation

This marine deposit is composed mostly of limestone that grades into sandstone and siltstone beds that are thicker in the eastern part of the Grand Canyon. The environment of deposition is interpreted to be a relatively shallow, low energy embayment. Plants reported from the Watahomigi Formation include the horsetail relative *Calamites* sp., the seed ferns *Neuropteris* sp., the conifer relative *Cordaites* sp., the conifer *Walchia* sp., *Taeniopteris* sp. (*incertae sedis*), and other unidentified plant fragments. These are found in the siltstone and sandstone beds in the eastern Grand Canyon near the base of the Watahomigi. These beds also preserve mud crack and rain drop surfaces, and their equivalent facies to the west are limestone (White 1929a; McKee 1982). Miscellaneous algal structures are also reported from the Watahomigi Formation, but without much accompanying information (White 1929a; McKee 1982).

Manakacha Formation

The Manakacha Formation was deposited during a second, larger transgression that flooded further inland to the northeast. Hence, it is mostly marine sandstone with some carbonate rock. The only identifiable plant foliage reported from the Manakacha Formation are ultimate branches of the conifer *Walchia* sp., and an undetermined fern (Filicopsida), which are found in shaly mudstone beds. Other reported plant fossils include algal limestone nodules, *Girvanella* bioclasts and calcisphere bioclasts (White 1929a; McKee 1982). The *Girvanella* and calcisphere bioclasts occur as bioclastic debris in carbonate facies in central and eastern Grand Canyon.

Wescogame Formation

The base of the Wescogame Formation lies on an unconformable surface marked by large (up to 24 m/80 ft deep) channel cuts (McKee 1982). The channel cuts are filled with conglomerate, and the overlying Wescogame Formation above consists largely of sand and mud in eastern Grand Canyon, that grades into limestone to the west. The Wescogame Formation, having been deposited in a higher energy marine environment, does not include well-preserved plant remains. Plants found in this formation are miscellaneous algae, including *Rivularites* (Spamer 1992), and possible algal and organic features (White 1927; McKee 1982). Additionally, some occurrences of *Walchia* sp. have been noted in the Grand Canyon area by Spamer (1992). "Fern or cycadofilice" fragments are reported from shaly beds with many mud crack and rain pit surfaces, but these may be from the overlying Esplanade Sandstone (Schuchert 1918; McKee 1982).

Esplanade Sandstone

The Esplanade is the thickest and most extensive member of the Supai Group. Its base is mostly mudstone and siltstone, coarsening upwards to cross bedded sandstone, which is especially prominent in central and eastern Grand Canyon (McKee 1982). The Esplanade Sandstone preserves rare shoots of *Walchia* sp., some undetermined plants, and plant fragments (White 1929a; McKee 1982).

Hermit Formation

The Hermit Formation, often referred to as the Hermit Shale, is early Permian in age (~280 Ma), and forms a soft, deep red slope near the top of Grand Canyon. It is composed of interbedded mudstones, shale, and fine sandstones, and represents sediments accumulating on a broad coastal plain in an environment that experienced strong seasonal drought (White 1929a). The Hermit Formation is one of very few fully terrestrial deposits in Grand Canyon and preserves by far the most detailed and diverse paleoflora from the park. The Permian represents a time of dramatic fluctuation in climate, in response to global cyclic glaciation (Fielding et al. 2008; Montañez and Poulsen 2013). In addition to these overall global-scale cycles, a more seasonal drought regime is interpreted for equatorial Pangea, in which dry periods became increasingly drier (Fielding et al. 2008; Montañez and Poulsen 2013). The Hermit Formation provides unique insight into less-often preserved plant communities that inhabited the continental basins during drier times.

The Hermit Formation flora is significant in that it is distinct from all other North American Permian floras, being a mix of cosmopolitan Permian species, and species with greater affinities to Gondwanan taxa than to North American taxa. Additionally, there are 24 type specimens from the David White collection (see Appendix 7-A).

The Hermit Formation lies directly above the Supai Group, filling erosional channels cut into the top of the Esplanade Sandstone, and forming an extensive deep red slope throughout the canyon. It was at one time included as part of the Supai Group, and was officially named a separate unit in 1922 by L. F. Noble, who removed the upper red shaley beds from the Supai Formation and named them the Hermit Formation (Noble 1922). The Supai Formation was later subdivided into the four current members of the Supai Group by McKee (1982). The Hermit Formation is composed mostly of thinly bedded silts and sandy silts, with thin discontinuous sandstone beds occurring in the basal and top

portions of the formation, and represents deposition in a broad fluvial environment. The Hermit Formation thickens to the west, measured as ~68.6 m (225 ft) thick at the South Kaibab trail and ~83.8 m (275 ft) thick in the Hermit Basin (near the Hermit Trail) (White 1929a).

This fossil flora is described only in a 1929 monograph by David White, but the record of known Permian floras extends to other localities in Arizona, Utah, Colorado, New Mexico, and Texas in the western United States, and also from China, France, Spain, and the Czech Republic (White 1929a; Gand et al. 1997; Wang 1997; DiMichele et al. 2005, 2006, 2007, 2012, 2013; Galtier and Broutin 2008; Opluštil et al. 2017). Of the 32 fully named plant fossil species listed by White (1929a) (with 10 additional species only identified to genus level, or of uncertain identification), about half of them were newly described species at the time he published them (see Appendix 7-A). However, it is likely that there are truly about half that many species in the Hermit Flora because many were poorly preserved and doubtfully described (Read and Mamay 1964). The major groups represented include Spermatopsida (seed plants, including seed ferns), conifers, horsetails, Ginkgoopsida, and algae (White 1929a).

Spermatopsida

The seed ferns (Spermatopsida) are a group of plants that were abundant in the Paleozoic and Mesozoic (Taylor et al. 2009), and they are an informal grouping of seed plants that otherwise cannot be firmly attributed to the seed plant lineages, conifers, cycadophytes, gnetaleans, or ginkgoes (Hilton and Bateman 2006). Their foliage superficially resembles that of ferns, but they produced seeds instead of spores as true ferns do. White listed 18 different species of seed fern (and five additional unidentified species) belonging to six different genera. The genera include *Autunia* or *Rhachiphyllum* (formerly *Callipteris* [Kerp and Haubold 1988]); *Supaia*, *Brongniartites*, *Yakia*, and *Neuropteridium* (Supaiaceae); and *Taeniopteris* (*incertae sedis*; *Taeniopteris* is included under seed ferns in White (1929a) but is truly a vast form genus that includes species of seed ferns and ferns) (Table 7-1).

The genus *Supaia* was established by White (1929a) to describe fern-like foliage dominant in the Hermit Formation flora (Figure 7-2). A key character of this genus is a single bifurcation in the lower frond that splits into two simply pinnate sections. The angle of the split is acute. White interpreted *Supaia* to be closely related to *Autunia* or *Rhachiphyllum* (what White referred to as *Callipteris*) (Kerp and Haubold 1988). DiMichele et al. (2018) commented specifically on *Supaia anomala*, and moved it into the genus *Auritifolia* Chaney, as *Auritifolia anomala*. The authors Kerp and Haubold (1988) commented specifically on the species *Callipteris conferta*, designating it *Autunia conferta*, and on the species *Callipteris raymondii*, designating it to the morphogenus *Sphenocallipteris raymondii*, as they argued that the name *Callipteris* should be abandoned. Kerp et al. (1991) then renamed *Sphenocallipteris* as *Gracilopteris*, meaning the most current name for *Callipteris raymondii* is *Gracilopteris raymondii*. The other species of *Callipteris* mentioned by White (1929) likely belong to the morphogenus *Rhachiphyllum* (Kerp and Haubold 1988), but no formal evaluation is presented here.

Table 7-1. List of all pteridosperm species reported in White (1929a), including locality information. Taxa marked with an asterisk are new species described by White (1929a). (Note: *Callipteris* is no longer a valid name, and these species likely belong to *Autunia* or *Rhachiphyllum* [Kerp and Haubold 1988]).

Fossil Plant Species	Family	Localities
Autunia conferta (*Callipteris conferta*)	*Incertae sedis*	Bright Angel Trail and South Kaibab Trail
*Callipteris arizonae**	*Incertae sedis*	Hermit Basin
Gracilopteris raymondii (*Callipteris raymondii*)	*Incertae sedis*	Bright Angel Trail and Hermit Trail
Callipteris ? sp.	*Incertae sedis*	Hermit Basin
*Supaia thinnfeldioides**	Supaiaceae	Hermit Basin
*Supaia rigida**	Supaiaceae	Hermit Basin
*Supai sturdevantii**	Supaiaceae	Bright Angel Trail
*Supaia merriami**	Supaiaceae	Hermit Trail
Supai compacta	Supaiaceae	Hermit Trail
Auritifolia anomala (*Supaia anomala**)	Supaiaceae	Hermit Basin and Bright Angel Trail
*Supaia linearifolia**	Supaiaceae	Bright Angel Trail
*Supaia breviloba**	Supaiaceae	Bright Angel Trail
*Supaia subgoepperti**	Supaiaceae	South Kaibab Trail
Supaia sp.	Supaiaceae	Hermit Basin
Supaia sp. indet	Supaiaceae	Hermit Basin
Supaia ? sp.	Supaiaceae	South Kaibab Trail
Brogniartites ? *yakiensis**	Supaiaceae	South Kaibab Trail and Hermit Trail
Brogniartites ? *aliena**	Supaiaceae	South Kaibab Trail
*Yakia heterophylla**	Supaiaceae	Hermit Basin and Bright Angel Trail
Neuropteridium sp.	Supaiaceae	Bright Angel Trail
Taeniopteris cf. *eckhardti*	*Incertae sedis*	Bright Angel Trail and Hermit Trail
*Taeniopteris angelica**	*Incertae sedis*	Bright Angel Trail
Taeniopteris coriacea	*Incertae sedis*	Bright Angel Trail, Hermit Basin

White reported several seeds potentially attached to different specimens of *Supaia* in both his 1929 monograph and in another article published in 1934 (which did not include any figures or references to specimens). Evidence for physical attachment of these seeds to foliage was unclear at best, was reviewed by Mamay and Watt (1971), and is currently interpreted to be some type of insect damage (William DiMichele and colleagues, pers. comm.).

Yakia was another genus newly established by White (1929a) and is defined by “closely spaced, relatively even and parallel ultimate divisions, and by the rather distant, narrow, crooked leaves, which have the appearance of being irregularly lobed…” (Figure 7-3). White also tentatively assigned a fructification that appears as five oblong bodies packed together in what White called “clusters of sporangia or (possibly seeds)”, located at the bases of the pinnules (White 1929a: Plate 40, Figures 1 and 2). The distinction between *Supaia* and the other genera included by White in Supaiaceae (*Brogniartites* and *Yakia*) is the size and overall shape (proportion) of the foliage—the

bifurcated fronds, pinnules, and venation are all otherwise very similar (it may be that they represent different leaf sizes produced by the same parent taxon).




Figure 7-2. *Supaia* specimens (CASSI KNIGHT). **A.** *Supaia* sp. specimen USNM 38040; **B.** *Supaia subgoepperti* specimen USNM 38038.




Figure 7-3. *Yakia heterophylla* (CASSI KNIGHT). **A.** USNM 312802 **B.** USNM 312805. The "clusters of sporangia or possibly seeds" mentioned by White (1929a) can be seen in the upper left corner of the specimen.

Coniferophyta

The conifers found in the Hermit Formation number ten different species (and two undetermined species) belonging to seven different families (Table 7-2). These include *Walchia, Ullmannia,* and *Voltzia* (Araucariaceae); *Paleotaxites* and *Taxites* (Paleotaxaceae); and *Brachyphyllum* and *Pagiophyllum* (Pinales?).

Table 7-2. List of all conifer species reported in White (1929a), including locality information. Taxa marked with an asterisk are new species described by White (1929a).

Fossil Plant Species	Family	Locality
Brachyphyllum arizonicum	?	Hermit Basin
Brachyphyllum tenue	?	Hermit Basin and Bright Angel Trail
Feysia sp. ? (*Ullmannia frumentaria*)	?	Hermit Basin and Bright Angel Trail
Hermitia dawsoni (*Walchia dawsoni**)	?	Hermit Basin
Hermitia sp. ? (*Walchia gracillima**)	?	South Kaibab Trail
Otovicia hypnoides ? (*Walchia hypnoides*)	?	South Kaibab Trail
Pagiophyllum dubium	?	Hermit Basin
Paleotaxites precursor	Paleotaxaceae	Hermit Basin and South Kaibab Trail
Taxites ? sp.	Paleotaxaceae	Hermit Basin
Voltzia dentiloba	?	Hermit Basin
Voltzia sp.	?	No locality given
Walchia piniformis	?	Hermit Basin, South Kaibab Trail, and Bright Angel Trails

Walchia is a morphogenus that describes leafy coniferous foliage, ultimate shoots and penultimate branch systems, that are all visually similar and typically too poorly preserved to allow certain identification (and with no preserved epidermal structure) (Visscher et al. 1986). There are several species of *Walchia* reported in White (1929a), and the author recognized the fact that there are multiple different species of conifer included under each name, noting that sometimes different cones are attached to branches of the same morphological form. *Walchia* is restricted to Euramerica during the Permian, and the species reported in the 1929 monograph were described based on their similarities to species of *Walchia* described from Europe by European authors. Visscher et al. (1986) established a new morphogenus (*Hermitia* Kerp and Clement-Westerhof) "to accommodate coniferous foliage without preserved epidermal structure; uncertain affinity at the family level" and commented on its relevance to some of the *Walchia* specimens described by White (1929a). Visscher et al. (1986) considered *Walchia piniformis* Sternberg sufficiently well known for the name to stand, placed *Walchia dawsoni* in *Hermitia* as *H. dawsoni* (White 1929a) Kerp and Clement-Westerhof. Kerp et al. (1990) transferred *Walchia hypnoides* to a new genus *Otovicia hypnoides*. *Walchia gracillima* fits the description of *Hermitia* (leafy shoots with needle-like leaves that lack preserved epidermal structures) and likely belongs to this morphogenus (Visscher et al. 1986; Broutin and Kerp 1994), but without preserved epidermal features or reproductive structures, no formal analysis will be presented here.

White reported some specimens of *Ullmannia frumentaria* with small, oval seeds or cones associated with foliage, and compared *U. frumentaria* to *Buriadica*, a genus found in India. His comparison is based on leaves that fork near their apex, but the description for *Ullmannia* does not include bifurcated leaves; this is common for penultimate leaves and cone bracts of Walchian conifers

(Schweitzer 1986). Furthermore, Broutin and Kerp (1994) designated a new morphogenus *Feysia* (which accommodates broad-leaved conifers), and strongly suggested that many specimens of *Ullmannia* likely could be referred to a species of *Feysia* (Broutin and Kerp 1994).

There are several conifer taxa that are only known from very few specimens or are particularly rare in the Hermit Formation flora. *Voltzia dentiloba* is known from the Hermit only by two moderately well-preserved cone bracts, and *Voltzia* sp. is known only from a single cone bract. *Paleotaxites* is a new morphogenus established by White (1929) of conifer with short, compact needles that are densely packed and spirally arranged. Some branches have cones closely associated, which are leathery in texture and have small scales at their base. The single species of *Paleotaxites* reported from the Hermit Formation, *P. precursor*, is rather rare.

White (1929a) also created a new species, *Pagiophyllum dubium*, to include fragmentary twigs and leaves that are not detailed enough to allow certain description, but all distinctly belong to a single species. These plants are loosely assigned to the genus *Pagiophyllum*, but more evidence is needed to confirm this assignment (White 1929a).

Equisetopsida

A single species of horsetail, *Sphenophyllum gilmorei* (Figure 7-4A), is reported from the Hermit Formation paleoflora, and is yet another new species established by White (1929a). The genus *Sphenophyllum* was originally assigned to Arthrophyta by White, but this division is an old name that is no longer used, and the genus *Sphenophyllum* is now attributed to Equisetopsida, which is the class commonly known as horsetails. *S. gilmorei* is similar to only one other species from this genus, *S. stoukenbergi* from the Ural Mountains, but has much larger, more elongate leaves.



Figure 7-4. Various taxa (CASSI KNIGHT). **A.** *Sphenophyllum gilmorei* specimen USNM 38025. **B.** *Psygmophyllum* sp. specimen USNM 38050. **C.** *Rivularites permiensis* specimen USNM 38024.

Sphenophyllum gilmorei is most commonly found in thin sandstone beds deposited within the channel cuts in the Esplanade Sandstone. It reportedly becomes much rarer outside the Hermit Basin, and a few specimens have been found near the Bright Angel and South Kaibab Trails. It can be found preserved in association with small vertebrate tracks on “slime mud” surfaces. Some specimens are preserved perpendicular to bedding, suggesting that *S. gilmorei* grew in shallow pools of water and were buried in situ. White interpreted the ecology of the plant to be one that grew in dense stands in sandy bottomed, intermittent pools, reaching a height of up to 1 m (3 ft) (White 1929a).

Ginkgoopsida

White reported one species of ginkgo in his 1929 monograph, and a second species of ginkgo is known from specimens in the Grand Canyon Museum collections. These plants are *Psygmophyllum* ? sp. and *Rhipidopsis* sp.?, respectively. *Psygmophyllum* ? sp. is known only from a single, poorly preserved leaf fragment found in the lower part of the Hermit Formation in the Hermit Basin (Figure 7-4B). The leaf fragment appears to be covered in a veil of silt and "slime", obscuring many of the detailed features. White also compared this specimen to the ginkgo *Rhipidopsis ginkgoides*. Both ginkgoes are known from the Asian portion of Gondwana (White 1929a).

Algae

One species of algae, *Rivularites permiensis* (Figure 7-4C), is found in the Hermit Formation. White described this as a new species in his 1929 monograph, and attributed *R. permiensis* to Thallophyta, which as a now abandoned term for algae. *R. permiensis* was identified based on its similarity to fossil *Rivularites* from France and living *Rivularites* from the French coast, but described as a new species because of some notable round protuberances that occurred in the Arizona specimens, but not those from Europe. There is, however, no firm evidence to suggest a direct relation between the fossil and modern blue-green algae.

Rivularites is common in the lower portion of the Hermit Formation, but has only been found in the channels that are cut into the underlying Esplanade Sandstone in the Hermit Basin. Colonies of the algae are found often on large ripple surfaces and may reach 0.5 m (nearly 2 ft) in diameter, but most tend to be smaller in size. White proposed that the algae grew in mats in moving water, based on the presence of associated ripple marks and the observation that all the protuberances were consistently pointed or pulled in a single direction (White 1929a).

Other Plants

A category for "fruits of uncertain affinities" includes three genera of different seed types, *Cyclocarpon*, *Carpolithus*, and *Eltovaria*, and gymnospermous ament. (Table 7-3). The different seed types are reportedly very rare in the Hermit Formation, and not many specimens have been found. The seeds *Cyclocarpon angelicum*, *C.* sp., and *Carpolithus* sp. both are small, round forms that are associated with pteridosperms, meaning they came from plants belonging to *Autunia*, *Rhachiphyllum*, or *Supaia*. There is only one known specimen of *Carpolithus* sp., and it is not particularly well-preserved.

Table 7-3. Fruits of uncertain affinities reported by David White (1929a) with locality information. Taxa marked with an asterisk are new species described by White (1929a).

Fossil Species	**Locality**
*Cyclocarpon angelicum**	Bright Angel Trail
Cyclocarpon sp.	No locality given
Carpolithus sp.	Hermit Basin
Eltovaria bursiformis	Bright Angel Trail and Hermit Basin

What is likely some type of fruitification formed by a modified leaf that is apparently folded longitudinally to create a type of pod is identified as *Eltovaria bursiformis*, *Eltovaria* being a new genus established by White for this form (White 1929a). The modified leaf most closely resembles *Taeniopteris* or *Supaia*, and White interpreted this structure as belonging to *Taeniopteris*. A small, single fragment of poorly preserved material is tentatively described as an ament (catkin-like structure) from a gymnosperm and, in this case, is likely some type of female pollen cone.

Acknowledgments

A sincere thank you to all who contributed to this report. I appreciate Klara Widrig and Diana Boudreau for spending their time to help track down needed references. I also greatly appreciate Klara, Diana, Anne Miller, Justin Tweet, and Vincent Santucci for their helpful comments and editorial eyes. Thank you to William DiMichele and Cindy Looy for reviewing this report and for their absolutely invaluable feedback. Finally, thank you to the Grand Canyon Association and Grand Canyon National Park for their support of this work, and many kind thanks to Vincent Santucci for giving me the opportunity to be involved in this project. You all made this possible. "Still the Grand Canyon is the sublimest thing on earth. It is so not alone by virtue of its magnitudes, but by virtue of the whole—its *tout ensemble*."—Clarence Dutton

Literature Cited

Beus, S. S. 1986. A geologic surprise in the Grand Canyon. Arizona Bureau of Geology and Mineral Technology Fieldnotes 16(3):1–4.

Beus, S. S. 1990. Redwall Limestone and Surprise Canyon Formation. Pages 115–135 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology. Oxford University Press, New York, New York.

Beus, S. S. 1995. Paleontology of the Surprise Canyon Formation (Mississippian) in Grand Canyon, Arizona. Proceedings – Fossils of Arizona Symposium 3:25–36.

Billingsley, G. H. 1979. Preliminary report of buried valleys at the Mississippian-Pennsylvanian boundary in western Grand Canyon. Pages 115–117 *in* S. S. Beus and R. R. Rawson, editors. Carboniferous stratigraphy in the Grand Canyon country, northern Arizona and southern Nevada. 9th International Congress of Carboniferous Stratigraphy and Geology, Field Trip 13. American Geological Institute Selected Guidebook Series 2.

Billingsley, G. H., and S. S. Beus. 1985. The Surprise Canyon Formation: An Upper Mississippian and Lower Pennsylvanian(?) rock unit in the Grand Canyon, Arizona. Pages A27–A33 *in* Stratigraphic notes, 1984. U.S. Geological Survey, Washington, D.C. Bulletin 1605-A.

Billingsley, H., and E. D. McKee. 1982. Pre-Supai buried valleys. Pages 137–153 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

Broutin, J., and H. Kerp. 1994. Aspects of Permian palaeobotany and palynology. XIV. A new form-genus of broad-leaved Late Carboniferous and Early Permian Northern Hemisphere conifers. Review of Palaeobotany and Palynology 83:241–251.

Cecil, C. B. 2003. Introduction: Allocyclic controls on Middle Pennsylvanian sedimentation in cratonic depositional systems (U.S.A.). Pages 69–72 *in* C. B. Cecil and N. T. Edgar. Climate controls in stratigraphy. Society of Economic Paleontologists and Mineralogists, Tulsa, Oklahoma. Special Publication 77.

DiMichele, W., and R. B. Aronson. 1992. The Pennsylvanian-Permian vegetational transition: A terrestrial analogue to the onshore-offshore hypothesis. Evolution 46(3):807–824.

DiMichele, W. A., H. Kerp, M. Krings, and D. S. Chaney. 2005. The Permian peltasperm radiation: Evidence from the southwestern United States. New Mexico Museum of Natural History and Science Bulletin 30:67–79.

DiMichele, W. A., N. J. Tabor, D. S. Chaney, and W. J. Nelson. 2006. Wetlands to wet spots: Environmental tracking and the fate of Carboniferous elements in Early Permian tropical floras. Pages 223–248 *in* S. F. Greb and W. A. DiMichele, editors. Wetlands through time. Geological Society of America, Boulder, Colorado. Special Paper 399.

DiMichele W. A., D. S. Chaney, W. J. Nelson, S. G. Lucas, C. V. Looy, K. Quick, and J. Wang. 2007. A low diversity, seasonal tropical landscape dominated by conifers and peltasperms: Early Permian Abo Formation, New Mexico. Review of Palaeobotany and Palynology 145:249–273.

DiMichele, W. A., S. G. Lucas, and K. Krainer. 2012. Vertebrate trackways among a stand of *Supaia* White plants on an early Permian floodplain, New Mexico. Journal of Paleontology 86:584–594.

DiMichele, W. A., D. S. Chaney, S. G. Lucas, H. Kerp, and S. Voigt. 2013. Flora of the Lower Permian Abo Formation redbeds, western equatorial Pangea, New Mexico. New Mexico Museum of Natural History and Science Bulletin 59:265–287.

DiMichele, W. A., S. G. Lucas, D. S. Chaney, M. P. Donovan, H. Kerp, R. A. Koll, and C. V Looy. 2018. Early Permian flora, Doña Ana Mountains, southern New Mexico, with special consideration of taxonomic issues and arthropod damage. New Mexico Museum of Natural History and Science Bulletin 79:165–205.

Fielding, C. R., and T. D. Frank, and J. L. Isbell. 2008. The late Paleozoic Ice Age—A review of current understanding and synthesis of global climate patterns. Pages 343–354 *in* C. R. Fielding, T. D. Frank, and J. L. Isbell, editors. Resolving the Late Paleozoic Ice Age in time and space. Geological Society of America, Boulder, Colorado. Special Paper 441.

Galtier, J., and J. Broutin. 2008. Floras from red beds of the Permian Basin of Lodève (Southern France). Iberian Journal of Geology 34:57–72.

Gand, G., H. Kerp, C. Passons, and E. Martinez-Garcia. 1997. Palaeoenvironmental and stratigraphic aspects of animal traces and plant remains in Spanish Permian red beds (Pena Sagra, Cantabrian Mountains, Spain). Geobios 30:295–318.

Gastaldo, R. A., and T. M. Demko. 2011. The relationship between continental landscape evolution and the plant-fossil record: Long term hydrologic controls on preservation. Pages 249–285 *in* P. A. Allison and D. J. Bottjer, editors. Taphonomy: process and bias through time. Springer, Dordrecht, Netherlands. Topics in Geobiology 32.

Hilton, J., and R. M. Bateman. 2006. Pteridosperms are the backbone of seed-plant phylogeny. The Journal of the Torrey Botanical Society 133(1):119–168.

Hodnett, J.-P. M., and D. K. Elliott. 2018. Carboniferous chondrichthyan assemblages from the Surprise Canyon and Watahomigi formations (latest Mississippian–Early Pennsylvanian) of the western Grand Canyon, northern Arizona. Journal of Paleontology 92, Memoir 77.

Kerp, H., and H. Haubold. 1988. Toward a reclassification of the West- and Central-European species of the form-genus *Callipteris* Brongniart. Zeitschrift für Geologische Wissenschaften, Berlin 16:865–876.

Kerp, J. H. F., S. V. Naugolnykh, and H. Haubold. 1991. *Gracilopteris*, a new name for *Sphenocallipteris* Haubold and Kerp. Review of Palaeobotany and Palynology 69(4):369–370.

Kerp, J. H. F., R. J. Poort, H. A. J. M. Swinkels, and R. Verwer. 2001. Aspects of Permian palaeobotany and palynology. IX. Conifer-dominated Rotliegend floras from the Saar-Nahe basin (? Late Carboniferous-early Permian; SW-Germany) with special reference to the reproductive biology of early conifers. Review of Palaeobotany and Palynology 62(3–4):205–248.

Mamay, S. H., and A. D. Watt. 1971. An ovuliferous callipteroid plant from the Hermit Shale (Lower Permian) of the Grand Canyon, Arizona. Pages C48–C51 *in* Geological Survey Research 1971. U.S. Geological Survey, Washington, D.C. Professional Paper 750–C.

McKee, E. D. 1982. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

Montañez, I. P., and C. J. Poulsen. 2013. The Late Paleozoic Ice Age: An evolving paradigm. Annual Review of Earth and Planetary Sciences 41:629–656.

Noble, L. F. 1922. A section of the Paleozoic formations of the Grand Canyon at Bass Trail. Pages 23–73 *in* Shorter contributions to general geology, 1922. U.S. Geological Survey, Washington, D.C. Professional Paper 131-B.

Opluštil, S., J. Jirasék, M. Schmitz, and K. Matýsek. 2017. Biotic changes around the radioisotopically constrained Carboniferous-Permian boundary in the Boskovice Basin (Czech Republic). Bulletin of Geosciences 92:95–122.

Read, C. B., and S. H. Mamay. 1964. Upper Paleozoic floral zones and floral provinces of the United States. U.S. Geological Survey, Washington, D.C. Professional Paper 545-K.

Schuchert, C. 1918. The Cambrian of the Grand Canyon of Arizona. American Journal of Science (4th series) 45:362–369.

Schweitzer, H.-J. 1986. The land flora of the English and German Zechstein sequences. Pages 31–54 *in* G. M. Harwood and D. B. Smith, editors. The English Zechstein and related topics. Geological Society of London, London, United Kingdom. Special Publication 22:31–54.

Spamer, E. E. 1992. The Grand Canyon fossil record: A source book in paleontology of the Grand Canyon and vicinity, northwestern Arizona and southeastern Nevada. Geological Society of America, Boulder, Colorado. Microform Publication 24.

Taylor, T. N., E. L. Taylor, and M. Krings. 2009. Paleobotany: The biology and evolution of fossil plants (2nd edition). Academic Press, Boston, Massachusetts.

Tidwell, W. D., J. R. Jennings, and S. S. Beus. 1992. A Carboniferous flora from the Surprise Canyon Formation in the Grand Canyon, Arizona. Journal of Paleontology 66(6):1013–1021.

Visscher, H., J. H. F. Kerp, and J. A. Clement-Westerhof. 1986. Aspects of Permian palaeobotany and palynology. VI. Towards a flexible system of naming Paleozoic conifers. Acta Botanica Neerlandica 35:87–99.

Wang, Z. Q. 1997. Permian *Supaia* fronds and an associated *Autunia* fructification from Shanxi, China. Palaeontology 40:245–277.

White, C. D. 1927. Study of the fossil floras in the Grand Canyon, Arizona. Carnegie Institution of Washington Year Book 26:366–369.

White, C. D. 1928. Algal deposits of Unkar Proterozoic age in the Grand Canyon, Arizona. Proceedings of the National Academy of Sciences of the United States of America 14(7):597–600.

White, C. D. 1929a. Flora of the Hermit shale, Grand Canyon, Arizona. Carnegie Institution of Washington Publication 405.

White, C. D. 1929b. Study of the fossil floras in the Grand Canyon, Arizona. Carnegie Institution of Washington Year Book 28:392–393.

White, C. D. 1934. The seeds of *Supaia*, a Permian pteridosperm. Science 79(2055):462.

Appendix 7-A. Paleobotanical Type Specimens

Appendix Table 7-A-1. Table of all known paleobotanical type specimens from GRCA, including name and specimen numbers. All of the newly created plant species were described by David White in his 1929 monograph (White 1929a),and were originally discovered in the Hermit Formation flora.

Species	Age	Lithology	Reference	Plant Group	Syntype Specimens
Brachyphyllum arizonicum	Permian	Hermit	White 1929a	Conifer	USNM 38060
Brachyphyllum tenue	Permian	Hermit	White 1929a	Conifer	USNM 38061, 312783, 312784
Brongniartites? *yakiensis*	Permian	Hermit	White 1929a	Pteridophyte	USNM 38042, 312807–312812, 324554–324559
Brongniartites? *aliena*	Permian	Hermit	White 1929a	Pteridophyte	USNM 38043, 312785–312789
Rhachiphyllum sp. (*Callipteris arizonae*)	Permian	Hermit	White 1929a	Pteridophyte?	USNM 38027
Cyclocarpon angelicum	Permian	Hermit	White 1929a	Seed/Fruit	USNM 38063
Eltovaria bursiformis	Permian	Hermit	White 1929a	Seed/Fruit	USNM 38066
Pagiophyllum dubium	Permian	Hermit	White 1929a	Conifer	USNM 38062, 312790–312792
Palaeotaxites praecursor	Permian	Hermit	White 1929a	Conifer	USNM 38058, 38060, 324560–324564
Sphenophyllum gilmorei	Permian	Hermit	White 1929a	Pteridophyte	USNM 38025, 324567–324575
Supaia anomala	Permian	Hermit	White 1929a	Pteridophyte	USNM 38035, 324581–324588
Supaia breviloba	Permian	Hermit	White 1929a	Pteridophyte	USNM 38037
Supaia compacta	Permian	Hermit	White 1929a	Pteridophyte	USNM 38034, 324590–324593
Supaia linearifolia	Permian	Hermit	White 1929a	Pteridophyte	USNM 38036 and 324594
Supaia merriami	Permian	Hermit	White 1929a	Pteridophyte	USNM 38033
Supaia rigida	Permian	Hermit	White 1929a	Pteridophyte	USNM 38031, 324595, 324596
Supaia sturdevantii	Permian	Hermit	White 1929a	Pteridophyte	USNM 38032, 38038, 324597
Supaia subgoepperti	Permian	Hermit	White 1929a	Pteridophyte	USNM 38038
Supaia thinnfeldioides	Permian	Hermit	White 1929a	Pteridophyte	USNM 38030, 324598–324603
Taeniopteris angelica	Permian	Hermit	White 1929a	Pteridophyte or Gingko	USNM 38048, 324565, 324566, 342587
Voltzia dentiloba	Permian	Hermit	White 1929a	Conifer	USNM 38056
Hermitia dawsoni (*Walchia dawsonil*)	Permian	Hermit	White 1929a	Conifer	USNM 38052, 312796–312798

Appendix Table 7-A-1 (continued). Table of all known paleobotanical type specimens from GRCA, including name and specimen numbers. All of the newly created plant species were described by David White in his 1929 monograph (White 1929a),and were originally discovered in the Hermit Formation flora.

Species	Age	Lithology	Reference	Plant Group	Syntype Specimens
Hermitia sp. (*Walchia gracillima*)	Permian	Hermit	White 1929a	Conifer	USNM 38053, 312793–312795
Yakia heterophylla	Permian	Hermit	White 1929a	Pteridophyte	USNM 38044, 312799–312804
Rivularites permiensis	Permian	Hermit	White 1929a	Cyanobacteria colonies	USNM 38024

Chapter 8. Paleozoic Invertebrate Ichnology of Grand Canyon National Park

By Anne E. Miller[1], Lorenzo Marchetti[2], Heitor Francischini[3], and Spencer G. Lucas[4]

[1]Grand Canyon National Park
1824 S Thompson St, Suite 200
Flagstaff, Arizona 86001

[2]Urweltmuseum GEOSKOP / Burg Lichtenberg (Pfalz)
Burgstraße 19
D-66871 Thallichtenberg, Germany

[3]Universidade Federal do Rio Grande do Sul (UFRGS)
Laboratório de Paleontologia de Vertebrados and Programa de Pós-Graduação em Geociências, Instituto de Geociências
Porto Alegre, Rio Grande do Sul, Brazil

[4]New Mexico Museum of Natural History and Science
1801 Mountain Road N.W.
Albuquerque, New Mexico, 87104

Introduction

Grand Canyon National Park (GRCA) encompasses an extensive stratigraphic record that typifies the evolution of diverse ecosystems and their inhabitants throughout the Paleozoic. The occurrence of invertebrate traces (ichnofossils) is not uncommon in these strata and represents direct evidence of interactions between the organisms and the substrates on which and/or in which they lived. During the Paleozoic, a diverse array of invertebrate groups were capable of producing trace fossils on or within the substrates where they lived, including arthropods (such as crustaceans, insects, arachnids, trilobites, and xiphosurans), mollusks (especially bivalves and mobile benthonic groups), brachiopods, cnidarians, echinoderms, various worm taxa (such as annelids and priapulids), and other rare taxa.

Some ichnogenera are specific to ichnofacies that reflect the bathymetry and/or the depositional features of marine or terrestrial depositional environments (Seilacher 1964; Hunt and Lucas 1998, 2007; MacEachern et al. 2010). Certain ichnofacies encompass many different types of ichnogenera, depending on the depositional context of the formation. These groupings of ichnogenera model the organisms' interaction with the sediment based on recurring behaviors and provide further interpretation of their depositional environments (MacEachern et al. 2010). They are reflected in most of the Grand Canyon's Paleozoic strata and provide valuable information on the paleoecology of invertebrate organisms. Thus, protection of such trace fossils is crucial to advancing the field of ichnology and to enabling interpretation of the ancient environments represented by the strata exposed in the Grand Canyon.

Trace Fossil Classifications

Trace fossils differ from body fossils in that they are classified by an organism's behavior, rather than the organism's anatomy (morphology). They are, quite literally, the record of interaction between an organism and its substrate. Thus, they are classified primarily by their trace morphology that reflects specific behavior, which can be related to the anatomy of the producer, especially for trackways (e.g., Bertling et al. 2006; Minter et al. 2007; Marchetti et al. 2019a).

The morphology of a trace fossil is not only a reflection of the behavior of an organism and the nature of preservation, but it also variably reflects the anatomy of the producer (e.g., Minter et al. 2007). In particular, trackways record the foot structure and some other aspects of producer anatomy, whereas burrows do so less, in part because different organisms with very different anatomy can more readily produce the same burrow structure than they can produce the same kinds of trackways. This is why vertebrate ichnology, which mostly focuses on trackways, often treats the morphology consistent with the anatomy of the producer, but not the taxonomy of the producer, as a key element in ichnotaxonomy (e.g., Marchetti et al. 2019a). In invertebrate taxonomy, what can be called the problem of homeomorphy—same trace fossil structure, different producers—is more prevalent, so the emphasis in ichnotaxonomy is on morphological features indicative of behavior, rather than of producer anatomy.

Invertebrate trace fossils are classified independent of the identity of their producer (and its taxonomy) using a binomial nomenclature (i.e., the ichnogeneric plus the ichnospecific epithets) analogous, but parallel to, the taxonomic classification of zootaxa (ICZN 1999). This is why the classification of trace fossils, or ichnotaxonomy, can be considered a parataxonomy. Ichnofamilies may be used to group ichnogenera into larger ichnotaxonomic groups, but they are generally less extensively studied and discussed than in the phylogenetic taxonomy of body fossils (Pemberton and Frey 1982). This is because ichnotaxonomy is separated from the taxonomic classification of zootaxa and has an ethological component (Pemberton and Frey 1982), therefore it is more difficult to group morphologies and assign categories that are a complex result of different behaviors and different anatomies.

The list of valid ichnotaxa in a defined unit and/or locality is named an ichnoassociation, and the composition of an ichnoassociation can be used to define ichnodiversity.

A further informal approach to trace fossil classification evaluates ichnodisparity (e.g., Buatois et al. 2017), which is the evaluation of the structural complexity of the ichnoassociations based on the number of architectural designs that are present. For such analysis, 79 architectural designs have been introduced that encompass 523 invertebrate-produced ichnogenera (Buatois et al. 2017).

Seilacher (1964) created a broad ethological classification consisting of five major behavioral (ethological) categories for invertebrate traces that has since been expanded into 13 categories, plus some subordinate subcategories (MacEachern et al. 2010; Vallon et al. 2016). Seilacher's (1964) original five comprise: Domichnia (dwelling), Fodinichnia (feeding), Pascichnia (grazing), Repichnia (locomotion), and Cubichnia (resting). The subsequently proposed additional groups comprise: Agrichnia (farming), Calichnia (brooding), Chemichnia (chemosymbiotic traces), Digestichnia

(digestion), Ecdysichnia (molting), Fixichnia (attachment), Fugichnia (escaping), and Praedichnia (predation) (MacEachern et al. 2010; Vallon et al. 2016). These behaviors seen in the trace fossil record reveal the way in which organisms respond to environmental changes such as climate, food resources, lithology, soil moisture, turbidity, water table fluctuations, and the energy of deposition (MacEachern et al. 2010). The dynamics of the depositional environment can be inferred based on the range of behavior present. However, the use of these ethological terms is rare among ichnologists, and behavior is more often described in simple descriptive terms (e.g., "feeding trace", "dwelling trace", etc.).

One can also informally classify trace fossils on a toponomic basis (relationship to the sediment) in order to understand their orientation and preservation within the sediment. Seilacher (1964) differentiated full-relief trace fossils, occurring within the sediment, from semi-relief traces, which are preserved along bedding surfaces. He grouped the semi-relief traces as epirelief (top surface of a bed) and hyporelief (bottom surface of a bed), in which they each can be concave (mold) or convex (cast). Martinsson (1970) went further and created four general groups of toponomic classification: endichnia, epichnia, exichnia, and hypichnia. Endichnia refers to traces within the medium, similar to Seilacher's (1964) full-relief traces. Epichnia refers to ridges or grooves on the top surface of a bed. The opposite is hypichnia, referring to ridges or grooves on the bottom surface of a bed. Exichnia refers to traces that have been filled in with sediment different from the surrounding medium.

Trace fossils reflect behavioral adaptations to specific environmental conditions. Some ichnotaxa are restricted to environmental settings or even to particular lithofacies. Therefore, ichnoassociations will be recurrent over geologic time whenever the respective environmental conditions recur. These ideas were used by Seilacher (1964) as the foundation for the concept of ichnofacies. Ichnofacies are distinct and recurrent (in space and time) associations of fossil traces (ichnoassociations) that reflect specific combinations of the organisms' responses to particular environmental conditions (MacEachern et al. 2007). In his seminal work, Seilacher (1964) recognized six archetypical ichnofacies, based on associations of invertebrate traces: *Skolithos* Ichnofacies, *Cruziana* Ichnofacies, *Zoophycos* Ichnofacies, *Nereites* Ichnofacies, *Glossifungites* Ichnofacies, and *Scoyenia* Ichnofacies. Among them, only the last one is present in continental settings. This framework was consequently expanded so that nine other invertebrate ichnofacies were proposed: *Celliforma* Ichnofacies, *Coprinisphaera* Ichnofacies, *Cubiculum* Ichnofacies, *Octopodichnus–Entradichnus* Ichnofacies, *Palaeoscolytus* Ichnofacies, *Psilonichnus* Ichnofacies, *Teredolites* Ichnofacies, *Termitchnus* Ichnofacies, and *Trypanites* Ichnofacies (MacEachern et al. 2007; Hunt and Lucas 2007; Lucas 2016). These new ichnofacies recognize greater ichnofaciological diversity in continental settings than was known to Seilacher (1964), and thus represent a more diverse array of environmental settings (paleosols, eolian dunes, supratidal deposits, etc.). The following will discuss the ichnological record of some units from the Grand Canyon in the light of the ichnofacies paradigm, but more complete and updated information on ichnofacies in general can be found in MacEachern et al. (2012).

Ichnology is essential in providing missing paleoecological information for those organisms whose fossilized bodies are frequently studied and even those that are never discovered, but it poses a

challenge in associating trace fossils with their producers. For example, trace fossils are often found in strata that lack body fossils, and thus provide the only available biogenic information for paleoenvironmental reconstructions. This phenomenon is very common in continental desert deposits, such as the Permian Coconino Sandstone, where invertebrate traces and tetrapod tracks comprise the only evidence of the occupation of that paleoenvironment by organisms (see discussion below). Fortunately, there is a relatively long history of Grand Canyon trace fossil research, so that this linkage has been achieved with some success (e.g., Elliott and Martin 1987; Lane et al. 2003). Currently, a project by one of the authors (AEM, pers. obs.) is addressing these challenges by studying the ichnology of the Bright Angel Shale, which holds the most diverse and well-preserved invertebrate trace fossils from Grand Canyon's Paleozoic strata. This study is intended to not only link potential producers to their traces, but also determine ichnofacies and ichnoassociations to better understand Cambrian paleoecology. The work should advance the field of invertebrate ichnology of the Cambrian, but it will also pose more questions and suggest important future work. Therefore, it is imperative that the significant trace fossil specimens in GRCA are preserved and protected because they provide opportunities for advancing our understanding of the history of life recorded in the Park.

Stratigraphic Distribution of Trace Fossils

Trace fossils have been reported from almost every Paleozoic geological unit of the Grand Canyon. Some of these reports lack comprehensive ichnotaxonomic lists, and this may be attributed to the poorly studied trace fossil record of some of the Paleozoic formations. The most extensively reported trace fossil record from Grand Canyon is from the Cambrian strata of the Tonto Group. Out of the three formations that comprise the Tonto Group, the Bright Angel Shale has yielded the majority of these reports. This is due to the sheer number and diversity of trace fossils preserved in the Bright Angel Shale relative to other units. Although the Coconino Sandstone is famous for its vertebrate trackways, it contains the second most reported invertebrate traces next to the Tonto Group. The Hermit Formation has also been the source of a few reports on invertebrate trace fossils, some of which have been considered to be of plant origin. These units represent completely different time periods and depositional environments and therefore contribute to a diverse invertebrate ichnological record for Grand Canyon. The Toroweap Formation has little to no published information on invertebrate trace fossils, so it is not included in this chapter, though trace fossils likely are present in that stratigraphic unit.

The following text is organized by geologic age and formation, including a list of collected museum specimens from GRCA (Table 8-3 for the Bright Angel Shale subchapter). The specimens only documented in the field and not collected were not numbered. For each formation, the ichnotaxa are listed by different architectural designs (according to Buatois et al. 2017), citing the relevant literature in chronologic order. For the most relevant ichnoassociations, a paragraph discussing ichnodiversity, ichnodisparity and ichnofacies of that specific unit was added. A list of valid ichnotaxa from GRCA is provided in Table 8-1. A list of invalid ichnotaxa from unpublished theses is also provided (Table 8-2).

Table 8-1. List of valid ichnogenera and their respective architectural designs (sensu Buatois et al. 2017), ordered by geological unit (from lower to upper).

Unit	Valid ichnogenera	Architectural designs
Tapeats Sandstone	*Cruziana*	5: Bilobate trails and paired grooves
	Diplichnites	6: Trackways and scratch imprints
	Monomorphichnus	6: Trackways and scratch imprints
	Rusophycus	8: Bilaterally symmetrical short, scratched impressions and burrows
	Planolites	11: Simple actively filled (massive) horizontal to oblique structures
	Palaeophycus	10: Passively filled horizontal burrows
	Treptichnus	17: Horizontal burrows with horizontal to vertical branches
	Teichichnus	22: Horizontal burrows with simple vertically oriented spreiten
	Rhizocorallium	26: Burrows with horizontal spreiten
	Skolithos	35: Vertical unbranched burrows
	Arenicolites	36: Vertical single U- and Y-shaped burrows
	Diplocraterion	36: Vertical single U- and Y-shaped burrows
Bright Angel Shale	*Cochlichnus*	1: Simple horizontal trails
	Cruziana	5: Bilobate trails and paired grooves
	Angulichnus	6: Trackways and scratch imprints
	Diplichnites	6: Trackways and scratch imprints
	Dimorphichnus	6: Trackways and scratch imprints
	Monomorphichnus	6: Trackways and scratch imprints
	Rusophycus	8: Bilaterally symmetrical short, scratched impressions and burrows
	Palaeophycus	10: Passively filled horizontal burrows
	Planolites	11: Simple actively filled (massive) horizontal to oblique structures
	Nereites	14: Complex actively filled horizontal structures
	Scolicia	14: Complex actively filled horizontal structures
	Phycodes	17: Horizontal burrows with horizontal to vertical branches
	Treptichnus	17: Horizontal burrows with horizontal to vertical branches
	Monocraterion	19: Radial to rosetted structures
	Teichichnus	22: Horizontal burrows with simple vertically oriented spreiten
	Spirophycus	23: Horizontal helicoidal burrows
	Bergaueria	34: Vertical plug-shaped burrows
	Skolithos	35: Vertical unbranched burrows
	Arenicolites	36: Vertical single U- and Y-shaped burrows
	Diplocraterion	36: Vertical single U- and Y-shaped burrows
	Belorhaphe	46: Uniramous meandering graphoglyptids
Muav Limestone	*Phycodes*	17: Horizontal burrows with horizontal to vertical branches
Temple Butte Formation	*Palaeophycus*	10: Passively filled horizontal burrows
Surprise Canyon Formation	*Conostichus*	34: Vertical plug-shaped burrows

Table 8-1 (continued). List of valid ichnogenera and their respective architectural designs (sensu Buatois et al. 2017), ordered by geological unit (from lower to upper).

Unit	Valid ichnogenera	Architectural designs
Wescogame Formation	*Helminthopsis*	1: Simple horizontal trails
	Diplichnites	6: Trackways and scratch imprints
Hermit Formation	*Lithographus*	6: Trackways and scratch imprints
	Stiaria	6: Trackways and scratch imprints
	Scoyenia	12: Simple actively filled (meniscate) horizontal to oblique structures
	Sphaerapus	13: Simple actively filled (pelletoidal) horizontal to oblique structures
Coconino Sandstone	*Diplopodichnus*	5: Bilobate trails and paired grooves
	Diplichnites	6: Trackways and scratch imprints
	Lithographus	6: Trackways and scratch imprints
Coconino Sandstone	*Paleohelcura*	6: Trackways and scratch imprints
	Stiaria	6: Trackways and scratch imprints
	Octopodichnus	6: Trackways and scratch imprints
	Taenidium serpentinum	14: Complex, actively filled horizontal structures

Table 8-2. List of invalid ichnotaxa, ordered by geological unit (from lower to upper).

Unit	Invalid ichnotaxa	Current interpretation
Tapeats Sandstone	*Isopodichnus*	Junior synonym of *Cruziana*
	Corophioides	Junior synonym of *Diplocraterion*
	Stipsellus	Junior synonym of *Skolithos*
Bright Angel Shale	*Cruziana grandcanyonensis*	*Nomen nudum*
	Cruziana rusiformis	*Nomen nudum*
	Isopodichnus problematicus	Junior synonym of *Cruziana problematica*
	Scalarituba	Junior synonym of *Nereites*
	Scalarituba alternolobus	*Nomen nudum*
	Scalarituba novum	*Nomen nudum*
	Flectostriatus imporcatus	*Nomen nudum*
	Fodichnites bitumulus	*Nomen nudum*
	Phycodes pedum	Synonym of *Treptichnus pedum*
	Frondichnus vespirolatus	*Nomen nudum*
	Bicavichnites lophoseparatus	*Nomen nudum*
	Bicavichnites martini	*Nomen nudum*
	Corophioides	Junior synonym of *Diplocraterion*
Hermit Formation	*Walpia hermitensis*	Junior synonym of *Sphaerapus larvalis*
Coconino Sandstone	*Triavestigia niningeri*	Junior synonym of *Paleohelcura tridactyla*
	Paleohelcura dunbari	Junior synonym of *Paleohelcura tridactyla*
	Mesichnum	Junior synonym of *Paleohelcura*

Table 8-2 (continued). List of invalid ichnotaxa, ordered by geological unit (from lower to upper).

Unit	Invalid ichnotaxa	Current interpretation
Coconino Sandstone (continued)	*Octopodichnus minor*	Junior synonym of *Octopodichnus didactylus*
	Octopodichnus raymondi	Regarded as *Paleohelcura* isp.
	Isopodichnus filiciformis	Regarded as *Diplichnites* isp.
	Unisulcus sinuosus	Probable junior synonym of *Diplopodichnus biformis*
	Scolecocoprus cameronensis	Junior synonym of *Taenidium serpentinum*
	Scolecocoprus arizonensis	Probable junior synonym of *Taenidium serpentinum*

Institutional Abbreviations—**GRCA**, Grand Canyon National Park Museum, Arizona, USA; **MNA**, Museum of Northern Arizona, Flagstaff, Arizona, USA; **MSP**, Museo di Serrapetrona, Macerata, Italy; **RAM**, Raymond M. Alf Museum of Paleontology, Claremont, California, USA; **SMM**, Science Museum of Minnesota, Saint Paul, Minnesota, USA; **UCMP**, University of California, Berkeley, California, USA; **USNM**, Smithsonian, National Museum of Natural History, Washington, D.C., USA

The museum collections of Grand Canyon National Park contain a robust record of invertebrate trace fossil specimens from several formations throughout GRCA's history. However, many have been deaccessioned or contain limited ichnotaxonomic, locality, and reference information because they were collected so long ago.

Cambrian

Tonto Group

The Tonto Group sits upon the Precambrian basement and forms a gradual slope onto what is known as the Tonto Platform in Grand Canyon. These rocks can be traced from the west along the Colorado River to the east midway down the canyon. The Tonto Group has recently been expanded to comprise five formations, in ascending order the Sixtymile Formation, Tapeats Sandstone, Bright Angel Shale, Muav Limestone, and Frenchman Mountain Dolostone (Karlstrom et al. 2020). The Tapeats Sandstone, Bright Angel Shale, and Muav Limestone are the historic core of the Tonto Group and form a typical transgressive sequence. The Frenchman Mountain Dolostone, previously known as the undifferentiated dolomites first mentioned by McKee and Resser (1945), is cut into by the paleochannels of the Devonian Temple Butte Formation.

The Tonto Group represents a time of global sea level rise with minor transgressive and regressive events, creating an ideal habitat for Cambrian invertebrate organisms. Approximately 500 million years ago, Cambrian strata were deposited along the western coast of Laurentia near the palaeoequator. Tonto Group sediment accumulated on a shallow marine shelf with various tidal channels, embayments, intertidal flats, and monadnocks from the underlying Precambrian surface (McKee and Resser 1945; Hereford 1977; Middleton 1989). The reaction between carbonate build-ups and the eroding, siliceous bedrock may have created nutrition for microbial mats, invertebrate grazers, and benthic feeders (Rose 2003). Although the remains of invertebrates are relatively sparse in the Tonto Group, the preserved behavior of these organisms is extremely abundant. In fact, the

Tonto Group contains the majority of invertebrate ichnogenera and ichnospecies reported from Grand Canyon. These trace fossils mainly consist of varied burrows, arthropod trackways, resting and feeding traces, dwelling structures, and other locomotive tracks (Martin 1985; Lane et al. 2003; Middleton and Elliott 2003).

Tapeats Sandstone

The Tapeats Sandstone, historically the basal-most unit of the Tonto Group, is known for its lower contact with the underlying Precambrian surface called "The Great Unconformity". Due to the uneven underlying surface, the Tapeats Sandstone varies in thickness from 30 to 120 m (98 to 394 ft) (Noble 1922; Middleton and Elliott 2003). The basal portion of the Tapeats Sandstone is composed of conglomeratic quartz and feldspar and commonly reflects the mineralogy of the Precambrian basement rocks. It typically forms a cliff of stacked, 1-m-thick (3 ft), cross-stratified sandstone beds. The upper unit consists of interbedded mudstone and fine- to medium-grained sandstone and grades into the upper Bright Angel Shale, via what is often referred to as the "transition zone" (McKee and Resser 1945; Middleton and Elliott 2003).

The Tapeats Sandstone was deposited on a shallow-marine shelf within intertidal areas of deposition seen in the older deposits toward the west, and fluvial regions of deposition seen in the younger deposits toward the east (Hereford 1977). The maximum depositional ages obtained by U/Pb dating from detrital zircons are 505.4 ± 8 Ma to 501.4 ± 3.8 Ma (Karlstrom et al. 2018), indicating that the Tapeats Sandstone represents a gradual advancement of the sea during the Cambrian Epoch 3. An evolving, braided delta plain emptied into a high-energy shoreface environment in which sediment was constantly reworked (Schuchert 1918; Rose 2003). Due to the high-energy environment in the basal portion of the Tapeats Sandstone, trace fossils are more common in the upper transition zone where the energy waned and the sea advanced inland. These traces commonly consist of vertical burrows and a variety of horizontal tracks and trails (Middleton and Elliott 2003) and were most likely created by gastropods and annelid worms (Spamer 1992) as well as arthropods (trilobites and crustaceans).

Trace Fossils

Worm burrows and trilobite trails have been reported from the Tapeats Sandstone (e.g., Noble 1922; Schenk and Wheeler 1942; McKee and Resser 1945), but detailed reports on the ichnotaxa are somewhat lacking. Many authors in the past used generalized identifications, such as annelid burrows or "fucoidal casts" (Schuchert 1918). The word "fucoid" (from the Greek *fýkos*, seaweed) originally referred to algae, because many paleontologists from the 1800s thought that trace fossils were algal structures (Osgood 1975). Thus, this word is only used here if it was the only identification in the literature or from collections in the past.

Passively Filled Horizontal Burrows

Schuchert (1918) first noted the presence of *Palaeophycus* in the Tapeats Sandstone and identified them as "seaweeds". *Palaeophycus* is now understood to be horizontally oriented, passively infilled burrows that record the dwelling behavior of predatorial or suspension-feeding organisms (Buatois et al. 2017).

Simple Actively Filled (Massive) Horizontal to Oblique Structures

Hagadorn et al. (2011) reported a few occurrences of *Planolites* in the Tapeats Sandstone. *Planolites* is often confused with *Palaeophycus* due to their similar morphology, but *Planolites* shows an actively infilled burrow of sediment different from its matrix that indicates it was made by a deposit feeder (Pemberton and Frey 1982).

Horizontal Burrows with Horizontal to Vertical Branches

Treptichnus was reported in eastern GRCA by Hagadorn et al. (2011). *Treptichnus* is a horizontal to oblique, zig-zag burrow sometimes branching or overlapping. This ichnogenus encompasses various morphology types and therefore contains several ichnospecies, one of which is discussed further in the Bright Angel Shale section.

Horizontal Burrows with Vertically Oriented Spreiten (Sedimentary Laminae Caused by a Moving Organism)

Hagadorn et al. (2011) noted an abundance of *Teichichnus* in the shale interbeds of the Tapeats Sandstone. *Teichichnus* is a straight to slightly curved, horizontal burrow with tapered ends and vertically stacked spreiten (or laminae) that record vertical shifting in the sediment (Knaust 2018).

Trackways and Scratch Imprints

A specimen collected by McKee in 1937 was assigned to *Diplichnites*, a well-known ichnogenus of arthropod trackway consisting of two parallel rows of closely spaced small tracks. Hagadorn et al. (2011) reported the occurrence of *Diplichnites* in eastern GRCA and central Arizona. *Monomorphichnus* was also reported in the Tapeats Sandstone (Hagadorn et al. 2011) and consists of parallel sets of ridges or grooves indicating dragging or swimming behavior, usually of arthropods.

Bilobate Trails and Paired Grooves

A specimen collected by McKee and Wheeler in 1937 has been assigned to *Isopodichnus*, which is a synonym of the more commonly used ichnogenus name *Cruziana*. *Cruziana* is a near-surface, horizontal burrow consisting of lobed furrows covered by striations and was most commonly made by trilobites or other arthropods. *Cruziana* typically represents feeding behavior within a shallow-marine, fluvial, or lacustrine environment. Seilacher (1970) reported *Cruziana arizonensis* from the upper portion of the Tapeats Sandstone. *C. arizonensis* mainly consists of shorter tracks with a 2.5 to 5.5 cm (0.98 to 2.2 in) width. *Cruziana* was also reported in the Tapeats Sandstone by Hagadorn et al. (2011).

Bilaterally Symmetrical, Short, Scratched Impressions and Burrows

Two specimens collected by McKee and Wheeler in 1937 at Grand Wash Cliffs have been assigned to *Rusophycus didymus* and *Rusophycus* isp. A specimen collected by Seilacher in 1955 is also assigned to *R. didymus*. *Rusophycus* is generally considered to be the resting trace made by the producer of *Cruziana* and thus has a similar morphology without the continuation of a trail. *Rusophycus* was reported in the Tapeats Sandstone by Hagadorn et al. (2011) and observed by one of the authors (AEM) in central GRCA (Figure 8-1).



Figure 8-1. *Rusophycus* in convex hyporelief within the Tapeats Sandstone, central Grand Canyon (NPS/ANNE MILLER).

Burrows with Horizontal Spreiten

A specimen collected by Seilacher in 1955 shows a U-shaped feeding burrow, assigned to *Rhizocorallium* or *Corophioides* (the latter is considered to be a junior synonym of *Diplocraterion*; see below). *Rhizocorallium* is a burrow with U-shaped spreiten parallel or oblique to the bedding plane and likely records dwelling or feeding behavior (Knaust 2013).

Vertical Unbranched Burrows

Skolithos is common in the Tapeats Sandstone (Frech 1893; McKee and Resser 1945; Middleton and Elliott 2003; Hagadorn et al. 2011). It consists of straight, vertical, unornamented cylindrical burrows made by a variety of invertebrates for dwelling, suspension feeding, and predation, and is found in almost every type of depositional environment throughout geologic history. Howell (1957) named *S. annulatus* for the *Skolithos* specimen collected by McKee and Resser (1945; Plate 8, Figure c). A specimen collected by Seilacher in 1955 has been assigned to *Skolithos* cf. *S. linearis.*

Howell (1957) applied the name *Stipsellus annulatus* to vertical, cylindrical burrows approximately 12 mm (0.47 in) in diameter with evenly spaced menisci (back-filled, annulate structures). These were found in the upper Tapeats Sandstone. Häntzschel (1975) considered *Stipsellus* a problematic

ichnotaxon, and the illustration of it in Häntzschel (1975: Figure 132.2) suggests that it may even be the cast of part of an eocrinoid column. Alpert (1974) considered *Stipsellus* a junior synonym of *Skolithos.*

Vertical Single U- and Y-Shaped Burrows

Schuchert (1918) mentioned the abundance of *Arenicolites,* which are vertically oriented, U- or J-shaped, cylindrical burrows that record feeding and dwelling behaviors made by a variety of invertebrate organisms, including polychaete worms and crustaceans. Hagadorn et al. (2011) also reported the abundance of *Arenicolites* in the Tapeats Sandstone creating highly bioturbated beds at outcrops within eastern and western GRCA. *Diplocraterion* (formerly *Corophioides*) is another vertical, U-shaped burrow common in the Tapeats Sandstone (Hereford 1977; Middleton and Elliott 2003). However, *Diplocraterion* differs from *Arenicolites* in that it contains spreiten, which indicates adjusting behavior to maintain a connection with the changing sediment-water interface. *Diplocraterion* is primarily a dwelling burrow of crustaceans and is usually found in shallow-water environments.

Discussion

A total of 12 valid ichnogenera (*Arenicolites*, *Cruziana*, *Diplichnites*, *Diplocraterion*, *Monomorphichnus*, *Palaeophycus*, *Planolites*, *Rhizocorallium*, *Rusophycus*, *Skolithos*, *Teichichnus,* and *Treptichnus*) have been reported from the Tapeats Sandstone, but only a few of them were classified to the ichnospecies level (*C. arizonensis*, *R. didymus*, and *S. annulatus*) (Table 8-1). The identification of *S. linearis* is still tentative, and its presence in the Tapeats Sandstone requires further evaluation. Other ichnotaxa proposed for this unit include *Isopodichnus* and *Corophioides*, which are currently interpreted as synonyms of *Cruziana* and *Diplocraterion*, respectively. Seven architectural designs can be recognized among the Tapeats Sandstone ichnological record. The ichnoassociation of the Tapeats Sandstone belongs to the transition between the *Cruziana* ichnofacies and the *Skolithos* ichnofacies of shallow marine settings; this should mean a more proximal, high-energy environment compared to the typical *Cruziana* ichnofacies (e.g., Buatois and Mángano 2011).

GRCA Museum Collections

A “fucoid” specimen from the Tapeats Sandstone was collected along Bright Angel Trail in 1928 by Glen Sturdevant (GRCA 2088). In October 1929, Edwin McKee collected specimens of trilobite tracks and trails from the upper Tapeats Sandstone (or basal Bright Angel Shale) on the Tonto Platform (GRCA 2059, GRCA 2062, GRCA 2063, GRCA 2069, and GRCA 2071). That same year, he also collected “fucoids” from the upper Tapeats Sandstone (or basal Bright Angel Shale), also on the Tonto Platform (GRCA 8640). Trilobite tracks originally identified as *Isopodichnus* (currently *Cruziana*) were found in the Tapeats Sandstone and collected by McKee and Wheeler in January 1937 (GRCA 2137). In that same month at Grand Wash Cliffs, they also collected an indeterminate *Rusophycus* ichnospecies (GRCA 2138) and *R. didymus* (GRCA 2143) from the upper Tapeats Sandstone. In November 1937, McKee collected *Diplichnites* from a Tapeats Sandstone cast in Fern Glen Canyon (GRCA 2146). In March 1939, McKee collected two fossil trails near the top of the Tapeats Sandstone (GRCA 8603 and GRCA 8604, both deaccessioned) and one from Plateau Point (GRCA 8605—deaccessioned), all of which he attributed to worms. On September 9, 1955, Adolph

Seilacher collected the vertical tubes of *Skolithos* cf. *S. linearis* (GRCA 8677), the resting traces of *Rusophycus didymus* (GRCA 8678), and the U-shaped feeding burrow *Rhizocorallium* or *Diplocraterion* (formerly *Corophioides*) (GRCA 8679) within the transition zone of the Tapeats Sandstone. On October 17, 1955, he also collected the U-shaped burrows of *Arenicolites* from the Tapeats Sandstone west of Plateau Point (GRCA 8700). In 1969, George Beck collected a Tapeats Sandstone specimen with "worm trails" (GRCA 21235—deaccessioned). A specimen with borings and numerous cylindrical structures was collected on August 2, 1963 (GRCA 17435—deaccessioned). *Skolithos annulatus* (formerly *Stipsellus annulatus*) was collected by Howell (1957) (GRCA 8538).

Bright Angel Shale

The Bright Angel Shale is a unit placed in the middle of the Tonto Group, forming the iconic light green slopes that grade onto the Tonto Platform. It is exposed throughout Grand Canyon, ranging from 99 m (325 ft) thick in the east and 82 m (269 ft) thick in central GRCA to more than 137 m (449 ft) thick in western GRCA (Middleton and Elliott 2003). The wide range of thicknesses is due to the transitional nature of the upper and lower contacts, which are often interpreted differently by researchers. This is because the Bright Angel Shale gradationally overlies the Tapeats Sandstone and has an intertonguing relationship with the overlying Muav Limestone. The Bright Angel Shale consists of interbedded siliciclastic sandstone, siltstone, and shale. These rocks are mostly composed of quartz with some amount of potassium feldspar and muscovite, as well as glauconite that imbues the characteristic green color seen in the deposits. McKee and Resser (1945) recognized one formal member in the Bright Angel Shale, the Flour Sack Member, which comprises the uppermost strata of the unit in western GRCA. This member appears as a massive bed and contains sandstone, siltstone, and shale, as well as limestone tongues towards the top. Many of the massive sandstones in the Bright Angel Shale contain cross-stratification, and the interbedded strata often contain wavy or lenticular heterolithic bedding, all of which indicate a changing tidal environment.

The heterogeneous lithology and stratigraphy of the Bright Angel Shale represent strandline migration of a shallow-marine shelf that provided a habitat for many invertebrates. This environment was generally within subtidal to intertidal areas in which fair-weather suspension settling and storm influence was frequent (Martin 1985; Middleton 1989). Estuaries may have also existed in certain areas (Rose 2006). Invertebrates such as trilobites, brachiopods, and gastropods flourished in this environment.

McKee and Resser (1945) used trilobite fossil assemblages to constrain the age of the Bright Angel Shale. The basal portion in western GRCA contains the *Olenellus–Antagmus* horizon, which dates the bottom as late Cambrian Epoch 2. The upper portion of the Bright Angel in the east contains the *Alokistocare–Glossopleura* horizon, which dates the upper contact as Cambrian Epoch 3. However, Karlstrom et al. (2018) noted that the Tapeats Sandstone is younger than previously thought and does not get younger to the east, as seen via recent U/Pb detrital zircon data from the basal portion matching correlative formations throughout the west. These detrital zircons from the basal Tapeats Sandstone were dated as 505–501 million years old. This means that the whole of the Bright Angel

Shale is most likely Cambrian Epoch 3 in age, but more work is necessary to solidify the age constraints of the Tonto Group.

Trace Fossils

Body fossils are relatively rare in the Bright Angel Shale, but an astounding record of their behavior is common. Most invertebrate trace fossil reports include arthropod tracks and trails, worm burrows, and dwelling structures identified at the ichnogenus and ichnospecies level. Other reports only briefly mention traces with very limited ichnotaxonomic identifications. Noble (1922) mentioned the presence of "worm markings" and "fucoidal casts" within the green, glauconitic laminations of the Bright Angel Shale. McKee and Resser (1945) reported circular trails, worm borings, and "trilobite trails in hematitic sandstone". Strother and Beck (2000) found wrinkle structures in the Sumner Butte area similar to the "interference tadpole ripples" that McKee and Resser (1945) identified and compared them to modern microbial mats from supratidal regions. Baldwin et al. (2004) used the same comparison and identified these structures as "wrinkle marks" formed when microbial mats bind with the sediment. Davies et al. (2016) addressed the enigma of wrinkle structures and classified them as "microbially induced sedimentary structures" (MISS).

For the following list of ichnotaxa, it is important to note that any new ichnotaxonomic names proposed by Martin (1985) or Rose (2003) must be considered as *nomina nuda* according to the International Code of Zoological Nomenclature (1999) because they were only introduced in an unpublished thesis.

Simple Horizontal Trails

Martin (1985) identified *Cochlichnus*, which is a sinuous, locomotive, surface burrow in hypichnial preservation most likely made by annelid worms and often found in association with *Belorhaphe, Palaeophycus, Cruziana, Rusophycus,* and *Teichichnus. Belorhaphe* and *Cochlichnus* were both found in eastern GRCA.

Bilobate Trails and Paired Grooves

White (1874) identified *Cruziana linnarssoni* and *C. rustica* from the "Tonto Shale" (currently interpreted as part of the Bright Angel Shale) in Grand Canyon (Figure 8-2). *C. linnarssoni* has slightly pointed ends with a distinct, median furrow and contains little to no rugae (ridges) on the surface. *C. rustica* has blunt ends with a shallow, median furrow and contains many distinct rugae on the surface. "*Cruziana grandcanyonensis" nomen nudum* (proposed holotype MNA N3857) was found in sandstone throughout the Bright Angel Shale in eastern GRCA (Martin 1985). This ichnospecies consists of horizontally oriented, elongated, ridged furrows in hypichnial preservation with medial grooves. It represents the crawling or feeding behavior of an arthropod and typically is associated with *Rusophycus*, which indicates that these traces were most likely made by the same organism. According to Rose (2003), "*Cruziana rusiformis" nom. nud.* (GRCA 76909) is his *nominen substitutum* for Martin's (1985) "*C. grandcanyonensis" nom. nud.*, but both are unpublished names, so they are *nomina nuda. "C. rusiformis" nom. nud.* was found in association with *Palaeophycus*, sometimes more specifically with *P. striatus*, and potentially showed predatory behavior in central Grand Canyon as well as outside the park (GRCA 76928). "*C. rusiformis*" *nom. nud.* often transitions to *Rusophycus dispar*. *Isopodichnus problematicus* (specimen MNA N3716,

currently *Cruziana problematica*) was identified near Grand Wash Cliffs in cross-stratified sandstone and consists of bilobate burrows with a medial groove oriented horizontally in hypichnial preservation (Martin 1985).



Figure 8-2. *Cruziana* isp. from the Bright Angel Shale preserved in convex hyporelief (NPS/GRCA).

Trackways and Scratch Imprints

Martin (1985) identified *Diplichnites* (MNA N3718), which represents the walking tracks of arthropods and consists of obliquely oriented, parallel sets of ridges or tracks, which suggests a possible trilobite producer. This specimen was found near Grand Wash Cliffs and is usually associated with *Cruziana*, *Rusophycus*, *Dimorphichnus*, and *Monomorphichnus*. *Monomorphichnus* (MNA N3861) and *Dimorphichnus* (MNA N3858) were found in hypichnial preservation and record the walking behavior of arthropods along bedding planes (Martin 1985). They both consist of parallel sets of ridges, but differ in that *Monomorphichnus* typically has one set of ridges that is more prominent, indicating swimming or dragging behavior along the sediment. They are both usually associated with similar arthropod traces, such as *Cruziana*, *Rusophycus*, and *Diplichnites*, as well as *Palaeophycus*, and are often found throughout the heterolithic facies of the Bright Angel Shale. In western GRCA, in situ *Monomorphichnus lineatus* var. *giganticus* preserved in convex hyporelief was identified by one of the authors (AEM, pers. obs.) (Figure 8-3). It occurred within a green, micaceous shale layer beneath a fine sandstone in Martin's (1985) heterolithic facies. The specimen contains one 25-cm-long (9.8 in) row of laterally repeating, obliquely parallel striations and is

distinguished from *M. lineatus* by its large size and is therefore a variation of the ichnospecies described by Crimes et al. (1977). The specimen was found in association with *Teichichnus*, *Palaeophycus*, and *Rusophycus* and was most likely produced by a large trilobite approximately 26 cm (10 in) long.



Figure 8-3. In situ *Monomorphichnus* aff. *lineatus* var. *giganticus* preserved in convex hyporelief from the lower Bright Angel Shale (NPS/ANNE MILLER). Scale bar is 10 cm (4 in).

Elliott and Martin (1987) identified *Angulichnus alternipes* from float material in the lower portion of the Bright Angel Shale in eastern GRCA (holotype MNA N3862). The trace consists of alternating tracks bordering a central zig-zag furrow in epichnial preservation and was most likely created by an arthropod walking across the sandy surface. The authors attributed the potential producer of *A. alternipes* to be the enigmatic arthropod *Habelia*, based on its body plan.

Bilaterally Symmetrical Short, Scratched Impressions and Burrows

"*Rusophycus biungis*" *nom. nud.* (proposed holotype MNA N3865) is similar to *Cruziana* in that it is a bilobate furrow, but *Rusophycus* is a resting trace and does not continue as a trail. Instead, it consists of a "coffee bean-shaped", horizontally oriented, hypichnial V-marking (Figure 8-4). "*R. biungis*" was found in the heterolithic facies throughout the Bright Angel Shale in hypichnial preservation in close ichnoassociation with "*C. rusiformis/C. grandcanyonensis*", which most likely had the same producer (Martin 1985). These invalid ichnospecies of *Cruziana* often transition to *R. dispar* (GRCA 76915). In some cases, *R. dispar* was found with superposed *Palaeophycus* and other scratch marks in western Grand Canyon (GRCA 76919). *R. dispar* was also found in association with *Teichichnus*, another example of predatory behavior from eastern Grand Canyon (GRCA76903).

Passively Filled Horizontal Burrows

In the heterolithic facies, *Palaeophycus* aff. *P. striatus* was identified as straight or curved, hypichnial, cylindrical burrows oriented horizontally and somewhat obliquely to the bedding plane (Figure 8-5) (Martin 1985; Baldwin et al. 2004). In marine settings, *Palaeophycus* is likely produced by predaceous polychaetes or other suspension-feeding organisms (Díez-Canseco et al. 2016) and is a common trace fossil in the Bright Angel Shale.



Figure 8-4. *Rusophycus* isp. from the Bright Angel Shale preserved in convex hyporelief (NPS/ANNE MILLER).

Simple, Actively Filled (Massive) Horizontal to Oblique Structures

Baldwin et al. (2004) reported the presence of *Planolites* in the heterolithic beds (F2B facies) of the middle Bright Angel Shale. *Planolites* is a trace similar to *Palaeophycus*, but it has a burrow fill different from the host rock, whereas the fill of *Palaeophycus* is the same as the host rock. Thus, *Planolites* was an actively filled burrow, whereas *Palaeophycus* was passively filled (Pemberton and Frey 1982).

Complex, Actively Filled Horizontal Structures

Martin (1985) identified *Scolicia* (MNA N3867), which was found in a cliff-forming sandstone and consists of horizontally oriented, sinuous crawling or grazing trails of gastropods in epichnial preservation. In the Bright Angel Shale, *Scolicia* tends to be associated with *Diplocraterion* within the *Cruziana* ichnofacies, even though it is more commonly associated with the *Nereites* ichnofacies elsewhere. Another ichnotaxon described from the Bright Angel Shale by Martin (1985) is *Scalarituba*, which is currently regarded as a junior synonym of *Nereites* (see ichnotaxonomic discussion in Uchman 1995 and Mángano et al. 2000). It originally included two ichnospecies ("*S. alternolobus"* and "*S. novum" nom. nuda*), but both are considered *nomina nuda*.



Figure 8-5. *Palaeophycus* isp. from the Bright Angel Shale preserved in convex hyporelief (NPS/ANNE MILLER).

"Flectostriatus imporcatus" nom. nud. (proposed holotype MNA N3860) was identified as a horizontally oriented, meandering burrow with lateral and medial ridges in epichnial preservation (Martin 1985). It was found in cross-stratified sandstone and often has an ichnoassociation with "*Scalarituba alternolobus*" *nom. nud.*, which has a similar morphology. "*S. alternolobus*" (proposed holotype MNA N3866) differs from "*F. imporcatus*" *nom. nud.* in that it has biserially opposed lobes and medial chevrons in epichnial preservation (Martin 1985). "*S. alternolobus*" *nom. nud.* was also found in cross-stratified sandstone and is interpreted as a backfilled feeding burrow made by worm-like organisms.

Rose (2003) identified "*Scalarituba novum*" in association with *Diplocraterion* in the central Grand Canyon (GRCA 76901; Figure 8-6). In addition, *Scalarituba* was associated with *P. striatus* and *Diplocraterion* outside of Grand Canyon (GRCA 76930). "*Scalarituba novum*" exhibits plowing activity of the producer with a step-like morphology similar to Martin's (1985) "*Scalarituba alternolobus*" *nom. nud.* and "*Fodichnites bitumulus*" *nom. nud.*, which could be transitional forms. Rose (2003) also identified *Taphrhelminthopsis* (a preservational variant of *Scolicia*), which is a

ribbon-like, spiral trail similar to *Scalarituba*, and was found to be associated with *Cruziana*, *Planolites,* and *Rusophycus*.



Figure 8-6. "*Scalarituba novum*" (GRCA 76901) collected by Eben Rose in September 1998. Note the paired, funnel-shaped depressions of *Diplocraterion* in the lower left (Rose 2003). Scale bar is 10 cm (4 in) (NPS/KLARA WIDRIG).

Horizontal Burrows with Horizontal to Vertical Branches

Phycodes aff. *P. circinnatum* (MNA N3709) was found in hypichnial preservation and appears as branched, cylindrical burrows oriented horizontally along one plane (Martin 1985). *Phycodes* is generally thought to have been produced by annelid worms moving along the sand-mud interface. Martin (1985) also reported *Phycodes pedum* (MNA N3864) from the lenticular heterolithic facies at several locations. Recently, Buatois (2018) discussed the ichnotaxonomy of *Phycodes pedum,* arguing that this ichnospecies belongs to the ichnogenus *Treptichnus*. Buatois (2018) also noted the presence of *T. pedum* (formerly *P. pedum)* in the lower part of the Bright Angel Shale, where it seems to occur in association with *Cruziana, Diplichnites, Monomorphichnus, Palaeophycus striatus* (besides other forms of *Palaeophycus*), *Rusophycus,* and *Teichichnus*. *T. pedum* (formerly *P. pedum)* is a curved, horizontal burrow with oblique branches and represents a feeding trace produced by priapulids (Figure 8-7) (Buatois 2018).



Figure 8-7. *Treptichnus pedum* (formerly *Phycodes pedum*) from the Bright Angel Shale preserved in convex hyporelief (NPS/ANNE MILLER).

Another ichnotaxon currently considered as a synonym of *T. pedum* (formerly *P. pedum)* is *Bicavichnites* (Lane et al. 2003; Seilacher et al. 2005). The proposed holotype (MNA N3242) of "*B. lophoseparatus*" *nom. nud.* was found in the dolomitic Meriwitica Tongue of the Bright Angel Shale near Grand Wash Cliffs (Martin 1985). The type material of a second ichnospecies (*B. martini*; MNA N9500) also came from the same stratigraphic level (Lane et al. 2003). It consists of two parallel sets of circular impressions in epichnial preservation. Initially, it was interpreted as crawling or walking behavior of arthropods and is often associated with "*Frondichnus*" *nom. nud*. The putative producer of *Bicavichnites* has been interpreted as the lobopod *Aysheaia*. It is important to note that this interpretation was not considered for the synonymous ichnogenus, *Treptichnus.*

Horizontal Burrows with Simple, Vertically Oriented Spreiten

Teichichnus rectus (MNA N3869) is a common ichnospecies found in the Bright Angel Shale, occurring as hypichnial, epichnial, and exichnial preservation in the heterolithic facies (Martin 1985). It consists of horizontally oriented burrows, probably produced by annelids, which are slightly curved and contain spreiten, exhibiting a stacked appearance (Figure 8-8). Ichnoassociations for *T. rectus* include *Bergaueria, Cruziana*, *Palaeophycus, Phycodes*, and *Rusophycus* (Martin 1985). According to Rose (2003), *Teichichnus* was found in association with *Palaeophycus* in central GRCA (GRCA 76925).



Figure 8-8. *Teichichnus* isp. from the Bright Angel Shale showing horizontal spreiten and tapering at one end (NPS/ANNE MILLER).

Two different potential morphotypes of *Teichichnus* were identified by one of the authors (AEM, pers. obs.) (Figure 8-9). Figure 8-9A represents horizontally oriented, slightly curved burrows with vertically stacked spreiten preserved in both convex hyporelief and epirelief of very fine sandstone and is found in situ, protruding into platy, green, micaceous shale layers. *Teichichnus* in Figure 8-9B has a similar morphology and occurrence, but is much smaller in size. The burrows in Figure 8-9A have 18 cm lengths (7 in) with 12 mm diameters (0.5 in), and the burrows in Figure 8-9B have 3 to 4 cm lengths (1 to 1.6 in) with 4 mm diameters (0.2 in). In addition, the larger burrows were found within a shale-dominated unit, and the smaller burrows were found within a rippled sandstone-dominated unit. Knaust (2018) interpreted *Teichichnus* as potentially produced by polychaetes, holothurians, bivalves, or crustaceans. However, more work is needed to narrow down the producers of *Teichichnus* and further resolve the ichnotaxonomy of different forms.




Figure 8-9. In situ *Teichichnus* from the lower Bright Angel Shale (NPS/ANNE MILLER). **A.** Large form of *Teichichnus* in shale-dominated facies. Scale is 7 cm (3 in). **B.** Small form of *Teichichnus* in sandstone-dominated facies. Scale is 10 cm (4 in).

Radial to Rosette Structures

Baldwin et al. (2004) reported *Monocraterion* in the heterolithic facies, where it occurs in association with *Diplocraterion*.

Horizontal Spiral Burrows

Martin (1985) identified *Spirophycus* (MNA N3868) in hypichnial preservation. *Spirophycus* consists of an asymmetrical, horizontally oriented, coiled, grazing burrow within medium-grained sandstone.

Vertical Plug-Shaped Burrows

Martin (1985) identified *Bergaueria* aff. *B. perata*, which consists of vertically oriented cylinders with radial grooves surrounding its smooth walls in hypichnial preservation (Figure 8-10). The specimen was found in the heterolithic facies and is often associated with *Teichichnus*, *Palaeophycus*, and *Cruziana*. *Bergaueria* is commonly thought to have been produced by actinarian, cerianthid, or pennatulacean cnidarians (Buatois and Mángano 2011).



Figure 8-10. *Bergaueria* isp. from the Bright Angel Shale (NPS/ANNE MILLER). **A.** and **B.** In situ specimens preserved in convex hyporelief. **C.** Float specimens of small burrows (black arrow) and large burrows with central, linear depression (white arrow). Tick marks and boxes on scale bars equal 1 cm.

Vertical Unbranched Burrows

Skolithos linearis is composed of simple, cylindrical, vertical burrows and is found in cross-stratified sandstone at Tanner Canyon (Martin 1985).

Vertical Single U- and Y-Shaped Burrows

Martin (1985) identified *Diplocraterion yoyo* (MNA N3859), which is a vertically oriented, U-shaped burrow containing spreiten, often with paired funnels on the surfaces of beds that connect to the burrow (Figure 8-11). *D. yoyo* is found in thickly bedded sandstone throughout the Bright Angel Shale and is associated with unspecified, epichnial burrows. *Diplocraterion* is interpreted to be a

dwelling burrow of suspension feeders such as crustaceans. Rose (2003) described the presence of *Corophioides* (GRCA 76924; a junior subjective synonym of *Diplocraterion*) in the lower Bright Angel Shale from the eastern Grand Canyon, where it occurs in association with *Bergaueria*. Baldwin et al. (2004) reported the presence of *Arenicolites*.



Figure 8-11. *Diplocraterion* isp. from the Bright Angel Shale showing vertical spreiten between limbs indicating downward shifting in the sediment by the tracemaker (NPS/ANNE MILLER).

Uniramous Meandering Graphoglyptids (Network of Dwelling Burrows)

Martin (1985) identified *Belorhaphe,* a zig-zag, horizontal burrow in hypichnial preservation, in association with *Cruziana*, *Palaeophycus*, *Rusophycus*, and *Teichichnus*. They tend to be found in Martin's (1985) lenticular heterolithic facies that represents a subtidal environment influenced by fair-weather suspension settling of mud, silt, and sand from periodic storm events.

Other Ichnotaxa Regarded as Nomina Nuda

"*Frondichnus versiprolatus*" *nom. nud.* (proposed holotype MNA N3856) is an epichnial trace composed of alternating, oblique furrows made by arthropods, but the behavior is unknown. It was also found in the dolomitic Meriwitica Tongue near Grand Wash Cliffs (Martin 1985). "*Pholetichnus circinatus*" *nom. nud.* (proposed holotype MNA N3863) occurs in the heterolithic facies. This specimen is an epichnial trace exhibiting crawling behavior and consists of circular depressions on the bedding plane oriented in a spiral pattern. This trace was most likely produced by an organism

with a proboscis that probed the sediment for food (Martin 1985). The complete list of invalid ichnotaxa found in the Bright Angel Shale can be found in Table 8-2.

Discussion

The ichnological record of the Bright Angel Shale is very diverse, including many valid ichnotaxa such as: *Angulichnus alternipes, Arenicolites* isp., *Belorhaphe* isp., *Bergaueria* aff. *B. perata, Cochlichnus* isp., *Cruziana linnarssoni, C. problematica, C. rusiformis, C. rustica, Dimorphichnus* isp., *Diplichnites* isp., *Diplocraterion yoyo, Monocraterion* isp., *Monomorphichnus lineatus* var. *giganticus, Nereites* isp., *Palaeophycus* aff. *Pa. striatus, Phycodes* aff. *Ph. circinnatum, Planolites* isp., *Rusophycus dispar, Scolicia* isp., *Spirophycus* isp., *Skolithos linearis, Teichichnus rectus,* and *Treptichnus pedum.* Other ichnotaxa ("*Bicavichnites lophoseparatus*" *nom. nud., B. martini*, "*Cruziana grandcanyonensis*" *nom. nud., "Isopodichnus problematicus, "Scalarituba alternolobus*" *nom. nud.,* and *S. novum*) were formerly proposed for this unit, but recent works have reinterpreted them. The ichnodisparity of this record is also considered high, with 15 different architectural designs.

Baldwin et al. (2004) classified ichnogenera from the Bright Angel Shale into three groups (or ichnoguilds) based on feeding behavior: sediment feeding, filter and suspension feeding, and surface feeding. Sediment-feeding refers to burrowing and furrowing horizontally along bedding planes and includes ichnogenera such as *Cruziana*, *Palaeophycus*, *Phycodes*, *Rusophycus*, and *Teichichnus*. Filter and suspension-feeding behavior creates vertical and "U"-shaped burrows and includes *Arenicolites*, *Diplocraterion*, *Monocraterion*, and *Skolithos*. Surface-feeding refers to trail casts on bedding surfaces and includes *Palaeophycus* and *Planolites*. *Teichichnus* and *Phycodes* were typically found densely packed together on the underside of resistant sandstone beds within highly bioturbated, heterolithic facies. The predominance of ichnotaxa related to deposit-feeding behaviors (e.g., *Cruziana, Palaeophycus, Phycodes, Planolites, Scolicia,* and *Teichichnus*) characterizes the *Cruziana* Ichnofacies. This ichnofacies is typical of shallow-marine settings with moderate to low energy, ranging from the offshore to most distal shoreface. Low rates of sedimentation and erosion favor the preservation of locomotion traces of mobile carnivorous or scavenger invertebrates (e.g., *Angulichnus*, *Diplichnites*, *Dimorphichnus,* and *Monomorphichnus*). The high ichnodiversity and disparity of traces found in the Bright Angel Shale is typical of the *Cruziana* Ichnofacies.

GRCA Museum Collections

A listing of the collected trace fossil specimens from the Bright Angel Shale in Grand Canyon, housed in the GRCA Museum Collections can be found in Table 8.3.

Table 8-3. List of collected trace fossil specimens from the Bright Angel Shale in Grand Canyon, housed in the GRCA Museum Collections.

Catalog #	Name	Collector	Collection Date
GRCA 2072	*Scolicia*	Edwin D. McKee	07/1930
GRCA 2078	"fucoid"	Edwin D. McKee	11/1930
GRCA 2079	*Rusophycus*	Unknown	Unknown

Table 8-3 (continued). List of collected trace fossil specimens from the Bright Angel Shale in Grand Canyon, housed in the GRCA Museum Collections.

Catalog #	Name	Collector	Collection Date
GRCA 2080	*Rusophycus*	Edwin D. McKee	07/1930
GRCA 2081	Worm trail	R. L. Nichols	08/1930
GRCA 2082	“fucoid”	Unknown	Unknown
GRCA 2083	*Bergaueria*	Edwin D. McKee	04/1930
GRCA 2084	“fucoid”	Unknown	Unknown
GRCA 2085	*Teichichnus*	Unknown	Unknown
GRCA 2086	“fucoid”	Edwin D. McKee	10/1929
GRCA 2087	“fucoid”	Edwin D. McKee	10/1929
GRCA 2090	*Rusophycus*	Edwin D. McKee	11/1930
GRCA 2091	*Teichichnus*	Edwin D. McKee	11/1930
GRCA 2092	*Teichichnus*	Edwin D. McKee	11/1930
GRCA 2093	*Teichichnus*	Edwin D. McKee	1930
GRCA 2097	“fucoid”	Barbara Hastings	1929
GRCA 2101	“fucoid”	Lloyd Davis	1931
GRCA 2102	“fucoid”	Lloyd Davis	1931
GRCA 2103	“fucoid”	Lloyd Davis	1931
GRCA 2104	“fucoid”	Lloyd Davis	1931
GRCA 2105	“fucoid”	Lloyd Davis	1931
GRCA 2106	“fucoid”	Edwin D. McKee	08/1928
GRCA 2107	“fucoid”	Barbara Hastings	05/1929
GRCA 2109	*Rusophycus*	Edwin D. McKee	08/1933
GRCA 2110	“fucoid”	Edwin D. McKee	08/1935
GRCA 2111	“fucoid”	Edwin D. McKee	08/1936
GRCA 2116	*Teichichnus*	Edwin D. McKee	07/1928
GRCA 2117	Trilobite trail	Glen Sturdevant	Unknown
GRCA 2119	“fucoid”	Edwin D. McKee	05/1936
GRCA 2120	Spiracle fossil	Edwin D. McKee	05/1936
GRCA 2124	*Rusophycus didymus*	Edwin D. McKee	1936
GRCA 2126	Trilobite trail	Edwin D. McKee	05/1936
GRCA 2144	*Rusophycus*	Frank Richardson	05/1938
GRCA 2516	*Rusophycus*	Frank Richardson	05/1938
GRCA 2538	*Teichichnus*	Unknown	Unknown
GRCA 2563	Trilobite trail	Edwin D. McKee	05/1936
GRCA 2579	“fucoid”	Edwin D. McKee	02/1937
GRCA 2580	“fucoid”	Edwin D. McKee	02/1937
GRCA 2654	“fucoid”	Edwin D. McKee	05/1936
GRCA 2655	“fucoid”	Edwin D. McKee	05/1936
GRCA 2656	“fucoid”	Edwin D. McKee	05/1936
GRCA 2657	“fucoid”	Edwin D. McKee	05/1936

Table 8-3 (continued). List of collected trace fossil specimens from the Bright Angel Shale in Grand Canyon, housed in the GRCA Museum Collections.

Catalog #	Name	Collector	Collection Date
GRCA 2658	“fucoid”	Edwin D. McKee	05/1936
GRCA 2659	“fucoid”	Edwin D. McKee	05/1936
GRCA 2660	“fucoid”	Edwin D. McKee	05/1936
GRCA 2661	“fucoid”	Edwin D. McKee	05/1936
GRCA 2662	“fucoid”	Edwin D. McKee	05/1936
GRCA 2663	“fucoid”	Edwin D. McKee	05/1936
GRCA 5266	“fucoid”	Edwin D. McKee	02/1936
GRCA 5267	“fucoid”	Edwin D. McKee	02/1936
GRCA 5268	“fucoid”	Edwin D. McKee	02/1936
GRCA 6386	“fucoid”	Edwin D. McKee	02/1936
GRCA 8528	*Teichichnus*	Edwin D. McKee	07/1928
GRCA 8596	burrow	Edwin D. McKee	04/1939
GRCA 8643	burrow	Dorothy Gardner	05/09/1943
GRCA 8645	“fucoid”	Bill Meadows	01/1947
GRCA 8667	*Cruziana*	Adolph Seilacher	09/22/1955
GRCA 8674	burrow	Adolph Seilacher	10/1955
GRCA 8675	burrow	Adolph Seilacher	10/1955
GRCA 8683	*Rusophycus didymus*	Adolph Seilacher	09/22/1955
GRCA 8684	*Phycodes pedum*	Adolph Seilacher	09/25/1955
GRCA 8685	*Phycodes pedum*	Adolph Seilacher	09/25/1955
GRCA 8686	*Phycodes pedum*	Adolph Seilacher	09/1955
GRCA 8687	*Phycodes pedum*	Adolph Seilacher	09/1955
GRCA 8688	*Phycodes pedum*	Adolph Seilacher	09/1955
GRCA 8689	*Phycodes pedum*	Adolph Seilacher	09/1955
GRCA 8691	Trilobite trail	Adolph Seilacher	10/1955
GRCA 8693	*Teichichnus*	Adolph Seilacher	09/25/1955
GRCA 8701	*Scolicia*	Adolph Seilacher	10/17/1955
GRCA 8702	*Corophioides*	Adolph Seilacher	10/17/1955
GRCA 8703	*Corophioides*	Adolph Seilacher	10/17/1955
GRCA 8704	*Bergaueria*	Adolph Seilacher	10/17/1955
GRCA 8706	*Sinusites*	Adolph Seilacher	10/17/1955
GRCA 8707	*Sinusites*	Adolph Seilacher	10/17/1955
GRCA 8708	*Sinusites*	Adolph Seilacher	10/17/1955
GRCA 8709	Ichthidian	Adolph Seilacher	10/17/1955
GRCA 11448	*Rusophycus*	Edwin D. McKee	07/1930
GRCA 11449	*Rusophycus*	Edwin D. McKee	07/1930
GRCA 11450	“fucoid”	Edwin D. McKee	04/1930
GRCA 11451	“fucoid”	Edwin D. McKee	04/1930
GRCA 11452	*Rusophycus*	Edwin D. McKee	11/1930

Table 8-3 (continued). List of collected trace fossil specimens from the Bright Angel Shale in Grand Canyon, housed in the GRCA Museum Collections.

Catalog #	Name	Collector	Collection Date
GRCA 14467	*Teichichnus*	Unknown	09/1966
GRCA 14468	*Teichichnus*	Unknown	09/1966
GRCA 14469	*Teichichnus*	Unknown	09/1966
GRCA 14471	“fucoid”	Unknown	09/1966
GRCA 15231	*Rusophycus*	Unknown	Unknown
GRCA 15232	*Rusophycus*	Unknown	Unknown
GRCA 17182	Trilobite trail	Unknown	Unknown
GRCA 17415	trail	Edwin D. McKee	02/1936
GRCA 21046	“fucoid”	Edwin Rothfuss	04/20/1963
GRCA 21047	“fucoid”	Edwin Rothfuss	04/20/1963
GRCA 21402	“fucoid”	Edwin D. McKee	10/1939
GRCA 39124	“fucoid”	Unknown	Unknown
GRCA 76901	*Scalarituba novum*	Eben C. Rose	09/1998
GRCA 76902	*Palaeophycus*	Eben C. Rose	05/1998
GRCA 76903	*Rusophycus dispar*	Eben C. Rose	09/1998
GRCA 76904	*Rusophycus dispar*	Eben C. Rose	09/1998
GRCA 76905	*Monomorphichnus*	Eben C. Rose	05/1999
GRCA 76909	*Cruziana rusiformis*	Eben C. Rose	05/1999
GRCA 76910	*Diplichnites*	Eben C. Rose	08/1999
GRCA 76911	*Teichichnus*	Eben C. Rose	09/1998
GRCA 76913	*Rusophycus dispar*	Eben C. Rose	08/1998
GRCA 76914	*Palaeophycus*	Eben C. Rose	09/1999
GRCA 76915	*Palaeophycus*	Eben C. Rose	09/1999
GRCA 76916	*Palaeophycus*	Eben C. Rose	09/1999
GRCA 76917	*Cruziana rusiformis*	Eben C. Rose	05/1999
GRCA 76918	*Scalarituba novum*	Eben C. Rose	08/1999
GRCA 76919	*Rusophycus dispar*	Eben C. Rose	05/1998
GRCA 76920	*Palaeophycus*	Eben C. Rose	05/1999
GRCA 76921	*Cruziana rusiformis*	Eben C. Rose	05/1998
GRCA 76922	*Scalarituba novum*	Eben C. Rose	03/1998
GRCA 76924	*Corophioides*	Eben C. Rose	08/1998
GRCA 76925	*Teichichnus*	Eben C. Rose	09/1998
GRCA 76926	*Treptichnus pedum*	Eben C. Rose	05/1999
GRCA 76928	*Cruziana rusiformis*	Eben C. Rose	05/1999
GRCA 76929	*Treptichnus pedum*	Eben C. Rose	05/1999
GRCA 76930	*Scalarituba*	Eben C. Rose	05/1999
GRCA 76931	*Palaeophycus*	Eben C. Rose	09/1999
GRCA 76932	*Treptichnus pedum*	Eben C. Rose	05/1999
GRCA 85217	Eocrinoidea	Kate Watters	10/25/2005

Muav Limestone

The Muav Limestone is mainly exposed as cliffs above the Tonto Platform and has an intertonguing relationship with the underlying Bright Angel Shale. According to Schuchert (1918), faunal assemblages date the Muav Limestone as late Cambrian (Furongian). He also noted the potential for finding strata younger than Cambrian, below the Devonian contact with the Temple Butte Formation, although no research on this possibility has occurred. Seven members were designated by McKee and Resser (1945) based on invertebrate faunal assemblages and lithology. The top three members can be traced throughout Grand Canyon, and the lowest four members are only present in the western half. The Muav Limestone is generally composed of mottled mudstone and packstone, both calcareous and dolomitic in composition, as well as conglomeratic beds (Middleton and Elliott 2003). Most of the Muav Limestone beds are structure-less or horizontally laminated, although many localities do contain cross-stratified beds. The thickness of the formation ranges from 42 m (138 ft) in eastern exposures to 252 m (827 ft) in the western Grand Canyon (Middleton and Elliott 2003). Beds are thicker in the west, but the siliciclastic content is higher in the east. These features are compatible with the overall perspective of eastward coastline advancement.

The Muav Limestone represents a series of transgressive and regressive events in an offshore marine environment, conducive to supporting a habitat for invertebrates. The transgressive events were recorded in five members of the Muav Limestone, namely (ascending): Rampart Cave, Sanup Plateau, Spencer Canyon, Peach Springs, and Kanab Canyon members. The intertonguing relationship with the underlying Bright Angel Shale reveals a step-like pattern in which the Muav Limestone diagonally crosses time planes. This is because the sea advanced in rapid, periodic stages, interrupted by at least four regressive events of a lesser extent. The regressive events are recorded in the various tongues of sediment that protrude into the limestone (McKee and Resser 1945). Although body fossils are sparse in the Muav Limestone, bioturbation is extremely common within calcareous-dominated sediment and is responsible for the mottled appearance (Schuchert 1918).

Trace Fossils

Trace fossils are common in the Muav Limestone, but there has never been a comprehensive ichnotaxonomic treatment. Schuchert (1918) stated, “The writer has not seen a Paleozoic marine deposit more bored into and consumed by mud eaters than this one”. Noble (1922) also noted that almost every carbonate bed in the Muav Limestone contains “fucoidal markings”. Clearly, this unit yields a substantial amount of traces, but those reported have not yet been placed within any particular ichnotaxon. Schuchert (1918) pointed out that, although most of the traces in the Muav Limestone are not easily recognizable, there is sufficient, observable evidence that annelids were very common. These annelid burrows were mostly preserved within the layers of shale as heaps or “circular sausage-like strings” (Schuchert 1918). Nonspecific worm borings and trails were also reported in the Muav Limestone by Schuchert (1918), Noble (1922), and McKee and Resser (1945).

Horizontal Burrows with Horizontal to Vertical Branches

The only ichnotaxon reported from the Muav Limestone was a specimen collected by Schellbach in 1936 and assigned to *Phycodes* isp.

GRCA Museum Collections

Unspecified “fucoids” were found in the Muav Limestone and collected by Edwin McKee in October and November 1930 (GRCA 2089, GRCA 2095, and GRCA 2096). Louis Schellbach collected several, unspecified “fucoid” traces from the Muav Limestone in September 1936 (GRCA 2127, GRCA 2128, GRCA 2129, GRCA 2130, GRCA 2131, GRCA 2132, and GRCA 2133). He also collected *Phycodes* in September 1936 (GRCA 2134). Edwin McKee collected two “fucoid” specimens from the base of the Muav Limestone near Grand Wash Cliffs in the fall of 1936 that he identified as “worm tubes” (GRCA 8541 and GRCA 8542). Adolph Seilacher collected specimens with vertical tubes and branching “worm trails” from the upper Muav Limestone near Indian Gardens on September 20, 1955 (GRCA 8676 and GRCA 8682). Four reticulate burrow specimens were collected from the upper Muav Limestone along the Kaibab Trail by Adolph Seilacher on September 22, 1955 (GRCA 8668, GRCA 8669, GRCA 8670, and GRCA 8671). On the same day, Seilacher also collected a specimen identified as *Eophyton*, which are actually inorganic drag marks or tool marks found in the upper Muav Limestone along Kaibab Trail (GRCA 8672). “Fucoids” were also collected from the Muav Limestone from an unknown location by an unknown collector (GRCA 2098, GRCA 2108, and GRCA 17153).

Frenchman Mountain Dolostone

The Frenchman Mountain Dolostone is an interval of dolomite overlying the Muav Limestone, creating an unconformity that is distinguished by a high-relief, erosional surface. Historically known as the “undifferentiated dolomites”, it was assigned to the Frenchman Mountain Dolostone by Karlstrom et al. (2020), who made it the uppermost unit of the Tonto Group. It is mostly exposed in the western portion of Grand Canyon and extends from Kanab Creek to Nankoweap Canyon (Noble 1922). This dolomitic sequence is up to 131 m (430 ft) thick and has its best exposure at Grand Wash Cliffs (Middleton and Elliott 2003). Three lithofacies were identified by McKee and Resser (1945) and were described as a white, structureless dolomite, a yellow, bedded dolomite, and a gray, bedded dolomite. Some facies contain ripples and cross-stratification. Carbonates with oolitic grainstones and stromatolites were documented by Brathovde (1986), which indicates a shallow subtidal to intertidal environment.

Trace Fossils

The most common trace fossils present in this unit are invertebrate trackways and horizontal burrows (Middleton and Elliott 2003). These burrows and trails were first reported by McKee (1945), but no comprehensive report has been completed.

Devonian

Temple Butte Formation

The Temple Butte Formation is exposed as scattered, inconspicuous lenses cutting into the Tonto Group. The lenses are relatively thin at 30 m (98 ft) thick and discontinuous in the eastern portion of Grand Canyon and are more than 220 m (722 ft) thick and continuous towards the central and western regions, often merging with the overlying Redwall Limestone cliffs (Beus 2003a). These lenses are considered to be paleochannels or paleo-valleys (up to 120 m or 394 ft wide) that had cut

into the underlying, undifferentiated dolomites as well as the Muav Limestone. The Temple Butte Formation mainly consists of dolomite with some sandstone, limestone, and conglomerate.

The Temple Butte Formation is considered to be early Late Devonian in age, potentially in part as old as the Middle Devonian based on conodont assemblages, making this formation the only Devonian unit in Grand Canyon (Beus 2003a). The discontinuous lenses in eastern GRCA are thought to represent tidal channels, and the fossils in the central and western portion of GRCA indicate a shallow, subtidal, open-marine environment. These conditions allowed for invertebrates to flourish (see Chapter 5 of this book).

Trace Fossils

Several reports have mentioned the occurrence of worm borings and trails in the Temple Butte Formation (Schuchert 1918; Noble 1922; Stoyanow 1936; Bond et al. 2018), but very little was identified and described, most likely due to the nature of preservation and difficulty of access.

Passively Filled Horizontal Burrows

Beus (2003a) noted the presence of certain unidentified, cylindrical traces that look similar to the ichnogenus *Palaeophycus*, found in dolomitic beds within the lower Temple Butte Formation. Beus (2003a) mentioned that they are sub-horizontal and straight or gently curved with a micritized core and could potentially resemble some form of algal activity.

GRCA Museum Collections

In September 1938, Edwin McKee collected a Devonian limestone slab with “worm borings” outside of Grand Canyon National Park along Sycamore Creek near Verde Valley, Arizona (GRCA 5443).

Mississippian

Redwall Limestone

The prominent, red-stained cliffs that are exposed nearly mid-way vertically down the canyon are outcrops of the Redwall Limestone. The unit sits horizontally on the Cambrian Tonto Group and the lenses of the Devonian Temple Butte Formation. Redwall Limestone cliffs are 150 to 250 m (492 to 820 ft) high, stained by the overlying, iron oxide-rich material drained down from the Supai Group (Beus 2003b). The unweathered surface, however, is mainly light to dark gray in color. The Redwall Limestone can be traced throughout GRCA, from the east in Marble Canyon all the way to the west at Grand Wash Cliffs. Its thickness increases to the northwest, from about 120 m (394 ft) thick near Tanner Trail to about 245 m (804 ft) thick at Iceberg Ridge (McKee and Gutschick 1969). The Redwall Limestone consists of four members, in ascending order: the Whitmore Wash Member, Thunder Springs Member, Mooney Falls Member, and Horseshoe Mesa Member. The two oldest members consist of limestone that transitions to mostly dolomite towards the central and eastern portion of GRCA, both thickening to the west. The thickest member is the overlying Mooney Falls Member, which is mostly limestone. The Redwall Limestone is topped by the Horseshoe Mesa Member, consisting of alternating beds of dark chert and light carbonate. The Redwall Limestone represents a shallow sea that contained a diverse assemblage of organisms such as mollusks, echinoderms, arthropods, bryozoans, and fish. These fossilized organisms date the Redwall Limestone as Early Mississippian in age (Beus 2003b).

Trace Fossils

Though trace fossils are scarce in the Redwall Limestone, Stoyanow (1936) noted “impressions of fucoids on bedding planes”. Edwin McKee collected several invertebrate trace specimens in 1952–1962 that mainly consist of horizontal trails.

GRCA Museum Collections

Edwin McKee collected three limestone slabs that contain “worm trails” on the surface. The slabs were collected in 1952–1962 (GRCA 17937, GRCA 17938, and GRCA 17939). He also collected “worm trail” specimens near the top of the Redwall Limestone on October 13, 1958 (GRCA 20017, GRCA 20018, and potentially GRCA 20019, which is listed as by an unknown collector from the same locality). One “fucoid” specimen associated with “worm trails” was collected near the top of the Redwall Limestone by George Beck in August 1968 (GRCA 21226).

Surprise Canyon Formation

The Surprise Canyon Formation is exposed as lenses of clastic and carbonate sediment that filled in the karst topography of the Redwall Limestone during the Late Mississippian. The paleo-valleys thicken to the west from 45 to 60 m (148 to 197 ft) and can be up to 1 km wide (0.6 mi). They consist of a sandstone conglomerate at the base, skeletal limestone in the middle unit, and slope-forming siltstone and limestone at the top (Beus 2003b). The Surprise Canyon Formation was deposited in a broad valley by dendritic streams that filled in collapsed sinkholes of the Redwall Limestone as the sea receded.

Trace Fossils

Although body fossils are more common in the Surprise Canyon Formation, there have been a few brief mentions of invertebrate trace fossils. Trace fossils typical of the *Skolithos* ichnofacies have been found in the basal sandstone/conglomerate. Trace fossils typical of the *Cruziana* ichnofacies were seen in the middle unit of Surprise Canyon Formation (Beus 2003b). Bonde et al. (2018) also noted the presence of unspecified trace fossils in the Surprise Canyon Formation.

Vertical Plug-Shaped Burrows

The basal sandstone/conglomerate unit contains vertical burrows and *Conostichus* (Beus 2003b), which are cone-shaped burrows most likely produced by sea anemones. *Conostichus* is typical of the *Skolithos* ichnofacies.

Pennsylvanian

Supai Group

The Supai Group is exposed as massive red beds and slopes forming a cliffy, stair-stepped profile that overlies the Redwall Limestone and the paleochannels of the Surprise Canyon Formation. The prominent red color comes from the ferritic pigment of shales washing through, while the unweathered surface is more of a light tan to pink color. The Supai Group mainly consists of sandstone, but some beds contain sandy shale and limestone (Noble 1922). The group is divided into four formations including, from oldest to youngest: the Watahomigi Formation, Manakacha Formation, Wescogame Formation, and Esplanade Sandstone.

Trace Fossils

Burrows, most frequently smooth and cylindrical (parallel, perpendicular, and oblique to the bedding plane), occur mostly in carbonates and silty sandstones of the Supai Group. Invertebrate trackways and plant root bioturbation have also been observed (Blakey 2003).

GRCA Museum Collections

Two limestone specimens with "fucoidal structures" were collected from indeterminate strata of the Supai Group by Edwin McKee near the Shivwits Plateau on an unknown date. The specimens were loaned to McKee on February 4, 1972 and never returned; thus, they were deaccessioned (GRCA 5669 and GRCA 6504). McKee also collected limestone slabs "covered with worm tubes" in 1952–1962 from the Supai Group near the Bright Angel Trail (GRCA 12753 and GRCA 12754).

Watahomigi Formation

The Watahomigi Formation is the oldest unit, thickening westward from 30 to 90 m (98 to 295 ft) and containing siltstone, mudstone, limestone, and dolomite (Blakey 2003).

Trace Fossils

McKee (1982) mentioned the presence of "worm borings" in siltstone and limestone from two locations. Bonde et al. (2018) also noted the presence of "worm tubes" and microbially induced sedimentary structures (MISS) in the Watahomigi Formation.

Manakacha Formation

The overlying Manakacha Formation is dominantly sandstone with a few mudstone beds and is thickest in the central portion of GRCA, averaging 90 m (295 ft) thick (Blakey 2003).

Trace Fossils

"Horseshoe-crab-like invertebrate tracks" were observed by McKee (1982), as well as "worm borings and trails".

Wescogame Formation

The Wescogame Formation ranges from 30 to 60 m (98 to 197 ft) in thickness and consists of lithologies similar to those of the underlying Manakacha Formation. The upper contact of the Wescogame Formation forms the boundary with the Permian rocks of the Esplanade Sandstone, the uppermost unit of the Supai Group (Blakey 2003).

Trace Fossils

Gilmore (1928) described two arthropod trackways (USNM 11693 and 11740) from the same lithofacies where vertebrate tracks were found, i.e., cross-bedded sandstones of eolian origin (Figure 8-12). McKee (1982) noted the presence of worm borings on a silty sandstone cliff unit from the lower portion of the Wescogame Formation. He also mentioned two different types of unknown invertebrate tracks were present.




Figure 8-12. Invertebrate traces from the Wescogame Formation (LORENZO MARCHETTI). **A.** *Helminthopsis* isp., indicated by arrows, convex hyporelief (USNM 11577). **B.** *Diplichnites* isp., concave epirelief (USNM 11740B). **C.** Partial arthropod trackway, undetermined, convex hyporelief (USNM 11740).

Simple Horizontal Trails

On the holotype slab of the vertebrate ichnotaxon *Anomalopus sturdevanti* Gilmore 1928 (USNM 11577; Figure 8-12A), thin, shallow, horizontal, slightly meandering and unornamented burrows can be observed (width 2 mm or 0.08 in). They can be assigned to *Helminthopsis* isp.

Trackways and Scratch Imprints

The specimen USNM 11740B (Figure 12B) shows an arthropod trackway with two parallel rows of closely spaced elongated tracks with a medial furrow, about 2 cm (0.8 in) wide. This is referable to *Diplichnites* isp.

Permian

Esplanade Sandstone

The Esplanade Sandstone is the most sandstone-dominated unit of all of the Supai units, with cross-stratified sandstone beds. It begins the Permian strata of Grand Canyon and was mostly deposited in eolian settings (Blakey 2003).

Trace Fossils

McKee (1982) mentioned the presence of “worm borings” in muddy films within siltstone and sandstone of the upper Esplanade Sandstone at several locations. He also noted that resistant sandstone from the basal part of the Esplanade Sandstone contains “worm borings”. A siltstone with these traces and mud pellets was found in 150-Mile Canyon (McKee 1982). McKee (1982) measured

vertical burrows that were 3.3 mm (0.13 in) in diameter, creating a resistant sandstone ledge in the Esplanade Sandstone. Some horseshoe-shaped tracks found in mudstone were also reported from the Esplanade Sandstone (McKee 1982).

Hermit Formation

The Hermit Formation is a poorly exposed, slope-forming, siliciclastic unit that gradationally overlies the Supai Group. It is mainly composed of red-to-brown mudstone, siltstone, and very fine sandstone deposited in a low-energy coastal environment rich in flora and fauna. The siltstone and sandstone often contain ripple marks and cross-stratification associated with meandering streams. These rocks can also be observed in outcrops outside of GRCA near Sedona, Arizona. Structure-less, 1-m-thick (3 ft) sandstone beds often form resistant ledges that may have been further lithified with calcitic concretions (Blakey 2003). Siliciclastic mudstone forms slopes and is usually structure-less with the occasional ripple mark. The thickness of the Hermit Formation ranges from 30 m (98 ft) in eastern Grand Canyon to over 270 m (886 ft) in western Grand Canyon (Blakey 2003). It is sharply overlain by the Coconino Sandstone throughout Grand Canyon. The Hermit Formation is thought to be Leonardian in age (Cisuralian), which was first assigned by White (1929) and later repeated by McKee (1982) based on the plant fossil *Callipteris arizonae*.

Trace Fossils

The invertebrate trace fossil record of the Hermit Formation mainly consists of trackways, burrows, and resting and feeding traces, although root bioturbation and microbial structures are more common (White 1929; Blakey 2003). McKee (1982a) reported the presence of “worm burrows” and “worm tubes”. Other invertebrate traces were reported by Spamer (1992). Unfortunately, many of these trace fossils are unidentifiable due to high levels of bioturbation. However, certain beds in the lower part of the formation reveal distinct traces that can be identified at the ichnogenus and ichnospecies level.

Trackways and Scratch Imprints

Two specimens from outside of GRCA in Gila County at Craddock Creek (UCMP-V 4012 and UCMP-V 4013) show arthropod trackways with two rows of regularly spaced sets of two to three elongated tracks perpendicular to each other, assignable to *Lithographus* isp. (Figure 8-13B). The specimen UCMP-V 4013 also includes a possible arthropod resting trace. A specimen from Yavapai County (UMCP-V 75214; Figure 8-13A) shows a trackway with two parallel rows of tracks arranged in series of three tracks with opposite symmetry; this is referable to *Stiaria* isp. These three specimens come from the Mogollon Rim area, from units correlated with the Hermit Formation of the Grand Canyon (Blakey 1979).




Figure 8-13. Arthropod trackways from the Hermit Formation of the Mogollon Rim area (LORENZO MARCHETTI). **A.** *Stiaria* isp. (UMCP-V 75214). **B.** *Lithographus* isp. (UCMP-V 4012). Arrow indicates a possible arthropod resting trace. Scale bar is 1 cm (0.4 in).

Simple, Actively Filled (Meniscate) Horizontal to Oblique Structures

White (1929) identified the new ichnospecies *Scoyenia gracilis* (page 115, Plate 4), found in the lower portion of the Hermit Formation (Figure 8-14). *Scoyenia* is a deposit feeding or dwelling structure with straight, wrinkly, overlapping burrow structures and is thought to be made by either a larval or adult insect. White (1929) noted that *S. gracilis* was somewhat different than other possible forms of *Scoyenia* based on the fasciculate morphology and concluded that it was the mold of an animal boring into the substrate.



Figure 8-14. *Scoyenia gracilis* (GRCA 8538) from the Hermit Formation (White 1929). Scale bar is 1 cm (0.4 in) (NPS/KLARA WIDRIG).

Simple, Actively Filled (Pelletoidal) Horizontal Burrows

The ichnospecies *Walpia hermitensis* (lectotype USNM 263675: Figure 8-15A; Paralectotype: USNM 263676; Figure 8-15B) was proposed by White (1929: 117, Plate 51) based on a specimen found in association with *Scoyenia gracilis*. White (1929) interpreted the nodules in *W. hermitensis* as possible fecal pellets produced by worms or crustaceans as they tunneled the substrate. Alternatively, Häntzschel (1975) interpreted the nodules to be sediment packets from arthropods excavating and backfilling as they burrowed along the surface. Lucas et al. (2011) interpreted *Walpia* as an arthropod feeding or locomotion structure (most likely a shallow compaction burrow made by a hexapod) consisting of slightly curved to sinuous burrows or trails lined with small, rounded nodules. Recently, the ichnospecies *W. hermitensis* was regarded as a junior synonym of *Sphaerapus larvalis* (Lucas et al. 2013). Early coleopterans could be the potential producers of *S. larvalis* due to the similarity to modern, beetle-produced burrows (Lucas et al. 2011, 2013).




Figure 8-15. *Walpia hermitensis* syntypes from the Hermit Formation (White 1929) (SPENCER LUCAS). **A.** Lectotype in concave epirelief (USNM 263675). **B.** Paralectotype in concave epirelief (USNM 263676). Scale bar is 3 cm (1.2 in).

GRCA Museum Collections

"Fucoid" specimens from the Hermit Formation with "worm trails and borings" were collected by Edwin McKee in 1929. One specimen was found along Yaki Trail (GRCA 3023) and the other was found in Hermit Basin and classified as "vermes" associated with *Scoyenia gracilis* (GRCA 3068; GRCA 8538, Figure 8-14; White 1929: 115, Plate 4). McKee collected another "fucoid" specimen with worm trails from the Hermit Formation (GRCA 3200). A slab of shale with many "worm borings" from the Hermit Formation was collected in the adjacent Lake Mead National Recreation Area by Edwin McKee (GRCA 5793). McKee also collected a siltstone specimen containing worm trails from the Hermit Formation (GRCA 5815). Lawrence Goebels collected another worm trail specimen from the Hermit Formation (GRCA 3203).

Discussion

The Hermit Formation presents only four valid ichnotaxa (*Lithographus* isp., *Scoyenia gracilis*, *Sphaerapus larvalis*, and *Stiaria* isp.) classified among three architectural designs. Arthropod trackways and horizontal to oblique burrows produced in continental settings characterize the *Scoyenia* ichnofacies, which represents low energy, inundated (at least periodically) continental settings, such as riverine point bars, channel banks, and crevasse splays, and lake margins and floodplains (MacEachern et al. 2012). Tetrapod tracks (Gilmore 1927; see the vertebrate ichnology chapter of this report) that occur in association with invertebrate traces corroborate this interpretation.

Coconino Sandstone

The Coconino Sandstone is a relatively thick unit (up to 300 m or 984 ft) exposed in several areas of the southern Colorado Plateau (e.g., Blakey and Knepp 1989), and especially in the GRCA and Mogollon Rim areas of Arizona. It is composed of large-scale cross-bedded sandstones of clear eolian origin (e.g., McKee 1979). The foreset dune beds are usually inclined to about 15–30°. The sandstones are pale-buff and fine-grained. The grains are well rounded and quartz is predominant. Interdunal deposits are rare. This unit usually forms evident and recognizable cliffs due to its massive structure. In GRCA, this unit overlies the Hermit Formation, interfingers with and is overlain by the Toroweap Formation, and is overlain by the Kaibab Formation in areas where the Toroweap Formation is not present. Because of this stratigraphic position, it is considered to be early Permian (Kungurian/Leonardian) in age (Blakey and Knepp 1989). This is consistent with the tetrapod footprint ichnoassociation, belonging to the *Erpetopus* footprint biochron (Marchetti et al. 2019b).

Trace Fossils

The only fossils known from the Coconino Sandstone are trace fossils (Spamer 1984). Invertebrate traces are known from GRCA, including as possible producers, worms, millipedes, isopods, spiders, scorpions (e.g., Gilmore 1926, 1927; Brady 1947; Middleton et al. 1990), and insects (Spamer 1992). The most common invertebrate ichnofossil is the possible scorpion track *Paleohelcura* (see Brady 1939). Other arthropod tracks are also common (Gilmore 1927; Brady 1947; Sadler 1993), and meniscate horizontal burrows (Brady 1947) and conical pits have also been reported (Elliott and Bartlett 2013). Invertebrate trace fossils from this unit have been the subject of numerous ichnotaxonomic studies (e.g., Gilmore 1926, 1928; Brady 1947, 1961; Alf 1968; Sadler 1993;

Kramer et al. 1995; Minter et al. 2007; Davies et al. 2007). They are also an important reference for paleoecological studies through ichnofacies: the dune-low diversity *Octopodichnus–Entradichnus* ichnofacies is largely based on the Coconino invertebrate ichnoassociation (Ekdale et al. 2007; Hunt and Lucas 2007; Buatois and Mángano 2011). These traces usually come from the lower half of the unit and are commonly preserved on the foreset surfaces of the eolian dunes (e.g., Gilmore 1926, 1927). The animal tracks in the Coconino Sandstone were probably made on dry sediments that were then moistened and covered by windblown dry sand before the sediments and tracks again dried out, or on dunes dampened by dew (Brady 1939; McKee 1944).

Trackways and Scratch Imprints

The ichnotaxon *Paleohelcura tridactyla* (Figure 8-16) is common in the Coconino Sandstone. It was erected by Gilmore (1926) based on a long continuous trackway with sets of three tracks arranged in lines oblique to the midline and with a continuous medial impression (holotype USNM 11145; Figure 8-16A). The ichnotaxon *Triavestigia niningeri* Gilmore 1927 (holotype USNM 11510) is considered a junior subjective synonym of *Paleohelcura tridactyla* because it represents an incomplete specimen of *P. tridactyla* with only one row and a medial impression shifted laterally (Braddy 1995). The ichnotaxon *Mesichnium benjamini* Gilmore 1926 (holotype USNM 11155) was assigned to *Paleohelcura* (*P. benjamini*) and differentiated from *P. tridactyla* because of the peculiar discontinuous preservation of the medial impression (Kozur et al. 1994). The ichnospecies *P. dunbari* Brady 1961 was erected based on the triangular arrangement of the three tracks preserved (holotype MNA-N 3694; Figure 8-16B), distinguished from the linear arrangement of the *P. tridactyla* holotype. However, transitions between these two patterns are observed along the same trackway (Sadler 1993), so this difference represents a minor extramorphological variant, and *P. dunbari* has to be considered a junior subjective synonym of *P. tridactyla* (e.g., Minter and Braddy 2009). According to Minter and Braddy (2009), the ichnogenus *Paleohelcura* is similar in morphology and pattern to the ichnogenus *Stiaria* Smith 1909. However, the ichnospecies *S. intermedia* can be distinguished by the maximum number of tracks (three) and the smaller maximum external width (18 mm or 0.71 in) (e.g., Minter and Braddy 2009); the ichnospecies *S. quadripedia* can be distinguished by the linear arrangement of the four tracks (Walker 1985), not observed in specimens of *Paleohelcura* showing four tracks per set (e.g., MNA-N 3669; Figure 8-16C). Therefore, although the incomplete specimens can be very similar, we suggest continuing to distinguish these two ichnogenera. Due to the two sets of four tracks, the most likely producers were arachnids (Brady 1947; Sadler 1993). Experiments with present-day scorpions produced trackways with this specific pattern (e.g., Brady 1947; Sadler 1993).




Figure 8-16. Arthropod trackways from the Coconino Sandstone, attributed to *Paleohelcura tridactyla* (LORENZO MARCHETTI). **A.** Holotype of *P. tridactyla*, concave epirelief (USNM 11145). **B.** Holotype of *P. dunbari*, concave epirelief (MNA-N 3694). **C.** Trackways showing sets of four tracks, convex hyporelief (MNA-N 3669). **D.** Artificial cast of the holotype of *Octopodichnus raymondi*, concave epirelief (MNA-N 9391). Scale bar is 2 cm (0.4 in).

The ichnotaxon *Octopodichnus didactylus* (Figure 8-17) is common in the Coconino Sandstone. It was introduced by Gilmore (1927) based on a trackway with two parallel rows of alternating sets of four, anteriorly bifurcated tracks. The proximal track is in a medial position, and the three distal tracks are arranged on a line oblique to the midline, distally closer (holotype USNM 11501; Figure 8-17A). This ichnotaxon was later amended by Sadler (1993). Based on a specimen from Seligman (holotype MNA-N 3654; Figure 8-17B), Brady (1947) erected a second ichnospecies, *O. minor*, characterized by smaller size and an opposite track arrangement in the series: the anterior track was medial and the three posterior tracks were oblique and progressively farther from the midline. However, the tracks can be oriented and bifurcated not only anteriorly, but also laterally and almost posteriorly due to the substrate inclination at the time of impression (similarly to the tetrapod tracks on foreset dune surfaces; Marchetti et al. 2019b). Therefore, the strong lateral bifurcation of the *O. minor* holotype probably caused an incorrect interpretation of the direction of progression, so that the pattern appeared to be the opposite of *O. didactylus*. As a consequence, *O. minor* may be a junior subjective synonym of *O. didactylus*.




Figure 8-17. Arthropod trackways from the Coconino Sandstone (LORENZO MARCHETTI). **A.** Holotype of *Octopodichnus didactylus*, convex hyporelief (USNM 11501). **B.** Holotype of *O. minor*, convex hyporelief (MNA-N 3654). **C.** *Lithographus* isp., holotype cast of *Permichnium coconinensis*. Trackway with change in direction and termination against a tetrapod trackway (arrow), convex hyporelief (MNA-N 9408). **D.** *Diplichnites* isp., holotype of *Isopodichnus filiciformis*, convex hyporelief. White scale bars in A and C are 2 cm (0.8 in).

A further ichnospecies was introduced by Sadler (1993) based on a specimen from Seligman (holotype RAM-JF 5905) first described by Alf (1968) as *O. raymondi*. However, the track arrangement and morphology of this trackway correspond to the diagnostic features of *Paleohelcura* from Gilmore (1926); therefore, this specimen is not considered to belong to *Octopodichnus*, but it is herein preliminarily assigned to *Paleohelcura tridactyla* (Figure 8-17D). Due to the two sets of four tracks, the most likely producers were arachnids (Brady 1947; Sadler 1993). Although experiments with present-day animals have not thus far produced trackways with this specific pattern (e.g., Brady 1947; Sadler 1993), the lack of a telson ("tail") impression and the slightly irregular pattern may indicate a spider as the producer.

Kramer et al. (1995) erected the ichnospecies *Permichnium coconinensis* based on a specimen from Seligman, in the Mogollon Rim area (holotype SMM-P 92.3.1). This specimen shows a trackway composed of two parallel rows of sets of two tracks perpendicular to each other. These are diagnostic features typical of *Lithographus* isp. (Figure 8-17C). The possible producers are insects such as blattoids (e.g., Kramer et al. 1995; Minter and Braddy 2009). It is noteworthy that this trackway ends and changes its pattern in correspondence with a tetrapod trackway proceeding transverse to it with a quite high pace. This was interpreted as predation behavior of the tetrapod trackmaker on the insect trackmaker (Kramer et al. 1995), but Hunt and Lucas (1998) questioned that interpretation. The fast pace of the tetrapod trackway is actually due to downslope locomotion with a lateral component (Marchetti et al. 2019b), therefore the different appearance of the *Lithographus* trackway termination could be simply due to the deformation provided by the tetrapod passage, although predation cannot be excluded.

Based on a specimen from Seligman, in the Mogollon Rim area, Brady (1947) erected the ichnospecies *Isopodichnus filiciformis* (holotype MNA-N 3653). However, *Isopodichnus* is characterized by extremely closely spaced tracks, probably due to a feeding behavior on the substrate, while the described material shows less closely spaced tracks arranged in two parallel rows, probably due to locomotion behavior. The latter morphology is typical of the ichnogenus *Diplichnites*. Therefore, this material is probably assignable to *Diplichnites* (Figure 8-17D). In addition, this form can be found in association with *Diplopodichnus*. Isopod crustaceans and millipedes are the most probable producers of *Diplichnites* (Brady 1947; Davis et al. 2007; Minter et al. 2007).

Bilobate Trails and Paired Grooves

The ichnotaxon *Diplopodichnus biformis* (Figure 8-18) is common in the Coconino Sandstone. It was introduced by Brady (1947) based on material from the Coconino Sandstone near Seligman in the Mogollon Rim area (holotype MNA-N 3657; Figure 8-18A). It is characterized by two parallel grooves that may be separated by a continuous ridge. The trace course is continuous and slightly sinuous, and is about 4 to 10 mm (0.2 to 0.39 in) wide. Gilmore (1927) introduced a morphologically similar ichnotaxon, *Unisulcus sinuosus* (holotype USNM 11498; Figure 8-18B). This is probably a poorly preserved *Diplopodichnus biformis*, because most of the trace is a single furrow, but occasionally two furrows and a ridge are recognizable. In some specimens and for small areas, it is possible to observe small, closely spaced tracks perpendicular to the medial ridge arranged in two parallel lines (GRCA 2882; Figure 8-18C). This suggests a possible transition with *Diplichnites* or similar morphotypes (Figure 8-18D). In three specimens, a possible transition to *Taenidium* cf. *T. serpentinum* has been observed (GRCA 2882, GRCA-NN 1 and MNA-N 3656). These compound traces suggest a single trackmaker capable of producing different morphologies and structures because of differing behavior. Brady (1947) considered this ichnotaxon as locomotion trails of myriapods, changing the morphology according to the trackway direction compared to the slope (the ridge seems to occur mostly in trackways with a downslope direction).




Figure 8-18. Bilobate trails and paired grooves from the Coconino Sandstone. **A.** Holotype of *Diplopodichnus biformis* associated with *Paleohelcura* and *Octopodichnus*, concave epirelief (MNA-N 3657) (LORENZO MARCHETTI). Scale bar is 2 cm (0.8 in). **B.** Holotype of *Unisulcus sinuosus*, concave epirelief (USNM 11498) (LORENZO MARCHETTI). **C.** Possible transition of *Taenidium* with *Diplopodichnus*, convex hyporelief (GRCA 2882) (HEITOR FRANCISCHINI). **D.** Detail of the latter specimen, showing closely spaced imprints (HEITOR FRANCISCHINI).

Complex, Actively Filled Horizontal Structures

Brady (1947) named *Scolecocoprus* (Figure 8-19), thinking it was a string of coprolites, and two ichnospecies have been named: *Scolecocoprus cameronensis* (holotype MNA-N 3707; MNA P3.129; Locality 168; Figure 8-19A) and *Scolecocoprus arizonensis* (holotype MNA-N 3655; Figure 8-19B). DeCourten (1978) later re-interpreted *Scolecocoprus* as a burrow with the behavior of a suspension-feeding organism responding to rapid sedimentation in a shallow-marine environment, thus creating "vertical and lateral displacements" in the burrow, as did D'Alessandro and Bromley (1987), who considered it a junior synonym of *Taenidium*. Indeed, the similarity of *Scolecocoprus cameronensis* (the type ichnospecies) to *Taenidium serpentinum* is strong and supports the conclusion of D'Alessandro and Bromley (1987). Nevertheless, *S. arizonensis* was still retained and it was not reallocated to any *Taenidium* ichnospecies (D'Alessandro and Bromley 1987; Keighley and Pickerill 1994). On the other hand, Buatois et al. (2017) considered *Scolecocoprus* a distinct valid ichnogenus. More detailed research is necessary to provide reliable information regarding this peculiar meniscate ichnogenus, but here we follow D'Alessandro and Bromley (1987) and consider *Scolecocoprus* as a junior synonym of *Taenidium*. The two ichnospecies *S. cameronensis* and *S. arizonensis* are herein assigned to *Taenidium serpentinum* and *Taenidium* cf. *T. serpentinum*, respectively.




Figure 8-19. Complex, actively filled horizontal structures from the Coconino Sandstone (LORENZO MARCHETTI). **A.** *Taenidium serpentinum*, holotype of *Scolecocoprus cameronensis*, convex hyporelief (MNA-N 3707). **B.** *Taenidium* cf. *serpentinum*, holotype of *Scolecocoprus arizonensis*, convex hyporelief (MNA-N 3655). **C.** *Scolecocoprus* isp. (arrow) in transition and associated with *Diplopodichnus*, convex hyporelief (GRCA-NN 1). White scale bar is 2 cm (0.8 in).

Discussion

A large number of invertebrate trace specimens, including long, complete, and beautifully preserved trackways and burrows, have been collected from the Coconino Sandstone. These specimens provide an ideal basis for ichnotaxonomy, because much extramorphological variation is registered. The effects of the trackmaker's locomotion on inclined planes and compound traces can also be recognized. Seven ichnogenera and 13 ichnospecies were erected based on Coconino Sandstone material. However, the number of valid ichnotaxa from this formation is relatively low, including six ichnogenera (*Diplichnites*, *Diplopodichnus*, *Lithographus*, *Octopodichnus*, *Palaeohelcura* and *Taenidium*). The number of architectural designs is even lower: only two types were recognized. This low-diversity ichnoassociation is less diverse and structurally complex than the typical continental ichnoassociations that are usually included in the *Scoyenia* ichnofacies. Because of this, a different ichnofacies based mostly on the Coconino ichnoassociation was proposed by Hunt and Lucas (2007), the *Octopodichnus* ichnofacies. Simultaneously, Ekdale et al. (2007) proposed the *Entradichnus* ichnofacies based mostly on traces from the Jurassic Navajo Sandstone. Buatois and Mángano (2011) unified the two ichnofacies into the *Octopodichnus–Entradichnus* ichnofacies, which includes low-diversity dune ichnoassociations with arthropod tracks and non-diverse burrows. Note that Lucas et al. (2010) and Lucas (2018) stated that the "*Entradichnus* ichnofacies" is supposedly characterized by an ichnofauna of *Arenicolites*, *Palaeophycus*, *Planolites*, *Skolithos*, and *Taenidium*, and ichnoassemblages that contain one or more of these ichnogenera are characteristic of interdunal facies in Permian–Jurassic eolianites of the American Southwest, although they can occur in what

must have been moist dunal settings. Therefore, the "*Entradaichnus* ichnofacies" is not a distinctive ichnofacies, but simply a synonym of the *Scoyenia* ichnofacies in interdunal settings (Hunt and Lucas 2016). The distinctive invertebrate ichnofacies of eolian paleoenvironments is the *Octopodichnus* ichnofacies, recognized by low diversity ichnofaunas of arthropod walking traces (*Octopodichnus*, *Paleohelcura*, etc.) and usually associated with footprints of the *Chelichnus* tetrapod ichnofacies.

GRCA Museum Collections

In July 1928, Edwin McKee collected a trackway specimen from the Coconino Sandstone within Hermit Basin that contained "worm trails" (GRCA 2883). In August 1928, a "fucoid" specimen from the base of the Coconino Sandstone with "worm trails showing body markings" was collected by McKee, but the specimen was never found (GRCA 2911; deaccessioned). In August 1931, McKee collected a trackway associated with "worm trails" from Marble Canyon (GRCA 2882). This specimen shows a transition between *Taenidium* and *Diplopodichnus*; the latter also includes transverse, closely spaced imprints. In November 1932, McKee collected invertebrate track specimens from the upper Coconino Sandstone outside of Grand Canyon in Apache County, Arizona at Kinlachee Creek (GRCA 2906 and GRCA 20652). An invertebrate track specimen, *Paleohelcura tridactyla*, was collected from the Coconino Sandstone in the Hermit Basin by an unknown collector at an unknown date; however, Gilmore (1926) named this specimen and may have been the collector (GRCA 2917). In October 1938, McKee collected a "worm trail" from the Coconino Sandstone (GRCA 2928). An invertebrate trackway identified as *Octopodichnus minor* (now regarded as *O. didactylus*), was collected from the Coconino Sandstone outside of Grand Canyon National Park near Seligman, Arizona, and is associated with Brady's collections in his 1947 report (GRCA 2930). An un-numbered specimen (GRCA-NN 1) includes several *Diplopodichnus* trails and a *Taenidium* burrow, possibly in transition with *Diplopodichnus*.

Besides specimens collected and housed in museum collections, it is worth noting the presence of invertebrate trackways and trails in situ in the Coconino Sandstone. Among them, *Paleohelcura tridactyla* and *Diplopodichnus* isp. can be recognized at the same level in which tetrapod tracks occur. *Diplopodichnus* isp. also occurs in association with *Ichniotherium sphaerodactylum* tracks in the fallen boulder of Coconino Sandstone described by Francischini et al. (2019). All these materials remain in situ.

Toroweap Formation

As noted earlier, we are not aware of any records of trace fossils from the Toroweap Formation, though we suspect that a careful search of that unit could reveal some.

Kaibab Formation

The youngest unit of Grand Canyon's Paleozoic stratigraphic section is the Kaibab Formation (Limestone). It forms the surface upon which visitors view the canyon from the rim at the top of the Kaibab and Coconino plateaus. The Kaibab Formation thickens westward from 90 to 120 m (295 to 394 ft) and sits upon the Woods Ranch Member of the Toroweap Formation in Grand Canyon (Hopkins and Thompson 2003). It consists of two conformable packages in GRCA: the Fossil Mountain Member and the overlying Harrisburg Member. The Fossil Mountain Member forms a cliff with hoodoo-like weathering that ranges from 75 to 105 m (246 to 344 ft) in thickness. It contains

cherty, fossiliferous limestone in the west and becomes siliciclastic towards the east. The Harrisburg Member forms receding ledges and cliffs towards the rim of the Grand Canyon and ranges from 25 to 90 m (82 to 295 ft) thick westward. It is composed of dolomite, gypsum, chert, sandstone, and limestone. Fossiliferous carbonate beds appear increasingly toward the west (Hopkins and Thompson 2003).

The Kaibab Formation was deposited in a subtidal, shallow-marine environment with mixed carbonates and siliciclastics affected by sea level changes. Invertebrate organisms such as brachiopods, bryozoans, mollusks, crinoids, sponges, and trilobites lived in this sea during the late Cisuralian. A diverse assemblage of selachian teeth is also known. Only a few invertebrate traces have been reported and collected.

Trace Fossils

"Fucoids, worm borings, and trilobite trails" have been reported from the Kaibab Formation (McKee 1938; Spamer 1992). McKee (1938) noted "small tubular structures which may represent worm borings" within inorganic chert layers. He also mentioned structures of algae within certain chert layers that may have made a soft substrate for worms to bore into.

GRCA Museum Collections

Specimens from the Kaibab Formation containing burrows and trails were collected by Louis Schellbach on October 7, 1941 east of Desert View (GRCA 2349, GRCA 2575, GRCA 2596, GRCA 10233, GRCA 11371, GRCA 14171, GRCA 14172). Invertebrate trackways were found in the Kaibab Formation and collected by Strickler in 1933 (GRCA 10043, GRCA 10045, and GRCA 14404). Cliff Carrol collected a marine "snail trail" from the Kaibab Formation near Yavapai Point (GRCA 10234 and GRCA 17336). In June 1936, a "worm trail" was collected by Everette Helman, found between Yaki and Grandview Points from near the top of the Kaibab Formation (GRCA 10044; Figure 8-20). Edwin McKee collected sandstone slabs covered with "fucoids" from the Kaibab Formation at Desert View (GRCA 6026 and GRCA 6598).




Figure 8-20. Worm trail specimen in the upper Kaibab Formation collected by Everette Helman in June 1936 (GRCA 10044). Scale bar is 1 cm (0.4 in) (NPS/KLARA WIDRIG).

Conclusions

The invertebrate trace fossil record of Grand Canyon is composed of trackways and traces produced in almost all geological units of this area, with the only exception being the Permian Toroweap

Formation due to a lack of research. Although the records of some units (i.e., the Frenchman Mountain Dolostone, the Redwall Limestone, the Watahomigi Formation, the Manakacha Formation and the Esplanade Sandstone) only include indeterminate or informally described traces, up to 33 valid ichnogenera occur among the Tapeats Sandstone, the Bright Angel Shale, the Muav Limestone, the Temple Butte Formation, the Surprise Canyon Formation, the Wescogame Formation, the Hermit Formation, the Coconino Sandstone, and the Kaibab Formation. The Cambrian Bright Angel Shale presents the most prolific record. With 21 ichnogenera classified in 15 architectural designs, it represents a very diverse and disparate association of trace fossils from the *Cruziana* Ichnofacies. Other marine and continental units also present reasonably high ichnodiversity and ichnodisparity: the Tapeats Sandstone, the Hermit Formation, and the Coconino Sandstone. Trace fossils often serve as the only indicators of past life in a particular geologic time interval and paleoenvironment. In the Grand Canyon region, trace fossils add significant clues to the diversity of past life and behaviors, especially within those formations that are mostly devoid of body fossils. Invertebrate trace fossils continue to be discovered throughout GRCA's Paleozoic units and can be regarded as significant, non-renewable resources for the park. Research on such specimens also continues with the help of park managers as they play an important role in protecting such invaluable resources.

Acknowledgments

Anne Miller was funded by the Grand Canyon Conservancy under a cooperative agreement between the U.S. Department of Interior, National Park Service and Northern Arizona University (Task Agreement P17AC00600 under cooperative agreement P14AC00921). Lorenzo Marchetti was funded by the Alexander von Humboldt Foundation. Heitor Francischini was funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES/PNPD; Grant Number: 88882.316166/2019-01). Spencer Lucas was funded by the New Mexico Museum of Natural History and Science.

Literature Cited

Alf, R. M. 1968. A spider trackway from the Coconino Formation, Seligman, Arizona. Southern California Academy of Sciences Bulletin 67:125–128.

Alpert, S. P. 1974. Systematic review of the genus *Skolithos*. Journal of Paleontology 48(4):661–669.

Baldwin, C. T., P. K. Strother, J. H. Beck, and E. Rose. 2004. Palaeoecology of the Bright Angel Shale in the eastern Grand Canyon, Arizona, USA, incorporating sedimentological, ichnological and palynological data. Pages 213–236 *in* D. McIlroy, editor. The application of ichnology to palaeoenvironmental and stratigraphic analysis. Geological Society of London, London, United Kingdom. Special Publications 228.

Bertling, M., S. J. Braddy, R. G. Bromely, G. R. Demathieu, J. Genise, R. Mikulaš, J. K. Nielsen, K. S. S. Kielsen, A. K. Rindsberg, M. Schlirf, and A. Uchman. 2006. Names for trace fossils: A uniform approach. Lethaia 39:265–286.

Beus, S. S. 2003a. Temple Butte Formation. Pages 107–114 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Beus, S. S. 2003b. Redwall Limestone and Surprise Canyon Formation. Pages 115–135 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Blakey, R. C. 2003. Supai Group and Hermit Formation. Pages 136–162 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Blakey, R. C., and R. Knepp. 1989. Pennsylvanian and Permian geology of Arizona. Pages 313–347 *in* J. P. Jenney and S. J. Reynolds, editors. Geologic evolution of Arizona. Arizona Geological Society Digest 17.

Bonde, A., V. L. Santucci, J. S. Tweet, E. Eichenberg, and B. Moore. 2018. Lake Mead National Recreation Area: Paleontological resources inventory. Natural Resource Report NPS/LAKE/NRR—2018/1618. National Park Service, Fort Collins, Colorado.

Braddy, S. J. 1995. A new arthropod trackway and associated invertebrate ichnofauna from the Lower Permian Hueco Formation of the Robledo Mountains, southern New Mexico. New Mexico Museum of Natural History and Science Bulletin 6:101–105.

Brady, L. F. 1939. Tracks in the Coconino Sandstone compared with those of small living arthropods. Plateau 12:32–34.

Brady, L. F. 1947. Invertebrate tracks from the Coconino Sandstone of northern Arizona. Journal of Paleontology 32:466–472.

Brady, L. F. 1961. A new species of *Paleohelcura* Gilmore from the Permian of northern Arizona. Journal of Paleontology 35:201.

Brathovde, J. E. 1986. Stratigraphy of the Grand Wash Dolomite (Upper? Cambrian), Western Grand Canyon, Mohave County, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Buatois, L. A. 2018. *Treptichnus pedum* and the Ediacaran–Cambrian boundary: Significance and caveats. Geological Magazine 155(1):174–180.

Buatois, L. A., and M. G. Mángano. 2011. Ichnology: Organism-substrate interactions in space and time. Cambridge University Press, Cambridge, United Kingdom.

Buatois, L. A., M. Wisshak, M. A. Wilson, and M. G. Mángano. 2017. Categories of architectural designs in trace fossils: A measure of ichnodisparity. Earth-Science Reviews 164:102–181.

Crimes, T. P., I. Legg, A. Marcos, and M. Arboleya. 1977. ?Late Precambrian–low Lower Cambrian trace fossils from Spain. Pages 91–138 *in* T. P. Crimes and J. C. Harper, editors. Trace Fossils 2: Proceedings from the 25th International Geological Conference. Geological Journal Special Issue 9.

D'Alessandro, A., and R. G. Bromley. 1987. Meniscate trace fossils and the *Muensteria-Taenidium* problem. Palaeontology 30:743–763.

Davies, N. S., A. G. Liu, M. R. Gibling, and R. F. Miller. 2016. Resolving MISS conceptions and misconceptions: A geological approach to sedimentary surface textures generated by microbial and abiotic processes. Earth-Science Reviews 154:210–246.

Davis, R. B., N. J. Minter, and S. J. Braddy. 2007. The neoichnology of terrestrial arthropods. Palaeogeography, Palaeoclimatology, Palaeoecology 255(3–4):284–307.

DeCourten, F. L. 1978. *Scolecocoprus cameronensis* Brady (1947) from the Kaibab Limestone of northern Arizona; a re-interpretation. Journal of Paleontology 52(2):491–493.

Díez-Canseco, D., L. A. Buatois, M. G. Mángano, M. Díaz-Molina, and M. I. Benito. 2016. Ichnofauna from coastal meandering channel systems (Upper Cretaceous Tremp Formation, South-Central Pyreneeds, Spain): Delineating the fluvial-tidal transition. Journal of Paleontology 90(2):250–268.

Ekdale, A. A., R. G. Bromley, and D. B. Loope. 2007. Ichnofacies of an ancient erg: A climatically influenced trace fossil association in the Jurassic Navajo Sandstone, southern Utah, USA. Pages 562–574 *in* W. Miller, editor. Trace fossils, concepts, problems, prospects. Elsevier, Amsterdam, Netherlands.

Elliott D. K., and R. W. Bartlett. 2013. A new trace fossil from the Permian Coconino Sandstone of Northern Arizona. New Mexico Museum of Natural History and Science Bulletin 60:79–80.

Elliott, D. K., and D. L. Martin. 1987. A new trace fossil from the Cambrian Bright Angel Shale, Grand Canyon, Arizona. Journal of Paleontology 61(4):641–648.

Francischini, H., S. G. Lucas, S. Voight, L. Marchetti, V. L. Santucci, C. L. Knight, J. R. Wood, P. Dentzien-Dias, and C. L. Schultz. 2019. On the presence of *Ichniotherium* in the Coconino Sandstone (Cisuralian) of the Grand Canyon and remarks on the occupation of deserts by non-amniote tetrapods. Paläontologische Zeitschrift online:1–19.

Frech, F. 1893. Section in Congress Canyon opposite Point Sublime. Pages 476–481 *in* Compte rendu. International Geologic Congress [Congres geologique international], 5th session. Government Printing Office, Washington, D.C.

Gilmore, C. W. 1926. Fossil footprints from the Grand Canyon. Smithsonian Miscellaneous Collections 77(9).

Gilmore, C. W. 1927. Fossil footprints from the Grand Canyon: Second contribution. Smithsonian Miscellaneous Collections 80(3).

Gilmore, C. W. 1928. Fossil footprints from the Grand Canyon: Third contribution. Smithsonian Miscellaneous Collections 80(8).

Hagadorn, J. W., J. L. Kirschvink, T. D. Raub, and E. C. Rose. 2011. Above the Great Unconformity: A fresh look at the Tapeats Sandstone, Arizona-Nevada, U.S.A. Museum of Northern Arizona Bulletin 67:63–77.

Häntzschel, W. 1975. Trace fossils and Problematica. Treatise on Invertebrate Paleontology, Part W: Miscellanea, Supplement 1. R. C. Moore, editor. Geological Society of America, Boulder, Colorado and University of Kansas Press, Lawrence, Kansas.

Hereford, R. 1977. Deposition of the Tapeats Sandstone (Cambrian) in central Arizona. Bulletin of the Geological Society of America 88:199–211.

Hopkins, R. L., and K. L. Thompson. 2003. Kaibab Formation. Pages 196–211 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Howell, B. F. 1957. *Stipsellus annulatus*, a *Skolithos*-like Cambrian fossil from Arizona. Bulletin of the Wagner Free Institute of Sciences of Philadelphia 32(2):7–20.

Hunt, A. P., and S. G. Lucas. 1998. Ichnological evidence for tetrapod predation in the Paleozoic: Is there any? New Mexico Museum of Natural History and Science Bulletin 12:59–62.

Hunt, A. P., and S. G. Lucas. 2007. Tetrapod ichnofacies: A new paradigm. Ichnos 14:59–68.

Hunt, A. P., and S. G. Lucas. 2016. The case for archetypal vertebrate ichnofacies. Ichnos 23(3–4):237–247.

International Commission on Zoological Nomenclature. 1999. International Code of Zoological Nomenclature (4th edition). International Trust for Zoological Nomenclature, London, United Kingdom.

Karlstrom, K., J. Hagadorn, G. Gehrels, W. Matthews, M. Schmitz, L. Madronich, J. Mulder, M. Pecha, D. Giesler, and L. Crossey. 2018. Cambrian Sauk transgression in the Grand Canyon region redefined by detrital zircons. Nature Geoscience 11:438–443.

Karlstrom, K. E., M. T. Mohr, M. D. Schmitz, F. A. Sundberg, S. M. Rowland, R. Blakey, J. R. Foster, L. J. Crossey, C. M. Dehler, and J. W. Hagadorn. 2020. Redefining the Tonto Group of Grand Canyon and recalibrating the Cambrian timescale. Geology in press. doi: https://doi.org/10.1130/G46755.1.

Keighley, D. G., and R. K. Pickerill. 1994. The ichnogenus *Beaconites* and its distinction from *Ancorichnus* and *Taenidium*. Palaeontology 37(2):305–337.

Knaust, D. 2013. The ichnogenus *Rhizocorallium*: Classification, trace makers, palaeoenvironments and evolution. Earth-Science Reviews 126:1–47.

Knaust, D. 2018. The ichnogenus *Teichichnus* Seilacher, 1955. Earth-Science Reviews 177:386–403.

Kozur, H., M. Löffler, and E. Sittig. 1994. First evidence of *Paleohelcura* (arthropod trackway) in the Rotliegend of Europe. Neues Jarhbuch für Geologie und Paläontologie, Monatshefte, 10:618–632.

Kramer, J. M., B. R. Erickson, M. G. Lockley, A. P. Hunt, and S. Braddy. 1995. Pelycosaur predation in the Permian: Evidence from *Laoporus* trackways from the Coconino Sandstone with description of a new species of *Permichnium*. New Mexico Museum of Natural History and Science Bulletin 6:245–249.

Lane, A. A., S. J. Braddy, D. E. G. Briggs, and D. K. Elliott. 2003. A new trace fossil from the Middle Cambrian of the Grand Canyon, Arizona, USA. Palaeontology 46:987–997.

Lucas, S. G. 2016. Two new, substrate controlled nonmarine ichnofacies. Ichnos 23(3–4):248–261.

Lucas, S. G. 2018. Re-evaluation of the original ichnoassemblage of the "*Entradaichnus* ichnofacies," Jurassic of Utah. Geological Society of America Abstracts with Programs 50(6): paper 259-3.

Lucas, S. G., J. A. Spielmann, H. Klein, and A. J. Lerner. 2010. Ichnology of the Upper Triassic (Apachean) Redonda Formation, east-central New Mexico. New Mexico Museum of Natural History and Science Bulletin 47.

Lucas, S. G., S. Voigt, A. J. Lerner, and W. J. Nelson. 2011. Late Early Permian continental ichnofauna from Lake Kemp, north-central Texas, USA. Palaeogeography, Palaeoclimatology, Palaeoecology 308:395–404.

Lucas, S. G., S. Voigt, A. J. Lerner, and E. C. Rainforth. 2013. *Sphaerapus*, a poorly known invertebrate trace fossil from nonmarine Permian and Jurassic strata of North America. Ichnos 20:142–152.

MacEachern, J. A., S. G. Pemberton, M. K. Gingras, and K. L. Bann. 2010. Ichnology and facies models. Pages 19–58 *in* N. P. James and R. W. Dalrymple, editors. Facies Models 4. Geological Association of Canada, St. John's, Canada.

Mángano, M. G., L. A. Buatois, C. G. Maples, and R. R. West. 2000. A new ichnospecies of *Nereites* from Carboniferous tidal-flat facies of eastern Kansas, USA: Implications for the *Nereites-Neonereites* debate. Journal of Paleontology 74(1):149–157.

Marchetti, L., M. Belvedere, S. Voigt, H. Klein, D. Castanera, I. Díaz-Martínez, D. Marty, L. Xing, S. Feola, R. N. Melchor, and J. O. Farlow. 2019a. Defining the morphological quality of fossil footprints: Problems and principles of preservation in tetrapod ichnology with examples from the Palaeozoic to the present. Earth-Science Reviews 193:109–145

Marchetti, L., S. Voigt, S. G. Lucas, H. Francischini, P. Dentzien-Dias, R. Sacchi, M. Mangiacotti, A. Gazzola, A. Ronchi, and A. Millhouse. 2019b. Tetrapod ichnotaxonomy in eolian

paleoenvironments (Coconino and De Chelly formations, Arizona) and late Cisuralian sauropsid radiation. Earth-Science Reviews 190:148–170.

Martin, D. L. 1985. Depositional systems and ichnology of the Bright Angel Shale (Cambrian), eastern Grand Canyon, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Martinsson, A. 1970. Toponomy of trace fossils. Pages 323–330 *in* T. P. Crimes and J. C. Harper, editors. Trace fossils. Geological Journal Special Issue 3.

MacEachern, J. A. K. L. Bann, M. K. Gingras, J.-P. Zonneveld, S. E. Dashtgrad, and S. G. Pemberton. 2012. The ichnofacies paradigm. Developments in Sedimentology 64:103–138.

MacEachern, J. A., S. G. Pemberton, M. K. Gingras, and K. L. Bann. 2010. Ichnology and facies models. Pages 19–58 *in* N. P. James and R. W. Dalrymple, editors. Facies Models 4. Geological Association of Canada, St. John's, Canada.

McKee, E. D. 1933. The Coconino Sandstone—its history and origin. Carnegie Institution of Washington Publication 440:77–115.

McKee, E. D. 1938. The environment and history of the Toroweap and Kaibab formations of northern Arizona and southern Utah. Carnegie Institution of Washington Publication 492.

McKee, E. D. 1944. Tracks that go uphill. Plateau 16:61–72.

McKee, E. D., and C. E. Resser. 1945. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

McKee, E. D. 1979. Ancient sandstones considered to be eolian. Pages 187–238 *in* E. D. McKee, editor. A study of global sand seas. U.S. Geological Survey, Washington, D.C. Professional Paper 1052.

McKee, E. D. 1982. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D., and R. C. Gutschick. 1969. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Middleton, L. T. 1989. Cambrian and Ordovician depositional systems in Arizona. Pages 273–286 *in* J. P. Jenney and S. J. Reynolds, editors. Geologic evolution of Arizona. Arizona Geological Society Digest 17.

Middleton, L. T., and D. K. Elliott. 2003. Tonto Group. Pages 90–106 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Middleton, L. T., D. K. Elliott, and M. Morales. 1990. Coconino Sandstone. Pages 352–368 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology. Oxford University Press, New York, New York.

Minter, N. J., and S. J. Braddy. 2009. Ichnology of an Early Permian intertidal flat: The Robledo Mountains Formation of southern New Mexico, USA. The Palaeontological Association, Special Papers in Palaeontology 82.

Minter, N. J., S. J. Braddy, and R. B. Davis. 2007. Between a rock and a hard place: Arthropod trackways and ichnotaxonomy. Lethaia 40:365–375.

Noble, L. F. 1922. A section of the Paleozoic formations of the Grand Canyon at Bass Trail. Pages 23–73 *in* Shorter contributions to general geology, 1922. U.S. Geological Survey, Washington, D.C. Professional Paper 131-B.

Osgood, R. G. 1975. The history of invertebrate ichnology. Pages 3–12 *in* R. W. Frey, editor. The study of trace fossils: A synthesis of principles, problems and procedures in ichnology. Springer-Verlag, New York, New York.

Pemberton, S. G., and R. W. Frey. 1982. Trace fossil nomenclature and the *Planolites-Paleophycus* dilemma. Journal of Paleontology 56:843–871.

Retallack, G. J. 2013. Ediacaran life on land. Nature Research Letters 493:89–92.

Rose, E. C. 2003. Depositional Environment and History of the Cambrian Tonto Group, Grand Canyon, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Rose, E. C. 2006. Nonmarine aspects of the Cambrian Tonto Group of the Grand Canyon, USA, and broader implications. Palaeoworld 15:223–241.

Sadler, C. J. 1993. Arthropod trace fossils from the Permian De Chelly sandstone, northeastern Arizona. Journal of Paleontology 67(2):240–249.

Schenk, E. T., and H. E. Wheeler. 1942. Cambrian sequence in western Grand Canyon, Arizona. Journal of Geology 50(7):822–899.

Schuchert, C. 1918. On the Carboniferous of the Grand Canyon of Arizona. American Journal of Science (4^{th} series) 45(269):347–361.

Seilacher, A. 1964. Sedimentological classification and nomenclature of trace fossils. Sedimentology 3:253–256

Seilacher, A. 1970. *Cruziana* stratigraphy of non-fossiliferous Paleozoic sandstones. Pages 447–476 *in* T. P. Crimes and J. C. Harper, editors. Trace fossils. Geological Journal Special Issue 3.

Seilacher, A. 2007. Trace fossil analysis. Springer, Berlin, Germany.

Smith, J. 1909. Upland fauna of the Old Red Sandstone of Carrick, Ayrshire. A. W. Cross, Kilwinning, United Kingdom.

Spamer, E. E. 1984. Paleontology in the Grand Canyon of Arizona: 125 years of lessons and enigmas from the late Precambrian to the present. The Mosasaur 2:45–128.

Spamer, E. E. 1992. The Grand Canyon fossil record: A source book in paleontology of the Grand Canyon and vicinity, northwestern Arizona and southeastern Nevada. Geological Society of America, Boulder, Colorado. Microform Publication 24.

Stoyanow, A. A. 1936. Correlation of Arizona Paleozoic formations. Geological Society of America Bulletin 37:459–540.

Strother, P. K., and J. H. Beck. 2000. Spore-like microfossils from Middle Cambrian strata: expanding the meaning of the term cryptospore. Pages 413–424 *in* M. M. Harley, C. M. Morton, and S. Blackmore, editors. Pollen and spores: Morphology and biology. Royal Botanic Gardens, Kew, United Kingdom.

Uchman, A. 1995. Taxonomy and paleoecology of flysch trace fossils: The Marnoso-arenacea Formation and associated facies (Miocene, Northern Apennines, Italy). Beringeria 15:1–115.

Vallon, L. H., A. K. Rindsberg, and R. G. Bromley. 2016. An updated classification of animal behavior preserved in substrates. Geodinamica Acta 28(1–2):5–20.

Walker, E. F. 1985. Arthropod ichnofacies of the Old Red Sandstone at Dunure and Montrose, Scotland. Transactions of the Royal Society of Edinburgh 76:287–297.

White, C. A. 1874. Preliminary report upon invertebrate fossils collected by the expeditions of 1871, 1872, and 1873, with descriptions of new species. *In* U.S. Army Engineer Department, Geographical and geological explorations and surveys west of the one hundredth meridian. U.S. Government Printing Office, Washington, D.C.

White, C. D. 1929. Flora of the Hermit Shale, Grand Canyon, Arizona. Carnegie Institution of Washington Publication 405.

Chapter 9. Paleozoic Vertebrate Ichnology of Grand Canyon National Park

By Lorenzo Marchetti[1], Heitor Francischini[2], Spencer G. Lucas[3], Sebastian Voigt[1], Adrian P. Hunt[4], and Vincent L. Santucci[5]

[1]Urweltmuseum GEOSKOP / Burg Lichtenberg (Pfalz)
Burgstraße 19
D-66871 Thallichtenberg, Germany

[2]Universidade Federal do Rio Grande do Sul (UFRGS)
Laboratório de Paleontologia de Vertebrados and Programa de Pós-Graduação em Geociências, Instituto de Geociências
Porto Alegre, Rio Grande do Sul, Brazil

[3]New Mexico Museum of Natural History and Science
1801 Mountain Road N.W.
Albuquerque, New Mexico, 87104

[4]Flying Heritage and Combat Armor Museum
3407 109th St SW
Everett, Washington 98204

[5]National Park Service
Geologic Resources Division
1849 “C” Street, NW
Washington, D.C. 20240

Introduction

Vertebrate tracks are the only fossils of terrestrial vertebrates known from Paleozoic strata of Grand Canyon National Park (GRCA), therefore they are of great importance for the reconstruction of the extinct faunas of this area. For more than 100 years, the upper Paleozoic strata of the Grand Canyon yielded a noteworthy vertebrate track collection, in terms of abundance, completeness and quality of preservation. These are key requirements for a classification of tracks through ichnotaxonomy. This chapter proposes a complete ichnotaxonomic revision of the track collections from GRCA and is also based on a large amount of new material. These Paleozoic tracks were produced by different tetrapod groups, such as eureptiles, parareptiles, synapsids and anamniotes, and their size ranges from 0.5 to 20 cm (0.2 to 7.9 in) footprint length. As the result of the irreversibility of the evolutionary process, they provide useful information about faunal composition, faunal events, paleobiogeographic distribution and biostratigraphy. Of note, these vertebrate trace fossils are always in situ (i.e., not transported before burial) and preserved in lithofacies representing different continental paleoenvironments (marginal marine, alluvial plain and desert), therefore they can provide useful paleoecological information. Also, the occurrence of surfaces with long trackways provides

fundamental insights into the locomotion of the trackmakers on inclined planes, such as the paleosurfaces of dune foresets.

Vertebrate Ichnology

Vertebrate ichnology is the discipline that principally studies the traces left on the substrate by vertebrates during their lives. These traces can be produced by different behaviors, such as: walking, running, hopping, resting, swimming, etc. The main subject of vertebrate ichnology is thus the study of tracks left on the sediment during locomotion. The tracks can be preserved as concave epirelief (natural mold) and convex hyporelief (natural cast) and can be preserved on multiple layers, forming undertracks, underprints and overtracks. On the actual trampled surface, these ichnofossils sometimes preserve expulsion rims (i.e., a marginal rim of displaced sediment produced by the footfall; Allen 1997). Their occurrence generally indicates subaerial exposure, the presence of a substrate adequate to record the footprint impression, and a non-erosive and rapid burial of the actual track surface. These conditions are common in intermittently wet environments such as lake margins, alluvial plains, tidal flats, fluvial channels, and/or in desert environments characterized by rapid sedimentary cover (dune foreset surfaces). In intermittently wet environments, they are often associated with mud cracks, rain drops, wave ripples, tool marks, and microbial structures, among other sedimentary structures, and with invertebrate tracks and burrows. In desert environments, they are often associated with sand avalanches, wind ripple crests and invertebrate tracks and burrows.

With regard to the Paleozoic, only footprints of quadrupedal vertebrates are known, which include front (manus) and hind (pes) foot imprints. Tail, body, and digit drag marks may be present, but are not relevant for footprint classification. These footprints are usually arranged in sequences of pes-manus couples, arranged in two parallel rows that form a trackway. From the footprint and trackway parameters, it is possible to infer the size, weight, gait and speed of the trackmakers (e.g., Leonardi 1987). Generally, the Paleozoic terrestrial vertebrates were not fast moving, and most of them adopted a sprawling gait as seen in modern lizards and salamanders. Their size was also not very large, especially before the Guadalupian (maximum footprint length of about 20 cm or 7.9 in).

Tetrapod footprints are classified by their morphology, and especially by the anatomy-consistent morphology (e.g., Marchetti et al. 2019a). This is the morphology that reflects the actual lower side of the trackmaker foot, and thus it is strongly connected with the trackmaker's anatomical structure, which is subject to irreversible evolution. Tracks showing these anatomy-consistent features are regarded as well-preserved. This kind of preservation, named morphological preservation, can be evaluated through a numerical scale (Marchetti et al. 2019a). In the study of tetrapod tracks, it is necessary to exclude from the analysis those tracks that are not anatomy-consistent due to loss of information or deformation during (e.g., footprint wall collapse) and after track registration in the sediment (e.g., superimposition of other footprints and sedimentary structures and partial erosion). The footprints preserved on the dune foreset surfaces show a peculiar process that causes loss of information of the anatomy: the oblique direction of the gravity force due to the dune surface inclination (usually more than 30°). This causes the sliding of the trackmaker digits during locomotion, and asymmetrical trackway patterns and footprint preservation in trackways oblique to the slope (Figure 9-1A–D). Also, the trackmaker speed was influenced, with very close pes-manus

couples and secondary overstep in trackways directed upslope (slow gait) and well-spaced pes-manus couples and primary overstep (fast gait) in trackways directed downslope (Figure 9-1A–D). Overstep is the preservation of the pes in front of the manus, different than what is usually seen (manus in front of the pes). This has been observed in trackways of the same track type (*Varanopus*), therefore these different gaits and morphologies are likely not due to different trackmakers (Marchetti et al. 2019b, 2019c).




Figure 9-1. Track formation and track measurements (LORENZO MARCHETTI & SPENCER LUCAS). **A.** Track formation on a dune foreset surface with different kind of progression: **B.** directly upslope, **C.** directly downslope and **D.** transversely upslope. **E.** Track and trackway measurements. FL=footprint length, FW=footprint width, psL=palm/sole length, psW=palm/sole width, DL=digit length, div=digit divergence, V=digit V, SLp=stride length of the pes, SLm=stride length of the manus, PLp=(oblique) pace length of the pes, PAp=pace angulation of the pes, PAm=pace angulation of the manus, PLm=(oblique) pace length of the manus, LPp=length of pace of the pes, LPm=length of pace of the manus, WPp=width of pace of the pes, WPm=width of pace of the manus, Divp=divarication from midline of the pes, Divm=divarication from midline of the manus, Dmp=manus-pes distance, BL=gleno-acetabular length.

After the selection of well-preserved material, the footprints and trackways are described and measured. The digit impressions are counted with Roman numbers, starting from the inner (medial) side of the trackway. Tracks with longer external digit impressions (commonly digit IV) are named ectaxonic. Common track measurements (Figure 9-1E) include the foot length and width, and the

sole/palm impression length and width; all compared to the digit III axis. Other measurements are the digit length and the interdigital angles. The distance between two consecutive tracks of the same kind (manus or pes) on the same trackway side is named stride length. The distance between two consecutive tracks of the same kind (manus or pes) on opposite trackway sides is termed (oblique) pace length, and the angle between two consecutive (oblique) pace lengths is termed pace angulation. The line equidistant from the manus-pes couples is the trackway midline. Other measurements are the trackway width, the manus-pes distance and the manus and pes divarication (i.e., its rotation), all compared to the trackway midline.

Ichnotaxonomy is the discipline that classifies tetrapod tracks and other traces. It is considered a parataxonomy because it is disentangled from the taxonomic classification of the producers. In fact, it is quite difficult to relate a track type directly to a trackmaker genus or species, with some exceptions (e.g., Voigt et al. 2007; Marchetti et al. 2017a). Tetrapod track ichnotaxa usually correspond to higher taxonomic levels than species and genera, because similar species and genera may produce very similar footprints (e.g., all of the "pelycosaur" synapsid genera, except for varanopids, may be attributed to a single ichnogenus, *Dimetropus*). The tetrapod tracks are classified with a binomial nomenclature (ichnogeneric epithet + ichnospecific epithet), following the procedures of the International Code of Zoological Nomenclature (1999). Ichnofamilies can be used as well, but are generally less extensively studied. The relevant criteria (ichnotaxobases) are the anatomy-consistent foot morphology and the trackway pattern, including measurements. The list of ichnotaxa from a formation and/or area is referred to as an ichnoassociation and helps to define the ichnodiversity. The ichnoassociations can be time-constrained, because ichnotaxa have limited stratigraphic distributions. Also, being geographically widespread, they are useful to define a tetrapod footprint biostratigraphy (e.g., Lucas 2007; Fillmore et al. 2012; Voigt and Lucas 2018). Other useful information that can be inferred from ichnoassociations is paleobiogeography and paleoecology. Recurrent tetrapod ichnoassociations in space and time, related to specific paleoenvironments, are named tetrapod ichnofacies (Hunt and Lucas 2006, 2007), similar to invertebrate ichnofacies. Although this approach has been criticized and must certainly be improved, the paleoecological signature of tetrapod ichnoassociations is evident from several studies (e.g., Voigt and Lucas 2017; Marchetti et al. 2017b).

History of Track Collection in Grand Canyon and Mogollon Rim Areas

The first discoveries of Paleozoic tetrapod footprints in the Grand Canyon were by Charles Schuchert in 1915 along the Hermit Trail (Coconino Sandstone and Hermit Shale), with material being stored at Yale University and described by Lull (1918). The most important excavations were led by John C. Merriam in 1924 and Charles W. Gilmore in 1926 and 1927 along the Hermit and Yaki trails. This material comes from the Coconino Sandstone, the Hermit Shale and the Wescogame Formation and is stored at the Smithsonian Institution's National Museum of Natural History (Spamer 1984). It was first described by Gilmore (1927, 1928) and later revised by several studies (e.g., Baird 1952, Haubold 1971; Haubold et al. 1995a, 1995b; Voigt 2005; Francischini et al. 2019; Marchetti et al. 2019b). Further finds from the Coconino Sandstone of the Grand Canyon were described by Brand (1979), Hunt and Santucci (2001), Francischini et al. (2019) and Marchetti et al. (2019b), including either specimens in the field and specimens stored in the GRCA, Smithsonian and Yale collections.

Some historic photos of in situ specimens are also stored in the GRCA collection (Figure 9-2). A slab bearing footprints, possibly coming from the Pennsylvanian Manakacha Formation (Supai Group), was reported by Rowland (2017). Other important finds come from the Coconino Sandstone of the Mogollon Rim area, and are stored at the University of California at Berkeley, the Museum of Northern Arizona and the Raymond Alf Museum of Paleontology (e.g., Brand and Tang 1991; Lockley 1992; Loope 1992; Haubold et al. 1995b; Francischini et al. 2019; Marchetti et al. 2019b). A significant part of the GRCA park museum collection, including undescribed tracks from the Wescogame Formation, Hermit Shale and Coconino Sandstone, is described for the first time in this chapter. This also includes the first description of a specimen in the field found by Thomas Martens along the Bright Angel Trail, and of tetrapod tracks from the Hermit Shale of the Mogollon Rim area.




Figure 9-2. Historic photos of in situ vertebrate tracks from the Coconino Sandstone at GRCA. The original photos are in the collection of the GRCA museum. **A.** Surface above a cliff showing a clear trackway. **B.** Surface with poorly preserved tracks, probably arranged in trackways. Hammer for scale. **C.** Surface on a fallen slab showing a clear trackway with a turn.

Stratigraphic Distribution of Tracks

The tetrapod tracks from the Grand Canyon come from four different stratigraphic units, dated between the late Carboniferous (Pennsylvanian) and the early Permian (Cisuralian). These units are, from the oldest to the youngest: the Manakacha Formation, the Wescogame Formation, the Hermit Shale and the Coconino Sandstone (Figures 9-3 and 9-4). The tetrapod ichnoassociations will be described and discussed separately for these four units, in stratigraphic order. Moreover, some specimens from the same units found in the Mogollon Rim area will be also described for completeness. All the ichnotaxa occurring in each unit will be described separately based on the material from that unit, and no diagnoses are provided. Lists with all the relevant specimens of the collections are also provided.




Figure 9-3. Localities and geological setting (LORENZO MARCHETTI & SPENCER LUCAS). **A.** Simplified geologic map and localities. 1) Grand Canyon, Hermit Trail/Dripping Springs Trail. 2) Grand Canyon, Bright Angel Trail. 3) Grand Canyon, Yaki Trail/South Kaibab Trail. 4) Grand Canyon, Tanner Trail. 5) Soap Creek. 6) Peach Springs. 7) Seligman. 8) Picacho Butte. 9) Ash Fork. 10) Williams. 11) Limestone Canyon Mouth. 12) Perkinsville. 13) Sycamore Pass/Sedona. 14) Carrizo Creek. **B.** Chronostratigraphic scheme of the fossiliferous area (modified from Blakey and Knepp 1989). GL=Guadalupian. Units including tetrapod footprints are indicated by a track outline.

Institutional Abbreviations—**GRCA**, Grand Canyon National Park Museum, Grand Canyon, Arizona; **MNA**, Museum of Northern Arizona, Flagstaff, Arizona; **NMMNH**, New Mexico Museum of Natural History and Science, Albuquerque, New Mexico; **RAM**, Raymond M. Alf Museum of Paleontology, Claremont, California; **UCMP**, University of California, Berkeley; **USNM**, Smithsonian, National Museum of Natural History, Washington, D.C.; **YPM**, Yale Peabody Museum of Natural History, New Haven, Connecticut.




Figure 9-4. Permian stratigraphy of the Grand Canyon. Panoramic view looking west at the wall of Hermit Canyon showing the track-bearing Permian units (LORENZO MARCHETTI & SPENCER LUCAS).

Manakacha Formation

Geology

In the Grand Canyon, the Supai Group consists of five formations (in ascending order of oldest to youngest): Watahomigi, Manakacha, Wescogame, Pakoon and Esplanade Sandstone formations (e. g. McKee 1982; Blakey and Knepp 1989; Blakey 1990). The unconformity-bounded Manakacha Formation is about 90 m (300 ft) thick and mostly interbedded quartzose sandstone and red-bed mudstone. Deposition was on a coastal plain, and Blakey (1990) has identified some of the cross-bedded sandstones of the Manakacha Formation as eolianites. Invertebrate fossils from the Supai Group (principally fusulinids and brachiopods) indicate it is mostly of Pennsylvanian age, and, based on these fossils, McKee (1982) assigned an Atokan (late Bashkirian–early Moscovian) age to the Manakacha Formation.

Previous Studies

The only mention of tetrapod footprints from the Manakacha Formation is the recent work of Rowland (2017) and Rowland and Caputo (2018), who reported a fallen slab along the Bright Angel Trail with a tetrapod trackway showing an unusual gait and assigned by them to *Chelichnus* isp. If the stratigraphic interpretation is correct, this is the oldest known tetrapod trackway from GRCA.

Trace Fossils

Undetermined Tracks

The slab reported by Rowland (2017) and Rowland and Caputo (2018) displays a trackway of a quadruped with closely spaced footprints showing all the characteristics of tetrapod locomotion on inclined planes (Marchetti et al. 2019b) (Figure 9-5). The digit impressions are all elongated in the same direction, meaning a trackmaker sliding due to gravity. The tracks on the right side of the trackway are more deeply impressed than those on the left side. According to the slope direction inferred from the digit slide and expulsion rim directions, the trackmaker proceeded upslope with a transverse component, and this caused the evident asymmetry among the two trackway sides. The closely spaced tracks show a complete secondary overstep, which is typical of very slow locomotion. Due to the absence of clear and complete footprints showing all the digit impressions, this material is not classifiable (grade 1.0 of the preservation scale, Marchetti et al. 2019a) but it certainly represents a small-sized tetrapod walking upslope on an inclined plane such as a dune foreset surface.




Figure 9-5. Tetrapod footprints from the Manakacha Formation (LORENZO MARCHETTI & SPENCER LUCAS). **A.** Fallen blocks including a natural cast of the trackway slab (convex hyporelief). **B.** The trackway slab preserved as a natural mold (concave epirelief). Strike and dip symbols indicate the supposed dip of the inclined bedding plane, dashed arrows indicate the direction of progression. p=pes imprint. m=manus imprint. The scale bar in B is 1 m (3 ft) long.

Wescogame Formation

Geology

The Wescogame Formation, the middle unit of the Supai Group, is about 30 to 60 m (100 to 200 ft) thick in the Grand Canyon and is lithologically similar to the unconformably underlying Manakacha Formation—mostly quartzose sandstone and red-bed mudstone (McKee 1982; Blakey and Knepp 1989; Blakey 1990). McKee (1982) assigned a Late Pennsylvanian (Virgilian/early Ghzelian) age to the Wescogame Formation (see above). Like the Manakacha Formation, it was deposited on a coastal plain by both fluvial and eolian processes (Blakey 1990).

Previous Studies

A few tetrapod track occurrences have been reported from the middle part of the Supai Group (now known as the Wescogame Formation). Gilmore (1927) erected three new ichnotaxa basing on poorly preserved material (grade 1.0–1.5 of the preservation scale, Marchetti et al. 2019a): *Anomalopus sturdevanti*, *Stenichnus yakiensis* and *Tridentichnus supaiensis*. Haubold (1971) considered *Tridentichnus* a junior subjective synonym of *Anomalopus* and proposed the new combination *Anomalopus supaiensis*. Gilmore (1928) erected a further ichnotaxon for similarly preserved material: *Ammobatrachus turbatans*. Haubold (1971) considered *Ammobatrachus turbatans* as a junior subjective synonym of *Anomalopus supaiensis*. Marchetti et al. (2019b) considered *Ammobatrachus turbatans* to be a *nomen dubium*.

Trace Fossils

Anamniote Tracks

The specimen USNM 11534 (Figure 9-6A–B) shows a trackway composed of four consecutive pes-manus couples that can be tentatively assigned to cf. *Amphisauropus* isp. The pes and manus imprints are pentadactyl and ectaxonic with rounded digit terminations. The manus track is about half the size of the pes track (pes length 60 mm or 2.4 in). The sole imprint is elliptical in shape. The pace angulation is quite low (65–80°). The digit drag marks and trackway asymmetry indicate locomotion on an inclined paleosurface (dune foreset). Due to poor preservation caused by digit sliding, the assignment is only tentative. This specimen is the holotype of *Ammobatrachus turbatans* Gilmore 1928, which is considered a *nomen dubium* (Marchetti et al. 2019b). *Amphisauropus* is generally regarded as the track of seymouriamorph anamniotes (Voigt 2005; Marchetti et al. 2017a).

The specimen USNM 11691 (Figure 9-6C–D) shows a trackway with 10 closely spaced consecutive pes-manus couples that can be tentatively assigned to cf. *Limnopus* isp. The tracks are plantigrade and ectaxonic. Only three digit impressions of the pentadactyl pes are clearly recognizable, and the manus imprint is apparently tetradactyl and smaller than the pes imprint (pes length is about 40 to 50 mm or 1.6 to 2.0 in). Digit impressions have rounded terminations. The pace angulation is very low (50–70°), and the trackway shows secondary overstep. These tracks are evidently deformed by digit sliding caused by upslope progression on an inclined paleosurface (dune foreset), so the assignment is tentative. This is the holotype of the ichnotaxon *Tridentichnus supaiensis* Gilmore 1927, which is herein considered a *nomen dubium* because of the poor preservation. *Limnopus* is generally regarded as the track of relatively large temnospondyls, such as eryopids (e.g., Voigt 2005).




Figure 9-6. Anamniote footprints from the Wescogame Formation (LORENZO MARCHETTI & SPENCER LUCAS). **A.** USNM 11534. cf. *Amphisauropus* isp., trackway, concave epirelief. Holotype of *Ammobatrachus turbatans* Gilmore 1928, *nomen dubium*. **B.** Enlargement of A, right pes–manus couple. **C.** USNM 11691. cf. *Limnopus* isp., trackway, concave epirelief. Holotype of *Tridentichnus supaiensis* Gilmore 1927, *nomen dubium*. **D.** Enlargement of C, three tracks. **E.** GRCA 4646. *Batrachichnus* isp., trackway, concave epirelief. Strike and dip symbols indicate the supposed dip of the inclined bedding plane, dashed arrows indicate the direction of progression. p=pes imprint. m=manus imprint.

The specimen GRCA 4646 (Figure 9-6E) displays a small-sized trackway (pes length 20 mm or 0.79 in) with tracks representing 10 consecutive pes-manus couples, that can be assigned to *Batrachichnus* isp. The tracks are ectaxonic, the pes imprint is pentadactyl and plantigrade, and the manus imprint is tetradactyl and semi-plantigrade; the manus imprint is about half the size of the pes imprint. The pace angulation is quite low (65–85°). The digit drag marks and trackway asymmetry indicate an inclined paleosurface (dune foreset). *Batrachichnus* is generally regarded as the track of small temnospondyl tetrapods (e.g., Voigt 2005).

Reptile Tracks

The specimen GRCA 11288 (Figure 9-7A) displays two consecutive pes-manus couples that can be assigned to *Varanopus* isp. These tracks are pentadactyl and ectaxonic, and the pes imprint is slightly larger than the manus imprint (pes length of about 40 mm or 1.6 in). Digit impressions are relatively long and thick and end in claw marks that can be bifurcated. The digit impression V of the pes is relatively long (about as long as the digit III impression) and straight. The footprints are semi-plantigrade, and the palm/sole imprint is relatively short. These tracks do not show typical features of

locomotion on inclined paleosurfaces, therefore they were probably impressed in an interdune paleoenvironment.




Figure 9-7. Reptile footprints from the Wescogame Formation (LORENZO MARCHETTI & SPENCER LUCAS). **A.** GRCA 11288. *Varanopus* isp., incomplete step cycle, concave epirelief. **B.** USNM 11533, cf. *Varanopus* isp., trackway, concave epirelief. Holotype of *Stenichnus yakiensis* Gilmore 1927, *nomen dubium*. Strike and dip symbols indicate the supposed dip of the inclined bedding plane, dashed arrows indicate the direction of progression. p=pes imprint. m=manus imprint.

The specimen USNM 11533 (Figure 9-7B) shows a trackway with tracks belonging to 11 consecutive pes-manus couples assignable to cf. *Varanopus* isp. The tracks are ectaxonic, probably pentadactyl and show long and thin digit impressions. The pes imprint is larger than the manus imprint (pes length of about 40 mm or 1.6 in). The pace angulation is low (about 70°), and partial secondary overstep is observed. The digit drag marks and trackway asymmetry indicate locomotion on an inclined paleosurface (dune foreset). Due to poor preservation caused by digit sliding, the assignment is only tentative. This is the holotype of the ichnotaxon *Stenichnus yakiensis* Gilmore 1927, which is herein considered a *nomen dubium* because of the poor preservation. *Varanopus* is generally regarded as a small captorhinomorph track (e.g., Voigt 2005).

Undetermined Tracks

An additional relevant specimen is USNM 11577, the holotype of *Anomalopus sturdevanti* Gilmore 1927, which is herein considered a *nomen dubium* because of the poor preservation caused by digit sliding on an inclined plane (dune foreset).

Discussion

The revised tetrapod ichnoassociation from the Wescogame Formation is the oldest from GRCA (the Manakacha Formation has not so far yielded classifiable footprints), and also the oldest worldwide tetrapod ichnoassociation from eolian paleoenvironments. It includes (Table 9-1): cf. *Amphisauropus* isp., *Batrachichnus* isp., cf. *Limnopus* isp., cf. *Varanopus* isp. from dune foreset paleoenvironments and *Varanopus* isp. from interdune paleoenvironments. All these ichnotaxa are described for the first time from this formation. The occurrence of *Amphisauropus* and *Varanopus* is consistent with the *Dromopus* footprint biochron, that began during the Late Pennsylvanian (Ghzelian) (e.g., Fillmore et al. 2012). This is in agreement with previous age interpretations of the Wescogame Formation.

It is noteworthy, as in the case of the Coconino and De Chelly sandstones (Marchetti et al. 2019b), that the tetrapod ichnoassociation is quite similar to floodplain, lacustrine and coastal tetrapod ichnoassociations of the same age. This ichnoassociation includes quite diverse anamniote (seymouriamorphs and temnospondyls) tracks and fewer reptile tracks. This is an important datum that confirms the hypothesis of anamniote occurrence in eolian paleoenvironments (Francischini et al. 2019), in this case possibly including batrachomorphs as well. Moreover, the first occurrence of tracks in eolian paleoenvironments is significantly extended temporally (from the late early Permian to the late Carboniferous). Because few specimens are known and nevertheless encompass important data, further prospecting for footprints in the Wescogame Formation of the GRCA is strongly recommended.

Table 9-1. Museum collections, Wescogame Formation.

Catalog #	Assignment	Type	Previous Ichnotaxon	Validity	Preservation	Description	Collector	Date	Locality
GRCA 4646	*Batrachichnus* isp.	N/A	N/A	N/A	concave epirelief	trackway	Thrasher, E.	1931	Grand Canyon, Yaki Trail
GRCA 11288	*Varanopus* isp.	N/A	N/A	N/A	concave epirelief	partial trackway	McKee, E. D.	1938	Grand Canyon, Tanner Trail
USNM 11533	cf. *Varanopus* isp.	holotype	*Stenichnus yakiensis* Gilmore 1927	*nomen dubium*	concave epirelief	trackway	Metzler, A.	1926	Grand Canyon, Yaki Trail
USNM 11534	cf. *Amphisauropus* isp.	holotype	*Ammobatrachus turbatans* Gilmore 1928	*nomen dubium*	concave epirelief	trackway	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11577	Undetermined tracks	holotype	*Anomalopus sturdevanti* Gilmore 1927	*nomen dubium*	concave epirelief	trackway	Sturdevant, G. E.	1926	Grand Canyon, Yaki Trail
USNM 11691	cf. *Limnopus* isp.	holotype	*Tridentichnus supaiensis* Gilmore 1927	*nomen dubium*	concave epirelief	trackway	Sturdevant, G. E.	1927	Grand Canyon, Bright Angel Trail

Hermit Shale

Geology

The Hermit Shale (Formation) is mostly slope-forming red beds of siltstone and very fine grained sandstone about 30 to 100 m (100 to 330 ft) thick in the Grand Canyon (Noble 1922; Blakey 1990) (Figure 9-8). The fossil plants and footprints from the Hermit Shale have long been the basis for assigning it an early Permian (Leonardian; late Artinskian–Kungurian) age (e.g., White 1929). Hermit Shale deposition was by fluvial processes on a seasonally arid coastal plain (Blakey 1990).

Previous Studies

Abundant and well-preserved tetrapod tracks have been found in the Hermit Shale of GRCA. The first report is by Lull (1918), who erected the ichnotaxa *Exocampe*? *delicatula* and *Megapezia? coloradensis* based on specimens found by Charles Schuchert in 1915 along the Hermit Trail. These specimens come from the uppermost levels of what was then called the Supai Formation, now considered the Hermit Shale. This locality is stratigraphically placed in the basal part of the Hermit Shale (Gilmore 1927), above an erosional surface on the underlying sandstones of the Esplanade Sandstone (Figure 9-8). A new excavation at the same locality was led by Charles W. Gilmore in 1926, and the material was described in Gilmore (1927), who erected the ichnotaxa: *Batrachichnus obscurus, Collettosaurus pentadactylus*, *Dromillopus parvus*, *Hyloidichnus bifurcatus* and *Hylopus hermitanus.* He also proposed the new combination *Batrachichnus delicatulus* (Lull 1918). After a further excavation in 1927, led by Charles W. Gilmore and David White along the Hermit and Yaki trails, Gilmore (1928) erected the ichnospecies *Hyloidichnus whitei* and described a large slab with several trackways of different forms (USNM 11707A, not relocated in Haubold et al. 1995a).

Although the preservation of these trace fossils is remarkable (commonly grade 2.0–3.0 of the preservation scale, Marchetti et al. 2019a), few additional finds and publications have appeared on the ichnology of the Hermit Shale. Haubold (1971) introduced the ichnogenus *Gilmoreichnus* with the new combination *Gilmoreichnus hermitanus* (Gilmore 1927), adding to the type material some specimens from Germany. He also proposed the new combination *Batrachichnus parvus* (Gilmore 1927). The only revision of the Hermit Shale material was proposed by Haubold et al. (1995a), who recognized: *Batrachichnus delicatulus*, *Dimetropus* isp., *Hyloidichnus bifurcatus*, *Gilmoreichnus hermitanus* and *Parabaropus* isp. More recent studies proposed the invalidity of the ichnotaxon *Gilmoreichnus hermitanus*, because it was based on poorly preserved material actually assignable to different ichnotaxa (Voigt 2005; Marchetti et al. 2019a).




Figure 9-8. Stratigraphic section of the Hermit Shale, measured by Noble (1922) near Bass Point in the Grand Canyon (LORENZO MARCHETTI & SPENCER LUCAS).

Trace Fossils

Anamniote Tracks

The specimen USNM 11707A is a large slab showing more than 100 footprints and different ichnotaxa (Figure 9-9). This slab includes a relatively long trackway (13 consecutive pes-manus couples) assignable to *Amphisauropus kablikae* (Figure 9-10A–C). The footprints are well-preserved, pentadactyl and ectaxonic with straight digit impressions and round digit terminations on both the pes and the manus imprints, and the pes imprint is larger than the manus imprint (pes length about 70 mm or 2.8 in). The pes imprint is plantigrade and shows an elliptical sole impression, and the manus imprint is semi-plantigrade and distinctly wider than long. The trackway is broad, the pace angulation is about 70–110°, and the manus imprint is rotated towards the midline. Gilmore (1928) assigned this trackway to *Parabaropus coloradensis*, which was considered by Haubold et al. (1995a) similar to *Amphisauropus latus*, later synonymized with *Amphisauropus kablikae* by Voigt (2005). However, the specimen USNM 11707A was not relocated. The type material of *Parabaropus coloradensis* (YPM 2145A-C) includes three syntype slabs from the Hermit Shale, showing: A) an isolated pes track, assignable to *Dimetropus* isp., B) an isolated pes track, assignable to cf. *Amphisauropus* isp., and C) several poorly preserved tracks, assignable to cf. *Amphisauropus* isp. Because this type material is incomplete and includes different morphotypes, we consider *Parabaropus* Gilmore 1927 and *Parabaropus coloradensis* (Lull 1918) to be *nomina dubia*. The specimen USNM 11707B includes a manus track and a partial pes track assigned to *Amphisauropus* isp. This ichnogenus is uncommon in the Hermit Shale and is generally regarded as the track of seymouriamorphs (Voigt 2005; Marchetti et al. 2017a).

The ichnotaxon *Batrachichnus salamandroides* is a common track type in the Hermit Shale. This material is characterized by small ectaxonic tracks with pentadactyl, plantigrade pes imprints and tetradactyl, semi-plantigrade manus imprints; the pes imprint is larger than the manus imprint (foot length of about 10 to 20 mm or 0.39 to 0.79 in). The trackway pattern shows broad trackways in a simple alternating arrangement, with low pace angulation (usually lower than 100°). The ichnotaxa (1) *Exocampe*? *delicatula* Lull 1918 (holotype YPM 3410), later assigned to *Batrachichnus* in the new combination *Batrachichnus delicatulus* by Gilmore (1927), and (2) *Dromillopus parvus* Gilmore 1927 (holotype USNM 11537, Figure 9-10D) are junior subjective synonyms of *Batrachichnus salamandroides* (Geinitz 1861), following Haubold (2000) and Voigt (2005). Other significant material includes: USNM 11518, showing a trackway with slightly sinuous tail impression; USNM 11519 and USNM 11520, which are completely covered by tracks; GRCA 3051, showing tracks associated with groups of parallel scratches interpreted as swimming traces; and GRCA 3218, GRCA 19408 and UCMP-V 75209A, showing incomplete step cycles. Other materials assignable to *Batrachichnus salamandroides* come from the Mogollon Rim area (Figure 9-10E), including the specimens: UCMP-V 75216A-C and UCMP-V 4013A-B. Some other material from GRCA is of dubious attribution. The ichnotaxon *Batrachichnus obscurus* Gilmore 1927 is assignable to cf. *Batrachichnus* isp. due to the poor preservation of the holotype trackway (USNM 11529), which shows a deep body/tail impression and unclear and incomplete tracks. The specimen USNM 11563 includes a rather poorly preserved trackway incorrectly assigned to *Gilmoreichnus hermitanus* by Haubold et al. (1995a). This material is instead assignable to cf. *Batrachichnus* isp., according to

Marchetti et al. (2019b). *Batrachichnus* is generally regarded as the track of small temnospondyls (e.g., Voigt 2005).



Figure 9-9. Tetrapod footprints from the Hermit Shale along Hermit Trail in USNM 11707A (LORENZO MARCHETTI & SPENCER LUCAS). This large slab shows more than 100 footprints and several different ichnotaxa (*Amphisauropus kablikae*, *Hyloidichnus bifurcatus*, cf. *Batrachichnus* isp.) in concave epirelief.

The ichnotaxon *Ichniotherium cottae* is known from a single specimen of the Hermit Shale (Figure 9-10F), discovered along the Bright Angel Trail in 2005 and left in place. Available photographs show an incomplete step cycle composed of a left pes, a right pes-manus couple and another left pes preserved in convex hyporelief. There is no information on the manual imprint morphology as the only referred track has been lost by weathering except for the terminal impression of digit I. Nevertheless, ichnospecific assignation is unambiguous due to the characteristic pedal morphology. *Ichniotherium* is characterized by ectaxonic, pentadactyl and plantigrade pes imprints, straight digit impressions with enlarged and distally rounded tips, as well as an oval-shaped heel impression (Voigt 2005). The ichnospecies *Ichniotherium cottae* is unique in its short impression of pedal digit V (about half of the length of pedal digit IV impression) and pedal tracks that are more or less parallel to the direction of the trackmaker's movement (Voigt 2005; Buchwitz and Voigt 2018). *Ichniotherium cottae* is most likely the track of advanced diadectid reptiliomorphs (Voigt et al. 2007; Buchwitz and Voigt 2018).




Figure 9-10. Anamniote footprints from the Hermit Shale (LORENZO MARCHETTI & SPENCER LUCAS). **A–C.** USNM 11707A, Hermit Trail. *Amphisauropus kablikae*: **A.** trackway, concave epirelief; **B.** Enlargement of A, left pes–manus couple; **C.** Right pes. **D.** USNM 11537, Hermit Trail. *Batrachichnus salamandroides*, trackway with thin tail impression, concave epirelief. Holotype of *Dromillopus parvus* Gilmore 1927. **E.** UCMP-V 4013A, Mogollon Rim. *Batrachichnus salamandroides*, right pes-manus couple and tail impression, concave epirelief. **F.** *Ichniotherium cottae*, incomplete step cycle. Specimen in situ in the Grand Canyon National Park, discovered by Thomas Martens. Dashed arrows indicate the direction of progression.

Synapsid Tracks

The ichnotaxon *Dimetropus* is uncommon in the Hermit Shale. This material is characterized by relatively large, pentadactyl, plantigrade and ectaxonic tracks (pes length of 50 to 120 mm or 2.0 to 4.72 in). The pes imprint is much larger than the manus imprint and has a postero-laterally elongated sole impression. The digit impressions are short and terminate in claw impressions, when preserved. The digit impression base is arranged in a semi-circle. These tracks are often collapsed (e.g., USNM 11527, UCMP-V 4013B) or preserved as undertracks (e.g., USNM 11598). The incomplete step cycles show a probable slow gait, with relatively short stride and pace lengths.

The first mention of clear synapsid tracks from the Hermit Shale was by Haubold et al. (1995a), who assigned to *Dimetropus* cf. *nicolasi* a single pes track (USNM 11528) and to *Dimetropus* isp., an incomplete step cycle previously assigned by Gilmore (1927) to *Parabaropus coloradensis* (USNM 11598). We assign USNM 11528 to *Dimetropus* isp. (Figure 9-11B) and USNM 11598 to cf. *Dimetropus* isp., because of its poor preservation. Notably, USNM 11528 shows a collapsed track, and its undertrack is on the back of the slab. We also assign the single pes track of specimen YPM 2145A (syntype of *Parabaropus coloradensis, nomen dubium*) to *Dimetropus* isp. (Figure 9-11A).

Also, some of the material assigned to *Gilmoreichnus hermitanus* by Haubold et al. (1995a), and notably the trackway of specimen USNM 11527, the holotype of *Collettosaurus pentadactylus* Gilmore 1927 (*nomen dubium* after Marchetti et al. 2019a), is assigned to cf. *Dimetropus* isp. (Figure 9-11D). Three specimens from the Mogollon Rim area (Figure 9-11C), each an isolated track (UCMP-V 4010A-B, UCMP-V 4013B), are assigned to *Dimetropus* isp. This ichnotaxon is usually attributed to non-therapsid synapsids ("pelycosaurs"), with the exception of the family Varanopidae (e.g., Voigt 2005).




Figure 9-11. Synapsid footprints from the Hermit Shale (LORENZO MARCHETTI & SPENCER LUCAS). **A.** YPM 2145A, Hermit Trail. *Dimetropus* isp., left pes cut by a mud crack, convex hyporelief. Syntype of *Parabaropus coloradensis* (Lull 1918), *nomen dubium*. **B.** USNM 11528, Hermit Trail. *Dimetropus* isp., collapsed pes track, concave epirelief. **C.** UCMP-V 4010B, Mogollon Rim. *Dimetropus* isp., right pes, undertrack preservation, convex hyporelief. **D.** USNM 11527, Hermit Trail. cf. *Dimetropus* isp., trackway with collapsed tracks, concave epirelief. Holotype of *Collettosaurus pentadactylus* Gilmore 1927, *nomen dubium*. Dashed arrows indicate the direction of progression.

Reptile Tracks

The ichnotaxon *Dromopus lacertoides* is uncommon in the Hermit Shale. This material is characterized by long, curved and tapering digit impressions ending in claw marks. Tracks are relatively small (pes length of 30 mm or 1.2 in), pentadactyl, semi-digitigrade and strongly ectaxonic. The impressions of digits I–IV are often superimposed and distally curved inwards, and the digit V impression is proximal and directed laterally. Often, only the digit III and IV impressions are preserved. Manus and pes tracks have the same morphology, but the manus imprint is smaller. The specimen GRCA 3171 (Figure 9-12A) shows several tracks assignable to *Dromopus lacertoides*. Some specimens from the Mogollon Rim area (Figure 9-12B) are also assignable to *Dromopus lacertoides* (UCMP-V 4010C and UCMP-V 4013C). This ichnotaxon is generally regarded as the track of araeoscelid diapsids or bolosaurid parareptiles (e.g., Voigt 2005).




Figure 9-12. Reptile footprints from the Hermit Shale (LORENZO MARCHETTI & SPENCER LUCAS). **A.** GRCA 3171, Yaki Trail. *Dromopus lacertoides*, several footprints, concave epirelief. **B.** UCMP-V 4010C, Mogollon Rim. *Dromopus lacertoides*, left footprint, convex hyporelief. **C.** USNM 11518, Hermit Trail. Holotype of *Hyloidichnus bifurcatus* Gilmore 1927, left pes–manus couple, convex hyporelief. **D.** USNM 11692, Yaki/South Kaibab Trail. *Hyloidichnus bifurcatus,* left pes–manus couple, concave epirelief. Holotype of *Hyloidichnus whitei* Gilmore 1928. **E.** UCMP-V 75216G, Mogollon Rim*. Hyloidichnus bifurcatus*, left pes–manus couple, concave epirelief. **F.** UCMP-V 75216D, Mogollon Rim. *Erpetopus* isp., partial trackway, convex hyporelief. **G.** UCMP-V 75216A, Mogollon Rim. *Erpetopus* isp., left pes–manus couple, convex hyporelief. **H.** UCMP-V 75209A, Hermit Trail. *Erpetopus* isp., partial trackway, convex hyporelief. Dashed arrows indicate the direction of progression. p=pes imprint. m=manus imprint.

The ichnotaxon *Hyloidichnus bifurcatus* Gilmore 1927 (Figure 9-12C) is the most common track type in the Hermit Shale. This material is characterized by small to relatively large (pes length of 20 to 70 mm, or 0.79 to 2.7 in), pentadactyl, ectaxonic, and semi-plantigrade tracks. Digit impressions are long, straight and end in flat claw marks that are often bifurcated. The digit IV impression is the longest, digit III and IV impressions are of similar length, and the digit V impression is about as long as the digit I–II impressions. The palm/sole impression is very short and shows a convex to flat proximal margin. The pes imprint is slightly larger than the manus imprint. The trackways are broad, with a relatively low pace angulation and manus tracks rotated inward. The ichnogenus holotype (USNM 11518) (Figure 9-12C) includes a trackway with five consecutive pes-manus couples.

The ichnospecies *Hyloidichnus whitei* Gilmore 1928 (holotype USNM 11692) (Figure 9-12D) is here considered a junior synonym of *Hyloidichnus bifurcatus*, in agreement with Haubold et al. (1995a). We confirm the assignment to *Hyloidichnus bifurcatus* of a pes-manus couple on specimen USNM 11522 (Haubold et al. 1995a). The step cycle on USNM 11519 and the several trackways and isolated tracks on USNM 11707A (Figure 9-9), assigned to *Hylopus hermitanus* by Gilmore (1927, 1928), and to *Gilmoreichnus hermitanus* by Haubold et al. (1995a), are herein assigned to

Hyloidichnus bifurcatus. The specimens YPM 304, showing differently impressed and superimposed trackways, and USNM 11711, showing a trackway, previously assigned to *Gilmoreichnus hermitanus* by Haubold et al. (1995a), are herein assigned to *Hyloidichnus bifurcatus*. Other significant specimens assigned to this ichnotaxon are: UCMP-V 75209B, showing a large-sized pes-manus couple with evident digit impression bifurcation, and USNM 11529 and GRCA 3174, showing an incomplete step cycle. Some specimens assignable to *Hyloidichnus bifurcatus* come from the Mogollon Rim area (Figure 9-12E): UCMP-V 75216F-G, showing an isolated pes and an isolated pes-manus couple, respectively. Because of the poor preservation, we consider the ichnogenus *Gilmoreichnus* Haubold 1971, and the ichnospecies *Gilmoreichnus hermitanus* (Gilmore 1927), based on the trackway holotype, USNM 11517, as *nomina dubia* and assign this material to cf. *Hyloidichnus* isp., in agreement with Marchetti et al. (2019a). *Hyloidichnus* is generally related to captorhinid trackmakers, such as the Moradisaurinae (e.g., Voigt et al. 2010).

A single specimen from the Hermit Trail (UCMP-V 75209A) (Figure 9-12F) and three specimens from the Mogollon Rim area (UCMP-V 75216A, D, E) (Figures 9-12G and 9-12H), each showing an incomplete step cycle, are assignable to *Erpetopus* isp. A specimen from the Hermit Trail (USNM 11518) shows an incomplete step cycle with partial overstep of the pes on the manus and a possible body impression. Because of the lack of the pedal digit impression V, this material is assigned to cf. *Erpetopus* isp. These specimens are characterized by small (pes length of about 10 to 40 mm, or 0.39 to 1.6 in), pentadactyl, ectaxonic, semi-plantigrade footprints with relatively long and thin digit impressions ending in small claw impressions. The digit I–IV impressions are distally curved inwards, and the digit V impression is in a proximal position and distally curved backwards. The pedal digit III–IV impressions can be parallel. The digit IV impression is the longest, the digit III and IV impressions are of similar length, and the digit V impression is about as long as the digit I–II impressions. The palm/sole impression is very short and can show a concave proximal margin. The manus imprint is smaller than the pes imprint and rotated inward. The step cycles show a low pace length. This track type has been related to small captorhinomorph or parareptile producers (e.g., Haubold and Lucas 2003; Marchetti 2016).

Undetermined Tracks

A specimen from the Hermit Shale along the Yaki Trail (GRCA 3172–3173) (Figure 9-13) displays a trackway with three small (pes length of about 30 mm or 1.2 in) pes-manus couples. These traces are pentadactyl and semi-plantigrade (although the pedal digit impression V is not impressed), with rounded digit terminations. The pes imprint is ectaxonic and more deeply impressed in its medial part, and the digit impressions are thick and straight; the imprint is about as long as wide. The manus imprint is wider than long and shows very short digit impressions. The impressions of digits III and IV are the longest and about equal in length, and the external digit impressions are reduced. The trackway is broad, with well-spaced pes-manus couples and a low pace angulation (70°). The pes imprint is somewhat similar to *Amphisauropus* and, to a lesser degree, *Limnopus*, but the manus imprint is different from these ichnotaxa. The manus imprint resembles some incomplete tracks of *Dimetropus*, but no claw marks were observed, and the pes imprint has a structure different from *Dimetropus.* In fact, the sole impression is shorter and not mediolaterally expanded, the digit impressions are relatively longer, and the footprints are more deeply impressed medially, whereas

they are more deeply impressed laterally in *Dimetropus*. These features are instead typical of the Guadalupian track *Lunaepes* and the Lopingian track *Pachypes*. Further studies are needed in order to correctly assign this material, and the discovery of middle–late Permian (Guadalupian–Lopingian) track morphotypes in the early Permian (Cisuralian) would be noteworthy. This material cannot be attributed to therapsid trackmakers because the pes imprint is more deeply impressed medially (see Marchetti et al. 2019d). An attribution to small, heavily built parareptiles such as the nycteroleterids is more probable.




Figure 9-13. Undetermined footprints from the Hermit Shale (LORENZO MARCHETTI & SPENCER LUCAS). **A.** GRCA 3172, Yaki/South Kaibab Trail. Trackway, concave epirelief. **B.** Enlargement of A, right pes–manus couple. **C.** GRCA 3173, counterpart of GRCA 3172. Right pes–manus couple, convex hyporelief. Dashed arrows indicate the direction of progression.

Discussion

The tetrapod ichnoassociation from the Hermit Shale is certainly the best-preserved and most diverse known from GRCA. It includes the following ichnotaxa (Table 9-2): *Amphisauropus kablikae, Batrachichnus salamandroides, Dimetropus* isp., *Dromopus lacertoides, Erpetopus* isp., *Hyloidichnus bifurcatus*, and *Ichniotherium cottae*. The ichnogenera *Amphisauropus, Dromopus, Erpetopus* and *Ichniotherium* are here recognized for the first time from this formation, notably increasing the ichnodiversity. The occurrence of *Dimetropus* and *Ichniotherium* suggests a Kungurian minimum age, and the occurrence of *Erpetopus* suggests a late Artinskian maximum age. Therefore, this ichnoassociation belongs to the *Erpetopus* tetrapod footprint biochron and suggests a late Artinskian to Kungurian age for the Hermit Shale, probably late Artinskian–early Kungurian because the Hermit Shale is overlain by the late Kungurian Coconino Sandstone (Blakey and Knepp 1989; Voigt and Lucas 2018; Marchetti et al. 2019b). The ichnoassociation includes abundant and relatively diverse reptile and anamniote tracks and rarer synapsid tracks. This is consistent with the ichnofaunal composition of the late Artinskian–Kungurian ichnoassociations (e.g., Haubold and

Lucas 2003; Gand and Durand 2006; Voigt et al. 2011; Marchetti et al. 2015a, 2015b; Voigt and Haubold 2015; Mujal et al. 2016; Voigt and Lucas 2017), characterized by an increased abundance and diversity of reptile tracks, therefore registering a low-latitude and possibly worldwide reptile radiation, whereas older ichnoassociations include more anamniote and synapsid tracks (Voigt and Lucas 2018; Marchetti et al. 2019b).

Table 9-2. Museum collections, Hermit Shale.

Catalog #	Assignment	Type	Previous Ichnotaxon	Validity	Preservation	Description	Collector	Date	Locality
GRCA 3051	*Batrachichnus salamandroides*	N/A	N/A	N/A	convex hyporelief	several tracks, swimming traces	–	–	Grand Canyon
GRCA 3171	*Dromopus lacertoides*	N/A	N/A	N/A	concave epirelief	several tracks	–	–	Grand Canyon
GRCA 3172	Undetermined	N/A	N/A	N/A	concave epirelief	trackway	Goebels, L. A.	1937	Grand Canyon, Yaki Trail
GRCA 3173	Undetermined	N/A	N/A	N/A	convex hyporelief	counterpart of GRCA 3172	Goebels, L. A.	1937	Grand Canyon, Yaki Trail
GRCA 3174	*Hyloidichnus bifurcatus*	N/A	N/A	N/A	concave epirelief	partial trackway	Goebels, L. A.	1937	Grand Canyon, Yaki Trail
GRCA 3218	*Batrachichnus salamandroides*	N/A	N/A	N/A	concave epirelief	partial trackway	Richardson, F.	1960	Grand Canyon, Hermit Trail
GRCA 19408	*Batrachichnus salamandroides*	N/A	N/A	N/A	concave epirelief	partial trackway	Hill, W. H.	1968	Grand Canyon, Hermit Trail
UCMP-V 4010A	*Dimetropus* isp.	N/A	N/A	N/A	concave epirelief	pes imprint	Camp, C. L.	–	Yavapai County, Sedona
UCMP-V 4010B	*Dimetropus* isp.	N/A	N/A	N/A	convex hyporelief	pes imprint	Camp, C. L.	–	Yavapai County, Sedona
UCMP-V 4010C	*Dromopus lacertoides*	N/A	N/A	N/A	convex hyporelief	manus imprint	Camp, C. L.	–	Yavapai County, Sedona
UCMP-V 4013A	*Batrachichnus salamandroides*	N/A	N/A	N/A	concave epirelief	pes-manus couple	Camp, C. L.	1940	Gila County, Carrizo Cr.
UCMP-V 4013 B	*Batrachichnus salamandroides*, *Dimetropus* isp.	N/A	N/A	N/A	concave epirelief	trackway (Bs), pes imprint (D)	Camp, C. L.	1940	Gila County, Carrizo Cr.
UCMP-V 4013C	*Dromopus lacertoides*	N/A	N/A	N/A	convex hyporelief	pes-manus couple	Camp, C. L.	1941	Gila County, Carrizo Cr.
UCMP-V 75209A	*Batrachichnus salamandroides*, *Erpetopus* isp.	N/A	N/A	N/A	concave epirelief	partial trackway (Bs), partial trackway (E)	Richardson, F.	1938	Grand Canyon, Hermit Trail
UCMP-V 75209B	*Hyloidichnus bifurcatus*	N/A	N/A	N/A	concave epirelief	pes-manus couple	Richardson, F.	1938	Grand Canyon, Hermit Trail
UCMP-V 75216A	*Batrachichnus salamandroides*, *Erpetopus* isp.	N/A	N/A	N/A	concave epirelief	trackway and tail impression (Bs), partial trackway (E)	Camp, C. L.	1940	Navajo County, Limestone Canyon
UCMP-V 75216B	*Batrachichnus salamandroides*	N/A	N/A	N/A	concave epirelief	pes-manus couple	Camp, C. L.	1940	Navajo County, Limestone Canyon
UCMP-V 75216C	*Batrachichnus salamandroides*	N/A	N/A	N/A	concave epirelief	manus	Camp, C. L.	1940	Navajo County, Limestone Canyon

Table 9-2 (continued). Museum collections, Hermit Shale.

Catalog #	Assignment	Type	Previous Ichnotaxon	Validity	Preservation	Description	Collector	Date	Locality
UCMP-V 75216D	*Erpetopus* isp.	N/A	N/A	N/A	convex hyporelief	incomplete step cycle	Camp, C. L.	1940	Navajo County, Limestone Canyon
UCMP-V 75216E	*Erpetopus* isp.	N/A	N/A	N/A	convex hyporelief	incomplete step cycle	Camp, C. L.	1940	Navajo County, Limestone Canyon
UCMP-V 75216F	*Hyloidichnus bifurcatus*	N/A	N/A	N/A	convex hyporelief	pes imprint	Camp, C. L.	1940	Navajo County, Limestone Canyon
UCMP-V 75216G	*Hyloidichnus bifurcatus*	N/A	N/A	N/A	concave epirelief	pes-manus couple	Camp, C. L.	1940	Navajo County, Limestone Canyon
USNM 11517	cf. *Hyloidichnus* isp.	holotype	*Gilmoreichnus hermitanus* (Gilmore 1927)	*nomen dubium*	concave epirelief and convex hyporelief	trackway	Gilmore, C. W., Metzler, A.	1926	Grand Canyon, Hermit Trail
USNM 11518	*Hyloidichnus bifurcatus*	holotype	*Hyloidichnus bifurcatus* Gilmore 1927	valid	convex hyporelief	trackway	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11518	*Batrachichnus salamandroides,* cf. *Erpetopus* isp.	N/A	N/A	N/A	concave epirelief	trackway and tail impression (Bs), partial trackway (E)	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11519	*Batrachichnus salamandroides*, *Hyloidichnus bifurcatus*	N/A	N/A	N/A	concave epirelief	several tracks (Bs), partial trackway (Hb)	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11520	*Batrachichnus salamandroides*	N/A	N/A	N/A	concave epirelief	several tracks	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11522	*Hyloidichnus bifurcatus*	N/A	N/A	N/A	concave epirelief	pes-manus couple	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11527	cf. *Dimetropus* isp.	holotype	*Collettosaurus pentadactylus* Gilmore 1927	*nomen dubium*	concave epirelief	trackway	Gilmore, C. W., Metzler, A.	1926	Grand Canyon, Hermit Trail
USNM 11528	*Dimetropus* isp.	N/A	N/A	N/A	concave epirelief	pes imprint	Sturdevant, G. E.	1926	Grand Canyon, Hermit Trail
USNM 11529	cf. *Batrachichnus* isp., *Hyloidichnus bifurcatus*	holotype	*Batrachichnus obscurus* Gilmore 1927	*nomen dubium*	concave epirelief	trackway and tail impression (B), partial trackway (Hb)	Gilmore, C. W., Metzler, A.	1926	Grand Canyon, Hermit Trail
USNM 11537	*Batrachichnus salamandroides*	holotype	*Dromillopus parvus* Gilmore 1927	junior synonym	concave epirelief	trackway and tail impression	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11563	cf. *Batrachichnus* isp.	N/A	N/A	N/A	concave epirelief	trackway	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11598	cf. *Dimetropus* isp.	N/A	N/A	N/A	convex hyporelief	partial trackway	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11692	*Hyloidichnus bifurcatus*	holotype	*Hyloidichnus whitei* Gilmore 1928	junior synonym	concave epirelief	partial trackway	Gilmore, C. W., White, D.	1927	Grand Canyon, Yaki Trail

Table 9-2 (continued). Museum collections, Hermit Shale.

Catalog #	Assignment	Type	Previous Ichnotaxon	Validity	Preservation	Description	Collector	Date	Locality
USNM 11707A	*Amphisauropus kablikae, Hyloidichnus bifurcatus*, cf. *Batrachichnus* isp.	N/A	N/A	N/A	concave epirelief	several trackways (Hb), trackway (Ak), trackway (B)	Gilmore, C. W., Metzler, A.	1927	Grand Canyon, Hermit Trail
USNM 11707B	*Amphisauropus kablikae*	N/A	N/A	N/A	concave epirelief	pes and manus imprints	Gilmore, C. W., Metzler, A.	1927	Grand Canyon, Hermit Trail
USNM 11711	*Hyloidichnus bifurcatus*	N/A	N/A	N/A	concave epirelief	trackway	Gilmore, C. W., Metzler, A.	1927	Grand Canyon, Hermit Trail
YPM 304	*Hyloidichnus bifurcatus*	N/A	N/A	N/A	concave epirelief	superimposed trackways	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
YPM 2145A	*Dimetropus* isp.	syntype	*Parabaropus coloradensis* (Lull 1918)	*nomen dubium*	convex hyporelief	pes track	Schuchert, C.	1915	Grand Canyon, Hermit Trail
YPM 2145B	cf. *Amphisauropus* isp.	syntype	*Parabaropus coloradensis* (Lull 1918)	*nomen dubium*	convex hyporelief	pes track	Schuchert, C.	1915	Grand Canyon, Hermit Trail
YPM 2145C	cf. *Amphisauropus* isp.	syntype	*Parabaropus coloradensis* (Lull 1918)	*nomen dubium*	convex hyporelief	partial trackway	Schuchert, C.	1915	Grand Canyon, Hermit Trail
YPM 3410	*Batrachichnus salamandroides*	holotype	*Exocampe? delicatula* Lull 1918	junior synonym	convex hyporelief	partial trackway	Schuchert, C.	1915	Grand Canyon, Hermit Trail

Coconino Sandstone

Geology

In the Grand Canyon, the Coconino Sandstone forms bold cliffs characterized by large-scale crossbeds between red bed slopes of the Hermit Shale (below) and the Toroweap Formation (above). The contact of the Coconino Sandstone with the underlying Hermit Shale is an unconformity, whereas the overlying contact with the Toroweap Formation is gradational and intertonguing (Middleton et al. 1990). In the Grand Canyon, the Coconino Sandstone is 20 to over 183 m (66 to 600 ft) thick and consists entirely of quartz sandstone (Figure 9-14). Deposition was in a large sand sea (erg) that covered much of northern Arizona and extended to the east to central New Mexico, where the equivalent lithosome is the Glorieta Sandstone (e.g., McKee 1979; Blakey and Knepp 1989; Middleton et al. 1990; Lucas et al. 2013). Coconino eolian deposits are primarily crossbedded sandstone that represent dunes, but there are some thin intervals of laminar and ripple laminar sandstone that are interdunal deposits (McKee 1979). The Coconino has long been recognized as of early Permian (Leonardian) age because it is bracketed by units of that age, Hermit below and lower Kaibab Formation above (Middleton et al. 1990). The recently revised tetrapod ichnoassociation of the Coconino Sandstone is consistent with a Kungurian age (Marchetti et al. 2019b).

Previous Studies

The most famous track-bearing formation of the Grand Canyon is the Coconino Sandstone, which is a reference unit for describing tetrapod tracks in Paleozoic eolian environments, because of the noteworthy abundance of tracks and trackways preserved on dune foreset paleosurfaces (e.g., Lockley et al. 1994; Hunt and Lucas 2006, 2007; Marchetti et al. 2019b). The first report of Coconino Sandstone tracks was by Lull (1918), who erected the ichnotaxa *Laoporus noblei* and *Laoporus schucherti* based on material collected by Charles Schuchert in 1915 along the Hermit Trail. The first large excavation along the Hermit Trail was led by John C. Merriam in 1924, and the material was described by Gilmore (1926). Based on this material, Gilmore erected the following ichnotaxa: *Agostopus matheri*, *Allopus*? *arizonae*, *Baropezia eakini*, *Barypodus palmatus*, *Dolichopodus tetradactylus*, *Nanopus merriami* and *Paleopus regularis*. The new combination *Baropezia arizonae* (Gilmore 1926) was proposed by Baird (1952). At the same locality, a permanent exhibit with tracks exposed on the foreset paleosurfaces on the side of the Hermit Trail was prepared in 1924 by the director of the National Park Service, Stephen F. Mather, with the aid of Charles W. Gilmore. These surfaces are still exposed today (Marchetti et al. 2019b) (Figure 9-15A). After a further excavation led by Charles W. Gilmore in 1926 at the same locality, he introduced the following ichnotaxa (in Gilmore 1927): *Agostopus medius*, *Amblyopus pachypodus*, *Baropus coconinoensis*, *Barypodus metszeri*, *Barypodus tridactylus* and *Nanopus maximus*. All the material from the Hermit Trail came from the lower half of the Coconino Sandstone (Gilmore 1926, 1927) (Figure 9-14).




Figure 9-14. Stratigraphic section of the Coconino Sandstone measured along the Hermit Trail (LORENZO MARCHETTI & SPENCER LUCAS).




Figure 9-15. Tetrapod footprints from the Coconino Sandstone (LORENZO MARCHETTI & SPENCER LUCAS). **A.** Trampled surface exposed along the Hermit Trail. **B.** Large slab RAM 244 showing more than 500 footprints, differently oriented trackways compared to the slope (upslope, downslope, transverse), and the longest trackway of the Paleozoic for number of footprints (almost 200), convex hyporelief.

Subsequent studies of Coconino Sandstone footprints have focused on ichnotaxonomy, locomotion and paleoecology, with the study of new material from the Mogollon Rim area and also through comparisons with modern equivalents (e.g., McKee 1944, 1947; Baird 1952; Brand 1979; Brand and Tang 1991; Lockley 1992; Loope 1992; McKeever and Haubold 1996; Haubold 2000; Hunt and Lucas 2006, 2007; Citton et al. 2012; Francischini et al. 2019; Marchetti et al. 2019b). The material from Ash Fork in the Mogollon Rim area comes from the middle part of the Coconino Sandstone, showing a similar stratigraphic distribution compared to the GRCA tracks (e.g., Hunt and Lucas 2005a). Some slabs coming from this locality show very long trackways, including the longest known trackway of the Paleozoic for number of tracks (Figure 9-15B). Trackways with closely spaced tracks were interpreted as subaerial upslope progression by McKee (1944, 1947), and

trackways with very asymmetric patterns were interpreted as subaqueous current-driven lateral progression (Brand and Tang 1991). This latter hypothesis was rejected by Loope (1992), Lockley (1992) and Marchetti et al. (2019b), also because of the clear eolian origin of the Coconino Sandstone lithofacies. Marchetti et al. (2019b, 2019c) proposed a locomotion model in eolian paleoenvironments mostly based on Coconino Sandstone material, distinguishing trackways showing upslope and downslope progression, with a possible transverse component. In this context, the track morphology is strongly influenced by gravity and thus by sliding of the digits during the trackmaker's locomotion. So, most of the previous ichnotaxonomic assignments were incorrect because they interpreted digit drag marks as actual digit impressions. This caused: (1) an overestimation of footprint diversity in older works (e.g., Gilmore 1926, 1927); and (2) an underestimation of footprint diversity in more recent works (McKeever and Haubold 1996; Haubold 2000). In fact, all the footprints from the Coconino Sandstone were assigned to the ichnogenus *Chelichnus* by McKeever and Haubold (1996) and Haubold (2000).

Based on this interpretation, Hunt and Lucas (2006, 2007) named the *Chelichnus* ichnofacies for low-diversity tetrapod ichnoassociations in eolian paleoenvironments, including the *Laoporus* ichnoassemblage of Lockley et al. (1994). However, the material assigned to *Chelichnus* and *Laoporus* is characterized by short and parallel "digit impressions," which are actually digit drag marks of different ichnotaxa. After a comprehensive revision, Marchetti et al. (2019b, 2019c) invalidated *Chelichnus*, *Laoporus* and several other ichnotaxa from the Coconino Sandstone and attributed the generally similar morphology of Paleozoic tetrapod footprints in desert paleoenvironments to the effects of locomotion on inclined planes. After excluding these effects, the Coconino Sandstone ichnoassociation is similar to the contemporary non-desert tetrapod ichnoassociations belonging to the *Erpetopus* footprint biochron, including: cf. *Amphisauropus* isp., cf. *Dromopus* isp., *Erpetopus* isp., *Ichniotherium sphaerodactylum,* cf. *Tambachichnium* isp. and *Varanopus curvidactylus*. Moreover, a possible facies-crossing, low-latitude and possibly worldwide reptile radiation was recognized in the tetrapod footprint record during the early Permian, and this is observable in the Coconino Sandstone as well (Marchetti et al. 2019b). Francischini et al. (2019) described new noteworthy specimens of *Ichniotherium sphaerodactylum* from the Coconino Sandstone of GRCA and discussed the presence of diadectomorphs in desert paleoenvironments.

Trace fossils

Anamniote Tracks

The ichnogenus *Amphisauropus* is rare in the Coconino Sandstone, and only one specimen (USNM 11135) (Figure 9-16A) showing an upslope-directed trackway has been tentatively assigned to cf. *Amphisauropus* isp. It is characterized by relatively large tracks (pes length of about 70 mm or 2.8 in) with plantigrade pes and semi-plantigrade manus imprints. The tracks are pentadactyl and ectaxonic. The pes imprint is about as long as wide, and the manus imprint is smaller and wider than long. The trackway is broad, and the pace angulation is very low (50–55°), a partial secondary overstep is observed, and the manus is directed inwards. This is the holotype of *Agostopus matheri* Gilmore 1926, considered a *nomen dubium* because of its poor preservation (Marchetti et al. 2019b). *Amphisauropus* is generally regarded as the track of seymouriamorphs (Voigt 2005; Marchetti et al. 2017a).




Figure 9-16. Anamniote footprints from the Coconino Sandstone (LORENZO MARCHETTI & SPENCER LUCAS). **A.** USNM 11135, Hermit Trail. cf. *Amphisauropus* isp., trackway, concave epirelief. Holotype of *Agostopus matheri* Gilmore 1926, *nomen dubium*. **B.** Specimen in situ, Dripping Springs Trail. *Ichniotherium sphaerodactylum*, trackway, convex hyporelief. **C.** Detail of B, four tracks. **D.** GRCA 3065, Kaibab Trail. *Ichniotherium sphaerodactylum*, partial trackway, concave epirelief. Strike and dip symbols indicate the supposed dip of the inclined bedding plane, dashed arrows indicate the direction of progression. p=pes imprint. m=manus imprint.

The ichnospecies *Ichniotherium sphaerodactylum* is more common in the Coconino Sandstone, being represented by 15 trackways found along the South Rim of GRCA (Dripping Springs, South Kaibab and Hermit trails) and the Mogollon Rim (Williams, Perkinsville and Seligman) (Francischini et al. 2019) (Figures 9-16B–D). This material is characterized by pentadactyl, plantigrade and ectaxonic manus and pes imprints with an oval-shaped palm/sole impression. The digit IV impression is the longest, and digit II–IV impressions can be bent inwards. The digit V impression is relatively longer in *Ichniotherium sphaerodactylum* than in other ichnospecies of the same ichnogenus. Pes imprints are slightly larger than manus imprints (about 80%). Digit impressions are clawless and have a typical drumstick shape, with expanded tips. Two specimens (USNM 11137 and USNM 11138), both collected by Charles W. Gilmore at the Hermit Trail in 1924, were first described by Gilmore (1926) as the holotype and paratype of *Baropezia eakini*. The ichnogenus *Baropezia* was erected by Matthew (1903) for Pennsylvanian tracks of the Joggins Formation of Nova Scotia, Canada. Although further analysis is necessary to understand the validity of this ichnogenus, we consider here the ichnospecies *Baropezia eakini* as a junior synonym of *Ichniotherium sphaerodactylum*, in

agreement with Francischini et al. (2019) and Marchetti et al. (2019b). The specimen USNM-V 11514, also from the Hermit Trail, is the holotype of *Baropus coconinoensis* (Gilmore 1927), but was regarded as a junior synonym of *Baropezia eakini* by Haubold (1971). As with *Baropezia*, more information is needed to evaluate the validity of the ichnogenus *Baropus* (whose type materials come from the Coal Measures of Kansas; Marsh 1894), but the ichnospecies *Baropus coconinoensis* is here regarded as a junior synonym of *Ichniotherium sphaerodactylum*, in agreement with Francischini et al. (2019) and Marchetti et al. (2019b). Some other *I. sphaerodactylum* specimens were previously assigned to *Baropus lentus* (UCMP 42945 and UCMP 42951), *Barypodus palmatus* (MNA-V 3386), *Chelichnus duncani* (USNM 11137 and USNM 11138), *Dolichopodus tetradactylus* (YPM 288), and *Laoporus noblei* (USNM-V 11508) (Gilmore 1926; McKeever and Haubold 1996). All of these ichnospecies are currently considered *nomina dubia* (Marchetti et al. 2019a). An incomplete specimen (GRCA 116801) found at the Hermit Trail was assigned to cf. *Ichniotherium* isp. (Francischini et al. 2019). *Ichniotherium* ichnospecies comprise one of the best examples of a strongly tied correlation between ichnotaxa and osteological taxa, representing tracks produced by diadectomorph reptiliomorphs. In particular, *I. sphaerodactylum* was assigned to the diadectid *Orobates pabsti*, based on complete trackways and skeletons from lower Permian (Rotliegend) of Germany (Voigt et al. 2007). The specimens from the Coconino Sandstone are the youngest/last record of *Ichniotherium*.

Synapsid Tracks

The ichnogenus *Tambachichnium* is uncommon in the Coconino Sandstone, and only a few specimens were assigned to cf. *Tambachichnium* isp. by Marchetti et al. (2019b). This material is characterized by relatively large tracks (foot length of about 50 to 100 mm, or 2.0 to 3.9 in) that are pentadactyl, ectaxonic and semi-plantigrade. The digit impressions are relatively thick and terminate in sharp claw marks. Digit I–IV impressions are superimposed and distally bent inwards. The digit V impression is proximal and oriented laterally. The manus imprint is smaller and less deeply impressed than the pes imprint. The trackways are broad and show a low pace angulation (50–95°) and possible secondary overstep. The most complete specimen is USNM 11506 (Figures 9-17A and B), including a trackway with complete pes and manus impressions. This is the holotype of *Nanopus maximus* Gilmore 1927, considered a *nomen dubium* by Marchetti et al. (2019b). Other specimens include the trackway holotypes of *Barypodus metszeri* Gilmore 1927 (USNM 11505) (Figure 9-17C) and *Barypodus tridactylus* Gilmore 1927 (USNM 11502) (Figure 9-17D), considered *nomina dubia* by Marchetti et al. (2019b) because of their poor preservation (incomplete tracks and digit drag marks). These three trackways are directed upslope, and those of specimens USNM 11505 and USNM 11506 also have a transverse component. An ichnotaxonomic revision of the ichnogenus *Tambachichnium* needs to be undertaken to definitively confirm the assignment of the Coconino material to this ichnogenus. This ichnotaxon is usually attributed to non-therapsid synapsids, such as the varanopids (e.g., Voigt 2005).




Figure 9-17. Synapsid footprints from the Coconino Sandstone, Hermit Trail (LORENZO MARCHETTI & SPENCER LUCAS). **A.** USNM 11506. cf. *Tambachichnium* isp., trackway, concave epirelief. Holotype of *Nanopus maximus* Gilmore 1927, *nomen dubium*. **B.** Detail of A, pes–manus couple. **C.** USNM 11505. cf. *Tambachichnium* isp., trackway, concave epirelief. Holotype of *Barypodus metszeri* Gilmore 1927, *nomen dubium*. **D.** USNM 11502. cf. *Tambachichnium* isp., trackway, concave epirelief. Holotype of *Barypodus tridactylus* Gilmore 1927, *nomen dubium*. Strike and dip symbols indicate the supposed dip of the inclined bedding plane, dashed arrows indicate the direction of progression. p=pes imprint. m=manus imprint.

Reptile Tracks

Only a few specimens from the Coconino Sandstone along the Mogollon Rim area are tentatively assigned to cf. *Dromopus* isp. (Marchetti et al. 2019b). This material is characterized by relatively small tracks (pes length of about 30 mm or 1.2 in) that are pentadactyl, markedly ectaxonic and semi-digitigrade. Digit impressions are long and slender and terminate in sharp claw marks. The digit I–IV impressions are proximally superimposed and distally curved inwards, and the impression of digit V is proximal and oriented outwards. The manus imprint is smaller than the pes imprint. The trackways are broad, with low pace angulation (60–100°), the pes oriented outward and the possible occurrence of a thin tail impression. The specimen UCMP-V 159262 includes a trackway with a thin tail impression and superimposed tracks, differently positioned at the two sides of the trackway. The specimen UCMP-V 159265 (Figure 9-18A) shows a trackway with superimposed pes-manus couples and superimposed digit impressions. Both trackways are directed upslope. This ichnotaxon is generally regarded as the track of araeoscelid diapsids or bolosaurid parareptiles (e.g., Voigt 2005).

The ichnogenus *Erpetopus* is relatively uncommon in the Coconino Sandstone (Marchetti et al. 2019b). This material is characterized by small tracks (pes length 10 to 15 mm, or 0.39 to 0.59 in) that are pentadactyl, ectaxonic and semi-plantigrade. Digit impressions are long and slender and terminate in sharp claw marks. The digit I–IV impressions are distally bent inwards, and the impression of digit V is relatively short and distally bent outward. The manus imprint is slightly smaller than the pes imprint and wider than long; the pes imprint is about as long as wide. The

trackways are broad or narrow, with variable pace angulation (60–125°); primary overstep can be observed, manus imprints are directed inward, and a thin tail impression may occur. The specimen USNM 11151 (Figure 9-18B) includes a partial trackway clearly assignable to this ichnogenus. Other specimens from the Mogollon Rim area (Figures 9-18C and D) include upslope-directed trackways assignable to this ichnogenus (MNA-V 3343 and MNA-V 3385). The upslope-directed trackway of USNM 11146, holotype of *Nanopus merriami* Gilmore 1926, *nomen dubium* after Marchetti et al. (2019b), is assignable to cf. *Erpetopus* isp. This track type is generally attributed to small captorhinomorph or parareptile producers (e.g., Haubold and Lucas 2003; Marchetti 2016).




Figure 9-18. Reptile footprints from the Coconino Sandstone (LORENZO MARCHETTI & SPENCER LUCAS). **A.** UCMP-V 159265, Mogollon Rim. cf. *Dromopus* isp., trackway, concave epirelief. **B.** USNM 11151, Hermit Trail. *Erpetopus* isp., pes imprint, convex hyporelief. **C.** MNA-V 3385, Mogollon Rim. *Erpetopus* isp., trackway, concave epirelief. **D.** Detail of C, left pes–manus couple. **E.** RAM-NN 1, Mogollon Rim. *Varanopus curvidactylus*, trackway, convex hyporelief. **F.** MNA-V 5193, Mogollon Rim. *Varanopus curvidactylus*, right pes–manus couple, convex hyporelief. **G.** GRCA 2914, Hermit Trail. *Varanopus curvidactylus*, trackway, concave epirelief. **H.** YPM 2143, Hermit Trail. cf. *Varanopus* isp., trackway, concave epirelief. Holotype of *Laoporus schucherti* Lull 1918, *nomen dubium*. Strike and dip symbols indicate the supposed dip of the inclined bedding plane, dashed arrows indicate the direction of progression.

The ichnotaxon *Varanopus curvidactylus* is by far the most common track type in the Coconino Sandstone (Marchetti et al. 2019b). This material is characterized by relatively small tracks (pes length 15 to 40 mm, or 0.59 to 1.6 in) that are pentadactyl, ectaxonic and semi-plantigrade. Digit impressions are long and slender and terminate in sharp claw marks. The digit I–IV impressions are distally bent inwards, and the pedal digit V impression is relatively long and straight. The manus imprint is slightly smaller than the pes imprint and wider than long; the pes imprint is about as long as wide. The trackways are broad to very narrow, with variable pace angulation (50–160°), can show primary overstep, and manus imprints are directed inward. Several specimens from the Mogollon

Rim area (Figures 9-18E and F) can be assigned to *Varanopus curvidactylus*: trackways directed upslope (MNA-V 3327, MNA-V 3470, MNA-V 5193, RAM-NN 1, UCMP-V 36852, UCMP-V 42944), trackways directed upslope with a transverse component (MNA-V 3376, RAM 236, RAM 422, RAM-NN 1, UCMP-V 42932), trackways transverse to the slope (RAM 420) and trackways directed downslope (RAM-NN 1, UCMP-V 42944). Among the material coming from the GRCA, the upslope-directed trackway of specimen GRCA 2914 (Figure 9-18G) and a pes-manus couple with partial overstep of specimen GRCA 2939 can be assigned to this ichnotaxon. Because of the poor and incomplete preservation of the holotypes, the ichnotaxa *Laoporus schucherti* Lull 1918 (YPM 2143, upslope-directed trackway) (Figure 9-18H) and *Laoporus noblei* Lull 1918 (YPM 2144, upslope-directed partial trackway) have been considered *nomina dubia,* and this material has been assigned to cf. *Varanopus* isp. (Marchetti et al. 2019b). *Varanopus curvidactylus* has generally been referred to relatively small captorhinomorph producers (e.g., Haubold and Lucas 2003), but small parareptile trackmakers are also possible.

Undetermined Tracks

Several specimens are not assignable to an ichnotaxon but are holotypes of invalid ichnotaxa (Marchetti et al. 2019b). These specimens include the holotypes of: *Dolichopodus tetradactylus* Gilmore 1926 (USNM 11123, downslope-directed trackway), *Palaeopus regularis* Gilmore 1926 (USNM 11143, upslope-directed trackway), *Barypodus palmatus* Gilmore 1926 (USNM 11134, possible pes-manus couple showing overstep), *Allopus*? *arizonae* Gilmore 1926 (USNM 11132, large-sized trackway directed upslope with transverse component), *Agostopus medius* Gilmore 1927 (USNM 11509, trackway directed upslope) and *Amblyopus pachypodus* Gilmore 1927 (USNM 11511, possible trackway).

Discussion

The Coconino Sandstone yields an extensive track collection. Because of the presence of very long trackways differently oriented compared to the paleoslope, it is the ideal unit for the reconstruction of vertebrate locomotion on desert dunes. Some specimens, such as RAM 244 from the Mogollon Rim area (Figure 9-15), are really remarkable: it includes the longest tetrapod trackway of the Paleozoic for the number of tracks (almost 200) and several trackways differently oriented compared to the slope. However, due to the effects of the locomotion on dune flanks, the track specimens useful for ichnotaxonomy are few, although significant (commonly grade 1.0–1.5 of the preservation scale, Marchetti et al. 2019a). According to Marchetti et al. (2019b), the tetrapod ichnoassociation from the Coconino Sandstone includes (Table 9-3): cf. *Amphisauropus* isp., cf. *Dromopus* isp., *Erpetopus* isp., *Ichniotherium sphaerodactylum*, cf. *Tambachichnium* isp., and *Varanopus curvidactylus*. The co-occurrence of *Ichniotherium* and *Erpetopus* suggests a late Artinskian–Kungurian age (*Erpetopus* biochron) for the Coconino Sandstone, more likely a Kungurian age because of its stratigraphic position (e.g., Blakey and Knepp 1989). The reptile tracks are very abundant and diverse, the anamniote tracks are less abundant and diverse, and the synapsid tracks are rare. This is in agreement with an early Permian low-latitude reptile radiation registered by the track record, that started during the Artinskian (Marchetti et al. 2019b).

Table 9-3. Museum collections, Coconino Sandstone.

Catalog #	Assignment	Type	Previous Ichnotaxon	Validity	Preservation	Description	Collector	Date	Locality
GRCA 2914	*Varanopus curvidactylus*	N/A	N/A	N/A	concave epirelief	trackway	Sturdevant, G. E.	1927	Grand Canyon, Hermit Trail
GRCA 2939	*Varanopus curvidactylus*	N/A	N/A	N/A	convex hyporelief	pes–manus couple	Sturdevant, G. E.	1926	Grand Canyon, Hermit Trail
GRCA 3065	*Ichniotherium sphaerodactylum*	N/A	N/A	N/A	concave epirelief	partial trackway	Byrd, J.	1934	Grand Canyon, South Kaibab Trail
GRCA 116801	cf. *Ichniotherium* isp.	N/A	N/A	N/A	convex hyporelief	partial trackway	GRCA Staff	2017	Grand Canyon, Hermit Trail
MNA-V 3327	*Varanopus curvidactylus*	N/A	N/A	N/A	convex hyporelief	trackway	–	–	Sycamore Pass
MNA-V 3343	*Erpetopus* isp.	N/A	N/A	N/A	concave epirelief	trackway and tail impression	–	–	Ash Fork
MNA-V 3376	*Varanopus curvidactylus*	N/A	N/A	N/A	convex hyporelief	trackway	–	–	Seligman, Aubrey Cliffs
MNA-V 3385	*Erpetopus* isp.	N/A	N/A	N/A	concave epirelief	trackway	–	–	Sandflat, between Bear Canyon Road and Perkinsville
MNA-V 3386	*Ichniotherium sphaerodactylum*	N/A	N/A	N/A	concave epirelief	pes–manus couple	–	–	Sandflat, between Bear Canyon Road and Perkinsville
MNA-V 3470	*Varanopus curvidactylus*	N/A	N/A	N/A	concave epirelief	trackway	–	–	Ash Fork
MNA-V 5193	*Varanopus curvidactylus*	N/A	N/A	N/A	convex hyporelief	trackway	–	–	Ash Fork
NMMNH-P 80306	*Ichniotherium sphaerodactylum*	N/A	N/A	N/A	convex hyporelief	trackway	Donated by a quarryman to Flagstaff Forest Service	1990s	Williams, commercial quarry
NMMNH-P 80307	*Ichniotherium sphaerodactylum*	N/A	N/A	N/A	concave epirelief	trackway	Donated by a quarryman to Flagstaff Forest Service	1990s	Williams, commercial quarry
RAM 236	*Varanopus curvidactylus*	N/A	N/A	N/A	convex hyporelief	two trackways	Alf, R. M.	N/A	Seligman
RAM 244	Undetermined tracks	N/A	N/A	N/A	convex hyporelief	several trackways	Alf, R. M.	N/A	Seligman
RAM 420	*Varanopus curvidactylus*	N/A	N/A	N/A	concave epirelief	two trackways	Alf, R. M.	N/A	Seligman
RAM 422	*Varanopus curvidactylus*	N/A	N/A	N/A	concave epirelief	trackway	Alf, R. M.	N/A	Seligman

Table 9-3 (continued). Museum collections, Coconino Sandstone.

Catalog #	Assignment	Type	Previous Ichnotaxon	Validity	Preservation	Description	Collector	Date	Locality
RAM-NN 1	*Varanopus curvidactylus*	N/A	N/A	N/A	concave epirelief	four trackways	Alf, R. M.	N/A	Seligman
UCMP-V 36852	*Varanopus curvidactylus*	N/A	N/A	N/A	concave epirelief	trackway	Peabody, F. E.	1940	Yavapai County, Peach Springs
UCMP-V 42932	*Varanopus curvidactylus*	N/A	N/A	N/A	concave epirelief	trackway	Kincade, S.	1940	Yavapai County, Peach Springs
UCMP-V 42944	*Varanopus curvidactylus*	N/A	N/A	N/A	concave epirelief	two trackways	–	–	Seligman
UCMP-V 42945	*Ichniotherium sphaerodactylum*	N/A	N/A	N/A	concave epirelief	trackway	–	–	–
UCMP-V 42951	*Ichniotherium sphaerodactylum*	N/A	N/A	N/A	convex hyporelief	trackway	Welles, S. P.	1949	Seligman
UCMP-V 159262	cf. *Dromopus* isp.	N/A	N/A	N/A	concave epirelief	trackway and tail impression	–	–	Yavapai County
UCMP-V 159265	cf. *Dromopus* isp.	N/A	N/A	N/A	concave epirelief	trackway	Richardson, F.	1938	Yavapai County
USNM 11123	Undetermined tracks	holotype	*Dolichopodus tetradactylus* Gilmore 1926	*nomen dubium*	concave epirelief	trackway	Gilmore, C. W.	1924	Grand Canyon, Hermit Trail
USNM 11132	Undetermined tracks	holotype	*Allopus? arizonae* Gilmore 1926	*nomen dubium*	concave epirelief	trackway	Gilmore, C. W.	1924	Grand Canyon, Hermit Trail
USNM 11134	Undetermined tracks	holotype	*Barypodus palmatus* Gilmore 1926	*nomen dubium*	concave epirelief	pes–manus couple	Gilmore, C. W.	1924	Grand Canyon, Hermit Trail
USNM 11135	cf. *Amphisauropus* isp.	holotype	*Agostopus matheri* Gilmore 1926	*nomen dubium*	concave epirelief	trackway	Gilmore, C. W.	1924	Grand Canyon, Hermit Trail
USNM 11137	*Ichniotherium sphaerodactylum*	holotype	*Baropezia eakini* Gilmore 1926	junior synonym	concave epirelief and convex hyporelief	trackway	Gilmore, C. W.	1924	Grand Canyon, Hermit Trail
USNM 11138	cf. *Ichniotherium* isp.	paratype	*Baropezia eakini* Gilmore 1926	junior synonym	concave epirelief	trackway	Gilmore, C. W.	1924	Grand Canyon, Hermit Trail
USNM 11143	Undetermined tracks	holotype	*Paleopus regularis* Gilmore 1926	*nomen dubium*	concave epirelief	trackway	Gilmore, C. W.	1924	Grand Canyon, Hermit Trail
USNM 11146	cf. *Erpetopus* isp.	holotype	*Nanopus merriami* Gilmore 1926	*nomen dubium*	concave epirelief	trackway	Gilmore, C. W.	1924	Grand Canyon, Hermit Trail
USNM 11151	*Erpetopus* isp.	N/A	N/A	N/A	convex hyporelief	partial trackway	Gilmore, C. W.	1924	Grand Canyon, Hermit Trail

Table 9-3 (continued). Museum collections, Coconino Sandstone.

Catalog #	Assignment	Type	Previous Ichnotaxon	Validity	Preservation	Description	Collector	Date	Locality
USNM 11502	cf. *Tambachichnium* isp.	holotype	*Barypodus tridactylus* Gilmore 1927	*nomen dubium*	concave epirelief and convex hyporelief	trackway	Gilmore, C. W., Metzler, A.	1926	Grand Canyon, Hermit Trail
USNM 11505	cf. *Tambachichnium* isp.	holotype	*Barypodus metszeri* Gilmore 1927	*nomen dubium*	concave epirelief	trackway	Gilmore, C. W., Metzler, A.	1926	Grand Canyon, Hermit Trail
USNM 11506	cf. *Tambachichnium* isp.	holotype	*Nanopus maximus* Gilmore 1927	*nomen dubium*	concave epirelief	trackway	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11508	*Ichniotherium sphaerodactylum*	N/A	N/A	N/A	concave epirelief	trackway	Gilmore, C. W., Metzler, A.	1927	Grand Canyon, Hermit Trail
USNM 11509	Undetermined tracks	holotype	*Agostopus medius* Gilmore 1927	*nomen dubium*	convex hyporelief	trackway	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11511	Undetermined tracks	holotype	*Amblyopus pachypodus* Gilmore 1927	*nomen dubium*	concave epirelief	trackway	Gilmore, C. W.	1926	Grand Canyon, Hermit Trail
USNM 11514	*Ichniotherium sphaerodactylum*	holotype	*Baropus coconinoensis* (Gilmore 1927)	junior synonym	concave epirelief	two pes–manus couples	Gilmore, C. W., Metzler, A.	1926	Grand Canyon, Hermit Trail
YPM 288	*Ichniotherium sphaerodactylum*	N/A	N/A	N/A	concave epirelief	partial trackway	Gidley, G. W., Wetmore, A.	–	Grand Canyon, Hermit Trail
YPM 2143	cf. *Varanopus* isp.	holotype	*Laoporus schucherti* Lull 1918	*nomen dubium*	concave epirelief	trackway	Schuchert, C.	1915	Grand Canyon, Hermit Trail
YPM 2144	cf. *Varanopus* isp.	holotype	*Laoporus noblei* Lull 1918	*nomen dubium*	concave epirelief	partial trackway	Schuchert, C.	1915	Grand Canyon, Hermit Trail

The finding of ichnoassociations typical of the *Dromopus* biochron in Ghzelian eolian units such as the Wescogame Formation (this work) and ichnoassociations typical of the *Erpetopus* biochron in Kungurian desert units such as the Coconino and De Chelly Sandstones (Marchetti et al. 2019b) suggests a facies-crossing reptile radiation and gives strength to the tetrapod footprint biostratigraphy. Some comments can be presented on the ichnofaunal composition of the Coconino Sandstone compared with the underlying Hermit Shale (this work). Both of the ichnoassociations include abundant and diverse reptile tracks, rare synapsid tracks and belong to the *Erpetopus* footprint biochron. Noteworthy, the most abundant track is *Hyloidichnus* in the Hermit Shale and *Varanopus* in the Coconino Sandstone, and both are lacking in the other formation. The reptile tracks *Erpetopus* and *Dromopus* are instead similarly distributed. The difference in anamniote tracks is noteworthy: small temnospondyl tracks (*Batrachichnus*) are very abundant in the Hermit Shale, while they are absent in the Coconino Sandstone; diadectomorph tracks (*Ichniotherium*) are relatively abundant in the Coconino Sandstone, but rare in the Hermit Shale (also, the *Ichniotherium* ichnospecies is different). *Amphisauropus* (seymouriamorph tracks) has a similar distribution in both units. With regard to the synapsid tracks, the ichnogenus is different: *Dimetropus* occurs only in the Hermit Shale, and *Tambachichnium* occurs only in the Coconino Sandstone. These differences are probably not related to stratigraphy, because all the Hermit Shale ichnogenera occur in younger units, and all the Coconino Sandstone ichnogenera occur in older units (e.g., Voigt and Lucas 2018; Marchetti et al. 2019b). Therefore, this seems to be due to a different paleoecology of the trackmakers that differentially occupied the floodplain (Hermit Shale) and desert (Coconino Sandstone) paleoenvironments. In this context, the non-occurrence of *Hyloidichnus*, *Batrachichnus* and *Dimetropus* in the Coconino Sandstone may have a paleoecological meaning, as does the non-occurrence of *Varanopus* and *Tambachichnium* in the Hermit Shale.

As mentioned above, the Coconino Sandstone is a unit deposited mainly by dunes of eolian origin intercalated with somewhat rare interdune deposits, representing a large erg located at low latitudes during the Kungurian. Since the Paleozoic, the tetrapod ichnological record of erg deposits is notoriously dominated by tracks of Amniota (i.e., eureptiles, parareptiles and synapsids) (e.g., Francischini et al. 2015; Krapovickas et al. 2015, 2016; Marchetti et al. 2019b, 2019c), but the Coconino Sandstone ichnoassociation is remarkable in containing two ichnotaxa (cf. *Amphisauropus* isp. and *Ichniotherium sphaerodactylum*) assigned to non-amniote tetrapods (Francischini et al. 2019; Marchetti et al. 2019b). *Amphisauropus* and *Ichniotherium* are considered to be seymouriamorph and diadectomorph (both reptiliomorph) tracks, and the presence of these ichnogenera in the Coconino Sandstone may represent the oldest record of anamniotes in desert environments and one of the oldest in eolian lithofacies (other occurrences are known from the Wescogame and De Chelly formations, Marchetti et al. 2019b and this work). This record suggests that the adaptations for living in arid settings could have appeared early in the Reptiliomorpha lineage, during the early Permian. Contrasting to this, the Coconino Sandstone lacks evidences of temnospondyl (Batrachomorpha lineage) tracks, such as *Batrachichnus* and *Limnopus*, which occur in the Wescogame Formation and Hermit Shale. Temnospondyl (including lissamphibian) tracks are still poorly known from eolian deposits (this work), and the colonization of arid settings by these tetrapods is an issue still to be investigated.

Hunt and Lucas (2005b, 2006, 2007, 2016) named the *Chelichnus* ichnofacies based on the now discredited interpretations of the ichnotaxonomy of the Coconino Sandstone and other Permian ichnofaunas of Arizona (De Chelly), and Colorado (Lyons) in the United States, Scotland, and Germany in Europe (as well as Late Triassic–Early Jurassic ichnofaunas of western North America which are in need of re-study: Hunt and Lucas 1998, 2005b, 2007). Ichnofacies retain their name even if the name-bearing ichnotaxon is shown to be a *nomen dubium*, although it would be preferable to use a valid ichnotaxon actually typical of that ichnofacies (e.g., *Varanopus*, *Procolophonichnium*, *Brasilichnium*). Does the *Chelichnus* ichnofacies of the Coconino still have utility? In a biotaxonomic sense it can now be recognized as a relatively depauperate ichnofauna lacking ichnotaxa, such as *Batrachichnus*, *Limnopus*, *Dimetropus* and *Hyloidichnus* and with a noteworthy abundance of *Varanopus*, relative to contemporary units (Marchetti et al. 2019b: Figure 13). It arguably has more significance as an ethoichnofacies (*sensu* Hunt and Lucas 2007), which reflects locomotor behavior such as digit drag marks and also as a taphofacies preserving features such as uni-directional ichnotaphonomic effects due to the slope direction (Marchetti et al. 2019b, 2019c). There is clearly a need to re-evaluate this ichnofacies.

Conclusions

The tetrapod ichnological record of the Grand Canyon and Mogollon Rim areas of Arizona is remarkable. Continental deposits are rare among the geological units of these areas and their Paleozoic vertebrate fossil record is restricted to tetrapod tracks of the Upper Pennsylvanian–lower Permian Manakacha Formation, Wescogame Formation, Hermit Shale and Coconino Sandstone (Figure 9-19). The copious and variably preserved tetrapod track record of the Wescogame Formation, Hermit Shale and Coconino Sandstone is composed of non-amniote, synapsid and reptile tracks and trackways produced in marginal marine, alluvial plain and desert settings. On the other hand, the Manakacha Formation record is still scant and indeterminate, but it represents a potential new window for future important discoveries.

The Grand Canyon tetrapod track record is unique in recording several facies-crossing ichnotaxa, leading to a re-interpretation of some vertebrate ichnofacies. Among these ichnotaxa, *Amphisauropus* and *Ichniotherium* may represent the oldest colonization of arid environments by non-amniote tetrapods. In addition, the Artinskian–Kungurian Hermit Shale and Coconino Sandstone ichnoassociations suggest that those paleoenvironments were dominated by reptiles (i.e., parareptiles, araeoscelids and captorhinomorphs), contrasting with older ichnoassociations found worldwide, where these trackmakers are rarer. These are only a few examples of how the tetrapod track record can contribute to biostratigraphic, paleobiological, and evolutionary problems of the fossil record. In spite of the fact that tracks have been collected in the Grand Canyon from over a century, new findings and re-interpretations make it an important area for understanding the composition and diversity of Paleozoic continental tetrapods.




Figure 9-19. Stratigraphic distribution of vertebrate tracks in the Paleozoic formations of GRCA (LORENZO MARCHETTI & SPENCER LUCAS). Gray lines and dots in the upper chart indicate uncertain occurrence. **1–10.** Tetrapod ichnogenera: **1.** *Batrachichnus*; **2.** *Limnopus*; **3.** *Amphisauropus*; **4.** *Ichniotherium*; **5.** *Dimetropus*; **6.** *Tambachichnium*; **7.** *Erpetopus*; **8.** *Varanopus*; **9.** *Hyloidichnus*; **10.** *Dromopus*. **I–VI.** Trackmaker groups: **I.** Temnospondyl anamniotes; **II.** Reptiliomorph anamniotes; **III.** Non-therapsid synapsids; **IV.** Parareptiles; **V.** Captorhinid eureptiles; **VI.** Diapsid eureptiles.

Acknowledgments

We are indebted to Thomas Martens, who kindly provided photographs and information on his discovery of *Ichniotherium cottae* in the Hermit Shale as well as permitted publication of the material. We extend additional thanks to Klara Widrig, Colleen Hyde and Kim Besom for assistance with photography of the vertebrate ichnofossil collections in the GRCA park museum collection. We also thank the staff of MNA, RAM, UCMP, USNM, and YPM for the help during the study of the collections. We thank the reviewers, Hendrik Klein and Andrew Milner, for their helpful comments. Heitor Francischini was funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES/PNPD; Grant Number: 88882.316166/2019-01). Lorenzo Marchetti was funded by the Alexander von Humboldt Foundation.

Literature Cited

Allen, J. R. L. 1997. Subfossil mammalian tracks (Flandrian) in the Severn Estuary, S.W. Britain: Mechanics of formation, preservation and distribution. Philosophical Transactions of the Royal Society of London B 352:481–518.

Baird, D. 1952. Revision of the Pennsylvanian and Permian footprints *Limnopus*, *Allopus* and *Baropus*. Journal of Paleontology 26:832–840.

Blakey, R. C. 1990. Supai Group and Hermit Formation. Pages 147–182 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology. Oxford University Press, New York, New York.

Blakey, R. C., and R. Knepp. 1989. Pennsylvanian and Permian geology of Arizona. Pages 313–347 *in* J. P. Jenney and S. J. Reynolds, editors. Geologic evolution of Arizona. Arizona Geological Society Digest 17.

Brand, L. R. 1979. Field and laboratory studies on the Coconino Sandstone (Permian) vertebrate footprints and their paleoecological implications. Palaeogeography, Palaeoclimatology, Palaeoecology 28:25–38.

Brand, L. R., and T. Tang. 1991. Fossil vertebrate footprints in the Coconino Sandstone (Permian) of northern Arizona: Evidence for underwater origin. Geology 19:1201–1204.

Buchwitz, M., and S. Voigt. 2018. On the morphological variability of *Ichniotherium* tracks and evolution of locomotion in the sister group of amniotes. PeerJ 6:e4346. Available at: https://doi.org/10.7717/peerj.4346 (accessed February 2020).

Citton, P., E. Sacchi, and U. Nicosia. 2012. Sometimes they come back: Recovery and reinterpretation of a trackway slab from the Permian Coconino Sandstone of the southwestern United States. Ichnos 19:165–174.

Fillmore, D. L., S. G. Lucas, and E. L. Simpson. 2012. Ichnology of the Mississippian Mauch Chunk Formation, eastern Pennsylvania. New Mexico Museum of Natural History and Science Bulletin 54.

Francischini, H., P. C. Dentzien-Dias, M. A. Fernandes, and C. L. Schultz. 2015. Dinosaur ichnofauna of the Upper Jurassic/Lower Cretaceous of the Paraná Basin (Brazil and Uruguay). Journal of South American Earth Sciences 63:180–190.

Francischini, H., S. G. Lucas, S. Voight, L. Marchetti, V. L. Santucci, C. L. Knight, J. R. Wood, P. Dentzien-Dias, and C. L. Schultz. 2019. On the presence of *Ichniotherium* in the Coconino Sandstone (Cisuralian) of the Grand Canyon and remarks on the occupation of deserts by non-amniote tetrapods. Paläontologische Zeitschrift online:1–19.

Gand, G., and M. Durand. 2006. Tetrapod footprint ichno-associations from French Permian basins. Comparisons with other Euramerican ichnofaunas. Pages 157–177 *in* S. G. Lucas, G. Cassinis, and J. W. Schneider, editors. Nonmarine Permian biostratigraphy and biochronology. Geological Society of London, London, United Kingdom. Special Publications 265.

Geinitz, H. B. 1861. Dyas oder die Zechsteinformation und das Rhotliegende-Die anamalischen Uberreste der Dyas: Wilhem Engelmann, Leipzig, Germany.

Gilmore, C. W. 1926. Fossil footprints from the Grand Canyon. Smithsonian Miscellaneous Collections 77(9).

Gilmore, C. W. 1927. Fossil footprints from the Grand Canyon: Second contribution. Smithsonian Miscellaneous Collections 80(3).

Gilmore, C. W. 1928. Fossil footprints from the Grand Canyon: Third contribution. Smithsonian Miscellaneous Collections 80(8).

Haubold, H. 1971. Ichnia amphibiorum et reptiliorum fossilium. Handbuch der Paleoherpetologie, Teil 18. Gustav Fischer Verlag, Stuttgart, Germany.

Haubold, H. 2000. Tetrapodenfährten aus dem Perm—Kenntnisstand und Progress 2000. Hallesches Jahrbuch fur Geowissenschaften B 22:1–16.

Haubold, H., and S. G. Lucas. 2003. Tetrapod footprints of the Lower Permian Choza Formation. Paläontologische Zeitschrift 77:247–261.

Haubold, H., A. P. Hunt, S. G. Lucas, and M. G. Lockley. 1995a. Wolfcampian (Early Permian) vertebrate tracks from Arizona and New Mexico. New Mexico Museum of Natural History and Science Bulletin 6:135–165.

Haubold, H., M. G. Lockley, A. P. Hunt, and S. G. Lucas. 1995b. Lacertoid footprints from Permian dune sandstones, Cornberg and De Chelly Sandstones. New Mexico Museum of Natural History and Science Bulletin 6:235–244.

Hunt, A. P., and S. G. Lucas. 1998. Implications of the cosmopolitanism of Permian tetrapod ichnofaunas. New Mexico Museum of Natural History and Science Bulletin 12:55–57.

Hunt, A. P., and S. G. Lucas. 2005a. Stratigraphic distribution of ichnofossils in the Coconino Sandstone (Permian: Leonardian) of Grand Canyon National Park and the Seligman/Ash Fork area, northern Arizona, USA. New Mexico Museum of Natural History and Science Bulletin 30:132.

Hunt, A. P., and S. G. Lucas. 2005b. Tetrapod ichnofacies and their utility in the Paleozoic. Pages 113–119 *in* R. J. Buta, A. K. Rindsberg, and D. C. Kopaska-Merkel, editors. Pennsylvanian footprints in the Black Warrior Basin of Alabama. Alabama Paleontological Society, Birmingham, Alabama. Monograph 1.

Hunt, A. P., and S. G. Lucas. 2006. Permian tetrapod ichnofacies. Pages 137–156 *in* S. G. Lucas, G. Cassinis, and J. W. Schneider, editors. Non-marine Permian biostratigraphy and biochronology. Geological Society of London, London, United Kingdom. Special Publication 265.

Hunt, A. P., and S. G. Lucas. 2007. Tetrapod ichnofacies: a new paradigm. Ichnos 14:59–68.

Hunt, A. P., and S. G. Lucas. 2016. The case for archetypal vertebrate ichnofacies. Ichnos 23:237–247.

Hunt, A. P., and V. L. Santucci. 2001. An unusual tetrapod track morphology from the Permian Coconino Sandstone, Grand Canyon National Park, Arizona. Pages 44–47 *in* V. L. Santucci and L. McClelland, editors. Proceedings of the 6th Fossil Resource Conference. NPS Geological Resources Division, Denver, Colorado. Technical Report NPS/NRGRD/GRDTR-01/01.

Krapovickas, V., C. A. Marsicano, A. C. Mancuso, M. S. de la Fuente, and E. G. Ottone. 2015. Tetrapod and invertebrate trace fossils from aeolian deposits of the lower Permian of central western Argentina. Historical Biology 27:827–842.

Krapovickas, V., M. G. Mángano, L. A. Buatois, and C. A. Marsicano. 2016. Integrated ichnofacies models for deserts: recurrent patterns and megatrends. Earth-Science Reviews 157:61–85.

International Commission on Zoological Nomenclature. 1999. International Code of Zoological Nomenclature (4th edition). International Trust for Zoological Nomenclature, London, United Kingdom.

Leonardi, G., editor. 1987. Glossary and manual of tetrapod footprint palaeoichnology. Departamento Nacional da Produção Mineral, Brasilia, Brazil.

Lockley, M. G. 1992. Comment and reply on "Fossil vertebrate footprints in the Coconino sandstone (Permian) of northern Arizona: evidence for underwater origin". Geology 20:666–670.

Lockley, M. G., A. P. Hunt, and C. Meyer. 1994. Vertebrate tracks and the ichnofacies concept: Implications for paleoecology and palichnostratigraphy. Pages 241–268 *in* K. Donovan, editor. The palaeobiology of trace fossils. John Wiley & Sons, Chichester, United Kingdom.

Loope, D. B. 1992. Comment and reply on "Fossil vertebrate footprints in the Coconino sandstone (Permian) of northern Arizona: evidence for underwater origin". Geology 20:666–670.

Lucas, S. G. 2007. Tetrapod footprint biostratigraphy and biochronology. Ichnos 14:5–38.

Lucas, S. G., K. Krainer, and R. J. Brose. 2013. The lower Permian Glorieta Sandstone in central New Mexico. New Mexico Museum of Natural History and Science Bulletin 59:201–211.

Lull, R. S. 1918. Fossil footprints from the Grand Canyon of the Colorado. American Journal of Science (4th series) 45:337–346.

Marchetti, L. 2016. New occurrences of tetrapod ichnotaxa from the Permian Orobic Basin (northern Italy) and critical discussion of the age of the ichnoassociation. Papers in Palaeontology 2:363–386.

Marchetti, L., A. Ronchi, G. Santi, and S. Voigt. 2015a. The Gerola Valley site (Orobic Basin, Northern Italy): A key for understanding late Early Permian tetrapod ichnofaunas. Palaeogeography, Palaeoclimatology, Palaeoecology 439:97–116.

Marchetti, L., A. Ronchi, G. Santi, P. Schirolli, and M. A. Conti. 2015b. Revision of a classic site for Permian tetrapod ichnology (Collio Formation, Trompia and Caffaro valleys, N. Italy), new

evidences for the radiation of captorhinomorph footprints. Palaeogeography, Palaeoclimatology, Palaeoecology 433:140–155.

Marchetti, L., E. Mujal, and M. Bernardi. 2017a. An unusual *Amphisauropus* trackway and its implication for understanding seymouriamorph locomotion. Lethaia 50:162–174.

Marchetti, L., A. Tessarollo, F. Felletti, and A. Ronchi. 2017b. Tetrapod footprint paleoecology: Behavior, taphonomy and ichnofauna disentangled. A case study from the Lower Permian of the southern Alps (Italy). Palaios 32:506–527.

Marchetti, L., M. Belvedere, S. Voigt, H. Klein, D. Castanera, I. Díaz-Martínez, D. Marty, L. Xing, S. Feola, R. N. Melchor, and J. O. Farlow. 2019a. Defining the morphological quality of fossil footprints. Problems and principles of preservation in tetrapod ichnology with examples from the Palaeozoic to the present. Earth-Science Reviews 193:109–145.

Marchetti, L., S. Voigt, S. G. Lucas, H. Francischini, P. Dentzien-Dias, R. Sacchi, M. Mangiacotti, S. Scali, A. Gazzola, A. Ronchi, and A. Millhouse. 2019b. Tetrapod ichnotaxonomy in eolian paleoenvironments (Coconino and De Chelly formations, Arizona) and late Cisuralian (Permian) sauropsid radiation. Earth-Science Reviews 190:148–170.

Marchetti, L., S. Voigt, and S. G. Lucas. 2019c. An anatomy-based study of the Lopingian eolian tracks of Germany and Scotland reveals the first evidence of the end-Guadalupian mass extinction at low paleolatitudes of Pangea. Gondwana Research 73:32–53.

Marchetti, L., H. Klein, M. Buchwitz, A. Ronchi, R. M. Smith, W. J. De Klerk, L. Sciscio, and G. H. Groenewald. 2019d. Permian-Triassic vertebrate footprints from South Africa: Ichnotaxonomy, producers and biostratigraphy through two major faunal crises. Gondwana Research 72:139–168.

Marsh, O. C. 1894. Footprints of vertebrates in the Coal Measures of Kansas. American Journal of Science (3rd series) 48:81–85.

Matthew, G. F. 1903. New genera of batrachian footprints of the Carboniferous System in Eastern Canada. Canadian Record of Science 9:100–111.

McKee, E. D. 1944. Tracks that go uphill. Plateau 16:61–73.

McKee, E. D. 1947. Experiments on the development of tracks in fine cross-bedded sand. Journal of Sedimentary Research 17:23–28.

McKee, E. D. 1979. Ancient sandstones considered to be eolian. Pages 187–238 *in* E. D. McKee, editor. A study of global sand seas. U.S. Geological Survey, Washington, D.C. Professional Paper 1052.

McKee, E. D. 1982. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKeever, P. J., and H. Haubold. 1996. Reclassification of vertebrate trackways from the Permian of Scotland and related forms from Arizona and Germany. Journal of Paleontology 70:1011–1022.

Middleton, L. T., D. K. Elliott, and M. Morales. 1990. Coconino Sandstone. Pages 352–368 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology. Oxford University Press, New York, New York.

Mujal, E., J. Fortuny, O. Oms, A. Bolet, À. Galobart, and P. Anadón. 2016. Palaeoenvironmental reconstruction and early Permian ichnoassemblage from the NE Iberian Peninsula (Pyrenean Basin). Geological Magazine 153:578–600.

Noble, L. F. 1922. A section of the Paleozoic formations of the Grand Canyon at Bass Trail. Pages 23–73 *in* Shorter contributions to general geology, 1922. U.S. Geological Survey, Washington, D.C. Professional Paper 131-B.

Rowland, S. M. 2017. Trackway of a sideways-walking pelycosaur in the Carboniferous Manakacha Formation in Grand Canyon National Park. Abstracts with Programs - Geological Society of America 49(6):abstract 136-12.

Rowland, S. M., and M. V. Caputo. 2018. Trackway of a sideways-walking basal tetrapod in the Pennsylvanian Manakacha Formation of Grand Canyon National Park. Journal of Vertebrate Paleontology, Program and Abstracts, 2018:206.

Spamer, E. E. 1984. Paleontology in the Grand Canyon of Arizona: 125 years of lessons and enigmas from the late Precambrian to the present. The Mosasaur 2:45–128.

Voigt, S. 2005. Die Tetrapodenichnofauna des kontinentalen Oberkarbon und Perm im Thüringer Wald – Ichnotaxonomie, Paläoökologie und Biostratigraphie. Cuvillier, Göttingen, Germany.

Voigt, S., and H. Haubold. 2015. Permian tetrapod footprints from the Spanish Pyrenees. Palaeogeography, Palaeoclimatology, Palaeoecology 417:112–120.

Voigt, S., and S. G. Lucas. 2017. Early Permian tetrapod footprints from central New Mexico. New Mexico Museum of Natural History and Science Bulletin 77:333–352.

Voigt, S., and S. G. Lucas. 2018. Outline of a Permian tetrapod footprint ichnostratigraphy. Pages 387–404 *in* S. G. Lucas and S. Z. Shen, editors. The Permian Timescale. Geological Society of London, London, United Kingdom. Special Publication 450.

Voigt, S., D. S. Berman, and A. C. Henrici. 2007. First well-established trackmaker association of Paleozoic tetrapods based on *Ichniotherium* trackways and diadectid skeletons from the Lower Permian of Germany. Journal of Vertebrate Paleontology 27:553–570.

Voigt, S., A. Hminna, H. Saber, J. W. Schneider, and H. Klein. 2010. Tetrapod footprints from the uppermost level of the Permian Ikakern Formation (Argana Basin, western High Atlas, Morocco). Journal of African Earth Sciences 57:470–478.

Voigt, S., A. Lagnaoui, A. Hminna, H. Saber, and J. W. Schneider. 2011. Revisional notes on the Permian tetrapod ichnofauna from the Tiddas Basin, central Morocco. Palaeogeography, Palaeoclimatology, Palaeoecology 302:474–483.

White, D. 1929. Flora of the Hermit Shale, Grand Canyon, Arizona. Carnegie Institute of Washington Publication 405.

Chapter 10. Mesozoic Paleontology of Grand Canyon National Park: Trace Fossils, Stratigraphy, and Regional Correlations

By Adam D. Marsh[1], William G. Parker[1], and Anne E. Miller[2]

[1]Petrified Forest National Park
Division of Science and Resource Management
1 Park Road #2217
Petrified Forest, Arizona 86028

[2]Grand Canyon National Park
Division of Science and Resource Management
1824 S Thompson St.
Flagstaff, Arizona 86001

Introduction

The only Mesozoic strata found within the boundary of Grand Canyon National Park (GRCA) are the Triassic Moenkopi Formation and Shinarump Member of the Chinle Formation (Figure 10-1). These Triassic rocks are best exposed at Cedar Mountain in eastern GRCA. An additional kilometer-thick section of later Mesozoic (Jurassic and Cretaceous) sedimentary rocks were once present in the region, but have eroded away as the Colorado Plateau continues to be uplifted (Flowers 2010; Liu and Gurnis 2010). Until this report, the only known paleontological resources of the Mesozoic section at GRCA were petrified logs in the Shinarump Member (Noble 1922).

The Shinarump Member of the Chinle Formation has always been recognized in the region as a cliff-forming conglomerate containing fossil logs (Powell 1873, 1876; Gilbert 1875; Dutton 1880), but other units have often been conflated or synonymized with it. The variegated finer beds above and below the "Shinarump Conglomerate" were first called the "Shinarump Group" (Powell 1876:54); the "lower portion" of Powell's Shinarump Group was later removed from that unit and named the "Moencopie beds" (Walcott 1880; Ward 1901:17). The spelling later changed to "Moenkopi Formation" (Gregory 1916:79), which has since been carried through the literature. Powell considered all of his "Shinarump Group" to be Triassic in age owing to the fossil logs, although once the "Moencopie beds" were removed from that group they were assigned to the Permian, because of the presence of supposed Permian-aged fossil plant impressions (Gregory 1916, 1917). The organization and formalization of the "Shinarump Conglomerate" and "Moencopie beds" is detailed below in the overview of Triassic strata at GRCA. A marked unconformity is present between these units throughout the region and represents the Tr-3 unconformity between the Middle Triassic and Upper Triassic series (Pipiringos and O'Sullivan 1978). The relatively hard Shinarump Member of the Chinle Formation "preserved the cliffs of the Moenkopi from destruction" (Gregory 1917:38) where the upper part of the Moenkopi Formation crops out from Houck, Arizona, through Holbrook and Radar Mesa, and along the Little Colorado River Valley towards Cedar Mountain.




Figure 10-1. Geographic location and stratigraphic section of Cedar Mountain, Grand Canyon National Park (NPS/MARSH ET AL.). **A.** Map of Arizona and southern Utah showing various locations discussed in the text. **B.** Stratigraphic section of the Mesozoic sedimentary rocks at Cedar Mountain, including the measured section of Noble (1922). **C.** Annotated view of the southwestern slope of Cedar Mountain. **D.** Geologic map of the Mesozoic section at Cedar Mountain (Billingsley et al. 2012).

Here we present a review of the Triassic sedimentary rocks of GRCA, their regional correlations, and the trace and body fossils found in the area. We also present the results of a two-day PaleoBlitz (an abbreviated paleontological inventory), which represents the second and most comprehensive paleontological inventory of the Mesozoic section at GRCA since shortly after the park was established (Noble 1922). These new data increase the fossil diversity of GRCA and shed light on an often-overlooked paleontological story in the park that chronicles an important change to the history of the area.

Institutional Abbreviations—**GRCA**, Grand Canyon National Park, Arizona; **MNA**, Museum of Northern Arizona, Flagstaff, Arizona; **PEFO**, Petrified Forest National Park, Arizona; **UCMP**, University of California Museum of Paleontology, Berkeley, California; **USNM**, National Museum of Natural History, Smithsonian Institution, Washington, D.C.; **WASO**, Washington D.C. Area Support Office, National Park Service, Geological Resources Division, Washington, D.C.; **WUPA**, Wupatki National Monument, Arizona.

Overview of Triassic Strata at GRCA

Moenkopi Formation

Geology

The "Moencopie beds" were separated from the former "Shinarump Group" based on differences in fossil invertebrates (Walcott 1880) within a succession of "dark reddish brown, soft, laminated, argillaceaous shales…highly charged with salt and gypsum" (Ward 1901:403) below the Shinarump Member of the Chinle Formation near Moenkopi Wash, Arizona (~37 km or 23 mi southeast of Cedar Mountain). The type section of the Moenkopi Formation is located in northern Arizona near Tanner Crossing in Cameron, Arizona and includes nearly 122 m (400 ft) of irregular beds of sandstone and shale with frequent horizontal and vertical gypsum beds and a calcareous base (Gregory 1917:23). The Moenkopi Formation at its type section sits unconformably over the Kaibab Formation and under the Shinarump Member of the Chinle Formation, just as it does at Cedar Mountain in GRCA (Gregory 1917; Noble 1922), although the section at Cedar Mountain is nearly 30 m (100 ft) thicker than the type section.

The Moenkopi Formation has different member-level divisions at various places in its outcrop belt in Colorado, New Mexico, Arizona, Utah, and Nevada. The formation thickens basinward to the northwest along the paleodrainage of a forearc basin east of the Cordilleran magmatic arc (McKee 1954; Dickinson and Gehrels 2008, 2009a). Exposures of the Moenkopi Formation north and west of the Canyon include the marginal marine Timpoweap Member, Virgin Limestone Member, Shnabkaib Member, and interspersed lower, middle, and upper "red members" (Gregory 1948, 1952; Reeside and Bassler 1922), and exposures south and east of the Canyon include prograding fluvial systems within the Wupatki Member, Moqui Member, and Holbrook Member (McKee 1954; Repenning et al. 1969; Stewart et al. 1972b).

The Moenkopi Formation section at Cedar Mountain is most similar to the eastern exposures along the Little Colorado River Valley. The Wupatki Member is best exposed east of the Black Point Monocline at Wupatki National Monument (~65 km or 40 mi southeast of Cedar Mountain; McKee

1954; Billingsley et al. 2007; Henderek et al. 2017). The Wupatki Member is characterized by "pale-reddish-brown, micaceous, horizontally and ripple-laminated siltstone that weathers to slopes" (Stewart et al. 1972b:7). The Moqui Member was named for the low badlands in the middle part of the Moenkopi Formation just west of Winslow, Arizona (McKee 1954). The Moqui Member in the eastern outcrops of the Moenkopi Formation is most similar to the Shnabkaib Member found in the western outcrops in that they are both very fine grained with numerous gypsum beds, but the Moqui Member lacks the sabkha-like cyclicity found in the type section of the Shnabkaib Member in Warner Valley, Utah (McKee 1954, 1964). The Moqui Member in the Little Colorado River Valley is characterized by "pale brown gypsiferous mudstone and siltstone beds" and is generally "a lighter shade of brown" than the rest of the section above and below it (Stewart et al. 1972b:8). The type section of the Holbrook Member occurs between Holbrook and Joseph City, Arizona (Hager 1922), where it crops out as "pale-red thin- to thick-bedded channel deposits of sandstone" that are very fine to medium-grained, poorly sorted, trough cross-bedded, and rich in siltstone (Stewart et al. 1972b:9). Closer to Cameron, the Holbrook Member contains thin beds of calcareous cross-bedded sandstones. The contacts between the three major members discussed here are defined lithologically; the lower contact of the Moqui Member with the Wupatki Member occurs just below the lowest gypsum bed, and the upper contact of the Moqui Member with the Holbrook Member occurs just above the highest gypsum bed (McKee 1954; Stewart et al. 1972b; Morales 1987).

Two prominent sandstone ledges within the Moenkopi Formation extend along the roughly east-to-west exposures throughout the Little Colorado River Valley. The "lower massive sandstone" occurs in the upper part of the Wupatki Member, and the "upper massive sandstone" occurs in the lower part of the Holbrook Member (McKee 1954). The lower massive sandstone near Cameron (i.e., Cedar Mountain) is a prominent ledge-forming, very fine grained, well-sorted sandstone that is 3 to 9 m (10 to 30 ft) thick, nearly 23 m (75 ft) from the base of the formation (Stewart et al. 1972b). This ledge-forming sandstone has been traced throughout the area from Wupatki National Monument north to the Echo Cliffs and south to Meteor Crater. The upper massive sandstone is not well documented in the Cameron area, but it is well reported just west of Holbrook, Arizona, where it forms a 13 m (43 ft) thick, fine-grained sandstone bench nearly 18 m (60 ft) above the base of the member (Welles 1947; McKee 1954).

Continental redbeds such as the Moenkopi Formation and Shinarump Conglomerate are notoriously difficult to date to specific Triassic geochronological time bins, because those are defined by marine biostratigraphy. However, a combination of vertebrate and trace fossil biostratigraphy and radiometric dating support the hypothesis that the Moenkopi Formation is Early Triassic to early Middle Triassic in age (see below). Marcou (1858) first estimated the age of the Moenkopi Formation to be Triassic owing to its lithological similarity to the Triassic Bunter sandstone in Germany, but Newberry (1861) later estimated the unit to be Permian, because the underlying sedimentary rocks were considered to be Carboniferous (Dutton 1882). Powell (1876) considered what is now the Moenkopi Formation in northwestern Arizona and southwestern Utah to be Triassic–Jurassic owing to the presence of brachiopods, crinoids, and bivalves in limestone beds in the lower part of the formation, but Walcott (1880) thought these fossils (and the strata) to be late Permian in age. Most researchers concurred with this designation (see discussion in McKee 1954:8), except for

Shimer (1919) who thought the fossils were Triassic in age, until vertebrate body and trace fossils were found throughout the formation along the Little Colorado River Valley (Welles 1947; McKee 1954).

Where the marine and marginal marine members of the Moenkopi Formation are exposed in eastern Nevada, they contain the Early Triassic *Meekoceras* ammonite zones (McKee 1954; Ogg 2012). The Moenkopi Formation between Holbrook and Cameron is quite fossiliferous in discrete stratigraphic horizons. Vertebrate fossils in the lower fossiliferous horizon occur in bonebeds of associated skeletons above the "lower massive sandstone" in the upper part of the Wupatki Member (exemplified by the UCMP locality V3835 near Meteor Crater; Nesbitt 2005a) and include actinopterygian fish, mastodonsauroid and trematosaurian temnospondyls, and non-archosaur archosauromorphs such as tanystropheids (Welles 1969, 1993; Nesbitt 2005a, 2005b). Fossils in the upper fossiliferous horizon occur as disarticulated skeletal elements or isolated partial skeletons just underneath the "upper massive sandstone" in the Holbrook Member (exemplified by the UCMP locality V3922 near Holbrook; Nesbitt 2000, 2005a, 2005b) and include sharks, actinopterygian and sarcopterygian fish, mastodonsauroid and brachyopid temnospondyls, and pseudosuchian archosaurs (Welles 1947, 1969; McKee 1954; Nesbitt 2000, 2002, 2005a, 2005b). Invertebrate and vertebrate trace fossils are found on bedding planes in fluvial sandstones, are often associated with sedimentary structures such as ripples, mud cracks, and flute casts, and are concentrated in similar horizons as vertebrate body fossils (McKee 1954; Klein and Lucas 2010b) in the Wupatki Member and Holbrook Member. Trace fossils within the Moenkopi Formation include invertebrate burrows, arthropod tracks, and tetrapod tracks, many of which were collected from northern Arizona and are housed at the UCMP and MNA.

The most recent vertebrate biostratigraphy suggests that at least the eastern terrestrial outcrops of the Holbrook Member are as young as the early Middle Triassic (Lucas 2010) based on the presence of the capitosaurid temnospondyl *Eocyclotosaurus*. This age estimate is supported by the presence of the tetrapod ichnogenera *Rotodactylus* and *Chirotherium* within the Holbrook Member; these ichnotaxa arguably only occur in the Middle Triassic (McKee 1954; Klein and Lucas 2010a; Henderek et al. 2017; see Brusatte et al. 2011 and Niedźwiedzki et al. 2013 for a possible Early Triassic record of cf. *Rotodactylus* isp.). Owing to the presence of Cordilleran magmatic arc-derived zircon grains incorporated into the prograding terrestrial Moenkopi–Chinle deposystems, U-Pb geochronology (either LA-ICP-MS or high-resolution CA/ID-TIMS) has been particularly effective in inferring the maximum depositional ages of these rocks. A detrital zircon sample from the Holbrook Member of the Moenkopi Formation (Dickinson and Gehrels 2009b) suggests that it may have been deposited as late as the early Late Triassic (Carnian, ~230 Ma), which is at least 10 million years later than what vertebrate biochronology suggests (Cohen et al. 2013).

Previous studies

The complete section of the Moenkopi Formation at Cedar Mountain in GRCA is important for determining the fluvial facies distribution across the aerial extent of the formation, because it is one of the only complete sections between the western marginal marine facies and the eastern fluvial facies. When it was first measured and described, the Moenkopi Formation exposed at Cedar

Mountain was not differentiated into member-level units owing to vegetative cover (Noble 1922); however, it was subsequently mapped as including the Wupatki Member, Shnabkaib Member, and Holbrook Member (Billingsley et al. 2012).

Noble (1922:72) noted a thin (0 to 1.2 m, 0 to 4 ft) "curious gnarly [conglomeratic] cherty bed" near the base of the section, which compares favorably to a similar bed found at the base of the Moenkopi Formation at Lee's Ferry (McKee 1954; Stewart et al. 1972b) that may represent the southeastern most expression of the Timpoweap Member (Blakey et al. 1993). However, similar cherty conglomeratic beds are also present at the top of the Kaibab Formation (McKee 1954), so the presence of the Timpoweap Member at Cedar Mountain is unlikely. Rather than being the Shnabkaib Member, the slope-forming, gypsiferous middle portion of Cedar Mountain represents the Moqui Member, as its lithology is inconsistent with the cyclic sabkha-like deposits of the type section of the Shnabkaib Member at St. George, Utah (Reeside and Bassler 1922). As Stewart et al. (1972b) noted, the Moqui Member of northern Arizona overlies the "lower massive sandstone" within the Moenkopi Formation (as it does at Cedar Mountain), but this sandstone occurs near the top of the Shnabkaib Member in southern Utah. Noble named a massive sandstone near the top of Cedar Mountain the "De Chelly sandstone" (1922:72), but this unit represents the "upper massive sandstone" of the Holbrook Member, and not the De Chelly Sandstone Member of the Permian Cutler Formation found within the Navajo Nation in northeastern Arizona (McKee 1933; Read and Wanek 1961).

GRCA Resources

There are no fossils previously reported from the Moenkopi Formation at Cedar Mountain, and Noble (1922) reported looking for, but not finding any fossils during his study. We collected a specimen referred to *Rotodactylus* isp. (GRCA 33186; Figure 10-2), located numerous sedimentary structures, and identified the ichnotaxon *Scoyenia* cf. *gracilis* from the Wupatki Member and Holbrook Member, respectively, from Cedar Mountain (see "Results" below).




Figure 10-2. GRCA 33186, *Rotodactylus* isp. from the Wupatki Member, Moenkopi Formation at Cedar Mountain (NPS/MARSH ET AL.). **A.** Trackway with arrows showing individual prints. **B.** Photo of locality where the *Rotodactylus* trackway was found (hammer at approximate location of fallen block).

Paleontological Potential

Medium; much of the Moenkopi Formation in northern and northeastern Arizona is fossiliferous if specific stratigraphic horizons are targeted (however, this may be a function of the lack of comprehensive surveying). For example, fossils are unknown in the slope-forming Moqui Member (McKee 1954 reported plant impressions found at Moqui Wash near Winslow, but it is unclear whether they actually came from the Moqui Member there). The two stratigraphic horizons that produce vertebrate fossils described above are present at Cedar Mountain, but are either largely covered by talus or vegetation (the upper part of the Wupatki Member) or are strongly cliff-forming (the "upper massive sandstone" of the Holbrook Member) along the section. Depending on if the Timpoweap Member is indeed present at the base of the Moenkopi Formation (see above), the ammonite genus *Meekoceras* may be present (McKee 1954; Stewart et al. 1972b), but the contact between the lowest part of the Moenkopi Formation and the uppermost Kaibab Formation is poorly exposed. Invertebrate and vertebrate trace fossils are possible in fallen sandstone blocks from the Wupatki Member and Holbrook Member, but finding these in situ is unlikely.

Chinle Formation (Shinarump Member)

Geology

What is now accepted as the Shinarump Member of the Chinle Formation has long been recognized as a significant sedimentary unit throughout the Colorado Plateau, though its definition and lithological diagnosis have undergone substantial revision since its first conception (Powell 1873, 1876; Gilbert 1875). Recognized as the basal unit of the Chinle Formation (Stewart 1957; Repenning et al. 1969; Stewart et al. 1972a), the Shinarump Member unconformably overlies the Moenkopi Formation and often deeply scours the top of that unit, eroding down into the Holbrook Member in large trough-stratified channels. Throughout the Little Colorado River Valley and near Cameron, the primary lithology of the Shinarump Member is usually characterized as a lightly colored sandstone with substantial conglomeratic lenses containing at least 50% clasts of 5 cm (2 in) in diameter (Gregory 1917) and large petrified conifer logs. The unit is approximately 8 m (25 ft) thick at Cedar Mountain (Nobel 1922), but its deposited thickness is nearly impossible to determine at a given outcrop unless overlying Chinle Formation strata are present. The typical conglomeratic facies of the Shinarump Member can grade laterally into finer-grained facies formally recognized as the Mesa Redondo Member (Cooley 1958), but these are likely syndepositional overbank deposits correlative to the channelized fluvial system of the Shinarump Member (Irmis et al. 2011; Riggs et al. 2016; Martz et al. 2012, 2017). Regionally, the upper portions of the Shinarump Member and Mesa Redondo Member have been heavily pedogenically modified, and this can be used to recognize these units locally (Irmis et al. 2011, supp.). This may be the same as the post-depositional "mottling" described by Stewart et al. (1972a), which is a pedogenic alteration of the top of the Moenkopi Formation and parts of the Shinarump Member. At Cedar Mountain the top of the Shinarump Member is weathered off and the upper mottled horizon is not preserved or was never present.

Whereas the Shinarump Member of the Chinle Formation lacks diagnostic vertebrate fossils, it contains wood fossils similar to "*Araucarioxylon*" (Figure 10-3) and it stratigraphically underlies lower Chinle Formation members that have been removed by erosion here. These removed overlying units, such as the Blue Mesa Member, are Late Triassic (Norian) in age owing to the presence of

non-pseudopalatine leptosuchomorph phytosaurs and the absence of the aetosaur *Typothorax* at PEFO and other regional localities (Lucas 2010; Martz and Parker 2017).




Figure 10-3. Plant fossils from the Shinarump Member of the Chinle Formation at Cedar Mountain (NPS/MARSH ET AL.). **A.** Impression of a stem or small log in a fallen block, exhibiting the chert pebble conglomerate and mud rip-up clasts of the member. **B.** Fragment of conifer log that eroded from the Shinarump Member.

Detrital zircon samples from the Shinarump Member of the Chinle Formation in that study (Dickinson and Gehrels 2009b) do not seem to contain the youngest deposition-age grains, but a high-resolution U-Pb date from the laterally equivalent Mesa Redondo Member supports a Late

Triassic age (Norian, ~228 Ma) for the Shinarump Member (Atchley et al. 2013; Riggs et al. 2016). Lithostratigraphic correlation of the Shinarump Member to the base of the Dockum Group of Texas also suggests an early Norian age based on the presence of basal phytosaurs in the latter unit (Martz and Parker 2017).

Previous studies

The Shinarump Member was originally recognized as the capping sandstone at Cedar Mountain (Noble 1922), but at the time the unit was not recognized as the basal unit of the Chinle Formation in the area. Noble (1922) described the Shinarump Member as a hard, poorly sorted, cross-bedded conglomerate with chert clasts and sedimentary clasts that do not resemble any units in the underlying Moenkopi Formation. At the time, the Shinarump Member was considered Late Triassic in age owing to the presence of large petrified logs (see below) and its correlation to other Shinarump Member outcrops in the region that underlie other sedimentary units with Late Triassic vertebrate fossils (i.e., the rest of the Chinle Formation).

GRCA Resources

Noble (1922) described in situ petrified logs from the Shinarump Member at Cedar Mountain, and several pieces of petrified wood were previously documented by GRCA interns in 2011 as weathered talus and in situ. We observed numerous pieces of petrified wood ex situ and in situ (Figure 10-3; see "Results" below).

Paleontological Potential

Low; the taxonomy of the petrified wood found in the Shinarump Member in northern Arizona has not been well studied, and the rare vertebrate bones that have been reported from the unit (Martz et al. 2017) are too fragmentary to determine identifications beyond major clades (e.g., Temnospondyli, Phytosauria). The finer-grained lateral equivalent of the Shinarump Member, the Mesa Redondo Member, is not fossiliferous where it crops out near PEFO (Martz et al. 2012, 2017).

Field Methods

The two-day PaleoBlitz at Cedar Mountain occurred on July 15 and 16, 2019 as part of the GRCA Paleontological Resource Inventory (study no. GRCA-00745, permit no. GRCA-2019-SCI-0002). Access to Cedar Mountain was provided by a difficult two-track dirt road that does not seem to have been improved since Noble first described it as "rather poor" (1922:71). Any future excursions to Cedar Mountain will require a high clearance, short-bed 4x4 vehicle to best navigate the patches of boulders, ruts, and narrow switchbacks on the road. Instead of surveying the slope of the mesa in a systematic fashion (which is largely made up of the unfossiliferous Moqui Member), we targeted the specific horizons in the Wupatki Member and Holbrook Member that have the best potential for fossil preservation, as described above (Figure 10-1D).

The locations of sedimentary structures and fossils were documented using a Garmin GPS MAP64S and a resource-grade GNSS receiver, and digital photographs were taken with a Canon Powershot ELPH 150 IS and iPhone camera. We surveyed for body fossils by inspecting split pieces of sandstone and looking for bones weathering out of softer units, and we surveyed for trace fossils by inspecting the underside of sandstone overhangs and flipping blocks of fallen sandstone. Field

consolidant was not used on any specimens. Copies of field notes, photographs, and GPS data are on file at GRCA and the WASO Paleontology Archives. The fossil potential for the Triassic units at GRCA described above in the overview of Triassic strata at GRCA include low (lacks fossils or contains fossils with little scientific significance), medium (contains fossils in equivalent sections regionally), and high potential (contains scientifically significant fossils).

Results

Moenkopi Formation

Wupatki Member

The Wupatki Member is best exposed in low relief along small drainages, which incise into the alternating layers of siltstone and sandstone that characterize the member. We could not locate an extensive "lower massive sandstone" in the unit and thus were not able to determine our exact stratigraphic position within the member at a given location. However, the incised outcrops in the drainages expose innumerable slabs of ripple-laminated sandstone; most of the ripples are asymmetrical and indicative of uni-directional flow. We were not able to discern the upper contact with the Moqui Member in outcrop owing to the coverage of juniper.

Rotodactylus *Peabody 1948*

Type species—*Rotodactylus cursorius* Peabody 1948

Type specimen—UCMP 37794, trackway of 11 pes-manus sets.

Type locality and horizon—UCMP V3835, near Meteor Crater, Arizona; Wupatki Member, Moenkopi Formation.

Diagnosis—Semi-bipedal, long-striding trackway of a small five-toed reptile; when present, the manus is closer to the midline than the pes (Peabody 1948); digit IV longer than digit III in the pes (Haubold 1967, 1971). Brusatte et al. (2011) suggested that the ichnogenera *Prorotodactylus* and *Rotodactylus* were made by dinosauromorphs, which would extend the origin of Dinosauromorpha at least into the Middle Triassic 5–10 million years before the appearance of the oldest dinosauromorph body fossil (Ezcurra et al. 2017).

Rotodactylus

Referred specimen—GRCA 33186 (Figure 10-2).

Location—Cedar Mountain, GRCA (Figure 10-1).

Description—This specimen was collected as a small sandstone slab (Figure 10-2A) that fell a short distance from its source horizon in the Wupatki Member at Cedar Mountain (Figure 10-2B). It includes three to four partial pes tracks that run obliquely along a rippled surface. The front two prints appear to include ungual drag marks, and the rest of the rippled surface contains flute casts and other drag marks. It is not possible to determine the ichnospecies because the prints are too subtle and do not preserve enough anatomy, but the presence of the ichnogenus *Rotodactylus* is characteristic of the Wupatki Member in northern Arizona (Peabody 1948; Klein and Lucas 2010), including nearby at WUPA (Henderek et al. 2017).

Moqui Member

We found a few sedimentary structures within the thin sandstone lenses in the Moqui Member at Cedar Mountain, but we did not locate any fossils. The fine-grained, gypsum-rich slopes that make up most of the outcrop of this member are mostly covered by juniper and are weakly bedded in the better exposures around the mesa. One location in the upper part of the Moqui Member included a local sandstone unit that protects the softer underlying beds in the member. We found mud cracks superposed on bi-directional ripples in a laminated sandstone from that channel (Figure 10-4). We were also able to locate primary gypsum beds below this prominent sandstone, indicating that the Wupatki Member–Moqui Member contact is lower than this horizon and that this sandstone is not the "lower massive sandstone" that is found near the top of the Wupatki Member elsewhere in the region.



Figure 10-4. Mud cracks superposed on bi-directional ripples in a sandstone from the Moqui Member of the Moenkopi Formation at Cedar Mountain. Hammer is 40 cm (16 in) long (NPS/MARSH ET AL.).

Holbrook Member

We were unable to locate a bench-forming conglomeratic layer within the Holbrook Member at Cedar Mountain, which would have been the most likely bed to yield vertebrate fossils. We determined that the second unit of Noble's measured section (1922:72) is the "upper massive sandstone" found throughout the Little Colorado River Valley; this cliff-former is difficult to navigate around the mesa, but provides a gentle slope above it containing interspersed exposures of orthogonal ripple-laminated sandstones (Figure 10-5). One location yielded internal molds of invertebrate burrows, but they were found ex situ and their orientation is impossible to determine. The burrows are straight, circular in cross section, have weak transverse striations on the outer surface, and a few appear to taper in one direction (Figure 10-6).




Figure 10-5. Hand sample of orthogonal ripples in alternating layers of laminated sandstone from the Holbrook Member of the Moenkopi Formation at Cedar Mountain (NPS/MARSH ET AL.).



Figure 10-6. Internal molds of burrows found ex situ in the Holbrook Member of the Moenkopi Formation at Cedar Mountain (NPS/MARSH ET AL.).

Scoyenia *White 1929*

Type species—*Scoyenia gracilis* White 1929

Type specimen—USNM 201869, straight to slightly curved, horizontally overlapping burrows considered to be a deposit feeding or dwelling structure, potentially made by insects (White 1929; see Paleozoic Invertebrate Ichnology chapter of this report).

Type locality and horizon—GRCA 8538, Grand Canyon National Park, Arizona; lower part of Hermit Formation (Permian).

Diagnosis—Horizontal, narrow, and curved burrows; unbranched (White 1929; Frey et al. 1984; MacEachern et al. 2010).

Scoyenia *cf.* gracilis

Referred specimen—Nothing collected; several hand samples observed but left in the field (Figure 10-7).

Location—Cedar Mountain, GRCA.

Description—These hand samples were observed as ex situ surface float in a talus pile of associated ripple-laminated sandstones above the "upper massive sandstone" in the Holbrook Member at Cedar Mountain. They include horizontal, gently curved burrows that generally trend in the same direction, but can overlap one another. The sedimentological and ichnological fabric of this portion of the Holbrook Member (and much of the Moenkopi Formation in northern Arizona) is consistent with the *Scoyenia* Ichnofacies (Buatois and Mángano 1995, 2004), which is characterized by horizontal feeding and locomotory traces, fish fin markings, vertical and horizontal dwelling structures, arthropod and vertebrate trackways, and plant root traces (MacEachern et al. 2010). These occur in low-energy terrestrial systems with alternating periods of flooding and subaerial conditions, which describes much of the depositional setting of the continental fluvial system of the Moenkopi Formation in northern Arizona (McKee 1954; Stewart et al. 1971b; MacEachern et al. 2010).




Figure 10-7. Hand samples of *Scoyenia* cf. *gracilis* found in the Holbrook Member of the Moenkopi Formation at Cedar Mountain (**A–D**) (NPS/MARSH ET AL.).

Chinle Formation

Shinarump Member

We found numerous fragments of petrified wood and impressions of wood in Shinarump Member clasts (Figure 10-3). The in situ logs within the Shinarump Member at Cedar Mountain are several meters in length and often have chert clasts silicified onto the outer surface of the wood. Mud rip-up clasts with concentric bands in cross section are commonly associated with the petrified wood near the base of fining-upward channel sequences in the Shinarump Member.

Conclusions

The Triassic sedimentary rocks at Cedar Mountain are the only Mesozoic units within the boundary of GRCA, and beneath the juniper cover are quite similar to equivalent units in the region. The Moenkopi Formation at GRCA includes the Wupatki Member, Moqui Member (but not the Shnabkaib Member), and Holbrook Member, all of which are similar with respect to sedimentary

architecture and sedimentary structures to the same units at Cameron, WUPA, Meteor Crater, and Holbrook, Arizona. We were able to match Noble's measured section to the outcrop at Cedar Mountain (1922), which allowed us to target specific areas for a two-day paleontological survey. The discovery of *Rotodactylus* in the Wupatki Member at GRCA is the first Mesozoic fossil specimen found in the park to be collected for the park's museum collections, and it confirms the presence of a small reptilian trackmaker (possibly a dinosauromorph; Brusatte et al. 2011) in the Middle Triassic similar to those reported from other sites in northern Arizona. Similarly, we confirm the presence of the *Scoyenia* ichnofacies in the Holbrook Member, which is typical of the fluvial and overbank depositional settings of the eastern exposures of the Moenkopi Formation. Large pieces of petrified wood are common in the Shinarump Member of the Chinle Formation, which caps Cedar Mountain and covers much of its slopes as talus.

Given the difficult access to Cedar Mountain as well as the low to medium potential to find exposed fossil resources, the Mesozoic section at GRCA needs very little monitoring in the future. More of the fallen sandstone slabs of the Holbrook Member and incised drainages in the Wupatki Member may be surveyed for additional trace and/or body fossils, but the likelihood of finding anything new is low. If future surveys or monitoring efforts are to take place, we suggest navigating the road from Desert View as few times as possible, camping at the base of Cedar Mountain for the duration of the trip, and contacting the Navajo Nation for permission to use the eastern road to access the site.

Acknowledgments

We thank Mark Nebel and Ronda Newton for providing logistical and financial support for the fieldwork described in this chapter, which was assisted by Diana Boudreau and Klara Widrig. Thanks to Vince Santucci, Justin Tweet, and Tim Connors of the Paleontology Program (Geologic Resource Division) of the National Park Service for inviting us on this project and for providing map and literature search assistance. We also thank Tracy Thomson (UC Davis) and Andrew Milner (St. George Dinosaur Discovery Site at Johnson Farm) for assistance with fossil identification. We appreciate the reviews by Justin Tweet, Diana Boudreau, Sterling Nesbitt and Jeff Martz.

Literature Cited

Atchley, S. C., L. C. Nordt, S. I. Dworkin, J. Ramezani, W. G. Parker, S. R. Ash, and S. A. Bowring. 2013. A linkage among Pangean tectonism, cyclic alluviation, climate change, and biologic turnover in the Late Triassic: the record from the Chinle Formation, southwestern United States. Journal of Sedimentary Research 83:1147–1161.

Billingsley, G. H., S. S. Priest, and T. J. Felger. 2007. Geologic map of Wupatki National Monument and vicinity, Coconino County, northern Arizona. U.S. Geological Survey, Reston, Virginia. Scientific Investigations Map 2958. Scale 1:24,000.

Billingsley, G. H., P. W. Stoffer, and S. S. Priest. 2012. Geologic map of the Tuba City 30' x 60' Quadrangle, Coconino County, Arizona. U.S. Geological Survey, Reston, Virginia. Scientific Investigations Map SIM-3227. Scale 1:50,000.

Blakey R. C., E. L. Basham, and M. J. Cook. 1993. Early and Middle Triassic paleogeography of the Colorado Plateau and vicinity. Museum of Northern Arizona Bulletin 59:13–26.

Brusatte, S. L., G. Niedźwiedzki, and R. J. Butler. 2011. Footprints pull origin and diversification of dinosaur stem lineage deep into Early Triassic. Proceedings of the Royal Society B: Biological Sciences 278:1107–1113.

Buatois, L. A., and M. G. Mángano. 1995. The paleoenvironmental and paleoecological significance of the lacustrine *Mermia* ichnofacies: An archetypical subaqueous nonmarine trace fossil assemblage. Ichnos 4:151–161.

Buatois, L. A., and M. G. Mángano. 2004. Animal-substrate interactions in freshwater environments: Applications of ichnology in facies and sequence stratigraphic analysis of fluvio-lacustrine successions. Geological Society of London Special Publication 228:311–333.

Cohen, K. M., S. C. Finney, P. L. Gibbard, and J.-X. Fan. 2013. The ICS International Chronostratigraphic Chart. Episodes 36:199–204.

Cooley, M. E. 1958. The Mesa Redondo Member of the Chinle Formation, Apache and Navajo Counties, Arizona. Plateau 31:7–15.

Dickinson, W. R., and G. E. Gehrels. 2008. U-Pb ages of detrital zircons in relation to paleogeography: Triassic paleodrainage networks and sediment dispersal across southwest Laurentia. Journal of Sedimentary Research 78:745–764.

Dickinson, W. R., and G. E. Gehrels. 2009a. Insights into North American paleogeography and paleotectonics from U-Pb ages of detrital zircons in Mesozoic strata of the Colorado Plateau, USA. International Journal of Earth Sciences 99:1247–1265.

Dickinson, W. R., and G. E. Gehrels. 2009b. Use of U-Pb ages of detrital zircons to infer maximum depositional ages of strata: a test against a Colorado Plateau Mesozoic database. Earth and Planetary Science Letters 288:115–125.

Dutton, C. E. 1880. Report on the geology of the high plateaus of Utah. U.S. Geographical and Geological Survey of the Rocky Mountain Region. Government Printing Office, Washington, D.C.

Dutton, C. E. 1882. Tertiary history of the Grand Cañon district, with atlas. U.S. Geological Survey, Washington, D.C. Monograph 2.

Ezcurra, M. D., L. E. Fiorelli, A. G. Martinelli, S. Rocher, M. B. von Baczko, M. Ezpeleta, J. R. A. Taborda, E. M. Hechenleitner, M. J. Trotteyn, and J. B. Desojo. 2017. Deep faunistic turnovers preceded the rise of dinosaurs in southwestern Pangaea. Nature Ecology and Evolution 1:1477–1483.

Flowers, R. M. The enigmatic rise of the Colorado Plateau. Geology 38:671–672.

Frey, R. W., S. G. Pemberton, and J. A. Fagerstrom. 1984. Morphological, ethological, and environmental significance of the ichnogenera *Scoyenia* and *Ancorichnus*. Journal of Paleontology 58:511–528.

Gilbert, G. K. 1875. Report on the geology of portions of Nevada, Utah, California, and Arizona, examined in the years 1871 and 1872. Pages 17–187 *in* Report on the geographical and geological explorations and surveys west of the One Hundredth Meridian (Wheeler). Government Printing Office, Washington, D.C. U.S. Geological and Geographical Survey, Publication of the Wheeler Survey 3(1).

Gregory, H. E. 1916. The Navajo country. U.S. Geological Survey, Washington, D.C. Water-Supply Paper 380.

Gregory, H. E. 1917. Geology of the Navajo country: a reconnaissance of parts of Arizona, New Mexico, and Utah. U.S. Geological Survey, Washington, D.C. Professional Paper 93:1–161.

Gregory, H. E. 1948. Geology and geography of Kane County, Utah. Geological Society of America Bulletin 59:211–248.

Gregory, H. E. 1952. The geology and geography of the Zion park region, Utah-Arizona. U.S. Geological Survey, Washington, D.C. Professional Paper 220.

Hager, D. 1922. Oil possibilities of the Holbrook area in northeastern Arizona. Mining and Oil Bulletin 8:23–27, 33–34, 71–75, 81, 94, 135–140.

Haubold, H. 1967. Eine Pseudosuchier-Fährten aus dem Buntsandstein Südthüringens. Hallesches Jahrbuch für Mitteldeutsche Erdgeschichte 8:12–48.

Haubold, H. 1971. Die Tetrapodenfährten des Buntsandsteins. Paläontologische Abhandlungen A 4:395–548.

Henderek, R. L., V. L. Santucci, B. Rizner, J. Tweet, and J. R. Wood. 2017. Wupatki National Monument paleontological resources inventory. Natural Resource Report NPS/WUPA/NRR-2017/1524.

Irmis, R. B., R. Mundil, J. W. Martz, and W. G. Parker. 2011. High-resolution U-Pb ages from the Upper Triassic Chinle Formation (New Mexico, USA) support a diachronous rise of dinosaurs. Earth and Planetary Science Letters 309:258–267.

Klein, H., and S. G. Lucas. 2010. Tetrapod footprints – their use in biostratigraphy and biochronology of the Triassic. Pages 419–446 *in* S. G. Lucas, editor. The Triassic timescale. Geological Society of London, London, United Kingdom. Special Publication 334.

Klein, H., and S. G. Lucas. 2010b. Review of the tetrapod ichnofauna of the Moenkopi Formation/Group (Early-Middle Triassic) of the American Southwest. New Mexico Museum of Nature and Science Bulletin 50:1–67.

Liu, L., and M. Gurnis. 2010. Dynamic subsidence and uplift of the Colorado Plateau. Geology 38:663–666.

Lucas, S. G. 2010. The Triassic timescale based on nonmarine tetrapod biostratigraphy and biochronology. Pages 447–500 *in* S. G. Lucas, editor. The Triassic timescale. Geological Society of London, London, United Kingdom. Special Publication 334.

MacEachern, J. A., S. G. Pemberton, M. K. Gingras, and K. L. Bann. 2010. Ichnology and facies models. Pages 19–58 *in* N. P. James and R. W. Dalrymple, editors. Facies Models 4. Geological Association of Canada, St. John's, Canada.

Marcou, J. 1858. Geology of North America. Zurcher and Furrer, Zurich, Switzerland.

Martz, J. W., and W. G. Parker. 2017. Revised formulation of the Late Triassic Land Vertebrate "Faunachrons" of western North America: Recommendations for codifying nascent systems of vertebrate biochronology. Pages 39–124 *in* K. E. Zeigler and W. G. Parker, editors. Terrestrial depositional systems: Deciphering complexities through multiple stratigraphic methods. Elsevier, Oxford, United Kingdom.

Martz, J. W., W. G. Parker, L. Skinner, J. J. Raucci, P. Umhoefer, and R. Blakey. 2012. Geologic map of Petrified Forest National Park. Arizona Geological Survey, Tucson, Arizona. Contributed Map CM-12-A. Scale 1:50,000.

Martz, J. W., J. I. Kirkland, A. R. C. Milner, W. G. Parker, and V. L. Santucci. 2017. Upper Triassic lithostratigraphy, depositional systems, and vertebrate paleontology across southern Utah. Geology of the Intermountain West 4:99–180.

McKee, E. D. 1933. The Coconino sandstone—its history and origin. Carnegie Institution of Washington Publication 440:77–115.

McKee, E. D. 1954. Stratigraphy and history of the Moenkopi Formation of Triassic age. Geological Society of America, New York, New York. Memoir 61.

McKee, E. D. 1964. Permian and Triassic cycles involving chemical sediments, northern Arizona. Kansas Geological Survey, Lawrence, Kansas. Bulletin 169:283–286.

Morales, M. 1987. Terrestrial fauna and flora from the Triassic Moenkopi Formation of the southwestern United States. Journal of the Arizona-Nevada Academy of Science 22:1–19.

Nesbitt, S. J. 2000. A preliminary report on new vertebrate fossil sites, including a microsite from the Holbrook Member of the Moenkopi Formation, Holbrook, Arizona. Mesa Southwest Museum Bulletin 7:17–30.

Nesbitt, S. J. 2002. An update on fossil reptile material from the upper Moenkopi Formation, Holbrook Member (Middle Triassic), northern Arizona. Mesa Southwest Museum Bulletin 8:3–7.

Nesbitt, S. J. 2005a. Stratigraphy and tetrapod fauna of major quarries in the Moenkopi Formation (Early-Middle Triassic) along the Little Colorado River of northern Arizona. Mesa Southwest Bulletin 11:18–33.

Nesbitt, S. J. 2005b. Osteology of the Middle Triassic pseudosuchian archosaur *Arizonasaurus babbitti*. Historical Biology 17:19–47.

Newberry, J. S. 1861. Geological report. *In* Ives, J. C., Report upon the Colorado River of the West, explored in 1857 and 1858. U.S. 36th Congress, 1st Session, House Executive Document 90, Serial 1058, Part 3.

Niedźwiedzki, G., S. L. Brusatte, and R. J. Butler. 2013. *Prorotodactylus* and *Rotodactylus* tracks: An ichnological record of dinosauromorphs from the Early–Middle Triassic of Poland. Pages 319–351 *in* S. J. Nesbitt, J. B. Desojo, and R. B. Irmis, editors. Anatomy, phylogeny and palaeobiology of early archosaurs and their kin. Geological Society of London, London, United Kingdom. Special Publication 379.

Noble, L. F. 1922. A section of the Paleozoic formations of the Grand Canyon at Bass Trail. Pages 23–73 *in* Shorter contributions to general geology, 1922. U.S. Geological Survey, Washington, D.C. Professional Paper 131-B.

Ogg, J. G. 2012. Triassic. Pages 681–730 *in* F. M. Gradstein, J. G. Ogg, M. D. Schmitz, and G. M. Ogg, editors. The geologic time scale 2012. Volume 2. Elsevier, Oxford, United Kingdom.

Peabody, F. E. 1948. Reptile and amphibian trackways from the Moenkopi Formation in Arizona and Utah. University of California Publications, Bulletin of the Department of Geological Sciences 27:295–468.

Pipiringos, G. N., and R. B. O'Sullivan. 1978. Principle unconformities in Triassic and Jurassic rocks, western interior United States, a preliminary survey. U.S. Geological Survey, Washington, D.C. Professional Paper 1035-A.

Powell, J. W. 1873. Some remarks on the geological structure of a district of country lying to the north of the Grand Canyon of the Colorado. American Journal of Science (3rd series) 5:456–465.

Powell, J. W. 1876. Report on the geology of the eastern portion of the Uinta Mountains and a region of country adjacent thereto. U.S. Geological and Geographical Survey of the Territories. Government Printing Office, Washington, D.C.

Read, C. B., and A. A. Wanek. 1961. Correlation of Permian rocks in northeastern Arizona and adjoining parts of New Mexico and Utah. Pages C156–160 *in* Short papers in the geologic and hydrologic sciences, articles 147-192. Geological Survey research 1961. U.S. Geological Survey, Washington, D.C. Professional Paper 424-C.

Reeside, J. B., and H. Bassler. 1922. Stratigraphic sections in southwestern Utah and northwestern Arizona. U.S. Geological Survey, Washington, D.C. Professional Paper 129-D.

Repenning, C. A., M. E. Cooley, and J. P. Akers. 1969. Stratigraphy of the Chinle and Moenkopi Formations, Navajo and Hopi Indian Reservations. U.S. Geological Survey, Washington, D.C. Professional Paper 521-B.

Riggs, N. R., Z. A. Oberling, E. R. Howell, W. G. Parker, A. P. Barth, M. R. Cecil, and J. W. Martz. 2016. Sources of volcanic detritus in the basal Chinle Formation, southwestern Laurentia, and implications for the Early Mesozoic magmatic arc. Geosphere 12:439–463.

Shimer, H. W. 1919. Permo-Triassic of northwestern Arizona. Geological Society of America Bulletin 30(4):471–498.

Stewart, J. H. 1957. Proposed nomenclature of part of Upper Triassic strata in southeastern Utah: American Association of Petroleum Geologists Bulletin 41:441–465.

Stewart, J. H., F. G. Poole, and R. F. Wilson. 1972a. Stratigraphy and origin of the Chinle Formation and related Upper Triassic strata in the Colorado Plateau region. U.S. Geological Survey, Washington, D.C. Professional Paper 690.

Stewart, J. H., F. G. Poole, and R. F. Wilson. 1972b. Stratigraphy and origin of the Moenkopi Formation and related strata in the Colorado Plateau region. U.S. Geological Survey, Washington, D.C. Professional Paper 691.

Walcott, C. D. 1880. The Permian and other Paleozoic groups of the Kanab Valley, Arizona. American Journal of Science (3rd series) 20:221–225.

Ward, L. F. 1901. Geology of the Little Colorado Valley. American Journal of Science (4th series) 12:401–413.

Welles, S. P. 1947. Vertebrates from the Upper Moenkopi Formation of northern Arizona. University of California Publication 27:241–294.

Welles, S. P. 1969. Collecting Triassic vertebrates in the Plateau Province. Journal of the West 8:231–246.

Welles, S. J. 1993. A review of the lonchorhynchine trematosaurs (Labrynthodontia), and a description of a new genus and species from the Lower Moenkopi Formation of Arizona. PaleoBios 14:1–24.

White, C. D. 1929. Flora of the Hermit Shale, Grand Canyon, Arizona. Carnegie Institute of Washington Publication 405.

Chapter 11. Pleistocene/Holocene Cave Fossils From Grand Canyon National Park

Ice Age (Pleistocene) Flora, Fauna, Environments, And Climate Of The Grand Canyon, Arizona

By Jim I. Mead[1], Justin S. Tweet[2], Vincent L. Santucci[3], Benjamin Tobin[4], Carol L. Chambers[5], Shawn C. Thomas[6], and Mary C. Carpenter[7]

[1]The Mammoth Site
1800 Hwy 18 ByPass
Hot Springs, South Dakota 577473

[2]National Park Service
Geologic Resources Division
1849 "C" Street, NW
Washington, D.C. 20240

[3]National Park Service
9149 79th St. S.
Cottage Grove, Minnesota 55016

[4]Kentucky Geological Survey
University of Kentucky
Lexington, Kentucky 40506

[5]Northern Arizona University
School of Forestry
Flagstaff, Arizona 86001

[6]Bat Conservation International
Olympia, Washington 98506

[7]National Park Service
Badlands National Park
Interior, South Dakota 57750

Introduction

The Colorado Plateau is a distinct physiographic province in western North America covering an area of roughly 337,000 km^2 (130,115 mi^2) across parts of Arizona, Colorado, New Mexico, and Utah. Elevations range from about 360 m (1,180 ft) in the overall Grand Canyon (GC; which includes the Grand Canyon National Park, GRCA) river corridor to an average at the eastern South Rim of 2,072 m (6,800 ft) to 3,850 m (12,630 ft) on the nearby San Francisco Peaks at Flagstaff, Arizona, with an average elevation of 1,525 m (5,000 ft). The Colorado River of Grand Canyon is located along the southwestern portion of the Colorado Plateau in Arizona and is renowned for its

dramatic display of geomorphic effects created by fluvial incision and its unique dry-preservation of fossils from the Ice Age (late Pleistocene and Holocene [Quaternary]; most recent 2.58 million years). Although there were at least 22 glacial-interglacial cycles during the Ice Age, this discussion is limited to the most recent episode (called the Wisconsinan Glaciation), which includes the transition to the modern climate (latest Pleistocene and Holocene; the most recent 50,000 years of geologic history).

Due to its range in elevations and physiographic position in western North America, the Colorado Plateau (CP; along with its GC river corridor) plays a key role in the continental monsoons much in the same way that the Tibetan Plateau affects the Southeast Asian monsoon climate (Tang and Reiter 1984; Adams and Comrie 1997). In general, precipitation decreases from high elevations to lower elevations. Summer precipitation decreases from the southern Colorado Plateau northward which correlates to the strength of the summer monsoon. These relationships have important consequences for modern and Ice Age biotic distributions in the Grand Canyon and on the surrounding plateaus (Mock 1996; Higgins et al. 1997; Anderson et al. 2000, among others).

Here we review the Ice Age floras and faunas found predominantly within the GC but also on adjacent rims made by the Coconino, Kaibab, Kanab, Uinkaret, Shivwits, and Hualapai plateaus in addition to the Marble Platform at the up-river end of GC. The Grand Canyon includes land administered by the Havasupai, Hualapai, and Navajo Indian tribes along with federal lands managed by National Park Service (Grand Canyon National Park, Lake Mead National Recreation Area), Bureau of Land Management (Grand Canyon-Parashant National Monument), and US Forest Service (Coconino and Kaibab National Forests). The GC and its Colorado River corridor extends nearly 448 km (278 mi) in length and encompasses an area of about 4,921 km^2 (1,900 mi^2).

The preservation potential of Ice Age fossil deposits within the confines of the GC is limited due to active downcutting, steep canyon walls, abundant mass wasting, and periodic catastrophic flooding. Occasionally one can find some perched Ice Age alluvium that has been spared being flushed by subsequent floods down to the Gulf of Mexico. What has been the “gold mine” for Ice Age preservation are the dry caves created in the numerous limestone rock formations coupled with the arid climate. For details about the geology and overall history of the region, see the Geology and Stratigraphy chapter in this report or Beus and Morales (2003).

There are a variety of geochemical analyses used to assess the age of the various deposits found throughout the GC. For the Ice Age fossils of preserved organic remains researchers typically use radiocarbon dating (^{14}C, radiometric dating). This isotopic dating technique is accurate for only about the most recent 50,000 years. Over the past couple of decades the technique has been refined and now researchers refer to ages in “calibrated” (or corrected) years before present (cal yr BP). “Before present” is expressed as pre-1950 (pre-atomic bomb) and can be thought of in terms of “years ago”. In the 1970s and 1980s a researcher would need enough organic remains to fill an old 35-mm film can in order to obtain an accurate radiocarbon age. Now, with a technique called accelerator mass spectrometry (AMS), a researcher can use a single seed or piece of wood the size of a pin head. In the review below of fossil localities, the established age range will be presented. The interested

reader who wants to know more details about the chronology will need to go to the original publications provided in the literature cited.

Preservation Scenarios (Taphonomy)

River Corridor Sediments

Much has been written about the Colorado River and its evolution in the GC region (e.g., chapters in Young and Spamer 2001; Beus and Morales 2003). Tobin et al. (in press) provide a detailed overview of the karst system model as one of the primary drivers of canyon development and stream piracy. Critical for the preservation of the fossil record is a preserved depositional environment. The fairly narrow river corridor offers few places where fluvial and alluvial sediments can persist for a long period of time. Periodic floods have occurred along the river corridor, especially related to the series of lava dams that were positioned in the west region of the GC (Hamblin 1994; Fenton et al. 2002). Damming with sediment infilling (along with outburst-flood deposits) within the GC has permitted select side canyons to preserve some of the perched alluvial deposits (although some of these observed units might be related to spring-fed deposition and not flood debris; see Kaufman et al. 2002 and references within). These rare sedimentary deposits hold a record of select Grand Canyon past environments.

Dry Caves

Cave morphology throughout the GC can be separated into two distinct groups: 1) caves formed under confined hydrogeologic conditions (i.e., phreatic zone, saturated, below water table) and 2) those formed under unconfined conditions (i.e., vadose, unsaturated zone; Hill and Polyak 2010). Those formed under confined conditions are typically older, dry today, and removed from current hydrologic processes and are typically assumed to have formed either during or prior to river corridor incision. Unconfined caves are being formed currently, are a part of the karst groundwater system of the region and are actively recharged from precipitation on the surrounding plateaus. These differing conditions have resulted in a dichotomy of cave morphology in GC (Huntoon 2000).

Some of the oldest and most impressive cave systems in the area are formed in the upper members of the Redwall Limestone (see overview in Tobin et al. in press). Dating of mammillary cave formations in some of these caves places their formation prior to 1.6 to 3.7 million years ago (Polyak et al. 2008). These caves form along regional fracture patterns, resulting in two-dimensional maze caves with minimal vertical development, except in rare cases. These “maze caves” formed under phreatic conditions (Hill 2010) and follow regional hydrologic gradients, ultimately emerging in the canyon. As the canyon incised, the water table dropped, resulting in the dewatering of these cave systems and exposing cave entrances along canyon walls. There are competing hypotheses on their formation with observations supporting both: epigenic processes (Huntoon 2000) versus hypogenic processes (Hill and Polyak 2010).

Due to the dry nature of these “confined” caves, they often provide the best environment for preservation of paleontological resources from the Pleistocene to recent times. The nature of the known entrances to these caves also plays a major role in what species are most likely to be preserved within them. Since these caves were exposed to the surface environment due to canyon incision, the entrances are typically in cliff faces with hundreds of feet of cliff above and below. This

results in a limited variety of species that can actively use them. These species typically include birds, bats, packrats, ringtails, and an extinct mountain goat. When these dry caves have easier access, evidence of a wider array of species can be found that utilized them, including ground sloths, carnivores of many types, tortoises, and other forms of Pleistocene fauna.

Ongoing cave development is evident throughout the region as well. While these "unconfined" caves are more sporadically distributed throughout the park, many large springs are tied to them. The speleogenesis of these caves is much simpler than the confined systems, following the typical model of epigenic karst development. As with caves at the top of the Redwall Limestone, these caves follow regional structural patterns. Water is sourced directly from precipitation that recharges the aquifer via sinkholes on the surrounding plateaus. These regional aquifers appear to have a distributary pattern (Jones et al. 2018), with individual sinkholes tied to multiple springs. Springs on either side of the Colorado River have distinct flow patterns (Tobin et al. 2018), resulting in vastly different spring morphologies, with only aquifers on the north side of the Colorado River having significant cave development. Due to the increased moisture in these caves, they typically have minimal paleontological resources, with the majority that have been noted tied to packrat (typically nearer the drier entrance) and bat use.

What makes the greater Grand Canyon region unique in the Southwest is the abundance of dry caves. The development of these confined caves helps keep the cave and its contents dry, but having the climate and environment outside arid is equally as critical for preservation of Pleistocene-age fossils. A case in point: a horizontal crevice a mere 20 cm (8 in) in from the dripline can preserve packrat middens (see below) for over 30,000 years.

Packrat Middens

Packrats (trade rats, woodrats) are a genus of rodent (Rodentia, Cricetidae, *Neotoma*) with about 20 species with distributions from Alaska to southern Mexico (Vaughan 1990). There are more species of packrats whose distributions include the greater Grand Canyon area than any other region for the genus. They all have the habit of making a nest and den along with the construction of debris piles over the living areas (Dial and Czaplewski 1990; Finley 1990). Materials for these debris piles include a wide diversity of plant remains (such as leaves, twigs, flowers, thorns, bark, and seeds), dung, bones, and rocks—typically all gathered from within 100 m (330 ft) of the nest (Spaulding et al. 1990). Faunal remains recovered from the debris piles are not all necessarily of local origin (especially when sourced from carnivore dung, regurgitated pellets, and raptor and vulturid nests; Figure 11-1) (Mead 2005). These nests and dens can be created in caves, crevices, overhanging rock ledges, and rockshelters. As long as the middens are protected from direct precipitation, the contents are preserved.

As the packrat uses the den and nest, it cleans the passageways and reassembles the pile of debris on top. In so doing, what looks like a mine tailing dump develops along the edge of the constructed mound. Over time this mound, containing all of the contents collected by the packrat along with its dung is incorporated into a mixed heap of debris and then is scent-marked with its urine. Repeated urination on the debris pile ultimately cements all the material into a rock-hard mass (concentrated urine is called "amberat"); the cemented debris pile is called a "midden" (Figure 11-2).




Figure 11-1. Steve Emslie (right) and Larry Coats (left) excavating a condor (*Gymnogyps*) nest and skeletal remains reclaimed by a packrat and made into a midden, Sandblast Cave, 1984 (EMILEE MEAD).

Some of the earliest paleoecological studies of the American Southwest using packrat middens began in the GC (Phillips and Van Devender 1974; Van Devender and Mead 1976). The entire GC has not been assessed for packrat middens and their paleoecological data; rather, research has been limited to select regions primarily in the eastern half. Major study areas are typically along the river corridor and in side canyons such as Stanton's Cave (Dryer 1994), Little Nankoweap Canyon (Mead et al. 2003), Chuar (Cole 1982, 1990), Hance Canyon and Horseshoe Mesa (Cole 1982, 1990), and Rampart Cave, Vulture Cave, and surrounding canyons (Phillips and Van Devender 1974; Mead and Phillips 1981; Phillips 1984; Spaulding 1990).



Figure 11-2. A juvenile extinct Harrington's mountain goat (*Oreamnos harringtoni*) skull in a packrat midden, Stevens Cave, 1984 (EMILEE MEAD).

These preserved middens are valuable for a number of reasons. Each can be accurately radiocarbon dated. They contain abundant macrobotanical remains typically from within the home range of less than 100 m (330 ft) (Spaulding et al. 1990). Multiple species of macrobotanical remains can be independently radiocarbon dated to produce accurate coeval habitat reconstructions. Microfaunal fossils found associated with botanical remains often include the delicate skeletal remains of the local herpetofauna, species often not found in typical alluvial localities due to the abrasive nature of deposition. Interestingly, radiocarbon-dated packrat middens were used to determine the rates of cliff retreat in the eastern GRCA (Cole and Mayer 1982).

Taxonomic Groups

Appendix 11-A presents a list of known biotic remains recovered from Pleistocene deposits in the GC region. The list presents taxa by primary publication resource and not by locality (which can be determined from the primary reference).

Plants

Due to the numerous dry cave and rockshelter localities coupled with the overall arid environment, Pleistocene-age plant remains are plentiful within each packrat midden unit. Packrats are basically browsers, over-selecting the woody and herbaceous plants but typically underrepresenting the graze species such as the grasses. Although equally as well-preserved as the macrobotanical fossil, pollen remains from the region have been less frequently utilized for paleoenvironmental reconstructions.

Martin et al. (1961) recovered pollen from Shasta ground sloth dung in Rampart Cave. Pollen frequencies recovered from sediments were compared to pollen counts from artiodactyl dung pellets and macrobotanical remains from the same sediments from test pits in Bida and Kaetan caves (see below; O'Rourke and Mead 1985). Pollen and microhistological remains from dung were also compared to fossils from test pits in Stanton's Cave (Robbins et al. 1984; see Hansen 1974 about technique). Pollen assemblages and associated plant macrofossils from packrat middens do reflect similar vegetation signals but this still needs to be further assessed for future paleoecological reconstructions (Anderson and Van Devender 1991). Ideally one would sample the macrobotanical remains and pollen from packrat middens and also assess the pollen and microhistological remains from dung also in the midden, hopefully selecting not only packrat pellets but also dung from potential grazers.

Invertebrates

The recovery and study of Ice Age and Holocene-age mollusks from the GC is still in its infancy. The extant taxa and their distributions are well studied from a few localities, primarily from the work of Pilsbry and Ferriss (1911). Spamer and Bogan (1993) provided a critical overview of the extant taxa. Spamer (1993) and Spamer and Bogan (1993) synthesized the known late Pleistocene malacofaunas (molluscan faunas) and emphasized that much is yet to be understood about the mollusks of the region dating to the last glacial and early Holocene. Kaufman et al. (2002) provide a few more records but still illustrate that the Pleistocene molluscan faunas are not adequately studied. Mollusks have also been incorporated into packrat middens (Cole and Mead 1981). Ostracodes are equally as poorly understood in the GC region (Kaufman et al. 2002). The record of Pleistocene arthropods is restricted to those fossils recovered from dry-preserved packrat midden and cave sediment localities (Elias et al. 1992), and again, the group as a whole for the Ice Age is poorly understood. The single discussion about nematodes comes from a study of dry-preserved sloth dung from Rampart Cave (Schmidt et al. 1992).

Vertebrates

The recovery and study of vertebrates from the GC is much more voluminous and well understood than the invertebrates. Appendix 11-A provides an extensive list of the fossils and relevant citations. The first study of the Pleistocene vertebrates in the GC occurred at Rampart Cave (see below) following the discovery of Shasta ground sloth dung (Evans 1936; Laudermilk and Munz 1938; Hansen 1978). Vertebrate remains include skeletal elements, dung, hair, dermal scales, and occasionally entire mummified animals (Figure 11-3). These fossils are recovered from cave sediments, dung mats (both bat and artiodactyl), packrat middens, owl pellets, raptor nests, and ringtail refuse areas (e.g., Mead and Van Devender 1981; Mead 2005).

A number of studies have been made of the Pleistocene vertebrate remains but there is only one comprehensive, albeit outdated, overview (Mead 1981; see review in Kenworthy et al. 2004). The Pleistocene record of amphibians (anurans and salamanders) is non-existent for the GC. The record of turtles is exceedingly rare and occurs only from cave and midden deposits in the far western river corridor (Rampart and Vulture caves; see below). Lizards and snakes are abundant in the record due to the presence of their remains in dry cave deposits and packrat middens from throughout the GC.

Some of the first or only fossil records of select squamate species come from packrat middens. Birds are fairly well represented due to the in-depth records from Rampart and Stanton's caves and localities in the Sandblast Cave area (see below; Emslie 1988), but there is a bias toward the river corridor avifauna.



Figure 11-3. A mummified myotid bat at its last perch among the gypsum crystals in Double Bopper Cave (NPS/SHAWN THOMAS).

Mammals are equally as well represented as the birds. Entrances to a number of the cave localities is inaccessible to most mammals due to their locations high on cliff faces (Figure 11-4). Only the best cliff climbers (mountain goats, packrats, ringtails) and fliers (bats) gain entrance to some of these caves. With all the caves in the GC, studies of extant and ancient bats and their guano (dung) deposits are abundant (e.g., Wurster et al. 2008; Pape 2014), with a number of mummified remains beginning to be studied in detail (e.g., Mead and Mikesic 2001; see Double Bopper Cave below). A number of medium to large mammals (some extinct) are reported from various caves throughout the GC region, both within the river corridor and above the Tonto Platform mid-canyon, including Shasta ground sloth (*Nothrotheriops shastensis*), camel (*Camelops* sp.)., Harrington's mountain goat (*Oreamnos harringtoni*), bighorn (*Ovis canadensis*), bison (*Bison* sp.), and horse (*Equus* sp.) (Mead 1981).




Figure 11-4. Ascending into Skylight Cave, 1984 (EMILEE MEAD).

Select Localities

Not all caves and packrat midden localities will be described here, only those with a more complex, unusual, and/or diverse fossil assemblage story. The descriptions below are arranged based on their approximate down-canyon/river location, beginning up-river.

Stanton's Cave and Skull Cave

Stanton's Cave (named after Robert Brewster Stanton) is a large Redwall Limestone (Mississippian) cavern along the Marble Canyon river corridor. Nearby is Vasey's Paradise (named after George Vasey, a friend of John Wesley Powell), a gushet spring flowing into the river (Springer et al. 2008) that appears to show how Stanton's Cave was formed in the distant past. The cave has a long history of use in part due to easy access from the river and from the canyon rim country above via nearby South Canyon. Euler (1984) provides a synthesis of the historic and prehistoric use of the cave. Much of the surface deposit is composed of Holocene-age sediments containing bighorn (*Ovis canadensis*) dung and archaeological artifacts. Excavations in 1969–1970 produced a wealth of archaeological and paleontological information (see chapters in Euler 1984). Dryer (1994) produced a research

project on the packrat middens recovered from the back room of the cave—an area not thoroughly studied previously. The dry environment within the cave provided a wealth of information about the extinct *Oreamnos harringtoni* with the preservation of horn sheaths and dung (Mead et al. 1986; Mead and Lawler 1994). Today, a large steel lattice keeps human visitors from entering the cave but bats and other animals can still utilize the cave.

Packrat middens from the back of the cavern produced ^{14}C ages ranging from about 11,000 to 35,000 years ago (Dryer 1994). Radiocarbon dates on dung, wood, and a bone of an extinct bird (*Teratornis*) produced ages ranging from as young as 1,500 to as old as about 17,000 (uncorrected) yr B.P.; drift wood at the base of the sediments dated to > 35,000 years old (Robbins et al. 1984).

Skull Cave is a rather small cavern that divides into three separate passages (Emslie 1988). Analysis of three test pits indicated that for the most part the cave was used by packrats and birds, producing an impressive avifauna (Emslie 1988). Only a few radiometric ages provide a preliminary chronology via a uranium series date on anhydrite from test pit sediments and AMS ages obtained from packrat pellets. All ages are less than 20,000 yr B.P.

Sandblast Cave and Nearby Caves

A series of cliff-entrance caves can be found in the exposed Redwall Limestone in the Marble Canyon river corridor. Probably the most significant locality is Sandblast Cave (Figure 11-5) which is a grouping of crevices and tunnels (Emslie 1988). Excavations produced important data about the condor (*Gymnogyps californianus*, including a preserved nest; Figure 11-1) in addition to specimens of the extinct mountain goat, bison, camel, and horse, along with the only reported mammoth remains from the GC; these large mammal remains are thought to be remnants of food items brought in by condors (Emslie 1987, 1988; Mead et al. 2003). Other caves with fossils in the corridor stretch include Skylight (Figure 11-4) and Hummingbird caves, among others (Emslie 1987). Radiocarbon ages on *Gymnogyps* skeletal remains range from about 10,000 to 13,000 (uncorrected) yr B.P. ^{14}C ages on packrat middens and *Oreamnos* dung are all older than 30,000 and wood dating in excess of 40,000 yr B.P. (Emslie 1987, Mead et al. 2003).



Figure 11-5. Close-up of the entrance to Sandblast Cave showing the series of openings, 1984 (EMILEE MEAD).

Little Nankoweap

The Little Nankoweap drainage is known to have countless caves, many containing important archaeological and paleontological remains (Emslie et al. 1987; Mead and Lawler 1994; Emslie et al. 1995). Crescendo (CC:5:1), Rebound (CC:5:5), Left and Right Eye, Five-Windows (CC:5:2), Shrine (CC:5:3), and Stevens (CC:5:4; Figure 11-6) caves have been the most intensely studied and described, but many chambers and passageways in these caves still contain numerous areas and deposits that remain to be fully analyzed. Besides data on condors (Emslie 1987, 1988), a series of packrat middens produced copious plant macrofossils (Coats et al. 2008) and faunal remains (Carpenter and Mead 2000; Mead et al. 2003). The entrances to most of these caves are on the sheer cliff face of the Redwall Limestone well out of the main river corridor. Some of the flora and fauna (e.g., extinct camel and the extinct shrubox *Euceratherium collinum*) recovered from these cave deposits likely represent inhabitants of the flat plateau surface immediately above (Figure 11-7),

which also provide access to the higher plateaus of the North Rim, and not the narrow, steep river corridor and side-canyons. A number of the caves have multiple packrat middens and *Oreamnos* dung remains dating from about 11,000 to 46,000 yr B.P. (Mead et al. 2003).



Figure 11-6. Stevens Cave entrance located high on the cliff face of the Redwall Limestone, 1984 (EMILEE MEAD).



Figure 11-7. An expansive flat region exists above the Redwall Limestone caves in the Little Nankoweap Canyon region. With access from the higher North Rim plateau region above, large artiodactyls such as camels (*Camelops*), shrubox (*Euceratherium*), and bison (*Bison*), not cliff-climbers, were able to easily enter into portions of the eastern Grand Canyon (JIM MEAD).

Hance Canyon and Horseshoe Mesa

A series of packrat middens from Bida Cave were described by Cole (1981, 1985). Faunal remains from the middens were reported in Cole and Mead (1981). Surface remains and a test pit produced a wealth of information about the skeleton, diet, and habitat of *Oreamnos harringtoni* (Mead 1983; O'Rourke and Mead 1985; Mead et al. 1986; Mead, Martin et al. 1986; Mead and Lawler 1994). The cave is extensive, with the lower entrance a large, gaping cavern. At the point where the main cavern turns and narrows into a tunnel and all outside light ceases, bedding depressions are littered with *O. harringtoni* dung (Mead 1983). The cave system goes through a series of small tubes and large rooms, many with additional fossil and subfossil remains that have yet to be fully documented and studied. Ultimately the cave emerges at the top of the Redwall Limestone as a small entrance providing rare access through the cliff to the Inner Gorge region.

Multiple packrat middens have been dated from Bida Cave ranging from 8,000 to about 13,000 (uncorrected) yr B.P. (Cole 1981). *Oreamnos* dung and keratinous horn sheaths range in ^{14}C age from 12,000 to about 25,000 (uncorrected) yr B.P. (Mead 1983; Mead et al. 1986).

The caves and dry crevices of Horseshoe Mesa up Hance and Cottonwood canyons have been studied speleologically and to a certain extent archaeologically (see Farmer and deSaussure 1955), but are

poorly known paleontologically. Ancient packrat middens were recovered from many areas in these two canyons and a few across the river (Cole 1985). Cole (1981, 1990) reviews all the radiocarbon ages from multiple packrat midden dating from about 10,000 to 35,000 (uncorrected) years old.

Cremation Creek Caves

The first realization that many caves preserve organic remains for a long period of time came with the report of split-twig figure artifacts at sites on Cremation Creek (Farmer and deSaussure 1955). The greater Cremation Creek provides a number of caves that were explored in the early 1950s and found to contain Ice Age fossils. Marmot (*Marmota*) remains (not found in the region today at these elevations) were recovered in Tse-an Olje, Cylinder, Tse-an Kaetan, and Tooth caves (Lange 1956; among other caves across the river). Caves being explored in the 1950s were to be kept secret so the names applied by Lange (1956) and deSaussure (1956) were given in the Navajo (Diné) language; tsé'áán refers to "rock cave".

Only Kaetan Cave ("prayer stick cave"; Figure 11-8) in this area is fairly well studied; it is a small cavern with abundant archaeological material in the entrance chamber (Schwartz et al. 1958). Although the cave has a dry entrance chamber, it becomes wet further in with pools. Because of the exploration in 1955 for split-twig figures, Schwartz et al. (1958) also provided the earliest Ice Age palynological analysis in the GC. This then pointed the way for further work in the cave in the late 1970s (Mead 1983; O'Rourke and Mead 1985). An owl roost deposit in the cave was sampled but never studied and remains to be examined. As with Bida Cave, *Oreamnos harringtoni* remains were abundant (Mead and Lawler 1994). Packrat middens with plant and micromammal remains are ^{14}C 15,000 to 19,000 (uncorrected) yr B.P. A stratified test pit produced uncorrected ages from 14,000 to 30,000 yr B.P. An owl roost deposit may date back to 21,000 yr B.P. (Mead 1983).




Figure 11-8. Entrance to Kaetan Cave, 1980 (JIM MEAD).

Double Bopper Cave

Double Bopper Cave is the longest known cave in GRCA. This cave developed in Redwall Limestone, and is located in a remote part of the north rim (Figure 11-9). Difficult to access and largely hidden from view, the cave was only discovered in 2008. Annual expeditions have mapped over 64 km (40 mi) of passage, making it the longest known cave in Arizona and among the 50 longest caves in the world. The cave is still being actively mapped, so the length will continue to increase with future exploration. Double Bopper Cave is a maze cave characterized by rectilinear, joint-controlled passage development (see Tobin et al. in press). Parallel primary passages are large and relatively easy to travel through. Between these primary corridors, smaller passages connect the parallels, sometimes with dense and complicated cave development. The cave is predominantly horizontal, though there is multi-level development in a few isolated areas with vertical passages connecting the levels. The massive extent of the cave and diversity of passage sizes provides a considerable amount of habitat for subterranean wildlife.



Figure 11-9. The two main entrances to Double Bopper Cave. Only the best of climbers and fliers can enter the cave via these entrances. Note that the green dot below the right entrance is a person entering the cave (NPS/SHAWN THOMAS).

One of the most unique features of Double Bopper Cave is the abundance of mummified bats, which are typically rare or altogether absent from other GRCA caves. Bats still actively use the cave with flyways indicated by fresh guano deposits. However, live bats are rarely seen in the cave aside from bats exiting the main entrance at dusk and occasional solitary individuals in torpor. Mummified bats occur throughout the cave, especially along the larger parallel passages. Mummies are typically found clinging to walls and secondary gypsum formations but are also found scattered on the ground. The stable microclimate conditions in the cave, with low relative humidity (typically 35–45%), have likely persisted for thousands of years, making for excellent preservation of specimens. Most mummified bats can be identified to species, possessing intact skin and fur, and many mummy specimens "roosting" on walls cannot immediately be distinguished from live bats without closer examination (Figure 11-3).

Though a complete census has not been conducted, estimates from survey teams suggest the cave contains many hundreds to thousands of bat mummies. At least eight bat species have been identified in the cave. Townsend's big-eared bat (*Corynorhinus townsendii*), a cave-obligate species, is the most common. Less abundant but still commonly observed are pallid bats (*Antrozous pallidus*), big

brown bats (*Eptesicus fuscus*), and myotid species (*Myotis* spp.). Canyon bats (*Parastrellus hesperus*) and free-tailed bats (*Tadarida brasiliensis* and possibly *Nyctinomops femorosaccus*) are also present. The rarest species, consisting of only a few specimens, include hoary bats (*Lasiurus cinereus*) and silver-haired bats (*Lasionycteris noctivagans*), which is unusual in that these species are typically considered tree-dwelling bats. A nearby cave, Leandras Cave, also contains abundant bat mummies but with a suspected higher proportion of hoary bats and silver-haired bats (future inventory work is being planned to answer this question). Radiocarbon dating of a subset of bat mummy tissues collected from Double Bopper Cave yielded ages ranging from 3,500 to 33,650 yr B.P., demonstrating long-term use of the cave by bats.

Double Bopper Cave also contains mummified remains of other mammals. Packrats (*Neotoma* spp.) are far outnumbered by bats but still common. Several ringtail (*Bassariscus astutus*) specimens have been found deep in the cave in excellent states of preservation (Figure 11-10) but have not been radiocarbon dated yet. Other skeletonized specimens have yet to be fully assessed, including one gray fox (*Urocyon cinereoargenteus*; Figure 11-11). Inventory and carbon dating continue for paleontological resources in Double Bopper Cave.



Figure 11-10. A mummified ringtail (*Bassariscus astutus*) from Double Bopper Cave. With the carcass still with all its hair and it only being slightly modified in color, the age of this individual is likely fairly recent. Other carcasses of this taxon in the cave will be radiocarbon dated (NPS/SHAWN THOMAS).



Figure 11-11. An articulated gray fox skeleton (not radiocarbon dated) from Double Bopper Cave (NPS/SHAWN THOMAS).

Rampart, Muav, and Vulture Caves

Rampart, Muav, and Vulture caves are located in the far western end of the GC not far from where the river exits the Colorado Plateau and heads across the Basin and Range Province. This series of caves and packrat midden studies are about 305 km (190 mi) west of the other well-known fossil localities mentioned above. Only a few packrat midden localities have been found in between (Van Devender and Mead 1976; Cole 1985).

The entrance to Rampart Cave is a fair distance up a long, steep talus slope from the Colorado River. Today there is a short inclined climb into the cave, but there may have been more of a subtle ramp into the cave during the Pleistocene based on the occurrence of Shasta ground sloth and tortoise remains in the deposit.

The cave was the first location studied in the GC for Pleistocene biotic remains. In 1936 the CCC (Civilian Conservation Corp) produced a few test pits (Figure 11-12). Laudermilk and Munz (1938) described the plant remains from the dry-preserved dung. Subsequent excavation in the cave was made by Remington Kellogg of the National Museum of Natural History in 1942. Preliminary description of the faunal remains was published by Wilson (1942). Martin et al. (1961) provided the

first detailed paleoenvironmental and chronological study of cave contents with the analysis of dung from the 1956 Shutler Profile. A detailed analysis of radiocarbon-dated sloth dung was produced by Long and Martin (1974; Long et al. 1974). Fortunately, the Kellogg test trench was excavated to the limestone cave floor, because in 1976–1977 the bulk (~70%) of the deposit was destroyed by a smoldering fire set by an unauthorized visitor. The test trench produced a fire break saving some of this non-renewable fossil deposit. An exhaustive history of the various field studies in the cave (including field notes, early photographs, and the relocation of field maps) and an assessment of the remaining, unburned deposit was presented in Carpenter (2004). The entrance today requires permission and a key to open the steel gate. Studies of packrat middens from Rampart Cave and a multitude of isolated limestone crevice localities in a number of nearby canyons were published by Phillips (Phillips and Van Devender 1974; Mead and Phillips 1981; Phillips 1984). Sloth dung from the 1.5 m (5 ft) deep deposit produced ages from as young at 11,000 (uncorrected) years old, back to greater than 40,000 (Long and Martin 1974).

Muav Caves is a series of small tubes just above the pre-dam river level and now just above the high-stand of Lake Mead. The caves are best known for the remains of Shasta ground sloth (Long and Martin 1974; Long et al. 1974). Although some test pits were made in the cave entrance long ago, very little is understood about the contents of the deposits. In some crevices below these caves are a series of packrat middens and ringtail refuse den deposits in the Whipple Cliffs, but these remains have not been published at this time (Mead et al. in progress).

Vulture Cave is primarily a low crawlway along short passageways, but all areas are heavily congested with a multitude of packrat middens, floor deposits, and ringtail den debris (Mead and Phillips 1981; Mead and Van Devender 1981). Besides remains of *Gymnogyps californianus*, *Cathartes aura* (turkey vulture), and *Camelops* sp. the deposits provided a wealth of information about the Ice Age desert and woodland herpetofauna (Mead and Phillips 1981). Packrat midden contents ranged in age from 1,100 to 33,000 (uncorrected) yr B.P. (Mead and Phillips 1981). Important herpetological data came from a ringtail den ^{14}C dated to 2,000 yr B.P. (Mead and Van Devender 1981).



Figure 11-12. Rampart Cave in the 1930s. **A.** Photograph by the CCC (Civilian Conservation Corp) of the Shasta sloth (*Nothrotheriops shastensis*) dung deposit prior to excavation in the 1930s and the fire in 1975–1976. **B.** Photograph of the CCC excavating a Shasta sloth skeleton from the dung deposit in the 1930s.

Environments and Climate Discussion

When one thinks of the GC, it is often visualized as an uncomplicated, sinuous, deep gorge east to west, with high elevations with forests at the rims, and low elevations with hot desert habitats along the roaring river at its spine down below. Conceptually this viewpoint may be good, but the real understanding is that the modern canyon is truly complex in all aspects, and this was equally true during the Pleistocene. The eastern region is distinctly different from the western end—climatically, ecologically, and geologically. The differences are in the details. To understand the record of the Pleistocene (Ice Age), preservation is at the core to the recovery of the details. Any cave or shelter with the occurrence of split-twig figures (e.g., Farmer and deSaussure 1955; Emslie et al. 1995) illustrates that preservation of these Archaic cultural remains in the chamber is ideal enough that Ice Age remains are more than likely also present and the cavern should be assessed for them as well. Split-twig figures are part of the Archaic culture that occurred throughout the GC region and elsewhere on the Colorado Plateau, all since the Ice Age and within the Holocene. Overviews of this cultural period as it relates to the GC region can be found in Geib (1995), Huckell (1986), and Janetski et al. 2012).

The CP is a distinct physiographic province, straddling the present transition between summer-wet climatic regimes to the south and summer-dry climatic regimes to the north. With the tremendous topographic diversity of the region, there are extremes in available habitats and plant communities today; and these attributes were certainly expressed with the Ice Age climate regimes. Anderson et al. (2000) provides a detailed synthesis of many southern CP late Ice Age localities for paleoclimatic and paleobotanic reconstructions.

The data about world-wide changes in temperature are derived from deep sea core samples. A multitude of fossil forms have been used to create a world-wide record of climate and temperature changes. Different phases are grouped into like clusters termed marine isotope stages (MIS) or oxygen isotope stages (OIS) (see Cronin 2010; Bradley 2015, 3rd edition). For the GC region, the preserved dataset permits one to examine MIS 1 (14,059 cal. yr BP to present), MIS 2 (27,500–14,060 cal yr BP, and MIS 3 (59,000–27,501 cal yr BP). Clearly much of the Pleistocene (~2.59 million to 11,000 years ago) is not understood. More is known for other regions on the CP, but these deposits are not to be recovered in the GC.

Some of the best high elevation paleoecological data comes from stratified sediment cores taken from lakes that are not found within the limestone region of the GC, but elsewhere on the southern CP (Potato Lake, Anderson 1993; Walker Lake, see Anderson et al. 2000, for detailed discussion of dataset; Hay Lake, Jacobs 1985).

Data about MIS 2 and MIS 3 plant communities above about 2,800 m (9,190 ft) elevation are presently not fully understood. Lake-core pollen records at about 2,700 m (8,860 ft) suggest that during MIS 3 high-elevation pine species (perhaps bristlecone and or limber) mixed with Engelmann spruce and subalpine fir created an open forest, possibly with sagebrush growing in open areas. Calculated average summer temperatures were about 3–4°C cooler during MIS 3 than at present (Anderson et al. 2000).

The knowledge of plant communities between 1,600–2,100 m (5,250–6,890 ft) elevation is minimal due to the lack of fossil data. One thing is consistent with the preserved macrobotanical record, there is a lack of ponderosa pine from MIS 3–2. Below 1,500 m (4,940 ft) elevation to about 450 m (1,380 ft) a juniper-desertscrub open woodland persisted, including sagebrush, prickly pear cactus, agave, and, lower down, saltbush. Looking down into the GC during the Ice Age, one would be overwhelmed with this open woodland desertscrub community growing clear to the river's edge. Side canyons would have stands of Douglas fir and white fir in the wetter habitats. A mixed-conifer forest (montane conifers) with limber pine, white fir, and Douglas fir occupied the upper slopes, buttes, and rim county.

The transition from MIS 2 to MIS 1 (~14,000 cal yr BP) occurred with a major reorganization of the plant communities from rim to river. Many species of mixed-conifer forest retreated upslope to attain their near-current elevation distributions. Ponderosa pine quickly became established across middle elevations, as they are today. At lower elevations desertscrub communities replaced the juniper woodlands. All species did not move in concert to these climate-induced changes; instead, what is observed is a mosaic change. Figure 11-13 shows diagrammatically the inferred elevational distribution of plant communities in and adjacent to the GC during the most recent approximately 50,000 years. The data seem to imply that during the last glacial, the southern position of the jet stream, a cooler tropical ocean, and a heavy spring snowpack over the CP and adjacent Colorado Rocky Mountains probably conspired to suppress the monsoonal flow pattern (Anderson et al. 2000). During this time the seasonality of precipitation appears to have been dampened to predominate during the winter storms. The change to a summer precipitation maximum at the beginning of MIS 1 may be recorded by the sudden appearance and rapid migration of ponderosa pine across the southern CP. Data suggest that the mean depression of late glacial temperatures was at least 5° C (~9° F) colder than today (Anderson et al. 2000).

The vertebrate species were likewise responding not only to the temperature and precipitation changes (i.e., they are affected directly by these parameters) as discussed above but also to the modifications to the local plant community changes (i.e., their food source and/or their habitat requirements). Some species were not directly affected by these parameters and have not changed their distribution within the greater GC region, such as the bighorn, possibly the bison, and Gila monster (*Heloderma suspectum*). Other species appear to have moved up in elevation but stayed within the overall region (some species of voles), or moved out of the GC region to other areas of the continent (*Gymnogyps californianus*), or died out completely (extinction occurred), such as *Oreamnos harringtoni*, *Euceratherium collinum*, *Nothrotheriops shastensis*, *Camelops* sp., and likely some carnivores.

The trends in climate and climate-induced biotic changes over the most recent 50,000 years are based on the data discussed above. It must be remembered that these statements are based on an incomplete fossil record, both through time and for the length of the GC. Packrat midden and dry-preserved cave data have at least a 360 km (190 mi) gap beginning in the eastern GC and going throughout much of the western GC. Most of the above data is really the eastern GC. Data are sparse to almost non-existent for the Hualapai Plateau on the southwest end of the GC. The topographic structure of the

northwestern GC (Shivwitts and Uinkaret plateaus) is completely different than the eastern GC so it should be expected that biotic communities and climate responses would have been different over the past 50,000 years. Much still needs to be recovered and assessed to understand the details surrounding this rapid and critical change in climate along with plant community distributional changes for the greater GC.




Figure 11-13. Inferred elevational distribution of plant communities within and adjacent to the Grand Canyon during the most recent 50,000 years. Cross-section is a line oriented from the San Francisco Peaks north through the Grand Canyon to the North Rim region (Anderson et al. 2000: Figure 8).

Acknowledgments

We thank Kim Besom and Colleen Hyde, Collections, GRCA for assisting us with identifying specimens in the collections that have not been fully published upon, and Janet Gillette, Museum of Northern Arizona, for assistance with Stanton's Cave material. We also recognize the contributions, comments, and reviews offered by Diana Boudreau, Robyn Henderek, Anne Miller, Gary Morgan, and Klara Widrig.

Literature Cited

Adams, D. K., and A. C. Comrie. 1997. The North American monsoon. Bulletin of the American Meteorological Society 78:2197–2214.

Anderson, R. S. 1993. A 35,000 year vegetation and climate history from Potato Lake, Mogollon Rim, Arizona. Quaternary Research 40:351–359.

Anderson, R. S., and T. R. Van Devender. 1991. Comparison of pollen and macrofossils in packrat (*Neotoma*) middens: A chronological sequence from the Waterman Mountains of southern Arizona, U.S.A. Review of Palaeobotany and Palynology 68:1–28

Anderson, R. S., J. L. Betancourt, J. I. Mead, R. H. Hevly, and D. P. Adam. 2000. Middle- and late-Wisconsin paleobotanic and paleoclimatic records from the southern Colorado Plateau, USA. Palaeogeography, Palaeoclimatology, Palaeoecology 155:31–57.

Beus, S. S., and M. Morales, editors. 2003. Grand Canyon geology (2nd edition). Oxford University Press, New York, New York.

Carpenter, M. C. 2002. *Vulpes vulpes* (red fox) remains from Stanton's Cave, Arizona: First known record from the Grand Canyon. Journal of Vertebrate Paleontology 22(supplement to 3):41A.

Carpenter, M. C. 2003. Late Pleistocene Aves, Chiroptera, Perissodactyla, and Artiodactyla from Rampart Cave, Grand Canyon, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Carpenter, M. C., and J. I. Mead. 2000. A mummified canid from a cave in the Grand Canyon, Arizona. Journal of Vertebrate Paleontology 20(supplement to 3):33A.

Coats, L. 1997. Middle to Late Wisconsinan vegetation change at Little Nankoweap, Grand Canyon National Park, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Coats, L. L., K. L. Cole, and J. I. Mead. 2008. 50,000 years of vegetation and climate history on the Colorado Plateau, Utah and Arizona, USA. Quaternary Research 70(2):322–338.

Cockerell, T. D. A. 1927. A large form of *Oreohelix yavapai* in the Grand Canyon. The Nautilus 40(3):101.

Cole, C. J., and T. R. Van Devender. 1976. Surface structure of fossil and Recent epidermal scales from North American lizards of the genus *Sceloporus* (Reptilia, Iguanidae). Bulletin of the American Museum of Natural History 156(4):451–514.

Cole, K. 1982. Late Quaternary zonation of vegetation in the eastern Grand Canyon. Science 217(4565):1142–1145.

Cole, K. 1985. Past rates of change, species richness, and a model of vegetational inertia in the Grand Canyon, Arizona. The American Naturalist 125(2):289–303.

Cole, K. L. 1990. Reconstruction of past desert vegetation along the Colorado River using packrat middens. Palaeogeography, Palaeoclimatology, Palaeoecology 76:349–366.

Cole, K. L. 1990. Late Quaternary vegetation gradients through the Grand Canyon. Pages 240–258 *in* J. L. Betancourt, T. R. Van Devender, and P. S. Martin, editors. Packrat middens: The last 40,000 years of biotic change. University of Arizona Press, Tucson.

Cole, K., and J. L. Mead. 1981. Late Quaternary animal remains from packrat middens in the eastern Grand Canyon, Arizona. Journal of the Arizona-Nevada Academy of Science 16(1):24–25.

Cole, K. L., and L. Mayer. 1982. Use of packrat middens to determine rates of cliff retreat in the eastern Grand Canyon, Arizona. Geology 10:597–599.

Cole, K. L., J. F. Fisher, K. Ironside, J. I. Mead, and P. Koehler. 2013. The biogeographic histories of *Pinus edulis* and *Pinus monophylla* over the last 50,000 years. Quaternary International 310:96–110.

deSaussure, R. E. 1956. Remains of the California Condor in Arizona caves. Plateau 29(2):44–45.

Dryer, J. E. 1994. Late Pleistocene vegetation change at Stanton's Cave, Colorado River, Grand Canyon National Park, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Elias, S. A., J. I. Mead, and L. D. Agenbroad. 1992. Late Quaternary arthropods from the Colorado Plateau, Arizona and Utah. Great Basin Naturalist 52(1):59–67.

Emslie, S. D. 1987. Age and diet of fossil California condors in Grand Canyon, Arizona. Science 237(4816):768–770.

Emslie, S. D. 1988. Vertebrate paleontology and taphonomy of caves in Grand Canyon, Arizona. National Geographic Research 4(1):128–142.

Emslie, S. D., R. C. Euler, and J. I. Mead. 1987. A desert culture shrine in Grand Canyon, Arizona, and the role of split-twig figurines. National Geographic Research 3:511–516.

Emslie, S. D., J. I. Mead, and L. Coats. 1995. Split-twig figurines in Grand Canyon, Arizona: New discoveries and interpretations. Kiva 61:145–173.

Euler, R. C. 1978. Archeological and paleobiological studies at Stanton's Cave, Grand Canyon National Park, Arizona; a report of progress. Research Reports - National Geographic Society 10(1969 Projects):141–162.

Euler, R. C., editor. 1984. The archaeology, geology, and paleobiology of Stanton's Cave, Grand Canyon National Park, Arizona. Grand Canyon Natural History Association, Grand Canyon, Arizona. Monograph 6.

Evans, W. 1936. Work progress report for the archeological project at Pierces Ferry. On file at Grand Canyon National Park, Grand Canyon, Arizona.

Farmer, M. F., and R. deSaussure. 1955. Split-twig animal figurines. Plateau 27:13–23.

Fenton, C. R., R. H. Webb, T. E. Cerling, R. J. Poreda, and B. P. Nash. 2002. Cosmogenic ^{3}He ages and geochemical discrimination of lava-dam outburst-flood deposits in western Grand Canyon, Arizona. Pages 191–215 *in* P. K. House, R. H. Webb, V. R. Baker, and D. R. Levish, editors. Ancient floods, modern hazards: Principles and applications of paleoflood hydrology. American Geophysical Union, Washington, D.C. Water Science and Application Series 5.

Geib, P. R. 1995. Radiocarbon record for Archaic occupation of the Central Colorado Plateau. Pages 89–136 *in* C. van Riper, III, editor. Proceedings of the second biennial conference on research in Colorado Plateau National Parks. Transactions and Proceedings Series NPS/NRNAU/NRTP-95-11.

Hamblin, W. K. 1994. Late Cenozoic lava dams in the western Grand Canyon. Geological Society of America, Boulder, Colorado. Memoir 183.

Hansen, R. M. 1974. Dietary of the chuckwalla, *Sauromalus obesus*, determined by dung analysis. Herpetologica 30:120–123.

Hansen, R. M. 1978. Shasta ground sloth food habits, Rampart Cave, Arizona. Paleobiology 4(3):302–319.

Harington, C. R. 1972. Extinct animals of Rampart Cave. Canadian Geographical Journal 85(5):178–183.

Higgins, R. W., Y. Yao, and X. L. Wang. 1997. Influence of the North American monsoon system on the United States summer precipitation regime. Journal of Climatology 10:2600–2622.

Hill, C. A, and V. J. Polyak. 2010. Karst hydrology of Grand Canyon, Arizona, USA. Journal of Hydrology 390:169–181.

Huckell, B. B. 1996. The Archaic prehistory of the North American Southwest. Journal of World Prehistory 10:305–373

Hunt, A. P., and S. G. Lucas. 2018. The record of sloth coprolites in North and South America: Implications for terminal Pleistocene extinctions. New Mexico Museum of Natural History and Science Bulletin 79:277–298.

Huntoon, P. W. 2000. Variability of karstic permeability between unconfined and confined aquifers, Grand Canyon region, Arizona. Environmental and Engineering Geoscience 6(2):155–170.

Jacobs, B. F. 1985. A middle Wisconsin pollen record from Hay Lake, Arizona. Quaternary Research 24:121–130.

Janetski, J. C., M. L. Bodily, B. A. Newbold, and D. T. Yoder. 2012. The Paleoarchaic to Early Archaic transition on the Colorado Plateau: the archaeology of North Creek Shelter. American Antiquity 77:125–159

Jones, C. J. R, A. E. Springer, B. W. Tobin, S. J. Zappitello, and N. A. Jones. 2018. Characterization and hydraulic behaviour of the complex karst of the Kaibab Plateau and Grand Canyon National Park, USA. Pages 237–260 *in* M. Parise, F. Gabrovsek, G. Kaufmann, and N. Ravbar, editors. Advances in karst research: Theory, fieldwork and applications. Geological Society of London, London, United Kingdom. Special Publication 466.

Kaufman, D. S., G. O'Brien, J. I. Mead, J. Bright, and P. Umhoefer. 2002. Late Quaternary spring-fed deposits of the Grand Canyon and their implication for deep lava-dammed lakes. Quaternary Research 58:329–340.

Kenworthy, J., V. L. Santucci, and K. L. Cole. 2004. An inventory of paleontological resources associated with caves in Grand Canyon National Park. Pages 211–228 *in* C. van Riper, III and K. L. Cole, editors. The Colorado Plateau: Cultural, biological and physical research. University of Arizona Press, Tucson, Arizona.

Kropf, M., J. I. Mead, and R. S. Anderson. 2007. Dung, diet, and the paleoenvironment of the extinct shrub-ox (*Euceratherium collinum*) on the Colorado Plateau, USA. Quaternary Research 67(1):143–151.

Lange, M. C. 1956. Woodchuck remains in northern Arizona caves. Journal of Mammalogy 37(2):289–291.

Laudermilk, J. D., and P. A. Munz. 1938. Plants in the dung of *Nothrotherium* from Rampart and Muav Caves, Arizona. Carnegie Institution of Washington Publication 487:271–281.

Lindsay, E. H., and N. T. Tessman. 1974. Cenozoic vertebrate localities and faunas in Arizona. Journal of the Arizona Academy of Science 9(1):3–24.

Long, A., and P. S. Martin. 1974. Death of American ground sloth. Science 186:638–640.

Long, A., R. M. Hansen, and P. S. Martin. 1974. Extinction of the Shasta Ground Sloth. Geological Society of America Bulletin 85(12):1843–1848.

Martin, J. M. 2014. Late Pleistocene and Holocene *Bison* of Grand Canyon and Colorado Plateau: Implications from the use of paleobiology for natural resource management policy. Thesis. East Tennessee State University, Johnson City, Tennessee.

Martin, J. M., R. A. Martin, and J. I. Mead. 2017. Late Pleistocene and Holocene *Bison* of the Colorado Plateau. The Southwestern Naturalist 62(1):14–28.

Martin, P. S., B. E. Sabels, and D. Shutler, Jr. 1961. Rampart Cave coprolite and ecology of the Shasta ground sloth. American Journal of Science 259(2):102–107.

Martin, T., and D. Whitis. 2018. Guide to the Colorado River in the Grand Canyon: Lees Ferry to South Cove. Vishnu Temple Press, Flagstaff, Arizona.

Mead, J. I. 1981. The last 30,000 years of faunal history within the Grand Canyon, Arizona. Quaternary Research 15(3):311–326.

Mead, J. I. 1983. Harrington's extinct mountain goat (*Oreamnos harringtoni*) and its environment in the Grand Canyon, Arizona. Dissertation, University of Arizona, Tucson.

Mead, J. I. 2005. Late Pleistocene (Rancholabrean) amphibians and reptiles of Arizona. New Mexico Museum of Natural History and Science Bulletin 29:137–152.

Mead, J. I., and T. R. Van Devender. 1981. Late Holocene diet of *Bassariscus astutus* in the Grand Canyon, Arizona. Journal of Mammalogy 62(2):439–442.

Mead, J. I., and A. M. Phillips, III. 1982. The late Pleistocene and Holocene fauna and flora of Vulture Cave, Grand Canyon, Arizona. The Southwestern Naturalist 26(3):257–288.

Mead, J. I., and M. C. Lawler. 1994. Skull, mandible, and metapodials of the extinct Harrington's mountain goat (*Oreamnos harringtoni*). Journal of Vertebrate Paleontology 14:562–576.

Mead, J. I., and D. G. Mikesic. 2001. First fossil record of *Euderma maculatum* (Chiroptera: Vespertilionidae), eastern Grand Canyon, Arizona. Southwestern Naturalist 46:380–383.

Mead, J. I., and S. L. Swift. 2012. Late Pleistocene (Rancholabrean) dung deposits of the Colorado Plateau, western North America. New Mexico Museum of Natural History and Science 57:337–342.

Mead, J. I., R. S. Thompson, and A. Long. 1978. Arizona radiocarbon dates IX: Carbon isotope dating of packrat middens. Radiocarbon 20(2):171–191.

Mead, J. I., M. K. O'Rourke, and T. M. Foppe. 1986. Dung and diet of the extinct Harrington's mountain goat (*Oreamnos harringtoni*). Journal of Mammalogy 67(2):284–293.

Mead, J. I., P. S. Martin, R. C. Euler, A. Long, A. J. T. Jull, L. J. Toolin, D. J. Donahue, and T. W. Linick. 1986. Extinction of Harrington's mountain goat. Proceedings of the National Academy of Science 83:836–839.

Mead, J. I., L. L. Coats, and B. W. Schubert. 2003. Late Pleistocene faunas from caves in the eastern Grand Canyon, Arizona. Pages 64–86 *in* B. W. Schubert, J. I. Mead, and R. W. Graham, editors. Ice Age cave faunas of North America. Indiana University Press, Bloomington and Indianapolis, Indiana.

Miller, L. 1960. Condor remains from Rampart Cave, Arizona. The Condor 62(1):70.

Miller, R. R., and G. R. Smith. 1984. Fish remains from Stanton's Cave, Grand Canyon of the Colorado, Arizona, with notes on the taxonomy of *Gila cypha*. Pages 61–65 *in* R. C. Euler, editor. The archaeology, geology, and paleobiology of Stanton's Cave, Grand Canyon National Park, Arizona. Grand Canyon Natural History Association, Grand Canyon, Arizona. Monograph 6.

Mock, C. J. 1996. Climatic controls and spatial variations of precipitation in the western United States. Journal of Climatology 9:1111–1125.

Olsen, J. W., and S. J. Olsen. 1984. Zooarcheological analysis of small vertebrates from Stanton's Cave, Arizona. Pages 47–57 *in* R. C. Euler, editor. The archaeology, geology, and paleobiology of Stanton's Cave, Grand Canyon National Park, Arizona. Grand Canyon Natural History Association, Grand Canyon, Arizona. Monograph 6.

O'Rourke, M. K., and Mead, J. I. 1985. Late Pleistocene and Holocene pollen records from two caves in the Grand Canyon of Arizona, USA. Pages 169–185 *in* B. F. Jacobs, P. L. Fall, and O. K. Davis, editors. Late Quaternary vegetation and climates of the American Southwest. American Association of Stratigraphic Palynologists, Houston, Texas. Contributions Series 16.

Pape, R. B. 2014. Biology and ecology of Bat Cave, Grand Canyon National Park, Arizona. Journal of Cave and Karst Studies 76:1–13.

Phillips, A. M., III, and T. R. Van Devender. 1974. Pleistocene packrat middens from the lower Grand Canyon of Arizona. Journal of the Arizona Academy of Science 9(3):117–119.

Phillips, A. M., III. 1984. Shasta ground sloth extinction: Fossil packrat midden evidence from the western Grand Canyon. Pages 148–158 *in* P. S. Martin and R. G. Klein, editors. Quaternary extinctions: A prehistoric revolution. University of Arizona Press, Tucson, Arizona.

Pilsbry, H. A., and J. H. Ferriss. 1911. Mollusca of the southwestern states, V: The Grand Canyon and northern Arizona. Proceedings of the Academy of Natural Sciences of Philadelphia 63:174–199.

Polyak, V. J., C. A. Hill, and Y. Asmerom. 2008. Age and evolution of the Grand Canyon revealed by U-Pb dating of water table-type speleothems. Science 319(5868):1377–1380.

Ray, C. E., O. J. Linares, and G. S. Morgan. 1988. Paleontology. Pages 19–30 *in* A. M. Greenhall and U. Schmidt, editors. Natural history of vampire bats. CRC Press, Boca Raton, Florida.

Rea, A. M., and L. L. Hargrave. 1984. The bird bones from Stanton's Cave. Pages 77–91 *in* R. C. Euler, editor. The archaeology, geology, and paleobiology of Stanton's Cave, Grand Canyon National Park, Arizona. Grand Canyon Natural History Association, Grand Canyon, Arizona. Monograph 6.

Robbins, E. I., P. S. Martin, and A. Long. 1984. Paleoecology of Stanton's Cave Grand Canyon, Arizona. Pages 115–130 *in* R. C. Euler, editor. The archaeology, geology, and paleobiology of Stanton's Cave, Grand Canyon National Park, Arizona. Grand Canyon Natural History Association, Grand Canyon, Arizona. Monograph 6.

Schmidt, G. D., D. W. Duszynski, and P. S. Martin. 1992. Parasites of the extinct Shasta Ground Sloth *Nothrotheriops shastensis*, in Rampart Cave, Arizona. Journal of Parasitology 78(5):811–816.

Schwartz, D. W., A. L. Lange, and R. deSausure. 1958. Split-twig figurines in the Grand Canyon. American Antiquity 23:264–274.

Spamer, E. E. 1988. Geology of the Grand Canyon: an annotated bibliography, with an annotated catalogue of Grand Canyon type fossils. Volume 3. Second supplement (to 1987) with an annotated bibliography of the world literature on the Grand Canyon type fossil *Chuaria circularis* Walcott, 1899, an index fossil for the late Proterozoic. Geological Society of America, Boulder, Colorado. Microform Publication 17.

Spamer, E. E. 1993. Late Pleistocene (?) land snails (Mollusca: Gastropoda) in "red earth" deposits of the Grand Canyon, Arizona. The Mosasaur 5:47–58.

Spamer, E. E., and A. E. Bogan. 1993. Mollusca of the Grand Canyon and vicinity, Arizona: New and revised data on diversity and distributions, with notes on Pleistocene-Holocene mollusks of the Grand Canyon. Proceedings of the Academy of Natural Sciences of Philadelphia 144:21–68.

Spaulding, W. G., J. L. Betancourt, L. K. Croft, and K. L. Cole. 1990. Packrat middens: Their composition and methods of analysis. Pages 59–84 *in* J. L. Betancourt, T. R. Van Devender, and P. S. Martin, editors. Packrat middens: The last 40,000 years of biotic change. University of Arizona Press, Tucson, Arizona.

Springer, A. E., L. E. Stevens, D. E. Anderson, R. A. Parnell, D. K. Kreamer, L. A. Levin, and S. P. Flora. Pages 49–75 *in* L. E. Stevens and V. J. Meretsky, editors. Aridland springs in North America: Ecology and conservation. University of Arizona Press, Tucson, Arizona.

Stevens, L. 2013. The Colorado River in the Grand Canyon: A river runner's map and guide to its natural and human history. Red Lake Books, Flagstaff, Arizona.

Tang, M., and E. R. Reiter. 1984. Plateau monsoons of the Northern Hemisphere: A comparison between North America and Tibet. Monthly Weather Review 112:617–637.

Tobin, B. W., A. E. Springer, J. Ballensky, and A. Armstrong. In press. Cave and karst of the Grand Canyon World Heritage Site.

Tobin, B. W., A. E. Springer, D. K. Kreamer, and E. R. Schenk. 2018. Review: The distribution, flow, and quality of Grand Canyon springs, Arizona (USA). Hydrogeology Journal 26:721–732.

Van Devender, T. R., and J. I. Mead. 1976. Late Pleistocene and modern plant communities of Shinumo Creek and Peach Springs Wash, lower Grand Canyon, Arizona. Journal of the Arizona Academy of Science 11(1):16–22.

Van Devender, T. R., and W. G. Spaulding. 1979. Development of vegetation and climate in the southwestern United States. Science 204(4394):701–710.

Van Devender, T. R., A. M. Phillips, III, and J. I. Mead. 1977. Late Pleistocene reptiles and small mammals from the lower Grand Canyon of Arizona. The Southwestern Naturalist 22(1):49–66.

Van Devender, T. R., P. S. Martin, R. S. Thompson, K. L. Cole, A. J. T. Jull, A. Long, L. J. Toolin, and D. J. Donahue. 1985. Fossil packrat middens and the tandem accelerator mass spectrometer. Nature 317:610–613.

Wilson, R. W. 1942. Preliminary study of the fauna of Rampart Cave, Arizona. Carnegie Institution of Washington Publication 530:196–185.

Wurster, C. M., W. P. Patterson, D. A. McFarlane, L. I. Wassenaar, K. A. Hobson, N. B. Athfield, and M. I. Bird. 2008. Stable argon and hydrogen isotopes from bat guano in the Grand Canyon, USA, reveal Younger Dryas and 8.2 ka events. Geology 36:683–686.

Young, R. A., and E. E. Spamer. 2001. Colorado River origin and evolution: Proceedings of a symposium held at Grand Canyon National Park in June, 2000. Grand Canyon Association, Grand Canyon, Arizona.

Appendix 11-A. Grand Canyon Ice Age Taxa

The following tables list plants and animals reported from various Ice Age localities in the greater Grand Canyon (GC), Arizona. Almost all reports are of localities within Grand Canyon National Park (GRCA) itself, although a few localities are just outside of the park (e.g., some mollusks from Kaufman et al. 1992). The tables are set up taxonomically and refer to the publication(s) that discuss each taxon. The publications are included in "Literature Cited". Taxa followed by an asterisk (*) were named from specimens found within GRCA.

Plants

Ice Age plant taxa reported from the Grand Canyon are listed in Appendix Table 11-A-1.

Appendix Table 11-A-1. Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Polypodiopsida	Equisetaceae	*Equisetum* sp	Hansen 1978, Robbins et al. 1984
	Pteridaceae	*Adiantum capillus-veneris*	Laudermilk and Munz 1938
	Pteridaceae	*Adiantum* sp	Hansen 1978, Robbins et al. 1984
Gnetophyta	Ephedraceae	*Ephedra nevadensis*	Laudermilk and Munz 1938, Hansen 1978
	Ephedraceae	*Ephedra nevadensis*/*viridis*	Cole 1985
	Ephedraceae	*Ephedra torreyana*	Robbins et al. 1984, Dryer 1994
	Ephedraceae	*Ephedra viridis*	Cole and Mead 1981
	Ephedraceae	*Ephedra* cf. *E. nevadensis*	Long et al. 1974, Phillips and Van Devender 1974
	Ephedraceae	*Ephedra* sp.	Euler 1978, Mead and Phillips 1981, Hevly 1984, Phillips 1984, Robbins et al. 1984, O'Rourke and Mead 1985, Cole 1990, Dryer 1994

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Pinophyta	Cupressaceae	*Juniperus communis*	Cole 1990, Coats 1997, Coats et al. 2008
	Cupressaceae	*Juniperus monosperma*	Coats et al. 2008
	Cupressaceae	*Juniperus monosperma* and/or *J. osteosperma*	Phillips and Van Devender 1974
	Cupressaceae	*Juniperus osteosperma*[1]	Hansen 1978, Cole and Mead 1981, Cole 1982, Cole 1990, Coats 1997, Coats et al. 2008, Cole et al. 2013
	Cupressaceae	*Juniperus scopulorum*	Dryer 1994, Coats 1997
	Cupressaceae	*Juniperus* cf. *J. monosperma*	Cole 1985, Cole 1990
	Cupressaceae	*Juniperus* cf. *J. osteosperma*	Cole 1985, O'Rourke and Mead 1985
	Cupressaceae	*Juniperus* sp.	Martin et al. 1961, Van Devender and Mead 1976, Van Devender et al. 1977, Euler 1978, Mead et al. 1978, Mead et al. 1986, Mead and Phillips 1981, Cole 1982, Ferguson 1984, Hevly 1984, Phillips 1984, O'Rourke and Mead 1985, Dryer 1994, Cole et al. 2013
	Cupressaceae	Cupressaceae undetermined	Hansen 1978
	Pinaceae	*Abies concolor*	Mead et al. 1978, Cole and Mead 1981, Cole 1985, Cole 1990, O'Rourke and Mead 1985, Cole et al. 2013
	Pinaceae	*Abies* sp	O'Rourke and Mead 1985, Mead et al. 1986, Dryer 1994
	Pinaceae	*Picea engelmannii*	Cole 1985
	Pinaceae	*Picea engelmannii*/*pungens*	Cole 1990
	Pinaceae	*Picea* cf. *P. pungens*	Dryer 1994
	Pinaceae	*Picea* sp.	Euler 1978, Cole and Mead 1981, Cole 1982, Hevly 1984, O'Rourke and Mead 1985, Mead et al. 1986
	Pinaceae	*Pinus* cf. *P. contorta*	O'Rourke and Mead 1985
	Pinaceae	*Pinus edulis*	Van Devender and Spaulding 1979, Cole and Mead 1981, Ferguson 1984, Cole 1990, Dryer 1994, Cole et al. 2013
	Pinaceae	*Pinus* cf. *P. edulis* (late Holocene)	O'Rourke and Mead 1985
	Pinaceae	*Pinus edulis* var. *fallax*	Cole et al. 2013

[1] *Juniperus californicus* var. *osteosperma* = *Juniperus osteosperma*

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Pinophyta (continued)	Pinaceae	*Pinus flexilis*	Cole and Mead 1981, Cole 1982, Cole 1985, Cole 1990, Van Devender et al. 1985, Coats 1997, Coats et al. 2008
	Pinaceae	*Pinus* cf. *P. flexilis*	O'Rourke and Mead 1985
	Pinaceae	*Pinus monophylla*	Van Devender and Spaulding 1979
	Pinaceae	*Pinus ponderosa*	Cole 1982, Cole 1985, Cole 1990, O'Rourke and Mead 1985
	Pinaceae	*Pinus* sp.	Euler 1978, Hevly 1984, O'Rourke and Mead 1985, Mead et al. 1986, Coats 1997
	Pinaceae	*Pseudotsuga menziesii*	Euler 1978, Mead et al. 1978, Cole and Mead 1981, Cole 1982, Cole 1985, Cole 1990, Ferguson 1984, O'Rourke and Mead 1985, Dryer 1994, Coats 1997, Coats et al. 2008
	Pinaceae	*Pseudotsuga* sp.	O'Rourke and Mead 1985, Mead et al. 1986
Magnoliophyta	Amaranthaceae	*Amaranthus* cf. *A. palmeri*	Coats 1997
	Amaranthaceae	*Amaranthus* sp.	Long et al. 1974
	Amaranthaceae	*Amaranthus* spp.	Hansen 1978
	Amaranthaceae	*Atriplex confertifolia*	Van Devender and Mead 1976, Van Devender et al. 1977, Van Devender and Spaulding 1979, Cole and Mead 1981, Mead and Phillips 1981, Cole 1982, Cole 1985, Cole 1990, Phillips 1984, Dryer 1994, Coats 1997
	Amaranthaceae	*Atriplex* cf. *A. confertifolia*	Long et al. 1974
	Amaranthaceae	*Atriplex* cf. *A. jonesi* (Holocene)	Robbins et al. 1984
	Amaranthaceae	*Atriplex* sp.	Laudermilk and Munz 1938, Robbins et al. 1984, Mead et al. 1986, Dryer 1994, Coats 1997
	Amaranthaceae	*Atriplex* spp.	Hansen 1978
	Amaranthaceae	*Chenopodium* sp. (Holocene only?)	Robbins et al. 1984
	Amaranthaceae	*Eurotia lanata*	Hansen 1978
	Amaranthaceae	*Tidestromia oblongifolia*	Long et al. 1974
	Amaranthaceae	*Tidestromia* spp.	Hansen 1978
	Amaranthaceae	Cheno-am pollen	Euler 1978, Hevly 1984, O'Rourke and Mead 1985, Mead et al. 1986
	Anacardiaceae	*Rhus trilobata*	Van Devender and Mead 1976, Mead and Phillips 1981, Phillips 1984, Robbins et al. 1984, O'Rourke and Mead 1985, Coats 1997

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Anacardiaceae	*Rhus* sp.	O'Rourke and Mead 1985, Mead et al. 1986, Emslie et al. 1987, Cole 1990, Dryer 1994, Coats 1997
	Apiaceae	*Caucalis microcarpa*	Phillips 1984
	Apiaceae	*Cymopterus* sp.	Mead et al. 1986
	Apiaceae	*Lomatium* sp.	Dryer 1994
	Apiaceae	Apiaceae (or "Umbelliferae") undetermined (pollen)	Mead et al. 1986
	Apocynaceae	*Amsonia eastwoodiana*	Van Devender et al. 1977
	Apocynaceae	*Amsonia tormentosa*	Phillips 1984
	Apocynaceae	*Amsonia* sp.	Mead and Phillips 1981
	Asparagaceae	*Agave utahensis*	Van Devender and Mead 1976, Mead et al. 1978, Cole and Mead 1981, Mead and Phillips 1981, Phillips 1984, Robbins et al. 1984, Cole 1985, Cole 1990, O'Rourke and Mead 1985, Van Devender et al. 1985, Cole et al. 2013
	Asparagaceae	*Agave* cf. *A. utahensis*	Long et al. 1974
	Asparagaceae	*Agave* sp.	Mead et al. 1978, Robbins et al. 1984, Dryer 1994, Coats 1997, Coats et al. 2008
	Asparagaceae	*Agave* spp.	Hansen 1978
	Asparagaceae	*Nolina microcarpa*	Van Devender et al. 1977, Hansen 1978, Mead et al. 1978, Van Devender and Spaulding 1979, Mead and Phillips 1981, Phillips 1984
	Asparagaceae	*Nolina* cf. *N. parryi*	Phillips and Van Devender 1974
	Asparagaceae	*Nolina* sp.	Laudermilk and Munz 1938, Long et al. 1974, Van Devender and Mead 1976
	Asparagaceae	*Yucca angustissima*	Robbins et al. 1984, Cole 1990, Dryer 1994
	Asparagaceae	*Yucca baccata*	Van Devender and Mead 1976, Van Devender et al. 1977, Mead and Phillips 1981, Phillips 1984, Robbins et al. 1984, Coats 1997
	Asparagaceae	*Yucca brevifolia*	Van Devender et al. 1977
	Asparagaceae	*Yucca mohavensis*	Laudermilk and Munz 1938
	Asparagaceae	*Yucca schidigera*	Mead and Phillips 1981
	Asparagaceae	*Yucca* cf. *Y. newberryi*	Long et al. 1974
	Asparagaceae	*Yucca* sp.	Mead et al. 1978, 1986, Hevly 1984, Robbins et al. 1984, Coats 1997
	Asparagaceae	*Yucca* spp.	Hansen 1978
	Asparagaceae	*Agave* and *Yucca* undifferentiated	Hevly 1984

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Asteraceae	*Ambrosia* sp.	O'Rourke and Mead 1985, Dryer 1994
	Asteraceae	cf. *Ambrosia* (pollen)	O'Rourke and Mead 1985, Mead et al. 1986
	Asteraceae	*Artemisia ludoviciana*	Cole 1990
	Asteraceae	*Artemisia tridentata*	Hansen 1978, Robbins et al. 1984, Cole 1985, Cole 1990, Coats 1997, Coats et al. 2008
	Asteraceae	*Artemisia* cf. *A. ludoviciana*	Van Devender and Mead 1976
	Asteraceae	*Artemisia* sp.	Euler 1978, Cole 1982, Hevly 1984, O'Rourke and Mead 1985, Mead et al. 1986
	Asteraceae	*Artemisia* spp.	Dryer 1994
	Asteraceae	*Aster* sp.	Robbins et al. 1984
	Asteraceae	*Aster* spp.	Hansen 1978
	Asteraceae	*Baccharis sergiloides*	Robbins et al. 1984
	Asteraceae	*Baccharis* sp.	Dryer 1994
	Asteraceae	*Bahia* sp.	Coats 1997
	Asteraceae	*Bahia* spp.	Hansen 1978
	Asteraceae	*Brickellia atractyloides*	Coats 1997
	Asteraceae	*Brickellia* cf. *B. scabra*	Coats 1997
	Asteraceae	*Brickellia* sp.	Mead and Phillips 1981, Hevly 1984, Robbins et al. 1984, Dryer 1994, Coats 1997
	Asteraceae	*Chrysopsis* cf. *C. hispida*	Robbins et al. 1984
	Asteraceae	*Chrysothamnus nauseosus*	Hansen 1978
	Asteraceae	*Chrysothamnus* cf. *C. viscidiflorus*	Coats 1997, Coats et al. 2008
	Asteraceae	*Chrysothamnus* sp.	Mead and Phillips 1981, Phillips 1984, Robbins et al. 1984, Dryer 1994, Coats 1997
	Asteraceae	*Cirsium* sp.	Mead and Phillips 1981, Phillips 1984
	Asteraceae	*Cirsium* spp.	Dryer 1994
	Asteraceae	cf. *Cirsium* (pollen)	Mead et al. 1986
	Asteraceae	*Coreopsis* sp.	Dryer 1994
	Asteraceae	*Dyssodia pentachaeta*	Robbins et al. 1984
	Asteraceae	*Dyssodia* sp.	Robbins et al. 1984
	Asteraceae	*Encelia farinosa*	Van Devender and Mead 1976, Van Devender et al. 1977, Mead and Phillips 1981, Phillips 1984, Cole 1990
	Asteraceae	*Encelia* sp.	O'Rourke and Mead 1985
	Asteraceae	*Encelia* spp.	Hansen 1978

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Asteraceae	*Erigeron* sp.	Mead and Phillips 1981
	Asteraceae	*Franseria confertifolia*	Phillips 1984
	Asteraceae	*Gutierrezia lucida*	Phillips and Van Devender 1974, Van Devender et al. 1977, Hansen 1978, Mead and Phillips 1981
	Asteraceae	*Gutierrezia microcephala*	Phillips 1984
	Asteraceae	*Gutierrezia* sp.	Long et al. 1974, Robbins et al. 1984, Dryer 1994
	Asteraceae	*Haplopappus* sp. (Holocene)[2]	Robbins et al. 1984
	Asteraceae	cf. *Helianthus* (pollen)	Mead et al. 1986
	Asteraceae	*Hofmeisteria pluriseta*	Hansen 1978
	Asteraceae	*Hofmeisteria* sp.	Long et al. 1974
	Asteraceae	*Laphamia congesta*	Robbins et al. 1984
	Asteraceae	*Lygodesmia exigua*	Robbins et al. 1984
	Asteraceae	*Peucephyllum schottii*	Hansen 1978
	Asteraceae	*Solidago* sp.	Robbins et al. 1984
	Asteraceae	*Tetradymia canescens*?	Robbins et al. 1984
	Asteraceae	cf. *Brickellia* and *Cirsium*	Hevly 1984
	Asteraceae	Asteraceae (or "Compositae") undetermined (pollen)	Euler 1978, Hevly 1984, O'Rourke and Mead 1985, Mead et al. 1986
	Berberidaceae	*Berberis repens* (Holocene)	Robbins et al. 1984
	Berberidaceae	*Berberis* sp.	O'Rourke and Mead 1985, Dryer 1994
	Betulaceae	*Alnus* sp. (pollen)	Euler 1978, Hevly 1984
	Betulaceae	*Betula* sp. (pollen)	Euler 1978, Hevly 1984
	Betulaceae	*Ostrya knowltoni*	Van Devender et al. 1977, Mead and Phillips 1981, Phillips 1984, O'Rourke and Mead 1985, Cole 1990, Coats 1997
	Betulaceae	*Ostrya* sp. (pollen)	O'Rourke and Mead 1985, Mead et al. 1986
	Boraginaceae	*Amsinckia intermedia*	Hansen 1978, Mead and Phillips 1981
	Boraginaceae	*Amsinckia* sp.	Van Devender and Mead 1976
	Boraginaceae	*Coldenia hispidissima*	Robbins et al. 1984
	Boraginaceae	*Coldenia* sp.	Robbins et al. 1984
	Boraginaceae	Cryptantha pterocarya	Mead and Phillips 1981
	Boraginaceae	*Cryptantha virginensis*	Mead and Phillips 1981, Phillips 1984
	Boraginaceae	*Cryptantha* cf. *C. confertiflora*	Coats 1997

[2] *Aplopappus = Haplopappus*

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Boraginaceae	*Cryptantha* cf. *C. recurvata*	Dryer 1994
	Boraginaceae	*Cryptantha* cf. *C. torreyana*	Coats 1997
	Boraginaceae	*Cryptantha* cf. *C. virginensis*	Coats 1997
	Boraginaceae	*Cryptantha* sp.	Robbins et al. 1984, Mead et al. 1986, Dryer 1994, Coats 1997
	Boraginaceae	*Cryptantha* spp.	Hansen 1978
	Boraginaceae	*Lappula occidentalis*	Dryer 1994, Coats 1997
	Boraginaceae	*Lappula redowskii*	Van Devender and Mead 1976
	Boraginaceae	*Lithospermum incisum*	Coats 1997, Coats et al. 2008
	Boraginaceae	*Pectocarya* spp.	Hansen 1978
	Boraginaceae	*Phacelia crenulata*	Mead and Phillips 1981, Phillips 1984
	Boraginaceae	*Phacelia* sp.	Long et al. 1974
	Boraginaceae	*Phacelia* spp.	Hansen 1978
	Boraginaceae	*Tiquilia* sp.	Coats 1997
	Boraginaceae	cf. *Trixis* (pollen)	Mead et al. 1986
	Boraginaceae	*Coldenia–Cryptantha*	Robbins et al. 1984
	Brassicaceae	*Descurainia pinnata*	Hansen 1978
	Brassicaceae	*Draba cuneifolia*	Hansen 1978
	Brassicaceae	*Lepidium* sp.	Long et al. 1974, Mead and Phillips 1981, Phillips 1984, Dryer 1994, Coats 1997
	Brassicaceae	*Lepidium* spp.	Hansen 1978
	Brassicaceae	*Lesquerella* sp.	Robbins et al. 1984, Mead et al. 1986
	Brassicaceae	cf. *Lesquerella* (pollen)	Mead et al. 1986
	Brassicaceae	*Streptanthella longirostris*	Dryer 1994
	Brassicaceae	*Thysanocarpus amplectens*	Phillips 1984
	Brassicaceae	Brassicaceae (or “Cruciferae”) undetermined	Hevly 1984, Mead et al. 1986
	Cactaceae	*Cylindropuntia* sp.	Dryer 1994, Coats 1997
	Cactaceae	*Echinocactus polycephalus*	Phillips and Van Devender 1974, Van Devender et al. 1977, Mead and Phillips 1981, Phillips 1984
	Cactaceae	*Echinocactus* sp.	Martin et al. 1961, Phillips 1984
	Cactaceae	*Echinocereus* sp.	Mead and Phillips 1981, Coats 1997
	Cactaceae	*Ferocactus acanthodes*	Van Devender and Mead 1976, Mead and Phillips 1981, Phillips 1984
	Cactaceae	*Ferocactus wislizeni*	Coats 1997
	Cactaceae	*Opuntia basilaris*	Phillips and Van Devender 1974, Van Devender et al. 1977, Mead and Phillips 1981, Phillips 1984, Robbins et al. 1984, Cole 1990

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Cactaceae	*Opuntia chorotica*	Van Devender and Mead 1976
	Cactaceae	*Opuntia erinacea*	Mead and Phillips 1981, Phillips 1984, Cole 1985, Cole 1990
	Cactaceae	*Opuntia macrorhiza*	Mead and Phillips 1981
	Cactaceae	*Opuntia phaeacantha*	Van Devender and Mead 1976, Cole 1990
	Cactaceae	*Opuntia whipplei*	Phillips and Van Devender 1974, Van Devender and Mead 1976, Van Devender et al. 1977, Mead and Phillips 1981, Phillips 1984, Dryer 1994
	Cactaceae	*Opuntia* cf. *O. whipplei*	Mead et al. 1978
	Cactaceae	*Opuntia* (*Platyopuntia*) sp.[3]	Coats 1997
	Cactaceae	*Opuntia* sp.	Laudermilk and Munz 1938, Long et al. 1974, Mead and Phillips 1981, Hevly 1984, Cole 1985, Cole 1990, O'Rourke and Mead 1985, Dryer 1994, Coats 1997, Coats et al. 2008
	Cactaceae	Cactaceae undetermined	Hansen 1978, Dryer 1994, Coats 1997
	Cannabaceae	*Celtis reticulata*	Van Devender and Mead 1976, Hansen 1978, Mead and Phillips 1981, O'Rourke and Mead 1985, Dryer 1994
	Cannabaceae	*Celtis* sp. (Holocene)	Hevly 1984
	Caprifoliaceae	*Lonicera* sp.	Mead et al. 1986
	Caprifoliaceae	*Symphoricarpos* sp.	Van Devender et al. 1977, Van Devender and Spaulding 1979, Mead and Phillips 1981, Phillips 1984, Mead et al. 1986, Cole 1990, Dryer 1994, Coats 1997, Coats et al. 2008
	Caryophyllaceae	*Arenaria fendleri*	Robbins et al. 1984
	Caryophyllaceae	*Arenaria* sp.	Robbins et al. 1984
	Caryophyllaceae	Caryophyllaceae undetermined (pollen)	O'Rourke and Mead 1985, Mead et al. 1986
	Celastraceae	*Mortonia scabrella*	Phillips and Van Devender 1974, Van Devender et al. 1977, Mead et al. 1978, Mead and Phillips 1981
	Celastraceae	*Mortonia scabrella* var. *utahensis*	Phillips 1984
	Celastraceae	*Pachystima myrsinites*	Cole 1990

[3] *Platypuntia* of Coats 1997 accepted as *Opuntia* (*Platyopuntia*)

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Convolvulacea	*Convolvulus* sp.	Robbins et al. 1984
	Crossosomaceae	*Crossosoma bigelovii*	Hansen 1978, Mead and Phillips 1981, Phillips 1984
	Crossosomaceae	*Crossosoma* sp.	Long et al. 1974
	Crossosomaceae	*Glossopetalon nevadense*	Hansen 1978
	Crossosomaceae	*Glossopetalon* sp.	Mead et al. 1986
	Cucurbitaceae	*Cucurbita* sp.	Van Devender and Mead 1976
	Cyperaceae	*Carex* sp.	Long et al. 1974, Robbins et al. 1984, Mead et al. 1986
	Cyperaceae	*Eleocharis* sp.	Robbins et al. 1984
	Cyperaceae	Cyperaceae undetermined	Hansen 1978
	Elaeagnaceae	*Shepherdia* sp.	Hevly 1984
	Elaeagnaceae	*Shepherdia* spp.	Hansen 1978
	Euphorbiaceae	*Argythamnia* sp.	Dryer 1994
	Euphorbiaceae	*Euphorbia* cf. *E. fendleri*	Coats 1997
	Euphorbiaceae	*Euphorbia sp.*	Van Devender and Mead 1976, Robbins et al. 1984, Dryer 1994
	Euphorbiaceae	*Tragia* sp.	Robbins et al. 1984
	Fabaceae	*Acacia greggii*	Long et al. 1974, Hansen 1978, Mead and Phillips 1981, Phillips 1984
	Fabaceae	*Acacia* sp.	Mead et al. 1986
	Fabaceae	*Astragalus nutallianus*	Phillips 1984
	Fabaceae	*Astragalus* sp.	Hansen 1978, Robbins et al. 1984, O'Rourke and Mead 1985, Dryer 1994
	Fabaceae	*Astragalus*-type pollen	Mead et al. 1986
	Fabaceae	*Cassia* sp.	Laudermilk and Munz 1938
	Fabaceae	*Cercis occidentalis*	O'Rourke and Mead 1985, Cole 1990
	Fabaceae	*Lotus* spp.	Hansen 1978
	Fabaceae	*Prosopis juliflora*	Long et al. 1974, Hansen 1978, Cole 1990
	Fabaceae	Fabaceae (or "Leguminosae") undetermined (pollen)	Mead et al. 1986
	Fabaceae	Fabaceae undetermined (driftwood)	Ferguson 1984
	Fabaceae	"Legume a and b"	Hevly 1984
	Fagaceae	*Quercus turbinella*	Mead and Phillips 1981, Cole 1982, Cole 1985, Cole 1990
	Fagaceae	*Quercus* sp.	Euler 1978, Hansen 1978, Hevly 1984, O'Rourke and Mead 1985, Mead et al. 1986

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Grossulariaceae	*Ribes montigenum*	Van Devender et al. 1977, Van Devender and Spaulding 1979, Mead and Phillips 1981, O'Rourke and Mead 1985
	Grossulariaceae	*Ribes pinetorum*	Coats et al. 2008
	Grossulariaceae	*Ribes* cf. *R. pinetorum*	Coats 1997
	Grossulariaceae	*Ribes* sp.	Cole 1985, Cole 1990, Coats 1997
	Grossulariaceae	cf. *Ribes* sp.	Dryer 1994
	Hydrangeaceae	*Fendlera rupicola*	Cole 1990
	Juglandaceae	*Juglans* sp. (Holocene body fossils)	Hevly 1984
	Juglandaceae	*Juglans* sp. (pollen)	Euler 1978, Hevly 1984
	Krameriaceae	*Krameria parvifolia*	Hansen 1978
	Lamiaceae	*Hedeoma diffusum*	Robbins et al. 1984
	Lamiaceae	*Hedeoma* sp.	Robbins et al. 1984
	Lamiaceae	*Salvia dorrii*	Mead and Phillips 1981, Phillips 1984
	Lamiaceae	Lamiaceae (or "Labiatae") undetermined (pollen)	Mead et al. 1986
	Liliaceae	Liliaceae undetermined (pollen)	Mead et al. 1986
	Linaceae	*Linum* sp. (pollen)	Mead et al. 1986
	Loasaceae	*Eucnide urens*	Hansen 1978
	Loasaceae	*Mentzelia puberula*	Mead and Phillips 1981
	Loasaceae	*Mentzelia* sp.	Phillips 1984, Robbins et al. 1984, Dryer 1994
	Malpighiaceae	*Janusia gracilis*	Van Devender and Mead 1976
	Malpighiaceae	*Janusia* sp.	Hansen 1978
	Malvaceae	*Malvastrum rotundifolium*	Hansen 1978
	Malvaceae	*Sida* sp.	Mead et al. 1986
	Malvaceae	*Sphaeralcea ambigua*	Laudermilk and Munz 1938, Van Devender et al. 1977, Hansen 1978
	Malvaceae	*Sphaeralcea* cf. *S. laxa*	Long et al. 1974
	Malvaceae	*Sphaeralcea* sp.	Martin et al. 1961, Van Devender and Mead 1976, Mead and Phillips 1981, Phillips 1984, Robbins et al. 1984, O'Rourke and Mead 1985, Mead et al. 1986, Dryer 1994
	Nyctaginaceae	*Abronia* sp.	Hansen 1978
	Nyctaginaceae	*Allionia incarnata*	Hansen 1978, Phillips 1984
	Nyctaginaceae	*Allionia* sp.	Long et al. 1974
	Nyctaginaceae	*Boerhavia coulteri*	Mead and Phillips 1981
	Nyctaginaceae	*Boerhavia* sp.	Van Devender and Mead 1976
	Nyctaginaceae	*Mirabilis multiflora*	Coats 1997

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Nyctaginaceae	*Oxybaphus* sp.	O'Rourke and Mead 1985
	Oleaceae	*Fraxinus anomala*	Long et al. 1974, Phillips and Van Devender 1974, Van Devender and Mead 1976, Van Devender et al. 1977, Hansen 1978, Mead et al. 1978, Van Devender and Spaulding 1979, Mead and Phillips 1981, Cole 1982, Cole 1985, Cole 1990, Phillips 1984, O'Rourke and Mead 1985, Coats 1997, Coats et al. 2008
	Oleaceae	*Fraxinus* cf. *F. anomala*	Dryer 1994
	Oleaceae	*Fraxinus* sp.	Laudermilk and Munz 1938
	Oleaceae	*Fraxinus* spp.	Hansen 1978
	Onagraceae	*Oenothera cavernae*	Phillips 1984
	Onagraceae	*Oenothera* sp.	Robbins et al. 1984, Mead et al. 1986
	Onagraceae	*Oenothera* spp.	Hansen 1978
	Orobanchaceae	*Castilleja* cf. *C. miniata*	Coats 1997
	Orobanchaceae	*Castilleja* sp.	Coats 1997
	Orobanchaceae	*Castilleja* spp.	Hansen 1978
	Orobanchaceae	*Castilleja* or *Orthocarpus*	Van Devender and Mead 1976
	Papaveraceae	*Argemone* sp.	Mead and Phillips 1981, Phillips 1984, O'Rourke and Mead 1985, Dryer 1994, Coats 1997
	Phrymaceae	*Mimulus* sp.	Robbins et al. 1984
	Plantaginaceae	*Penstemon eatonii*	Phillips 1984
	Plantaginaceae	*Penstemon* sp.	Van Devender and Mead 1976, Mead and Phillips 1981
	Plantaginaceae	*Plantago* sp.	Dryer 1994
	Poaceae	*Agropyron* sp.	Robbins et al. 1984, Mead et al. 1986
	Poaceae	*Agropyron* spp.	Hansen 1978
	Poaceae	*Andropogon* sp.	Robbins et al. 1984
	Poaceae	*Aristida glauca*	Robbins et al. 1984
	Poaceae	*Aristida longiseta*	Robbins et al. 1984
	Poaceae	*Aristida* sp.	Laudermilk and Munz 1938, Long et al. 1974, Robbins et al. 1984, Mead et al. 1986, Dryer 1994
	Poaceae	*Aristida* spp.	Hansen 1978
	Poaceae	*Bouteloua eripoda* (Holocene)	Robbins et al. 1984
	Poaceae	*Bouteloua gracilis*	Robbins et al. 1984
	Poaceae	*Bouteloua simplex*	Robbins et al. 1984
	Poaceae	*Bouteloua trifida*	Robbins et al. 1984
	Poaceae	*Bouteloua* sp.	Robbins et al. 1984, Mead et al. 1986
	Poaceae	*Bouteloua* spp.	Hansen 1978

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Poaceae	*Bromus* sp.	Long et al. 1974, Robbins et al. 1984, Mead et al. 1986
	Poaceae	*Bromus* spp.	Hansen 1978
	Poaceae	*Enneapogon desvauxii*	Robbins et al. 1984
	Poaceae	*Festuca arizonica*	Hansen 1978
	Poaceae	*Festuca* sp.	Robbins et al. 1984, Mead et al. 1986
	Poaceae	*Hilaria rigida*	Hansen 1978
	Poaceae	*Lycurus phleoides* (Holocene)	Robbins et al. 1984
	Poaceae	*Muhlenbergia* sp.	Long et al. 1974, Hansen 1978
	Poaceae	*Panicum huachucae*	Robbins et al. 1984
	Poaceae	*Oryzopsis hymenoides*	Mead and Phillips 1981, O'Rourke and Mead 1985, Coats 1997, Coats et al. 2008
	Poaceae	*Oryzopsis* sp.	Hansen 1978, Mead et al. 1986
	Poaceae	*Phragmites communis*	Laudermilk and Munz 1938, Hansen 1978
	Poaceae	*Phragmites* sp.	Long et al. 1974, Robbins et al. 1984
	Poaceae	*Poa* sp.	Mead et al. 1986
	Poaceae	*Poa* spp.	Hansen 1978
	Poaceae	*Schedonnardus paniculatus*	Robbins et al. 1984
	Poaceae	*Schizachyrium scoparium*	Dryer 1994
	Poaceae	*Sporobolus flexuosus*	Robbins et al. 1984
	Poaceae	*Sporobolus texanus*	Robbins et al. 1984
	Poaceae	*Sporobolus* cf. *S. cryptandrus*	Robbins et al. 1984
	Poaceae	*Sporobolus* sp.	Robbins et al. 1984, Mead et al. 1986
	Poaceae	*Sporobolus* spp.	Hansen 1978
	Poaceae	*Stipa arida*	Robbins et al. 1984
	Poaceae	*Stipa hymenoides*	Dryer 1994
	Poaceae	*Stipa* cf. *neomexicana*	Dryer 1994
	Poaceae	*Stipa* sp.	Long et al. 1974, Mead et al. 1986
	Poaceae	*Stipa* spp.	Hansen 1978
	Poaceae	*Tridens pulchellus*	Robbins et al. 1984
	Poaceae	*Tridens* sp	Long et al. 1974, Hansen 1978
	Poaceae	*Tridens* spp.	Hansen 1978
	Poaceae	*Zea mays* (late Holocene)	Cole 1982
	Poaceae	Gramineae A	Robbins et al. 1984
	Poaceae	Poaceae (or "Gramineae") undetermined	Euler 1978, Hevly 1984, Robbins et al. 1984, O'Rourke and Mead 1985, Mead et al. 1986
	Polemoniaceae	*Gilia* sp.	Hevly 1984
	Polemoniaceae	cf. *Leptodactylon* (pollen)	Mead et al. 1986
	Polemoniaceae	*Linanthus demissus*	Hansen 1978
	Polemoniaceae	*Phlox* sp. (pollen)	Mead et al. 1986
	Polemoniaceae	*Phlox*/*Leptodactylon*	Mead et al. 1986
	Polemoniaceae	Polemoniaceae undetermined (pollen)	O'Rourke and Mead 1985

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Polygonaceae	*Eriogonum deflexum*	Robbins et al. 1984
	Polygonaceae	*Eriogonum* sp.	Robbins et al. 1984, Mead et al. 1986
	Polygonaceae	*Eriogonum* spp.	Hansen 1978
	Polygonaceae	*Polygonum* sp.	Dryer 1994
	Polygonaceae	*Rumex* sp.	Dryer 1994
	Polygonaceae	Polygonaceae undetermined (pollen)	O'Rourke and Mead 1985, Mead et al. 1986
	Ranunculaceae	*Anemone tuberosa*	Hansen 1978, Phillips 1984
	Ranunculaceae	*Anemone* sp.	Van Devender and Mead 1976
	Ranunculaceae	*Aquilegia chrysantha*	Hansen 1978
	Ranunculaceae	*Caltha* sp. (pollen)	Mead et al. 1986
	Ranunculaceae	*Clematis ligusticifolia*	Robbins et al. 1984
	Ranunculaceae	*Ranunculus* sp.	Dryer 1994
	Ranunculaceae	Ranunculaceae undetermined (pollen)	O'Rourke and Mead 1985, Mead et al. 1986
	Rhamnaceae	*Rhamnus betulaefolia*	Phillips 1984
	Rhamnaceae	*Rhamnus* sp. (late Holocene)	O'Rourke and Mead 1985
	Rhamnaceae	Rhamnaceae undetermined (pollen)	O'Rourke and Mead 1985
	Rosaceae	*Amelanchier* sp.	Hevly 1984, Robbins et al. 1984, Coats 1997
	Rosaceae	*Cercocarpus intricatus*	Van Devender et al. 1977, Van Devender and Spaulding 1979, Phillips 1984, Robbins et al. 1984, O'Rourke and Mead 1985, Cole 1990, Coats 1997, Coats et al. 2008
	Rosaceae	*Cercocarpus montanus*	Hansen 1978
	Rosaceae	*Cercocarpus* sp. (pollen)	Mead et al. 1986
	Rosaceae	cf. *Cercocarpus* (pollen)	O'Rourke and Mead 1985
	Rosaceae	*Chamaebatiaria millefolium*	Cole 1990, Coats 1997, Coats et al. 2008
	Rosaceae	*Coleogyne ramosissima*	Phillips and Van Devender 1974, Van Devender et al. 1977, Mead and Phillips 1981, Phillips 1984, O'Rourke and Mead 1985, Cole 1990, Coats 1997, Coats et al. 2008
	Rosaceae	*Cowania mexicana*	O'Rourke and Mead 1985, Cole 1990, Coats 1997
	Rosaceae	*Fallugia paradoxa*	Van Devender and Mead 1976, Robbins et al. 1984
	Rosaceae	*Geum* spp.	Hansen 1978
	Rosaceae	*Holodiscus dumosus*	Cole 1990, Coats 1997
	Rosaceae	*Potentilla* spp.	Hansen 1978
	Rosaceae	*Prunus fasciculata*	Phillips and Van Devender 1974, Van Devender et al. 1977, Hansen 1978, Mead et al. 1978, Phillips 1984, Robbins et al. 1984, Cole 1990

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Rosaceae	*Prunus virginiana*	Dryer 1994, Coats 1997
	Rosaceae	*Prunus* sp.	Laudermilk and Munz 1938, Long et al. 1974, Mead et al. 1986
	Rosaceae	*Purshia mexicana*	Dryer 1994
	Rosaceae	*Purshia* sp.	Hevly 1984
	Rosaceae	*Rosa* cf. *R. arizonica*	O'Rourke and Mead 1985
	Rosaceae	*Rosa stellata*	Coats et al. 2008
	Rosaceae	*Rosa* cf. *R. stellata*	Cole 1985, Cole 1990, Dryer 1994, Coats 1997
	Rosaceae	*Rosa* sp.	O'Rourke and Mead 1985, Dryer 1994, Coats 1997
	Rosaceae	*Rubus* sp.	Cole 1990, Dryer 1994, Coats 1997
	Rosaceae	Rosaceae undetermined (pollen)	Mead et al. 1986
	Rubiaceae	*Galium* sp.	Van Devender and Mead 1976, Mead and Phillips 1981, Phillips 1984, Robbins et al. 1984
	Rubiaceae	*Galium* spp.	Hansen 1978
	Rutaceae	*Ptelea trifoliata*	Coats 1997, Coats et al. 2008
	Rutaceae	*Ptelea trifoliata* var. *pallida*	Cole 1985, Cole 1990, O'Rourke and Mead 1985
	Rutaceae	*Thamnosma montana*	Dryer 1994
	Rutaceae	*Thamnosma* sp.	Mead and Phillips 1981, Hevly 1984
	Salicaceae	*Populus fremontii*	Hansen 1978
	Salicaceae	*Populus fremontii*?	Euler 1978
	Salicaceae	*Populus* sp.	Laudermilk and Munz 1938, Ferguson 1984, Mead et al. 1986
	Salicaceae	*Salix* sp. (pollen)	Euler 1978, Hevly 1984, Mead et al. 1986
	Salicaceae	*Populus* and *Salix* (Holocene body fossils)	Hevly 1984
	Santalaceae	*Phoradendron californicum*	Phillips 1984
	Santalaceae	*Phoradendron* sp.	Long et al. 1974, Hansen 1978, Mead and Phillips 1981, Robbins et al. 1984
	Sapindaceae	*Acer glabrum*	Coats 1997, Coats et al. 2008
	Sapindaceae	*Acer* sp. (pollen)	Euler 1978, Hevly 1984
	Sarcobataceae	*Sarcobatus vermiculatus*	Robbins et al. 1984
	Sarcobataceae	*Sarcobatus* sp. (pollen)	Euler 1978, Hevly 1984
	Saxifragaceae	*Mitella* sp.	Dryer 1994
	Saxifragaceae	cf. *Saxifraga* (pollen)	Mead et al. 1986
	Saxifragaceae	Saxifragaceae undetermined (pollen)	O'Rourke and Mead 1985
	Solanaceae	*Datura metaloides* (GRCA: Holocene)	Van Devender and Mead 1976, Robbins et al. 1984
	Solanaceae	*Lycium andersonii*	Phillips 1984
	Solanaceae	*Lycium* sp.	Dryer 1994
	Solanaceae	cf. *Lycium* (pollen)	Mead et al. 1986

Appendix Table 11-A-1 (continued). Ice Age plant taxa reported from the Grand Canyon. Those records that are pollen-only are reported as such; otherwise, records may be macrofossil-only or mixed macrofossil and palynomorph.

Category	Family	Taxa Observed	Sources
Magnoliophyta (continued)	Solanaceae	*Physalis fendleri*	Robbins et al. 1984
	Solanaceae	*Physalis* sp.	Laudermilk and Munz 1938, Van Devender and Mead 1976, Mead and Phillips 1981, Phillips 1984
	Solanaceae	Solanaceae undetermined (pollen)	Mead et al. 1986
	Urticaceae	*Urtica* sp.	Robbins et al. 1984
	Verbenaceae	*Aloysia wrightii*	Robbins et al. 1984, Cole 1990
	Verbenaceae	*Verbena* sp.	Van Devender and Mead 1976
	Vitaceae	*Vitis arizonica*	Mead and Phillips 1981, Phillips 1984
	Vitaceae	*Vitis* sp.	Mead et al. 1978
	Zygophyllaceae	*Kallstroemia* sp.	Van Devender and Mead 1976
	Zygophyllaceae	*Larrea divaricata*	Laudermilk and Munz 1938
	Zygophyllaceae	*Larrea tridentata*	Martin et al. 1961, Hansen 1978, Mead and Phillips 1981, Phillips 1984
	Zygophyllaceae	*Ceanothus*/*Cercocarpus*	Mead et al. 1986
	Zygophyllaceae	Undetermined wood	Emslie 1988, Dryer 1994, Kaufman et al. 2002
	Zygophyllaceae	Undetermined plants	Phillips and Van Devender 1974, Mead et al. 1978, Mead et al. 1986, Hevly 1984, Emslie et al. 1987, Emslie 1988, Kaufman et al. 2002

Invertebrates

Ice Age invertebrate taxa reported from the Grand Canyon are listed in Appendix Table 11-A-2.

Appendix Table 11-A-2. Ice Age invertebrate taxa reported from the Grand Canyon. Taxa followed by an asterisk (*) were named from specimens found within GRCA.

Phylum	Class	Order	Family	Taxon Observed	Sources
Mollusca	Bivalvia	Sphaeriida	Sphaeriidae	*Pisidium* cf. *P. casertanum*	Kaufman et al. 2002
	Bivalvia	Sphaeriida	Sphaeriidae	*Pisidium* cf. *P. nitidum*	Kaufman et al. 2002
	Bivalvia	Sphaeriida	Sphaeriidae	*Pisidium* cf. *P. subtruncatum*	Kaufman et al. 2002
	Bivalvia	Sphaeriida	Sphaeriidae	*Pisidium* cf. *P. walkeri*	Kaufman et al. 2002
	Bivalvia	Sphaeriida	Sphaeriidae	*Pisidium* sp.	Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Succineidae	*Catinella* cf. *C. vermeta*[4]	Spamer 1993, Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Succineidae	*Catinella* sp.	Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Cionellidae	*Cionella lubrica*[5]	Spamer 1993, Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Lymnaeidae	*Fossaria dalli*	Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Lymnaeidae	*Fossaria* sp.	Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Planorbidae	*Gyraulus parvus*	Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Oreohelicidae	*Oreohelix yavapai*[6]	Spamer and Bogan 1993, Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Succineidae	*Oxyloma* cf. *O. haydeni kanabensis*	Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Succineidae	*Oxyloma* sp.	Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Physidae	*Physella* cf. *P. humerosa*	Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Physidae	*Physella* cf. *P. virgata*	Kaufman et al. 2002
	Gastropoda	“Pulmonata”	Xanthonychidae	cf. *Sonorella* sp.	Cole and Mead 1981

[4] *Catinella* cf. *C. avara = Catinella* cf. *C. vermeta*

[5] *Cochlicopa lubrica = Cionella lubrica*

[6] Subspecies of *Oreohelix yavapai*, such as *O. y. fortis** Cockerell 1927, are now generally rolled into *Oreohelix yavapai*

Appendix Table 11-A-2 (continued). Ice Age invertebrate taxa reported from the Grand Canyon. Taxa followed by an asterisk (*) were named from specimens found within GRCA.

Phylum	Class	Order	Family	Taxon Observed	Sources
Mollusca (continued)	Gastropoda	"Pulmonata"	Vertiginidae	*Vertigo ovata*	Kaufman et al. 2002
	–	–	–	Mollusca undetermined	Hevly 1984
Nematoda	–	–	–	*Agamofilaria oxyura**	Schmidt et al. 1992
	–	–	–	*Strongyloides shastensis**	Schmidt et al. 1992
	–	–	–	Nematoda unspecified	Laudermilk and Munz 1938
Arthropoda	Arachnida	Ixodida	Ixodidae	*Dermacentor andersoni*	Elias et al. 1992
	Arachnida	Ixodida	Ixodidae	*Dermacentor* sp.	Elias et al. 1992
	Arachnida	Scorpiones	Buthidae	*Centruroides* sp.	Elias et al. 1992
	Diplopoda	–	–	Diplopoda undetermined	Elias et al. 1992
	Insecta	Coleoptera	Carabidae	*Agonum* (*Rhadine*) *perlevis* (late Holocene)	Elias et al. 1992
	Insecta	Coleoptera	Carabidae	*Agonum (Rhadine*) sp.	Elias et al. 1992
	Insecta	Coleoptera	Carabidae	*Calosoma* cf. *C. scrutator* (late Holocene)	Elias et al. 1992
	Insecta	Coleoptera	Chrysomelidae	*Chaetocnema* sp. (late Holocene)	Elias et al. 1992
	Insecta	Coleoptera	Chrysomelidae	*Lema trilinea*	Elias et al. 1992
	Insecta	Coleoptera	Chrysomelidae	Chrysomelidae undetermined	Elias et al. 1992
	Insecta	Coleoptera	Cleridae	*Acanthoscelides* sp.	Elias et al. 1992
	Insecta	Coleoptera	Curculionidae	*Apleurus angularis*	Elias et al. 1992
	Insecta	Coleoptera	Curculionidae	*Cleonidus trivittatus* or *C. quadrilineatus*	Elias et al. 1992
	Insecta	Coleoptera	Curculionidae	*Ophryastes* sp.	Elias et al. 1992
	Insecta	Coleoptera	Curculionidae	*Orimodema protracta* (late Holocene)	Elias et al. 1992
	Insecta	Coleoptera	Curculionidae	*Sapotes* sp.	Elias et al. 1992
	Insecta	Coleoptera	Curculionidae	*Scyphophorus acupunctatus*	Elias et al. 1992
	Insecta	Coleoptera	Curculionidae	Cleridae undetermined	Elias et al. 1992
	Insecta	Coleoptera	Dermestidae	Dermestidae undetermined	Elias et al. 1992

Appendix Table 11-A-2 (continued). Ice Age invertebrate taxa reported from the Grand Canyon. Taxa followed by an asterisk (*) were named from specimens found within GRCA.

Phylum	Class	Order	Family	Taxon Observed	Sources
Arthropoda (continued)	Insecta	Coleoptera	Elateridae	Elateridae undetermined	Elias et al. 1992
	Insecta	Coleoptera	Histeridae	Histeridae undetermined	Elias et al. 1992
	Insecta	Coleoptera	Melandryidae	*Anaspis rufa*	Elias et al. 1992
	Insecta	Coleoptera	Meloidae	Meloidae undetermined	Elias et al. 1992
	Insecta	Coleoptera	Nitidulidae	Nitidulidae undetermined	Elias et al. 1992
	Insecta	Coleoptera	Ptinidae	*Niptus* cf. *N. ventriculus*	Elias et al. 1992
	Insecta	Coleoptera	Ptinidae	*Ptinis* sp.	Elias et al. 1992
	Insecta	Coleoptera	Ptinidae	Ptinidae undetermined	Cole and Mead 1981
	Insecta	Coleoptera	Scarabaeidae	*Aphodius* near *A. ruficlarus*	Elias et al. 1992
	Insecta	Coleoptera	Scarabaeidae	*Aphodius* near *A. ruficlarus*	Elias et al. 1992
	Insecta	Coleoptera	Scarabaeidae	*Aphodius* sp.	Elias et al. 1992
	Insecta	Coleoptera	Scarabaeidae	*Diplotaxis* sp.	Elias et al. 1992
	Insecta	Coleoptera	Scarabaeidae	*Onthophagus* sp.	Elias et al. 1992
	Insecta	Coleoptera	Scarabaeidae	*Phyllophaga* sp.	Elias et al. 1992
	Insecta	Coleoptera	Scarabaeidae	*Serica* sp.	Elias et al. 1992
	Insecta	Coleoptera	Scarabaeidae	Scarabaeidae undetermined	Elias et al. 1992
	Insecta	Coleoptera	Scotylidae	Scotylidae undetermined	Elias et al. 1992
	Insecta	Coleoptera	Silphidae	*Thanatophilus truncatus* (late Holocene)	Elias et al. 1992
	Insecta	Coleoptera	Tenebrionidae	*Coniontis* sp.	Elias et al. 1992
	Insecta	Coleoptera	Tenebrionidae	*Eleodes* cf. *E. nigrina*	Elias et al. 1992
	Insecta	Coleoptera	Tenebrionidae	*Eleodes* spp.	Elias et al. 1992
	Insecta	Coleoptera	Tenebrionidae	Tenebrionidae undetermined	Hevly 1984
	Insecta	Diptera	–	Diptera undetermined	Elias et al. 1992
	Insecta	Hemiptera	–	Hemiptera undetermined	Elias et al. 1992
	Insecta	Homoptera	Cicadidae	Cicadidae undetermined	Elias et al. 1992
	Insecta	Hymenoptera	Superfamily Apoidea	Apoidea undetermined	Elias et al. 1992

Appendix Table 11-A-2 (continued). Ice Age invertebrate taxa reported from the Grand Canyon. Taxa followed by an asterisk (*) were named from specimens found within GRCA.

Phylum	Class	Order	Family	Taxon Observed	Sources
Arthropoda (continued)	Insecta	Lepidoptera	–	Lepidoptera undetermined	Elias et al. 1992
	Insecta	Neuroptera	Myrmelodontidae	Myrmelodontidae undetermined	Elias et al. 1992
	Insecta	Orthoptera	Acrididae	Acrididae undetermined	Elias et al. 1992
	Insecta	–	–	Insecta undetermined	Cole and Mead 1981, Mead and Phillips 1981
	Ostracoda	Podocopida	Candonidae	*Candona* sp. (late Holocene)	Kaufman et al. 2002
	Ostracoda	Podocopida	Cyprididae	*Cypridopsis okeechobei*	Kaufman et al. 2002
	Ostracoda	Podocopida	Cyprididae	*Cypridopsis vidua*	Kaufman et al. 2002
	Ostracoda	Podocopida	Darwinulidae	*Darwinula stevensoni*	Kaufman et al. 2002
	Ostracoda	Podocopida	Cyprididae	*Heterocypris incongruens*	Kaufman et al. 2002
	Ostracoda	Podocopida	Cyprididae	*Ilyocypris bradyi*	Kaufman et al. 2002
	Ostracoda	Podocopida	Cyprididae	*Strandesia meadensis*	Kaufman et al. 2002
	–	–	–	Arthropoda undetermined	Mead and Van Devender 1981, Hevly 1984

Vertebrates

Ice Age vertebrate taxa reported from the Grand Canyon are listed in Appendix Table 11-A-3.

Appendix Table 11-A-3. Ice Age vertebrate taxa reported from the Grand Canyon.

Class	Order	Taxa Observed	Sources
Osteichthyes	–	*Catostomus discobolus*	Miller and Smith 1984
	–	*Catostomus latipinnis* (possibly Holocene)	Miller and Smith 1984
	–	*Gila cypha* (probably Holocene)	Miller and Smith 1984
	–	*Gila elegans* (probably Holocene)	Miller and Smith 1984
	–	*Ptychocheilus lucius* (probably Holocene)	Miller and Smith 1984
	–	Osteichthyes undetermined	Hevly 1984, Emslie 1988, Dryer 1994
Amphibia	Anura	*Bufo* sp.	Spamer 1988
	Anura	*Hyla* sp.	Spamer 1988
	Anura	*Scaphiopus* sp.	GCM
	Urodela	*Ambystoma tigrinum* (late Holocene)	Mead 2005
Reptilia	Testudines	*Gopherus agassizii*	Wilson 1942, Van Devender et al. 1977, Mead 1981, Mead 2005, Hunt et al. 2018
	Testudines	*Gopherus morafkai*	Hunt et al. 2018
	Squamata	*Cnemidophorus tigris*	Mead 2005
	Squamata	*Cnemidophorus* cf. *C. tigris*	Van Devender et al. 1977, Mead 1981
	Squamata	*Cnemidophorus* sp.	Cole and Van Devender 1976, Van Devender et al. 1977, Cole and Mead 1981
	Squamata	*Coleonyx variegatus*	Van Devender et al. 1977, Mead 1981, Mead 2005
	Squamata	*Crotaphytus collaris*	Van Devender et al. 1977, Mead 1981, Mead 2005, Mead and Phillips 1981
	Squamata	*Crotaphytus* cf. *C. collaris*	Cole and Mead 1981
	Squamata	*Crotaphytus* sp.	Cole and Van Devender 1976
	Squamata	*Heloderma suspectum* (uncertain age)	Mead 2005
	Squamata	*Phrynosoma hernandesi*	Mead et al. 2003, Mead 2005
	Squamata	cf. *Phrynosoma*	Mead et al. 2003
	Squamata	*Sauromalus ater*[7]	Wilson 1942, Van Devender et al. 1977, Mead 1981, Mead 2005, Hunt et al. 2018
	Squamata	*Sceloporus magister*	Hunt et al. 2018

[7] *Sauromalus obesus = Sauromalus ater*

Appendix Table 11-A-3 (continued). Ice Age vertebrate taxa reported from the Grand Canyon.

Class	Order	Taxa Observed	Sources
Reptilia (continued)	Squamata	*Sceloporus tristichus*	Hunt et al. 2018
	Squamata	*Sceloporus undulatus*	Mead 2005
	Squamata	*Sceloporus* cf. *S. magister*	Van Devender et al. 1977, Mead 1981, Mead and Phillips 1981
	Squamata	*Sceloporus* cf. *S. undulatus*	Van Devender et al. 1977, Cole and Mead 1981, Mead 1981, Mead and Phillips 1981, Hunt et al. 2018
	Squamata	*Sceloporus* sp.	Van Devender et al. 1977, Cole and Mead 1981
	Squamata	*Sceloporus* spp.	Cole and Van Devender 1976
	Squamata	*Uta stansburiana*	Van Devender et al. 1977, Mead 1981, Mead 2005, Mead and Phillips 1981
	Squamata	Undetermined lizard	Emslie 1988, Dryer 1994, Emslie et al. 1995
	Suborder Serpentes	*Coluber* or *Masticophis* (late Holocene)	Mead and Phillips 1981
	Suborder Serpentes	*Crotalus mitchelli* or *C. viridis*	Van Devender et al. 1977, Mead 1981, Mead 2005, Mead and Phillips 1981
	Suborder Serpentes	*Crotalus* sp.	Van Devender et al. 1977
	Suborder Serpentes	*Diadophis punctatus*	Mead et al. 2003, Mead 2005
	Suborder Serpentes	*Hypsiglena torquata*	Van Devender et al. 1977, Mead 1981, Mead 2005, Mead and Phillips 1981
	Suborder Serpentes	*Lampropeltis getula*[8]	Van Devender et al. 1977, Mead 1981, Mead 2005, Mead and Phillips 1981, Olsen and Olsen 1984, Hunt et al. 2018
	Suborder Serpentes	*Lampropeltis pyromelana*	Mead 1981, Mead 2005, Mead and Phillips 1981
	Suborder Serpentes	*Lampropeltis triangulum* (late Holocene)	Mead and Phillips 1981, Mead 2005
	Suborder Serpentes	cf. *Lampropeltis*	Mead et al. 2003
	Suborder Serpentes	*Pituophis catenifer*	Hunt et al. 2018
	Suborder Serpentes	*Pituophis melanoleucus*	Van Devender et al. 1977; Mead 1981, 2005
	Suborder Serpentes	*Rhinocheilus lecontei*	Van Devender et al. 1977; Mead 1981, Mead 2005, Mead and Phillips 1981; Hunt et al. 2018
	Suborder Serpentes	*Salvadora* cf. *S. hexalepis* (late Holocene)	Mead and Phillips 1981

[8] *Lampropeltis getulus = Lampropeltis getula*

Appendix Table 11-A-3 (continued). Ice Age vertebrate taxa reported from the Grand Canyon.

Class	Order	Taxa Observed	Sources
Reptilia (continued)	Suborder Serpentes	*Sonora semiannulata*	Van Devender et al. 1977; Mead 1981, Mead 2005, Mead and Phillips 1981
	Suborder Serpentes	*Thamnophis* sp.	Mead 1981
	Suborder Serpentes	*Trimorphodon bisulcatus* (late Holocene)	Mead and Phillips 1981
	Suborder Serpentes	Serpentes undetermined	Emslie 1988, Dryer 1994, Mead et al. 2003
	–	Reptilia undetermined	Hevly 1984
Aves	Accipitriformes	*Accipiter striatus* (Holocene general)	Mead 1981, Hevly 1984
	Accipitriformes	*Aquila chrysaetos*	Hevly 1984, Carpenter 2003, Hunt et al. 2018
	Accipitriformes	*Buteo jamaicensis*	Hevly 1984, Hunt et al. 2018
	Accipitriformes	*Buteo regalis* (Holocene general)	Mead 1981
	Accipitriformes	*Buteo* sp.	Emslie 1988
	Accipitriformes	cf. *Buteo jamaicensis*	Carpenter 2003
	Accipitriformes	*Buteogallus anthracinus* (Holocene general)	Mead 1981, Hevly 1984
	Accipitriformes	*Circus cyaneus*	Emslie 1988
	Accipitriformes	cf. *Circus cyaneus* (Holocene general)	Hevly 1984
	Accipitriformes	Hawk similar to *Buteo jamaicensis*	Miller 1960
	Anseriformes	*Aix sponsa* (Holocene general)	Mead 1981, Hevly 1984
	Anseriformes	*Anas acuta* (Holocene general)	Mead 1981, Hevly 1984
	Anseriformes	*Anas americana*	Hevly 1984, Emslie 1988
	Anseriformes	*Anas clypeata*	Hevly 1984, Emslie 1988
	Anseriformes	*Anas crecca*	Emslie 1988
	Anseriformes	*Anas crecca carolinensis*?	Mead 1981
	Anseriformes	*Anas crecca* cf. *carolinensis* (Holocene general)	Hevly 1984
	Anseriformes	*Anas cyanoptera* (Holocene general)	Mead 1981, Hevly 1984
	Anseriformes	*Anas discors*	Rea and Hargrave 1984
	Anseriformes	*Anas platyrhynchos*	Rea and Hargrave 1984, Emslie 1988
	Anseriformes	*Anas strepera* (Holocene general)	Mead 1981, Hevly 1984
	Anseriformes	*Anas* sp.	Emslie 1988, Emslie et al. 1995
	Anseriformes	*Aythya americana* (Holocene general)	Mead 1981, Hevly 1984
	Anseriformes	*Aythya affinis*	Hevly 1984, Emslie 1988
	Anseriformes	*Aythya marila* (Holocene general)	Mead 1981, Hevly 1984

Appendix Table 11-A-3 (continued). Ice Age vertebrate taxa reported from the Grand Canyon.

Class	Order	Taxa Observed	Sources
Aves (continued)	Anseriformes	*Aythya valisineria* (Holocene general)	Mead 1981, Hevly 1984
	Anseriformes	*Aythya* sp.	Emslie 1988
	Anseriformes	*Aythya* sp.? (Holocene general)	Hevly 1984
	Anseriformes	*Branta canadensis* (Holocene general)	Mead 1981, Hevly 1984
	Anseriformes	*Bucephala albeola* (Holocene general)	Mead 1981, Hevly 1984
	Anseriformes	*Bucephala clangula* (Holocene general)	Mead 1981
	Anseriformes	*Chen caerulescens*	Hevly 1984, Emslie 1988
	Anseriformes	cf. *Clangula hyemalis* (Holocene general)	Hevly 1984
	Anseriformes	*Mergus cucullatus* (Holocene general)	Mead 1981, Hevly 1984
	Anseriformes	*Mergus merganser* (Holocene general)	Mead 1981, Hevly 1984
	Anseriformes	*Olor columbianus* (Holocene general)	Mead 1981
	Anseriformes	cf. *Olor columbianus* (Holocene general)	Hevly 1984
	Anseriformes	*Oxyura jamaicensis* (Holocene general)	Mead 1981, Hevly 1984
	Apodiformes	*Aeronautes saxatalis*	Emslie 1988
	Cathartiformes	*Cathartes aura*	Mead 1981, Hevly 1984, Emslie 1988, Mead and Phillips 1981
	Cathartiformes	*?Cathartes aura*	Carpenter 2003
	Cathartiformes	*Coragyps atratus*	Carpenter 2003, Hunt et al. 2018
	Cathartiformes	*Coragyps occidentalis*	Hunt et al. 2018
	Cathartiformes	*Gymnogyps amplus*?	deSaussure 1956
	Cathartiformes	*Gymnogyps californianus*	*see text*
	Cathartiformes	*Gymnogyps* sp.	Rea and Hargrave 1984
	Cathartiformes	*Teratornis merriami*	Mead 1981, Rea and Hargrave 1984
	Cathartiformes	*Teratornis* cf. *T. merriami*	Dryer 1994
	Cathartiformes	*Teratornis* sp.	Lindsay and Tessman 1974
	Charadriiformes	*Actitis macularia*	Rea and Hargrave 1984
	Charadriiformes	*Calidris melanotos*	Rea and Hargrave 1984
	Charadriiformes	*Capella gallinago*	Mead 1981, Rea and Hargrave 1984
	Charadriiformes	*Larus pipixcan* (Holocene general)	Hevly 1984
	Charadriiformes	*Larus* sp.	Emslie 1988
	Charadriiformes	*Numenius americanus* (Holocene general)	Mead 1981, Hevly 1984
	Charadriiformes	*Phalaropus fulicarius*	Rea and Hargrave 1984

Appendix Table 11-A-3 (continued). Ice Age vertebrate taxa reported from the Grand Canyon.

Class	Order	Taxa Observed	Sources
Aves (continued)	Charadriiformes	*Phalaropus lobatus*	Rea and Hargrave 1984, Emslie 1988
	Charadriiformes	*Phalaropus* cf. *fulicarius*	Mead 1981
	Charadriiformes	*Recurvirostra americana*	Hevly 1984, Emslie 1988
	Charadriiformes	*Tringa semipalmata*[9]	Emslie et al. 1995
	Columbiformes	*Zenaida macroura*	Hevly 1984, Emslie 1988, Emslie et al. 1995
	Falconiformes	*Falco femoralis*	Miller 1960, Carpenter 2003, Hunt et al. 2018
	Falconiformes	*Falco mexicanus*	Hevly 1984, Emslie 1988
	Falconiformes	*Falco peregrinus*	Emslie 1988
	Falconiformes	*Falco sparverius*	Rea and Hargrave 1984, Emslie 1988, Emslie et al. 1995
	Galliformes	*Centrocercus urophasianus* (Holocene general)	Mead 1981, Hevly 1984
	Galliformes	cf. *Colinus virginianus*	Emslie 1988
	Galliformes	*Meleagris crassipes* (Holocene general)	Mead 1981, Hevly 1984
	Gruiformes	*Fulica americana*	Emslie 1988
	Gruiformes	*Gallinula chloropus* (Holocene general)	Mead 1981, Hevly 1984
	Gruiformes	cf. *Porzana carolina*	Emslie 1988
	Passeriformes	*Agelaius phoeniceus*	Emslie 1988
	Passeriformes	*Aphelocoma caerulescens* (Holocene general)	Mead 1981
	Passeriformes	cf. *Aphelocoma caerulescens* (Holocene general)	Hevly 1984
	Passeriformes	*Catherpes mexicanus* (Holocene general)	Mead 1981, Hevly 1984
	Passeriformes	*Cinclus mexicanus*	Rea and Hargrave 1984
	Passeriformes	*Contopus sordidulus* (Holocene general)	Mead 1981, Hevly 1984
	Passeriformes	*Corvus corax*	Hevly 1984, Emslie 1988
	Passeriformes	*Corvus corax sinuatus* (Holocene general)	Mead 1981
	Passeriformes	*Corvus* sp.	Emslie 1988, Emslie et al. 1995
	Passeriformes	*Dendroica coronata* (Holocene general)	Mead 1981, Hevly 1984
	Passeriformes	*Empidonax* sp.? (Holocene general)	Hevly 1984
	Passeriformes	*Eremophila alpestris*	Mead 1981, Hevly 1984
	Passeriformes	*Hirundo* sp. (Holocene general)	Mead 1981

[9] *Catoptrophorus semipalmatus = Tringa semipalmata*

Appendix Table 11-A-3 (continued). Ice Age vertebrate taxa reported from the Grand Canyon.

Class	Order	Taxa Observed	Sources
Aves (continued)	Passeriformes	*Hirundo* sp.? (Holocene general)	Hevly 1984
	Passeriformes	*Icterus galbula* (Holocene general)	Hevly 1984
	Passeriformes	*Icterus* sp. (Holocene general)	Mead 1981
	Passeriformes	*Junco hyemalis*	Mead 1981, Rea and Hargrave 1984
	Passeriformes	cf. *Junco* sp.	Emslie 1988
	Passeriformes	*Lanius excubitor* (Holocene general)	Mead 1981, Hevly 1984
	Passeriformes	*Loxia* cf. *L. curvirostra*	Mead 1981
	Passeriformes	*Myadestes townsendi* (Holocene general)	Mead 1981, Hevly 1984
	Passeriformes	*Passerella iliaca* (Holocene general)	Hevly 1984
	Passeriformes	*Passerina* sp. (Holocene general)	Hevly 1984
	Passeriformes	*Pica hudsonia* (Holocene general)[10]	Hevly 1984
	Passeriformes	*Salpinctes obsoletus* (Holocene general)	Hevly 1984
	Passeriformes	*Salpinctes obsoletus*?	Mead 1981
	Passeriformes	*Sayornis nigricans* (Holocene general)	Hevly 1984
	Passeriformes	*Sayornis saya* (Holocene general)	Mead 1981, Hevly 1984
	Passeriformes	*Sialia currucoides* (Holocene general)	Mead 1981, Hevly 1984
	Passeriformes	*Turdus grayi* (Holocene general)	Hevly 1984
	Passeriformes	*Turdus migratorius* (Holocene general)	Hevly 1984
	Passeriformes	*Turdus migratorius*?	Mead 1981
	Passeriformes	*Turdus* sp.?	Mead 1981
	Passeriformes	*Zonotrichia* cf. *Z. leucophrys* (Holocene general)	Hevly 1984
	Passeriformes	cf. Fringillidae (late Holocene)	Mead and Van Devender 1981
	Passeriformes	Passeriformes undetermined	Emslie 1988, Dryer 1994, Emslie et al. 1995
	Pelecaniformes	*Ardea herodias* (Holocene general)	Mead 1981, Hevly 1984
	Pelecaniformes	*Nycticorax nycticorax*	Hunt et al. 2018
	Pelecaniformes	cf. *Nycticorax nycticorax*	Carpenter 2003

[10] *Pica pica hudsonica = Pica hudsonia*

Appendix Table 11-A-3 (continued). Ice Age vertebrate taxa reported from the Grand Canyon.

Class	Order	Taxa Observed	Sources
Aves (continued)	Pelecaniformes	Ardeidae undetermined	Carpenter 2003
	Piciformes	*Colaptes auratus*	Emslie 1988
	Piciformes	*Sphyrapicus varius*	Emslie 1988
	Piciformes	Picidae undetermined	Emslie 1988
	Podicipediformes	*Aechmophorus occidentalis*	Hevly 1984, Emslie 1988
	Podicipediformes	*Podiceps auritus* (Holocene general)	Mead 1981
	Podicipediformes	*Podiceps nigricollis* (Holocene general)	Mead 1981, Hevly 1984
	Podicipediformes	cf. *Podiceps nigricollis*	Emslie 1988
	Podicipediformes	*Podilymbus podiceps*	Emslie 1988
	Strigiformes	*Bubo virginianus*	Mead 1981, Mead and Van Devender 1981, Rea and Hargrave 1984
	Strigiformes	*Otus asio* (Holocene general)	Mead 1981, Hevly 1984
	Strigiformes	*Tyto alba*	Miller 1960, Carpenter 2003, Hunt et al. 2018
	–	Aves undetermined	Cole and Mead 1981, Mead and Phillips 1981, Hevly 1984, Emslie 1988, Dryer 1994
Mammalia	Pilosa	*Nothrotheriops shastensis*[11]	*see text*
	Eulipotyphla	*Notiosorex crawfordi*	Mead 1981, Mead and Phillips 1981, Mead and Van Devender 1981, Emslie 1988
	Rodentia	*Ammospermophilus* cf. *A. leucurus* (late Holocene)	Mead and Van Devender 1981
	Rodentia	*Castor canadensis* (Holocene general)	Mead 1981, Olsen and Olsen 1984
	Rodentia	*Dipodomys* sp.	Lindsay and Tessman 1974, Mead 1981, Mead and Van Devender 1981, Hunt et al. 2018
	Rodentia	*Erethizon dorsatum*	Van Devender et al. 1977, Mead 1981, Mead and Phillips 1981, Hunt et al. 2018
	Rodentia	*Eutamias* sp.	Lindsay and Tessman 1974, Cole and Mead 1981, Hunt et al. 2018
	Rodentia	*Lemmiscus curtatus*	Mead et al. 2003
	Rodentia	*Marmota flaviventris*	Lange 1956, Van Devender et al. 1977, Hunt et al. 2018
	Rodentia	*Marmota flaviventris* cf. *M. f. engelhardti*	Wilson 1942
	Rodentia	*Marmota* cf. *M. flaviventris*	Mead 1981, Mead and Phillips 1981

[11] *Nothrotherium shastense = Nothrotheriops shastensis*

Appendix Table 11-A-3 (continued). Ice Age vertebrate taxa reported from the Grand Canyon.

Class	Order	Taxa Observed	Sources
Mammalia (continued)	Rodentia	*Marmota* sp.	Lindsay and Tessman 1974, Emslie et al. 1995
	Rodentia	*Microtus* sp.	Cole and Mead 1981, Mead 1981, Mead and Phillips 1981, Emslie 1988, Mead et al. 2003
	Rodentia	*Neotoma cinerea*	Mead et al. 2003
	Rodentia	*Neotoma devia* or *N. lepida*	Hunt et al. 2018
	Rodentia	*Neotoma lepida*	Van Devender et al. 1977, Mead 1981, Hunt et al. 2018
	Rodentia	*Neotoma mexicana*	Mead 1981, Hunt et al. 2018
	Rodentia	*Neotoma stephensi*	Van Devender et al. 1977, Mead 1981, Hunt et al. 2018
	Rodentia	*Neotoma* cf. *N. cinerea*	Cole and Mead 1981
	Rodentia	*Neotoma* cf. *N. lepida*	Cole and Mead 1981
	Rodentia	*Neotoma* cf. *N. mexicana*	Van Devender et al. 1977
	Rodentia	*Neotoma* sp.	Lindsay and Tessman 1974, Van Devender et al. 1977, Cole and Mead 1981, Olsen and Olsen 1984, Emslie 1988, Dryer 1994, Mead et al. 2003
	Rodentia	*Neotoma* spp.	Mead 1981, Mead and Phillips 1981
	Rodentia	*Ondatra zibethicus* (Holocene general)	Mead 1981, Olsen and Olsen 1984
	Rodentia	*Perognathus* cf. *P. intermedius*	Mead 1981, Mead and Phillips 1981
	Rodentia	*Perognathus* sp.	Emslie 1988
	Rodentia	*Peromyscus* sp.	Van Devender et al. 1977, Cole and Mead 1981, Mead and Phillips 1981, Olsen and Olsen 1984, Emslie 1988, Dryer 1994, Mead et al. 2003, Hunt et al. 2018
	Rodentia	*Peromyscus* spp.	Mead 1981, Mead and Van Devender 1981
	Rodentia	cf. *Reithrodontomys*	Cole and Mead 1981
	Rodentia	*Sciurus* sp. (Holocene?)	Olsen and Olsen 1984
	Rodentia	cf. *Sciurus* sp.	Emslie 1988
	Rodentia	*Spermophilus variegatus*	Mead 1981, Mead and Phillips 1981
	Rodentia	*Spermophilus* sp.	Lindsay and Tessman 1974, Mead et al. 2003, Hunt et al. 2018
	Rodentia	*Tamias* sp.	Hunt et al. 2018
	Rodentia	*Thomomys* sp.	Lindsay and Tessman 1974, Cole and Mead 1981, Hunt et al. 2018
	Rodentia	Sciuridae undetermined	Emslie et al. 1995
	Rodentia	Rodentia undetermined	Hevly 1984
	Lagomorpha	*Lepus californicus*	Mead 1981, Olsen and Olsen 1984
	Lagomorpha	*Lepus* near *L. californicus*	Wilson 1942

Appendix Table 11-A-3 (continued). Ice Age vertebrate taxa reported from the Grand Canyon.

Class	Order	Taxa Observed	Sources
Mammalia (continued)	Lagomorpha	*Lepus* sp.	Lindsay and Tessman 1974, Emslie 1988, Emslie et al. 1995, Hunt et al. 2018
	Lagomorpha	*Sylvilagus* cf. *S. audubonii*	Olsen and Olsen 1984
	Lagomorpha	*Sylvilagus* sp.	Lindsay and Tessman 1974, Van Devender et al. 1977, Mead 1981, Emslie 1988, Emslie et al. 1995, Mead et al. 2003, Hunt et al. 2018
	Lagomorpha	Lagomorpha undetermined	Hevly 1984
	Chiroptera	*Antrozous pallidus*	Emslie 1988
	Chiroptera	*Desmodus stocki*	Ray et al. 1988, Carpenter 2003, Hunt et al. 2018
	Chiroptera	*Eptesicus* cf. *E. fuscus*	Olsen and Olsen 1984
	Chiroptera	*Euderma maculatum*	Mead and Mikesic 2005
	Chiroptera	*Eumops* sp.	Carpenter 2003, Hunt et al. 2018
	Chiroptera	*Lasiurus cinereus*	*see text*
	Chiroptera	*Lasionycteris noctivagens*	*see text*
	Chiroptera	*Myotis* sp. (possibly Holocene)	Olsen and Olsen 1984
	Chiroptera	cf. *Myotis* sp.	Emslie 1988
	Chiroptera	*Pipistrellus hesperus* [*Parastrellus*]	Emslie 1988
	Chiroptera	*Plecotus townsendi* [*Corynorhinus*]	Emslie 1988
	Chiroptera	*Tadarida brasiliensis*	Carpenter 2003, Hunt et al. 2018
	Chiroptera	Undetermined Chiroptera	Van Devender et al. 1977
	Carnivora	*Bassariscus astutus*	Wilson 1942, Mead 1981, Mead and Phillips 1981, Hunt et al. 2018
	Carnivora	*Bassariscus* sp.	Lindsay and Tessman 1974
	Carnivora	*Canis latrans* (Holocene general)	Mead 1981, Olsen and Olsen 1984
	Carnivora	*Canis* sp. (wolf)	Emslie 1988
	Carnivora	*Lontra canadensis*	Mead 1981, Olsen and Olsen 1984
	Carnivora	*Lynx rufus*	Mead 1981
	Carnivora	*Lynx* sp.	Wilson 1942, Hunt et al. 2018
	Carnivora	*Mustela* sp.	Lindsay and Tessman 1974, Hunt et al. 2018
	Carnivora	*Procyon lotor* (Holocene general)	Mead 1981, Olsen and Olsen 1984
	Carnivora	*Puma concolor*[12]	Mead 1981, Mead et al. 2003, Hunt et al. 2018
	Carnivora	*Puma concolor*?	Wilson 1942

[12] *Felis concolor = Puma concolor*

Appendix Table 11-A-3 (continued). Ice Age vertebrate taxa reported from the Grand Canyon.

Class	Order	Taxa Observed	Sources
Mammalia (continued)	Carnivora	*Spilogale putorius*	Emslie 1988
	Carnivora	*Spilogale gracilis* or *S. putorius*	Hunt et al. 2018
	Carnivora	*Spilogale* sp.	Lindsay and Tessman 1974
	Carnivora	*Urocyon cinereoargenteus* (Holocene general)	Mead 1981, Olsen and Olsen 1984
	Carnivora	*Vulpes vulpes*	Carpenter 2002
	Carnivora	Canidae undetermined	Carpenter and Mead 2000
	Proboscidea	*Mammuthus* sp.	Emslie 1987, Emslie 1988
	Perissodactyla	*Equus* sp. (*E. conversidens*?)	Harington 1984
	Perissodactyla	*Equus* sp.	Wilson 1942, Mead 1981, Emslie 1987, Emslie 1988, Carpenter 2003, Mead et al. 2003, Hunt et al. 2018
	Artiodactyla	*Antilocapra americana*	Mead 1981, Mead and Phillips 1981
	Artiodactyla	*Bison* sp.	Harington 1984, Emslie 1987, Emslie 1988, Mead et al. 2003, Martin 2014, Martin et al. 2017
	Artiodactyla	*Camelops hesternus*	Mead et al. 2003
	Artiodactyla	*Camelops* cf. *C. hesternus*	Mead 1981, Mead and Phillips 1981
	Artiodactyla	*?Camelops* sp.	Emslie 1987
	Artiodactyla	cf. *Camelops* sp.	Emslie 1988
	Artiodactyla	*Euceratherium collinum*	Mead et al. 2003, Kropf et al. 2007
	Artiodactyla	cf. *Euceratherium collinum*	Mead et al. 2003
	Artiodactyla	*Odocoileus hemionus* (Holocene general)	Mead 1981, Olsen and Olsen 1984
	Artiodactyla	*Odocoileus* sp.	Mead 1981, Mead and Phillips 1981
	Artiodactyla	*Oreamnos harringtoni*	*see text*
	Artiodactyla	*Oreamnos* sp. (probably Holocene)	Harington 1984
	Artiodactyla	*Oreamnos* or *Ovis*	Emslie 1988
	Artiodactyla	*Ovis canadensis*	Mead 1981, Mead and Phillips 1981, Olsen and Olsen 1984, Harington 1984, Carpenter 2003, Mead et al. 2003, Hunt et al. 2018
	Artiodactyla	*Ovis* sp.	Wilson 1942
	Artiodactyla	Bovidae undetermined	Carpenter 2003
	Artiodactyla	Artiodactyla undetermined	Cole and Mead 1981, Harington 1984, Dryer 1994, Carpenter 2003
	–	Undetermined large mammal	Emslie 1988, Emslie et al. 1995
	–	Mammalia undetermined	Hevly 1984
	–	Vertebrata undetermined	Hevly 1984

Ichnofossils and Reproductive Traces

Ice Age ichnofossils and reproductive traces reported from the Grand Canyon can be seen in Appendix Table 11-A-4.

Appendix Table 11-A-4. Ice Age ichnofossils and reproductive traces reported from the Grand Canyon. Taxa followed by an asterisk (*) were named from specimens found within GRCA.

Category	Traces Observed	Sources
Invertebrates	Dipteran pupal case	Hevly 1984
	"Helminth" eggs	Schmidt et al. 1992
	Nematode eggs	Laudermilk and Munz 1938
Vertebrates	Artiodactyl dung	Mead and Swift 2012, Hunt et al. 2018
	Bassariscus astutus dung	Mead and Swift 2012, Hunt et al. 2018
	Bat guano	*see text*
	Bison sp. dung	Mead and Swift 2012
	Equus sp. dung	Mead and Swift 2012
	Erethizon dorsatum dung	Mead and Swift 2012
	Large felid dung	Mead and Swift 2012, Hunt et al. 2018
	cf. *Lepus* sp. dung	Mead and Swift 2012, Hunt et al. 2018
	Neotoma spp. dung	abundant in *Neotoma* middens; *see text*
	Nothrotheriops shastensis dung (*Castrocopros martini**)	*see text*
	Oreamnos harringtoni dung	*see text*
	Ovis canadensis dung	Robbins et al. 1984, O'Rourke and Mead 1985, Mead and Swift 2012, Hunt et al. 2018
	Peromyscus sp. dung	Emslie 1988
	Rabbit dung	Hevly 1984
	Rodent dung	Hevly 1984, Mead and Swift 2012, Hunt et al. 2018
	Sauromalus dung (age not stated)	Mead and Swift 2012
	cf. *Sylvilagus* sp. dung	Mead and Swift 2012, Hunt et al. 2018
	Bird regurgitation pellets	Emslie et al. 1995, Mead and Swift 2012
	Bassariscus astutus middens (late Holocene)	Mead 1981, Mead and Phillips 1981
	Neotoma spp. *middens*	*see text*
	Cathartes aura eggshells	Miller 1960, Harington 1975
	Gymnogyps californianus eggshells	Emslie 1987
	Gymnogyps nest	Martin 2014

Other Fossils

Other fossil Ice Age taxa reported from the Grand Canyon are listed in Appendix Table 11-A-5.

Appendix Table 11-A-5. Other Ice Age fossil taxa reported from the Grand Canyon. Taxa followed by an asterisk (*) were named from specimens found within GRCA.

Phylum	Class	Class or Subclass	Traces Observed	Sources
Apicomplexa	Conoidasida	Coccidia	*Archaeococcidia antiquus**	Schmidt et al. 1992
	Conoidasida	Coccidia	*Archaeococcidia nothrotheriopsae**	Schmidt et al. 1992
–	–	–	Fungal spores	Robbins et al. 1984, Schmidt et al. 1992

Chapter 12. Grand Canyon National Park Paleontological Resources Management and Protection

By Diana Boudreau[1], Vincent Santucci[2], Klara Widrig[1], Mark Nebel[1], Anne Miller[1], Ronnie Colvin[1], Kim Besom[1], and Colleen Hyde[1]

[1]Grand Canyon National Park
Grand Canyon, Arizona 86023-0129

[2]National Park Service
Geologic Resources Division
1849 C Street, NW, Room 2644
Washington, DC 20240

Introduction

Paleontological resources are an integral part of Grand Canyon National Park (GRCA). As stated in the park's foundation statement, one of the purposes of GRCA is to preserve and protect Grand Canyon's unique geologic, paleontologic, and other natural and cultural features for the benefit and enjoyment of the visiting public. Furthermore, paleontological resources are considered a fundamental resource, warranting primary consideration during planning and management, contributing to the park's significance, and helping achieve the park's purpose (NPS 2017).

The foundation document also highlights fossil resources of particular interest, such as the diverse invertebrate and vertebrate trace fossils in the Coconino Sandstone, Precambrian stromatolites, Pleistocene vertebrate fossils found in cave deposits, and the large assemblages of marine invertebrate fossils in the Redwall Limestone, Surprise Canyon Formation, and Kaibab Formation. While it is known that GRCA contains an extremely diverse and complete fossil record, visitor use trends and site conditions for heavily visited fossil localities are lacking. The foundation statement supports the development of this paleontology inventory report and paleontological resource protection plan in order to preserve paleontological resources and support the park's purpose (NPS 2017).

In previous years, there have been a few dedicated paleontologists studying fossil tracks, cave fossils, vertebrates, and other fossil resources found at GRCA; however, they are often outside researchers who do not work directly for the NPS. During the 2019 Paleontology Inventory, GRCA preserved and protected paleontological resources by building a park-based team and partnering with paleontologists from the NPS Geologic Resources Division. Therefore, 2019 has been an important year that demonstrated that establishing a team of dedicated paleontologists can greatly advance the knowledge and protection of fossil resources in NPS units and provide valuable outreach opportunities for the public.

GRCA Paleontological Resource Inventory

The inventory of paleontological resources is one of the most fundamental resource management activities that can be undertaken by a park. Baseline paleontological resource inventories help

identify the scope, significance, distribution, and resource management issues associated with fossils. The information gained through the inventory of park fossils enables park management to incorporate this information into park planning, programming, and decision-making.

Paleontological resource inventories are specifically identified in Section 6302 of the Paleontological Resources Preservation Act of 2009 (16 USC 470aaa–1), in Section 4302 of the Federal Cave Resources Protection Act of 1988 (16 USC 4301), and in NPS policy. The NPS Paleontology Program provides assistance to parks, including GRCA, by documenting and preserving paleontological resource information in the NPS Paleontology Archives and Library.

Between 1970 and 2010, most of the paleontology related activities undertaken at GRCA were primarily related to research and collecting by outside academic paleontologists and geologists working in the park. Jim Mead and students from the Quaternary Studies Program at Northern Arizona University coordinated paleontological research and collecting at Rampart Cave and other caves in GRCA (Mead 1981; Mead and Van Devender 1981; Mead and Phillips 1982; Mead et al. 2003; Carpenter 2003). In 2001, an inventory of paleontological resources associated with NPS caves, including those in GRCA, was undertaken by the NPS Geologic Resources Division (Santucci et al. 2001). This work later led to a GRCA-focused inventory of cave paleontological resources (Kenworthy et al. 2004). A more comprehensive inventory of paleontological resources at GRCA was included in a report documenting the fossils of the Southern Colorado Plateau Inventory and Monitoring Network in 2009 (Tweet et al. 2009). Between 2012 and 2018 a trans-boundary and collaborative project known as the Greater Grand Canyon Landscape Assessment (GGCLA) was undertaken to assess cave resources, including cave fossils, in and around the park. This project was expanded to include a Resource Condition Assessment (RCA) for the cave resources evaluated in the GGCLA project (Stortz 2018).

A number of paleontology interns were hired at GRCA after Deanna Greco was hired as the GRCA Physical Science Program Manager in 2010. Cassi Knight, Jeff Dobbins, James Super, Robyn Henderek, and Anne Miller served as Geoscientists-in-the-Parks (GIP) paleontology interns at GRCA and assisted with a variety of paleontological resource inventory projects. The work completed by these paleontology interns, along with their field notes, are incorporated into the GRCA museum and resource management archives.

In preparation for GRCA's centennial in 2019, the NPS Paleontology Program (Vincent Santucci) initiated communication with GRCA staff (Jeanne Calhoun) in early 2018 proposing the development of a park specific paleontological resource inventory for GRCA in conjunction with the park's centennial. This proposal was met with support from GRCA leadership and is the basis for this report.

In addition to the publication of this paleontology resource inventory report, GRCA staff conducted several surveys to document fossil localities as part of the second ever NPS PaleoBlitz (the first was conducted at CHIC in 2016). A few stratigraphic units at GRCA were the focus of these PaleoBlitz activities; Chinle Formation, Moenkopi Formation, Kaibab Formation, Coconino Sandstone, and Bright Angel Shale. Paleontologists Adam Marsh and Bill Parker from Petrified Forest National Park

(PEFO) assisted in surveying the Mesozoic strata. JP Hodnett, program coordinator for the Maryland-National Capital Park and Planning Commission (M-NCPPC) Dinosaur Park (MD) and National Fossil Day coordinator, conducted surveys for vertebrate material in the Kaibab Formation on the North and South Rims. The GRCA Paleontology Team (Mark Nebel, Anne Miller, Diana Boudreau, Klara Widrig, and Jered Hansen) conducted targeted surveys of, primarily, the Coconino Sandstone and, to a lesser degree, the Kaibab and Toroweap Formations and the Bright Angel Shale, throughout the 2019 season, and discovered a number of previously unknown track localities (Figure 12-1). Each survey used a combination of field notebooks, paper locality forms, and digital data collection devices leveraging the ArcGIS Online and Collector technologies to physically and digitally collect locality and site information, setting the stage for faster and more efficient data entry for future paleontology surveys.



Figure 12-1. The GRCA paleontology team documenting a new fossil track locality in the Coconino Sandstone during the 2019 PaleoBlitz (Left to Right: Klara Widrig, Anne Miller, Diana Boudreau, and Jered Hansen) (NPS/MARK NEBEL).

GRCA Paleontological Resource Monitoring

The monitoring of in situ paleontological resources is another important resource management tool for the NPS. Paleontological resource monitoring enables the long term assessment of fossil stability

through the use of a variety of techniques (Santucci and Koch 2003; Santucci et al. 2009). Monitoring strategies may be developed for each fossil locality to consider the site specific variables, either natural or anthropogenic, which may threaten the paleontological resources and their values. Paleontological monitoring prescriptions are developed to determine the site specific monitoring activities such as 1) how frequently to monitor the fossil locality; 2) which monitoring methods are to be used; 3) when conditions warrant the collection of fossils from the locality; and other considerations.

The most common technique to monitor paleontological sites is repeat photography. GRCA archives contain many images of known fossil sites, many of which would benefit from monitoring. Many of these images have recently been linked to the paleontology geodatabase for ease of access and ability to spatially analyze paleontology site and locality data. Recreating these images would allow GRCA staff to assess the vulnerability of those sites due to natural or anthropogenic effects. Photogrammetry is another great option for monitoring. By taking a series of photographs and stitching them together to create 3D models, it is possible to preserve large specimens, such as trackways, to establish a baseline condition. However, neither of these techniques are quantitative in nature. Miller et al. developed a methodology, based on paleontological sites from GRCA, that uses clearly defined indices to quantitatively identify high-priority sites for monitoring (2018). Both quantitative and qualitative methods should be utilized to monitor paleontological resources.

There are a number of natural processes, environmental conditions, and other natural factors beyond human management that affect the stability of paleontological sites at GRCA. Fossils are primarily affected by weathering and erosion, particularly at GRCA where unvegetated outcrops and seasonal climate are common. Freeze-thaw cycles contribute to increased fracturing and monsoonal precipitation can increase rates of erosion. Winds transporting sediments can also gradually abrade fossils already exposed. Fossils found near creeks and streams at GRCA are threatened by flash flood activity that could abrade exposed fossils or wash fossils out of geologic context and deeper into the canyon. Sudden geologic processes, such as rock falls, may also reveal, cover, or damage fossil resources. Biological activity, such as lichen growth, plant growth, or trampling by deer, elk, or bighorn, could impact the visibility or quality of those fossils.

In addition to natural processes, anthropogenic or human caused impacts, both intentional and unintentional, can lead to degradation of fossil sites. Unintentional harm could be done by individuals that are not aware of the fossils or those who don’t understand that their actions negatively impact fossil stability. Despite the fossil protection provided by the Paleontological Resources Preservation Act (PRPA), fossils are still intentionally disturbed, removed, or vandalized. Due to these impacts, determining the monitoring plan for a fossil site should factor in ease of public access, visitor use near or at the site, potential for any ground disturbing activities, such as trail maintenance or construction work, and other factors that may contribute to theft or vandalism (Miller et al. 2018: Table 4).

GRCA Paleontological Resource Protection

Paleontological resources are threatened by natural processes such as erosion, as well as anthropogenic impacts, such as unauthorized collecting and vandalism. These threats impact the

fossils' inherent scientific and educational value, as removing fossils from their geologic context impairs scientists' ability to accurately understand a fossil's significance in the geologic record. For resource management and protection, patrols of known paleontological sites and resources should be conducted to identify any indirect or direct human or natural impacts.

In 2009, the Paleontological Resources Preservation Act (PRPA) was signed into law as part of the Omnibus Public Land Management Act of 2009. New paleontology legislation is currently being drafted by an interagency coordination team including representatives from the Bureau of Land Management (BLM), Bureau of Reclamation (BOR), National Park Service (NPS), and U.S. Fish & Wildlife Service (FWS) within the Department of the Interior (DOI). This legislation will provide guidance related to fossil inventories, monitoring, public education, research and collecting permits, curation, and criminal prosecution on DOI lands. For more information regarding this act, visit https://www.nps.gov/subjects/fossils/fossil-protection.htm.

History of Paleontological Resource Protection at GRCA

Fossil theft and vandalism at GRCA have not been frequently reported. Few incident reports have been filed regarding paleontology theft and vandalism. However, anecdotal evidence from Law Enforcement Officers (LEO) in South Rim, Inner Canyon, and North Rim districts indicate that damage to fossils may occur more frequently than the incident report records show (Figures 12-2 and 12-3). Most fossil incidents occur within the Canyon District, because more fossils are exposed in that district.



Figure 12-2. Track blocks have been chiseled out and removed from the Coconino Sandstone along Hermit Trail (left). Individuals have attempted to remove other trackways (solid blue oval), but did not complete the task, leaving behind chisel marks (dashed red oval) (NPS/VINCENT SANTUCCI).



Figure 12-3. A nail was used by a visitor attempting to pry this brachiopod fossil from the Kaibab Formation (NPS/VINCENT SANTUCCI).

The first record of fossil vandalism was in 1946. Superintendent Harold Bryant saw threats to park resources, including fossils, and compiled a list including abused signs, interpretive displays, trees, shrubbery, and graffiti, as well as noting damaged or stolen binoculars and a fossil specimen from the Yavapai Observation Station (now renamed Yavapai Geology Museum). Bryant's efforts emboldened staff to prevent, as he called them, "such vicious acts of destruction." In addition, this was the start of littering fines and proactive measures to lessen harm to park resources (Anderson 2000).

More recently, there have been continued reports of fossil theft and vandalism. In 2007 or 2008, law enforcement investigated a report of a fossil track block that was stolen off of North Kaibab Trail. During Spring Break of 2017, the Fossil Fern Exhibit along South Kaibab Trail at Cedar Ridge and fern impressions surrounding the exhibit were heavily vandalized (Figures 12-4 and 5). Most of the damage involved scratching rocks and the plexiglass covering the exhibit. Part of the plexiglass was broken, exposing the fossils housed inside the exhibit. Fossil specimens may have been removed from the area by visitors, but this has not been confirmed.



Figure 12-4. Scratch marks on the Fossil Fern Exhibit structure along the South Kaibab Trail in 2017 (NPS/ANNE MILLER).




Figure 12-5. Photographs from 2013 **(A)** and 2017 **(B)** of the same fossil fern impressions near the exhibit showing vandalism that occurred during Spring Break in 2017 (NPS).

On May 20, 2019 two individuals were caught collecting approximately 13.5 kg (30 lbs.) of specimens from the fossil beds along the western Rim Trail (Figure 12-6). They were reported by a visitor, and law enforcement responded in time to apprehend the individuals. The couple appeared to be avid rock collecting hobbyists and used small hammers and picks to collect specimens. All fossil specimens were confiscated and are being held as evidence.



Figure 12-6. Rock and fossil specimens found in suspect's backpack after GRCA law enforcement apprehended them removing specimens from park land in 2019 (NPS).

Paleontological Resource Protection Recommendations

Canyon Rangers, law enforcement, and GRCA staff should be attentive during their regular duties for individuals attempting to take or deface specimens. Many well-preserved specimens are close to the main trails in the park and are therefore at a higher risk of damage or theft due to ease of access. Regular patrols should be conducted near heavily visited paleontological localities.

Fossil theft and vandalism trainings can better prepare staff to identify, report, and manage fossil theft and vandalism incidents. For more information on Paleontology Resource Protection trainings, contact Vincent Santucci (vincent_santucci@nps.gov). Regular training of LEOs and interpretive staff can help reduce the number of fossil theft or vandalism incidents. However, active recruitment of paleontological research scientists should also be considered a management strategy.

If fossil specimens are found during regular patrols by staff or visitors, follow these steps:

- Do not collect, remove, or disturb the fossils without consulting the Chief of Resources or individual responsible for paleontological resources at GRCA.
- Record locality information, including, but not limited to, geographic location, nearest trail or natural feature, GPS coordinates, and description of fossil within the rock.

- Take photos of the fossil specimen itself with a scale, a wider view of the fossil in the surrounding rock, and images looking up trail, down trail, and/or with identifiable landmarks in frame. Use an item for scale in each photo (e.g., another hiker, backpack, water bottle, coin).
- Reach out to individuals responsible for paleontological resources at GRCA to notify them of the resource and provide them with the associated information.

GRCA Paleontological Resource Research and Permits

Internal Permits

Several internal blanket permits were issued to past Chiefs of Resources to facilitate study and interpretation of geological and paleontological resources within the park. Permit GRCA-2011-SCI-0034 was issued to Deanna Greco to facilitate a paleontological inventory of Grand Canyon National Park with Geoscientists-in-the-Parks (GIP) interns Jeff Dobbins and James Super. Judy Helmich was issued permit GRCA-2003-SCI-0085 that allowed interpretive staff to collect specimens while in the field to use in their programs. Both of these permits are now expired.

On October 18, 2019, the GRCA Paleontology Program received a programmatic Categorical Exclusion (CE) (PEPC 90271) to guide ongoing paleontology work in the Park. The CE describes routine, ongoing paleontology inventory and monitoring work and resource mitigations for performing this work.

Previous Research Permits

Jim Mead and Mary Carpenter were issued permit GRCA-1999-SCI-0001 for their project *"Late Pleistocene Vertebrate Communities in the Lower Grand Canyon, Arizona: Rampart and Muav Caves."* This work supported Mary Carpenter's thesis (Carpenter 2003) and an abstract (Carpenter and Mead 1999).

John Foster was issued permit GRCA-2009-SCI-0032 in an attempt to relocate historic trilobite quarry locations. The permit was for in situ research only. He was unsuccessful in relocating the Cameron-Walcott quarry site, but the McKee Quarry was relocated in 2009. His findings were published in the Museum of Northern Arizona bulletins (Foster 2011).

Jessica Metcalfe was issued permit GRCA-2013-SCI-0001 for her dissertation project, "*Late Pleistocene Paleoecology of the Colorado Plateau*." The overall goal of this study was to reconstruct late Pleistocene paleoecology on the Colorado Plateau using isotopic analysis to examine feeding and migratory patterns. No collections were made as part of this project, which instead relied on GRCA and MNA museum collections (Metcalfe 2018).

Permit GRCA-2017-SCI-0054 was issued to Stephen Rowland for the study of fossil trackways in the Supai Formation. Research was focused on a trackway along the Bright Angel Trail, which may represent the oldest vertebrate trackway in the park. A paper on this trackway is currently in peer review.

Anne Miller was issued permit GRCA-2017-SCI-0059 for her Master's thesis at Northern Arizona University "*Ichnology of the Bright Angel Shale Formation, Grand Canyon, AZ: Indicators for Middle Cambrian Paleoecology."* This research is still ongoing.

A complete list of paleontology research permits issued in Grand Canyon National Park from 1999 to 2019 can be found in Appendix 12-A.

Current Permits

Permit GRCA-2019-SCI-0002 was issued to Vincent Santucci for the Grand Canyon National Park 2019 Paleontological Resource Inventory and PaleoBlitz. All field activities related to this report were authorized by this permit. Field activity is focused on the documentation and assessment of paleontological localities; however, surface collection of exceptional specimens is permitted.

GRCA Paleontological Resource Curation and Museum Collections

Museum collections at GRCA are stored at 2C Albright Avenue, Grand Canyon, Arizona, 86023. This storage facility was completed in 1999. It contains 6,000 ft^2 (557 m^2) of climate-controlled storage and houses over 1.6 million objects. Items are cataloged using the Interior Collections Management System (ICMS), version 8.20.48.3660 as of April 29, 2019.

Scope of Paleontological Collections

Grand Canyon National Park's paleontology collection includes a total of 13,428 catalog numbers. Of these, the vast majority were found within the park. Specimens collected from outside the park were once brought into the collections for teaching, research, exhibit, or other purposes, with the majority being from Arizona. This practice was discontinued in 1978, when the service's management policies stated "a scope of collection statement, in which the limits of museum collection are detailed, must be prepared and approved for every park." Fossils from other NPS units that remain in the GRCA collections include: 12 specimens from Harpers Ferry Center, one of which was returned to the Harpers Ferry Center museum storage facility; 38 specimens from Lake Mead National Recreation Area (LAKE); 45 specimens from Petrified Forest National Park (PEFO); and 239 specimens from Grand Canyon-Parashant National Monument (PARA). Most of the non-GRCA fossils are from areas in the vicinity of the park in northern Arizona. It should be noted that some Grand Canyon fossil specimens come from lands formerly part of GRCA, but now within the Havasupai Indian Reservation, which may lead to some confusion, because the original locality information may report that they are from the park. With 2,857 cataloged specimens, brachiopods are the most common fossil type in the collection, followed by mammals (2,177), trilobites (1,438), and bivalves (1,006). When considering the rock unit of origin, the greatest number of fossils in the collection were found in Quaternary deposits (4,994 catalog numbers), followed by the Kaibab Formation (3,302) and the Bright Angel Shale (1,669).

At least 167 fossil taxa have been named from specimens (holotypes, syntypes, etc.) collected within the modern boundaries of GRCA (Appendix A). An additional 13 taxa are based on specimens potentially collected within GRCA, but the provenance information is too vague to know for certain (Appendix B) (Tweet et al. 2016). Four holotype specimens are stored in a locked cabinet separate from the rest of the paleontology collections for added security.

Notable Contributors to Paleontological Collections

Many of the fossils housed at Grand Canyon were collected by notable geologists and paleontologists including: Raymond Alf, Merrill Beal, Stan Beus, Major L. F. Brady, Jim Mead, Kenneth L. Cole, Arthur L. Lang, Raymond deSaussure, Steve Emslie, Bob Euler (archeologist), Lawrence Goebels, Paul Martin, Edwin D. McKee, Larry Coates, Charles Merriam, Arthur Phillips, Allen Phillips, Larry Powers, Louis Schellbach, Edward T. Schenk, Adolph Seilacher, Glen Sturdevant, and David White.

Paleontological Collections in Outside Repositories

Numerous GRCA specimens are housed at outside institutions. The exact number of these institutions will probably never be known. There are several reasons for this. Among the most important factors are the following: the Grand Canyon region has long been a popular area for geological and paleontological field trips; undocumented collecting frequently occurred before the permitting system was implemented; the park boundaries have changed several times; standards for reporting locality information were formerly more relaxed, so that fossils collected from within GRCA are undoubtedly lost behind provenance information limited to “Grand Canyon” or “Arizona”; provenance information has been partially or entirely lost when material from smaller collections has been absorbed by other institutions; and many fossil specimens are easily accessible, portable, and therefore prone to illegal collecting. Because of these and other factors, it would not be surprising to find a small quantity of Grand Canyon fossils in any large paleontological collection.

A list of institutions containing GRCA specimens has been compiled from the literature. This list is based on mentions of definite GRCA specimens in specific collections (particularly type and figured specimens). This is not an exhaustive list. Contact information is provided in Appendix D.

- Academy of Natural Sciences of Drexel University (ANSP, formerly the Academy of Natural Sciences of Philadelphia; Philadelphia, Pennsylvania)
- American Museum of Natural History (AMNH; New York, New York)
- Brigham Young University (BYU; Provo, Utah)
- California Academy of Sciences, including material from the former collections of Stanford University (CAS; San Francisco, California)
- Carnegie Museum of Natural History (CMNH; Pittsburgh, Pennsylvania)
- Cincinnati Museum Center, including material from the former collections of the University of Minnesota (CMC; Cincinnati, Ohio)
- Columbia University (CU; New York, New York)
- Field Museum of Natural History (FMNH; Chicago, Illinois)
- Harvard University Herbaria (HUH; Cambridge, Massachusetts)
- Lake Mead National Recreation Area (LAKE; Boulder City, Nevada)
- Lund University (UL; Lund, Sweden)
- Museum of Comparative Zoology (MCZ; Harvard, Massachusetts)
- Museum of Northern Arizona (MNA; Flagstaff, Arizona)

- Natural History Museum of Los Angeles County, including material from the former collections of the California Institute of Technology and University of California at Los Angeles (LACM; Los Angeles, California)
- Raymond Alf Museum (RAM; Claremont, California)
- Smithsonian National Museum of Natural History (USNM; Washington, D.C.)
- University of Arizona Laboratory of Paleontology (UALP; Tucson, Arizona)
- University of California Museum of Paleontology (UCMP; Berkeley, California)
- University of Notre Dame (UND; Notre Dame, Indiana)
- University of Tübingen (Eberhard Karls University) (UT; Tübingen, Germany)
- Western Archeological and Conservation Center (WACC; Tucson, Arizona)
- Yale Peabody Museum (YPM; New Haven, Connecticut)

USNM houses significant material from GRCA, including the holotypes of 138 of the 167 fossil species named from GRCA material. Notable non-holotype specimens include a complete vampire bat skull (*Desmodus stocki*) found in Rampart Cave (USNM V 25478). The Smithsonian also houses the GRCA collections of Charles Doolittle Walcott, Charles Gilmore's track collections, David White's fossil plants, and Rampart Cave collections.

Photographic Archives

The GRCA Museum Collection maintains an image index database of historical photos related to paleontology in the park. This collection includes 247 photos, the oldest of which is an 1858 portrait of J. S. Newberry, the first geologist to describe fossils from Grand Canyon. Other notable geologists and paleontologists in the image database include David White, John Merriam, George Hesemann, Edwin McKee, and Charles Doolittle Walcott. Many of the photographs in the database document fieldwork and ichnofossils throughout the canyon. As such, they are useful for relocating historic sites and providing comparative photos for site monitoring and management. The oldest of these field photographs was taken in 1913, by Francois Matthes, and documents vertebrate tracks in the Coconino Sandstone. In 1930, a series of photographs of ichnofossils in the Coconino Sandstone were taken by George Grant. Civilian Conservation Corps (CCC) projects in the late 1930s were well documented via photography. These projects included the construction of the Fossil Fern Exhibit on the South Kaibab (Yaki) Trail, construction of trilobite exhibits near Indian Garden, and work at Rampart Cave. The CCC project at Rampart Cave was photographed by Michael Bobko and includes photographs of sloth dung filling the cave and the CCC crew excavating fossil bones. Kaibab fossils within the Museum Collection were also extensively photographed in the 1930s.

The image index database also includes a handful of photographs from outside the park. Photos of dinosaur tracks near Moenkopi, Cameron, and Tuba City can be found in the archives. Two photos of modern animal tracks in a sand dune north of Kanab, UT, were added for comparison with fossil trackways. Other notable photos include petrified wood from PEFO and a human skeleton at Willow Beach photographed during the CCC expedition to Rampart Cave. Western portions of GRCA were

once within the boundary of Lake Mead National Recreation Area (LAKE), so additional archival items may be found within LAKE's archives.

Digital Archives

3D models produced by photogrammetry techniques are an excellent way to make fossils available to the general public and remote researchers. A total of 15 photogrammetry models have been created from GRCA specimens using this technique. Two models can be found on the Smithsonian 3D Digitization webpage. Both specimens were originally from GRCA and are now housed at USNM.

In April 2019, a project with Geology and Photogrammetry Specialist Jack Wood (GRD) was initiated to digitize additional museum specimens to display as online 3D models for National Fossil Day. These models have been uploaded to the National Park Service Geologic Resource Division Sketchfab page (https://sketchfab.com/grd_nps/models), GRCA photogrammetry series website (https://www.nps.gov/articles/series.htm?id=A9E62040-AC6F-A6D7-BE564A036F1D6146), and photos and files associated with each 3D model have been archived on IRMA. For a complete list of specimens with photogrammetry 3D models see Appendix 12-B.

Very few photographic records exist for fossils within the collections. Although it would be a lengthy undertaking, photographing each item in the collection would be extremely useful for archival and research purposes. It would also be valuable to digitize more museum objects via 3D photogrammetry. This would make the museum collections far more accessible to visitors and researchers, the vast majority of whom do not have the opportunity to tour or visit the museum collection.

Outreach

The GRCA Museum Collection is a storage and research facility that is open to the public by appointment only. All tours of the collections must be requested in advance. Museum curators normally give 50 to 60 tours each year. The main audiences for these tours are school groups, park staff, and members of other affiliated organizations, such as Xanterra and the Grand Canyon Conservancy. A yearly art exhibition is held in mid-September, where the museum hosts a three-hour open house to display the art collection.

GRCA Paleontological Resource Interpretation, Education, and Outreach

The fossils preserved in the rock layers of GRCA are an integral part of explaining the geologic history of the canyon to the public. Their presence helps scientists better understand past depositional environments and how the landscape and lifeforms of this region have changed over time. There are many ranger-led programs, self-guided hikes, brochures, booklets, and educational programs at GRCA that focus on fossils and paleontological resources. All of the programs, handouts, and available resources are compiled here to better prepare interpretation and education staff. While most events are localized to the South Rim, any large interpretive event or program that is developed by the park should extend to the North Rim, Desert View, and Inner Canyon districts as often as possible. This can be accomplished by distributing flyers, cards, stickers, posters, or by adapting programs for use at these other locations.

Current Long Range Interpretive Plan

There is no current or proposed long range interpretive plan for paleontology at GRCA.

Current Paleontological Interpretive Programs

Ranger-Led Programs

Fossil Walk—Summer Only

The Fossil Walk program takes visitors on a journey back in time to discover what fossils can tell us about this part of the continent 270-million-years ago. Visitors join a Park Ranger for a hike along the western Rim Trail from Bright Angel Trailhead to visit the fossil beds preserved in the Kaibab Formation. Fossils are abundant in this area including *Meekella* and productid brachiopods, crinoid stem segments, stick and lacy bryozoans, burrows, horn corals, and sponges. This locality is also very close to the rim, providing an excellent opportunity to discuss other fossiliferous units in the canyon and the significance of the fossil record.

Geo-Glimpse—Year-Round

This 20-minute ranger-led program explains how Grand Canyon formed while exploring the canyon rim near Yavapai Geology Museum. Although fossils are not a central part of this program, paleontological topics are woven into the geologic history of the Grand Canyon to better understand paleoenvironments and the canyon we see today. A 45-minute version of this program is offered spring to fall as a Geo-Walk Ranger Program.

Junior Ranger Family Program—Summer Only

During summer months, in Grand Canyon Village on the South Rim, special family-friendly programs are offered with youth and Junior Rangers in mind. Two of these programs, Family Adventure Hike and Natural Wonders, highlight the paleontological resources found within the layered rocks of Grand Canyon. Family Adventure Hike guides individuals down the Hermit Trail for a two-hour hike to explore the canyon's resources first hand. Natural Wonders is a 30-minute Park Ranger presentation on the historic El Tovar Hotel Canyon-side porch.

Evening Programs—Summer Only

Evening programs presented by Park Rangers cover a wide variety of topics. Occasionally, programs discuss paleontology and geology of GRCA.

Self-Guided Hikes

Trail of Time Geology Walk—South Rim, Year-Round

The Trail of Time is a self-guided walking timeline trail along the South Rim that prompts visitors to understand the complex and long geologic history of Grand Canyon without having to leave the rim (Figure 12-7). Examples of each of the major rock layers are displayed along the trail, some of which contain fossil resources such as stromatolites, trace fossils, and invertebrate body fossils. A number of wayside signs explain paleontological concepts such as evolution, deep time, depositional environments, and erosion of past fossil-bearing strata. A Trail of Time companion booklet was compiled to provide visitors with a more in-depth interpretive experience. Self-guided materials for school groups or other larger groups are also available for teachers who could not schedule a ranger-led field trip.



Figure 12-7. Entrance portal along the Trail of Time near the Grand Canyon Headquarters building (NPS/DIANA BOUDREAU).

Widforss Trail—North Rim, April through October

The Widforss Trail provides access to many views of the canyon along the North Rim. The interpretive brochure available at the trailhead or Visitor Center provides visitors with paleontology information at stops 2 and 9. Stop 2 encourages visitors to examine the fossils preserved in the Kaibab Formation beneath their feet including crinoids and other small marine creatures (Figure 12-8). Stop 9 identifies the cliff forming Coconino Sandstone from a particularly scenic overlook. Visitors are informed of the wealth of fossil tracks preserved in this prominent geologic unit and the brochure includes an image of fossil vertebrate tracks.



Figure 12-8. Crinoid stems, stick and lacy bryozoans, and shell fragments from exposures of the Kaibab Formation can be found along the rim of the canyon (NPS/DIANA BOUDREAU).

Curriculum-Based Programs

Ranger-Led Field Trips—Spring and Fall

Each year the Environmental Education Branch of Grand Canyon National Park offers many ranger-led field trips for grades 3–12. Students from Arizona, and across the world, sign up for these free, curriculum-based programs. “Stories in Stone” is a two-and-a-half-hour trip focused on teaching introductory geology and paleontology, how fossils form, how to interpret ancient environments, and the importance of field work and making observations. “Grand Canyon Rocks” is a similar field trip to “Stories in Stone”, but it includes more exercises related to geology rather than paleontology. “Dynamic Earth” is a five-hour field trip that combines the information presented in these two programs. During each of these field trips, students fill out a journal with guiding questions and exercises to introduce new geologic and paleontological terms and concepts (Figure 12-9).



Figure 12-9. Students identify fossils in the Kaibab Formation at the fossil beds during the “Stories in Stone” field trip (NPS/RONNIE COLVIN).

Classroom Ranger Visits—Winter Only

During the winter months, Park Rangers make classroom visits to nearby schools in Arizona and surrounding states to educate students about a variety of topics, including paleontology and geology of GRCA.

Distance Learning—Year-Round

These programs are available year-round to classrooms or anyone with an internet connection. One of the distance learning programs is the “Ancient Life Program” which aims to teach 3rd–5th grade students about fossils. This program introduces students to the National Park Service, orients them to the geographic location of GRCA, familiarizes students with examples of GRCA fossils and rock units that contain fossils, relates rock types to depositional environments, explains environmental change over geologic time, and introduces students to the vast amount of time represented by the rock layers in the Grand Canyon. Students also better understand fossil bias in the rock record by playing a card game in which students are either turned into fossils or not, depending on the cards they are dealt.

Junior Paleontologist Program—Year-Round

This nationwide program targets kids age 5–12 and encourages them to explore, learn about, and protect fossil resources in National Parks, including those at Grand Canyon. The NPS Geologic Resources Division can provide Junior Paleontologist Activity Program supplies including activity booklets, badges, posters, and other fossil-related educational resources. Contact Vincent Santucci for more details (vincent_santucci@nps.gov).

Grand Canyon Field Institute and School Programs

There are a number of outside companies that offer guided trips into the canyon, such as the Grand Canyon Field Institute that is run by the park's non-profit partner, Grand Canyon Conservancy. Some of these guided trips take time to educate their groups on the geology and fossil resources found within the park.

Canyon Field School is a partnership between the Grand Canyon Conservancy and the National Park Service and offers unique opportunities for youth to experience the great outdoors at Grand Canyon National Park. They offer overnight camps that center around scientific discovery and learning at Grand Canyon National Park.

National Fossil Day Event

National Fossil Day is an annual event that takes place nationwide on Wednesday of the second full week in October, which is also Earth Science Week. Grand Canyon National Park has hosted a National Fossil Day event on the South Rim for the past five years (Figure 12-10). In the past, Park Rangers have led a fossil walk program, scheduled fossil themed games and activities for kids, hosted a Q&A session with fossil experts, remotely toured the Kaibab Formation fossil beds and interacted with rangers and fossil experts using "Facebook Live", and presented a fossil themed evening program focused on National Fossil Day and GRCA fossil resources. The NPS Geologic Resources Division can assist the park with planning for National Fossil Day activities and provide Junior Paleontologist Activity Program supplies including activity booklets, badges, posters, and other fossil-related educational resources. Contact Vincent Santucci for more details (vincent_santucci@nps.gov).



Figure 12-10. Park Rangers talk with the public about fossils at the 2016 National Fossil Day celebration at Grand Canyon National Park (NPS).

The 2019 National Fossil Day Celebration was a larger event compared to previous years, as an opportunity to commemorate the park's 100th anniversary (Figure 12-11). In 2019, Grand Canyon National Park hosted a number of outside researchers and partners on the South Rim on September 27 and 28. The festivities started off with a mini Paleontology Symposium on Friday, September 27, 2019. Four guest speakers shared their research on GRCA paleontology and specimens from the museum were on display during the symposium (Figure 12-12). During the National Fossil Day Celebration on September 28, visitors could participate in ranger-led programs to the fossil beds, create fossil-themed crafts during children's activities at the Yavapai Geology Museum, and were even able to ask paleontologists questions virtually through a special Facebook Live event. Twelve partner organizations participated with informational booths at the Grand Canyon Visitor Center plaza, including a green-screen photobooth provided by the American Geosciences Institute (Figure 12-13). A welcome ceremony for the event took place at the Mather Amphitheater with guest speakers Bruce MacFadden (President of the Paleontological Society) and Vincent Santucci (NPS Senior Paleontologist), Science and Resource Management's Chief of Resources, Jeanne Calhoun, and an Arizona State Greeting for the National Fossil Day event (Figure 12-14). The National Fossil Day Celebration was capped by a special paleontology evening program given by NPS Senior Paleontologist and Paleontology Program Coordinator Vincent Santucci. For more in-depth information on the National Fossil Day Celebration, refer to the 2019 Paleontology Project After Action Report (Boudreau 2020).



Figure 12-11. NPS paleontologists and educators who assisted with the 10th Annual National Fossil Day activities and programs on the South Rim of Grand Canyon National Park on September 28, 2019. Left to right (front): Jeremy Childs, Jennifer Glennon, Anne Miller, Hazel Wolfe, Celia Dubin, Kevin Garcia, John-Paul Hodnett, Vincent Santucci, Mary Carpenter, Jim Mead, Joel Despain, Adam Blankenbicker, Chris Symons, Jason Kenworthy, and Justin Tweet. Left to right (back): Grace Lilly, Janet Gillette, David Gillette, Tom Olson, Richard McMichael, Don Weeks, Anne Scott, Eleanour Snow, Andy Grass, Sequoyah McGee, Robyn Henderek, Sandy Croteau, Erin Eichenberg, Diana Boudreau, Sherman Mohler, Mary Ontiveros, Doug Wolfe, Maria Rodriguez, Veronica Colvin, Joel Kane, and Bryan Maul (NPS/MICHAEL QUINN).



Figure 12-12. Advertisement for the four speakers at the Paleontology Symposium on Friday September 27, 2019 at the Shrine of the Ages (NPS).



Figure 12-13. Advertisement for the National Fossil Day Celebration event at the South Rim of Grand Canyon National Park (NPS).



Figure 12-14. Speakers at the Welcome Ceremony at Mather Amphitheater at the National Fossil Day Celebration on September 28, 2019. Left to right: Jeanne Calhoun, Ronnie Colvin, Bruce MacFadden, and Vincent Santucci (JOHN-PAUL HODNETT).

Below are some statistics concerning visitation during the National Fossil Day Celebration:

- Approximately 110 visitors attended the Paleontology Symposium on Friday, September 27, 2019
- 85 visitors attended the Paleontology Evening Program given by Vincent Santucci on Saturday, September 28, 2019
- Ask-A-Scientist social media livestream reached 7,617 individuals, a maximum of 104 individuals viewed simultaneously, and 116 viewers submitted comments or questions
- Fossil Bed social media livestream event reached 23,788 individuals, over double the viewership compared to last year which reached 10,756 individuals. This livestream peaked at 290 viewers and received 116 comments and questions.

A number of new interpretive materials were developed specifically for the 2019 National Fossil Day event. Interpretation and education staff designed a portable pop-up exhibit, highlighting fossils from GRCA with paleo environments, to use for events hosted at GRCA, such as National Fossil Day, or for classroom ranger visits (Figure 12-15). This pop-up exhibit was unveiled on September 27, 2019,

and was on display at the Grand Canyon Visitor Center for two months following the event. A set of nine GRCA fossil-themed trading cards and National Fossil Day informational postcards were also developed for distribution at fossil events.



Figure 12-15. A large pop-up exhibit highlighting Grand Canyon National Park fossils was developed for the 2019 National Fossil Day Celebration at Grand Canyon (NPS/DIANA BOUDREAU).

Interpretive Signs and Exhibits

Permanent Signs and Exhibits

South Rim

There are a number of fossil interpretive signs on the South Rim in Grand Canyon Village. The largest concentration of paleontological interpretation is at the Yavapai Geology Museum. There is a small wall case that contains real and replica fossil specimens, such as brachiopods, stromatolites, trilobites, worm burrows, and a cast of a dragonfly wing. Many interpretive signs within the museum explain past depositional environments at Grand Canyon and have models of fossils from the canyon, including a trilobite with tracks, brachiopods, and *Chelichnus* tetrapod trackway (Figure 12-16). A cast of the Shasta Ground Sloth skull and dung are on exhibit in the main lobby of the Grand Canyon Visitor Center. A regularly scheduled program about the formation of the Grand Canyon titled “The Canyon World” plays in the Science on the Sphere exhibit at the Grand Canyon Visitor Center and highlights the fossil record.



Figure 12-16. Paleozoic exhibit at the Yavapai Geology Museum (NPS/DIANA BOUDREAU).

The Rim Trail between the Grand Canyon Visitors Center and the Yavapai Geology Museum is void of interpretive signage; however, traveling west along the Rim Trail from the Yavapai Geology Museum, visitors explore the "Trail of Time." See the Trail of Time section above for more information. The Rim Trail is often bordered by a small stone wall that contains large chunks of petrified wood. Most notable is the section of wall across from Verkamp's Visitors Center (Figure 12-17); it also has two large sections of petrified wood placed by the entrance ramp.



Figure 12-17. Two large chunks of petrified wood used as building materials for the wall along the Rim Trail outside the Verkamp's Visitor Center on the South Rim (NPS/DIANA BOUDREAU).

A number of other localities on the South Rim expose the public to GRCA's rich fossil assemblage. Bright Angel Lodge has a historic geology fireplace designed by Mary Colter (Figure 12-18A). Visitors can see stromatolites and a coiled nautiloid embedded in the rock fireplace. In the central courtyard of GRCA Headquarters, a few track blocks have been placed in the benches and stone floor (Figure 12-18B). In the main lobby, a small exhibit, compiled by a previous GIP, contains paleo environment images, fossil specimens, and an explanation of the fossil history of GRCA. These are examples of paleontological resources in cultural contexts; see Kenworthy and Santucci (2006) for further information on similar occurrences.




Figure 12-18. A. The geology fireplace in the Bright Angel Lodge designed by Mary Colter exhibits geology and paleontology of the Grand Canyon (NPS/MICHAEL QUINN). **B.** Fossil trackway slab used as building material for a bench in the Grand Canyon Headquarters courtyard. (NPS/DIANA BOUDREAU).

North Rim

The North Rim has one interpretive sign that outlines the paleoenvironments of the Kaibab, Coconino, Hermit, and Supai formations. The sign is located along a paved path between the North Rim Lodge and Bright Angel Point, providing an opportunity for visitors to read about past depositional environments of the canyon. Visitors can also learn about erosion, geologic time, and lack of Mesozoic strata and dinosaurs in the park on an interpretive sign at Point Imperial.

Desert View

There is no paleontological signage at Desert View. However, a coiled nautiloid that was found in the Kaibab Formation during the construction of the Desert View Watchtower was placed in a small glass wall case with other cultural and historic artifacts on the 2nd floor of the watchtower with a small descriptive label (Figure 12-19).




Figure 12-19. A coiled nautiloid found in the Kaibab Formation during construction of the Desert View Watchtower on display at Desert View (NPS/SALLY CARTTAR).

Canyon District

Most of the paleontological interpretation within the canyon district occurs via ranger programs. However, interpretive staff are planning to restore and add interpretive signage to the historic Fossil Fern Exhibit along the South Kaibab Trail at Cedar Ridge. For more information on this exhibit see the Historic Interpretive Sites section.

Future Signs and Exhibits

There are a few ongoing projects to update and add to paleontological interpretation at Grand Canyon. A proposed wayside exhibit for the fossil beds site would provide visitors with information for that site year-round, however there are concerns about fossil theft and vandalism at the site. Staff also plan to restore the historic Cedar Ridge CCC Fossil Fern Exhibit along the South Kaibab Trail, produce an interactive fossil component for the GRCA park app, and develop a mini program to display at the Grand Canyon Visitor Center's "Science on the Sphere". Construction on the Maswick South Lodging Complex began in 2019 and each of the four new lodging buildings will be named for fossils (Trilobite, Shasta, Fern, and Burrow). Xanterra plans to incorporate paleontological interpretive and outreach elements into the construction and landscaping of these buildings.

Historic Interpretive Sites

Cedar Ridge CCC Fossil Fern Exhibit

Located along the South Kaibab Trail at Cedar Ridge approximately 2.4 km (1.5 mi) down trail, the CCC Fossil Fern Exhibit resides at the edge of a cliff (Figure 12-20). During the construction of the South Kaibab Trail (formerly called the Yaki Trail) in the 1930s, trail workers uncovered well-preserved fossil ferns in the Hermit Formation. Recognizing their importance, the trail crew constructed an exhibit case to display and protect the well-preserved fossil plant materials. Since that time, little has been done to maintain and protect this exhibit. Therefore, the paleontology, interpretation, and outreach staff will work along with volunteers to rebuild this historic site in future years.




Figure 12-20. Fossil Fern Exhibit at Cedar Ridge on South Kaibab Trail in the 1930s **(A)** and in the spring of 2019 **(B)** (NPS).

"Lost" Trilobite Exhibit

A trilobite exhibit was built near Indian Garden in 1935 along the Bright Angel Trail by the CCC. There are no indicators on the trail today that this site ever existed. A team from the Museum of Western Colorado relocated one quarry using historic images in 2011 (Foster 2011). This exhibit was developed near a known fossil locality along the Tonto Trail. A total of three quarries were excavated; however, only the third quarry proved suitable for an exhibit. Unfortunately, the exhibit case did not protect the fossils from water and erosional processes. The Bright Angel Shale is very friable in nature, which caused the overlying shales to fall on top of the exhibit, obscuring the fossil trilobites. Noticing this threat to fossil resources, the park superintendent requested the exposed fossils be removed from the exhibit before it was completely buried. These specimens are now housed in GRCA museum collections. The location of the trilobite quarries is not entirely known by GRCA staff.

Interpretive Handouts, Site Bulletins, Books, and Brochures

2015. Grand Canyon National Park. Journey Through Time: Grand Canyon Geology. PDF File.

Thayer, D. 2009. An Introduction to Grand Canyon Fossils. Grand Canyon Association. 64 p.

Karlstrom, K, and L. Crossey. 2019. The Grand Canyon Trail of Time Companion: Geology Essentials for your Canyon Adventure. Four Colour Printing, Korea. 142 p.

Interpretive Staff Training and Resources

Paleontology Trainings

No formal paleontological training is provided to seasonal or permanent interpretive staff. However, Ronnie Colvin, Interpretive Park Ranger, provides a "Fossil 101" training upon request for incoming interpretive staff with an interest in fossils and paleontology. There are also teacher workshops available upon request that are taught by Ronnie Colvin and the Education Branch staff (Figure 12-21). In addition, interpreters and others interested in paleontological training may contact the park paleontologist to explore participating in field work.



Figure 12-21. Teacher workshop at the fossil beds site along the Rim Trail (NPS).

Paleontological Interpretation Resources

There are a number of tools and resources available for the interpretive staff. A cache of fossil specimens from GRCA are available for use in public programs and outreach events. The education and interpretation department has a few items located within the park. These specimens were deaccessioned from the museum collections and made available to the interpretation division.

Contact GRCA Museum Specialists Kim Besom (x7766) or Colleen Hyde (x7769) for more information. In previous years, a collection permit was provided for the interpretation staff to acquire more educational materials; however the permit has since expired and should be updated if more materials are needed.

As technology advances, examples of fossil resources are also available in digital forms. During the 2019 paleontology inventory, a number of important museum specimens were imaged to create 3D models which could then be 3D printed or viewed online as interactive models. See Appendix 12-B for a detailed list of available 3D specimens. Contact Vincent Santucci (vincent_santucci@nps.gov) or GRCA Museum Specialists Kim Besom (x7766) and Colleen Hyde (x7769) for more information.

Park websites were produced by the Geologic Resources Division to compile information about the 2019 Paleontology Inventory, PaleoBlitz, National Fossil Day Celebration, and links to available 3D specimens. See https://www.nps.gov/subjects/fossils/100-years-of-paleontological-discoveries-at-the-grand-canyon.htm, https://www.nps.gov/articles/park-paleo-fall-2019-grand-canyon.htm, and https://www.nps.gov/grca/learn/nature/fossils.htm for more information.

All paleontology programs developed by Park Rangers should instruct visitors on how to be paleontologically aware while in the park. The following topics should be highlighted when mentioning fossils in interpretive programs:

- When paleontologists survey for fossils, an important tool is a geologic map. Paleontological resources are more common in certain geologic units, so knowing where those units are exposed is important for a successful survey. Other field paleontology tools include small picks and brushes, appropriate glues and consolidants, GPS, camera, topographic maps, and appropriate First Aid and safety equipment. It might be helpful to provide examples of these items for visitors during an interpretive talk, if available.
- If fossils are found in the park by a visitor, they should photograph it, note the coordinates, and notify a ranger of where the resource was found, but most importantly they should leave the fossil where they found it. It is extremely important for scientific and resource management purposes for original location information to be preserved. Visitors should be informed that park fossils are non-renewable resources that are protected by federal law (Paleontological Resources Preservation Act, 2009).

GRCA Paleontological Resource Data Management

GRCA has been a focal point of paleontology research even before its designation as a National Park 100 years ago, leading to a wealth of data and resources that need to be properly managed. In addition, the Paleontological Resources Preservation Act (PRPA) mandates the management of fossils and associated data using scientific principles and expertise. Information regarding paleontological resources at GRCA, such as field notes, publications, and paleontology locality datasheets, are managed within the Park's internal network. Field data collected at paleontological sites and localities are maintained within a geospatial database. NPS Paleontology Archives and Library has a copy of all these files. These databases and archives have been fully updated as part of the 2019 Paleontology Inventory.

GRCA Paleontology Archives

GRCA maintains a paleontology archive and library to organize all documents related to paleontology resources at the park. Items such as field notes, publications, historic photos, maps, research permits, and museum collection documents can be found within the archives.

Due to the sensitive nature of paleontological resources, paleontology archives are restricted access only. To request access to information within the archives, contact Mark Nebel (mark_nebel@nps.gov) or Anne Miller (anne_miller@nps.gov).

NPS Paleontology Program Archives (WASO)

The National Park Service Paleontology Program (WASO) maintains a copy of all digital and hardcopy field notes, sketches, photographs, maps, reports, publications, and lists of archived fossil specimens at GRCA (Santucci et al. 2018). This ensures the safety of paleontological resource data in the event either GRCA or WASO archives are lost. To request access to materials in the WASO Paleontology Archives contact Vincent Santucci (vincent_santucci@nps.gov) or Justin Tweet (justin_tweet@nps.gov).

E&R Files

E&R files (from "Examination and Report on Referred Fossils") are unpublished internal USGS documents. For more than a century, USGS paleontologists identified and prepared informal reports on fossils sent to the survey by other geologists, for example to establish the relative age of a formation or to help correlate beds. The system was eventually formalized as a two-part process including a form sent by the transmitting geologist and a reply by the survey geologist. Sometimes the fossil identifications were incorporated into publications, but in many cases this information is unpublished. These E&R files include documentation of numerous fossil localities within GRCA, including from L. F. Noble's and Edwin McKee's work. Extensive access to the original files was granted to the NPS by the USGS beginning in 2014 (Santucci et al. 2014).

McKee and Walcott Archives

Paleontological research has a long history at GRCA. Two individuals, Edwin McKee and Charles Walcott, were heavily involved in recording and studying fossils in the canyon and created extensive archives. The McKee Archives, housed at USGS in Denver, include hand drawn stratigraphic columns, specimen lists, and correspondence with outside researchers. The Walcott Archives, housed at the Smithsonian Institute, are almost entirely composed of his field notes and sketches from 1930. To request access to these archives, contact Vincent Santucci (vincent_santucci@nps.gov), Justin Tweet (justin_tweet@nps.gov), Mark Nebel (mark_nebel@nps.gov), or Anne Miller (anne_miller@nps.gov).

Geospatial Database

Earle E. Spamer originally created, for his own personal and professional use, a *Paradox* database in the 1980s to track published data, and some unpublished theses, on paleontological resources of the Grand Canyon region and their related taxonomy. This database was later migrated to an *Access* database format that was presented by Spamer to the National Park Service (NPS) in 2003.

GRCA GIS Program Manager Mark Nebel and Geologist Steve Rice initially developed a spatial paleontology geodatabase for GRCA in 2011, adapting and significantly modifying the framework of Spamer's *Access* database, specifically for use at GRCA. The primary motivation for this effort was the imminent arrival, after many years without a paleontology program, of two Geoscientists-in-Parks (GIP) paleontology interns at GRCA. This initial GRCA geodatabase consisted of a single point feature class for paleontology localities. This was the primary database used and populated by a series of GIP interns at GRCA from 2011 into 2017.

Beginning in early 2017, GRCA Paleontology (Anne Miller) and GIS (Mark Nebel) staff recognized many shortcomings of the existing database and the need for development of a more comprehensive and systematic paleontology geodatabase for tracking data on paleontological resources in the park. After more than a year of development, including a needs assessment, database schema design, development, and field testing, the new GRCA paleontology geodatabase was deployed for park use in 2019. This database was accompanied by the development and testing of new field forms. GIS layer files (.lyr) have also been developed to provide a consistent map symbology for all feature classes in the geodatabase. This geodatabase can be adapted for use at other NPS units. For more information, contact Mark Nebel (mark_nebel@nps.gov).

The GRCA paleontology geodatabase is an ESRI file geodatabase that currently contains five feature classes (Paleontology Localities, Sites, Photopoints, Survey Lines, and Survey Polygons) and two related geodatabase tables (Paleontology Photos and Collected Specimens) (Figure 12-22). Photograph files are managed separately through a GRCA internal file system. All features in the paleontology database and all photographs have unique record IDs and are subject to naming conventions.




Figure 12-22. Screenshot from ArcCatalog showing the structure of the GRCA Paleontology geodatabase (NPS).

A ***PaleontologyLocality*** is a contiguous fossiliferous area (polygon) of a known minimum extent, based on field observations, and characterized by a unique geographic location, geology, and fossil assemblage. “In Context” ***PaleontologyLocality*** polygons include fossils that are in place within bedrock (“In Situ”), or situated loose on underlying bedrock of the formation from which the fossils are directly derived. Minor weathering out or local transport may have occurred. “Float” ***PaleontologyLocality*** polygons include fossils that are out of context, i.e., derived from a geologic formation different from that on which they are situated or located, having been transported to their current location.

A ***PaleontologySite*** point can be either a ***PaleontologyLocality*** discovery point, a specific fossil or fossil assemblage to be noted or monitored, or a fossil specimen collection point. ***PaleontologySite*** features are always within a ***PaleontologyLocality***.

PaleoPhotopoints are points that represent the locations from which ***PaleontologyLocality*** photographs are taken, including photos of representative fossils, outcrop or outcrop-scale photos, and photos of setting or location. They are referenced to a ***PaleontologyLocality*** (as opposed to a ***PaleontologySite***), but do not need to be within a locality polygon (Figure 12-23). Photographs provide scientific documentation of the resource and provide a baseline for monitoring their condition over time. Photographs are hot-linked in the geodatabase, so that they can be viewed directly from an ArcGIS map application connected to the GRCA network file system.




Figure 12-23. Screenshot from ArcMap showing an example of a Paleontology Locality with associated Sites and Photopoints (NPS).

PaleontologySurvey (line and polygon) features delineate areas where systematic paleontology surveys have been undertaken.

PaleoPhotos and ***CollectedSpecimen*** geodatabase tables record any photographs or collected specimens, respectively, that are associated with the Localities, Photopoints, and Sites, as appropriate, and are linked to the respective features via geodatabase relationship classes. Photographs are hot-linked to their respective ***PaleoPhotopoints*** via the GRCA network file system.

The geodatabase was designed initially to be compatible with Trimble GPS data collection devices, data dictionaries, and workflows. GRCA is currently developing and migrating to new protocols that leverage the ArcGIS Online and Collector software environments and GNSS data collection devices. The GRCA Paleontology geodatabase, definitions, protocols, naming conventions, and schema, field forms, etc. are evolving documents. To obtain the most current version of these documents contact Mark Nebel (GIS Program Manager; mark_nebel@nps.gov).

The following evolving GRCA paleontology documents are available upon request:

1. Geodatabase Documentation (.docx)
2. Geodatabase Schema and Domains (.xlsx)
3. Paleontology Field Form and Guide (.docx or .pdf)

Conclusions

As illustrated in the many chapters of this paleontological resources inventory report and the after-action report (Boudreau 2020), 2019 has been an important year in establishing a comprehensive baseline for paleontological resource stewardship and science at Grand Canyon National Park. Collectively, the paleontology-focused projects undertaken and accomplishments achieved at GRCA during 2019 represents the largest collaborative effort for paleontology in National Park Service history. This unprecedented work will help guide and inform future paleontological resource planning and activities for other NPS areas. In addition, the concurrence of this 2019 project with the celebration of GRCA's centennial, 10th anniversary of National Fossil Day, and the Department of Interior's final rulemaking for the Paleontological Resources Preservation Act (2009) bolstered each management action and supports continued affirmative paleontological resource management and stewardship within the park.

Despite the paleontology program at Grand Canyon being understaffed for many decades, great strides have been made, building upon 150 years of dedicated individuals who observed, recorded, preserved, and promoted the paleontological resources found within the Grand Canyon. In recent years, the park's paleontology projects have been primarily accomplished through the supervision of Geoscientists-in-the-Parks interns and as collateral duties by physical science or GIS staff members, with guidance and assistance from the WASO NPS Paleontology Program. The establishment of a committed, multi-disciplinary team for the 2019 paleontology project provided an opportunity for the development of goals and projects to advance the paleontological knowledge of the park on behalf of science, public education, and resource stewardship. The GRCA Paleontological Resource Inventory provides the essential foundation for current and future park leaders to ensure that the non-renewable

paleontological resources at GRCA will continue to be monitored, conserved, and protected into the future.

Literature Cited

Anderson, M. F. 2000. World War and its wake, 1942–1945. Pages 41–56 *in* M. F. Anderson. Polishing the jewel: An administrative history of Grand Canyon National Park. Grand Canyon Association, Grand Canyon, Arizona. Monograph 11.

Boudreau, D. M. 2020. Grand Canyon National Park: 2019 paleontology project after action report. Natural Resource Report NPS/GRCA/NRR—2020/2054. National Park Service, Fort Collins, Colorado.

Carpenter, M. C. 2003. Late Pleistocene Aves, Chiroptera, Perissodactyla, and Artiodactyla from Rampart Cave, Grand Canyon, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Carpenter, M. C., and J. I. Mead. 1999. Late Pleistocene vertebrate communities of the lower Grand Canyon: Rampart and Muav Caves. Journal of Vertebrate Paleontology 19(supplement to 3):36A.

Foster, J. R. 2011. Trilobites and other fauna from two quarries in the Bright Angel Shale (Middle Cambrian, Series 3; Delamaran), Grand Canyon National Park, Arizona. Pages 99–120 *in* J. S. Hollingsworth, F. A. Sundberg, and J. R. Foster, editors. Cambrian stratigraphy and paleontology of northern Arizona and southern Nevada. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 67.

Kenworthy, J. P., and V. L. Santucci. 2006. A preliminary investigation of National Park Service paleontological resources in cultural context: Part 1, general overview. New Mexico Museum of Natural History and Science Bulletin 34:70–76.

Kenworthy, J. P., V. L. Santucci, and K. L. Cole. 2004. An inventory of paleontological resources associated with caves in Grand Canyon National Park. Pages 211–228 *in* C. van Riper, III and K. L. Cole, editors. The Colorado Plateau: Cultural, biological, and physical research. University of Arizona Press, Tucson, Arizona.

Mead, J. I. 1981. The last 30,000 years of faunal history within the Grand Canyon, Arizona. Quaternary Research 15(3):311–326.

Mead, J. I., and A. M. Phillips, III. 1982. The late Pleistocene and Holocene fauna and flora of Vulture Cave, Grand Canyon, Arizona. The Southwestern Naturalist 26(3):257–288.

Mead, J. I., and T. R. Van Devender. 1981. Late Holocene diet of *Bassariscus astutus* in the Grand Canyon, Arizona. Journal of Mammalogy 62(2):439–442.

Mead, J. I., L. L. Coats, and B. W. Schubert. 2003. Late Pleistocene cave faunas in the eastern Grand Canyon, Arizona. Pages 64–86 *in* B. W. Schubert, J. I. Mead, and R. W. Graham, editors. Ice Age cave faunas of North America. Indiana University Press, Bloomington and Indianapolis, Indiana.

Metcalfe, J. Z. 2018. Pleistocene hairs: Microscopic examination prior to destructive analysis. Paleoamerica 4(1):16–30.

Miller, A. E., N. Steele, and B. W. Tobin. 2018. Vulnerability and fragility risk indices for non-renewable resources. Journal of Environmental Monitoring and Assessment 190:article 373.

National Park Service. 2017. Foundation Document: Grand Canyon National Park, Arizona. National Park Service, Denver, Colorado.

Santucci, V. L., and A. L. Koch. 2003. Paleontological resource monitoring strategies for the National Park Service. Park Science 22(1):22–25.

Santucci, V. L., J. Kenworthy, and R. Kerbo. 2001. An inventory of paleontological resources associated with National Park Service caves. NPS Geological Resources Division, Denver, Colorado. Technical Report NPS/NRGRD/GRDTR-01/02.

Santucci, V. L., J. P. Kenworthy, and A. L. Mims. 2009. Monitoring in situ paleontological resources. Pages 189–204 *in* R. Young and L. Norby, editors. Geological monitoring. Geological Society of America, Boulder, Colorado.

Santucci, V. L., J. M. Ghist, and R. B. Blodgett. 2014. Inventory of U.S. Geological Survey paleontology collections to identify fossil localities in National Park Service areas. Proceedings of the 10th Conference on Fossil Resources. Dakoterra 6:215–218.

Santucci, V. L., J. S. Tweet, and T. Connors. 2018. The Paleontology Synthesis Project and establishing a framework for managing National Park Service paleontological resource archives and data. New Mexico Museum of Natural History and Science Bulletin 79:589–601.

Stortz, S. D., C. E. Aslan, T. D. Sisk, T. A. Chaudhry, J. M. Rundall, J. Palumbo, L. Zachmann, and B. Dickson. 2018. Natural resource condition assessment: Greater Grand Canyon landscape assessment. Natural Resource Report NPS/GRCA/NRR—2018/1645. National Park Service, Fort Collins, Colorado.

Tweet, J. S., V. L. Santucci, J. P. Kenworthy, and A. L. Mims. 2009. Paleontological resource inventory and monitoring—Southern Colorado Plateau Network. Natural Resource Technical Report NPS/NRPC/NRTR—2009/245. National Park Service, Fort Collins, Colorado.

Tweet, J. S., V. L. Santucci, and H. G. McDonald. 2016. Name-bearing fossil type specimens and species named from National Park Service areas. New Mexico Museum of Natural History and Science Bulletin 74:277–288.

Appendix 12-A. GRCA Paleontology Research Permits 1999–2019

Appendix Table 12-A-1. List of paleontological research permits issued from 1999 to 2019.

Permit Number	Year	Investigator(s)	Study Title	IAR/Report Published
GRCA-1999-SCI-0001	1999–2004	Jim Mead, Mary Carpenter	Late Pleistocene Vertebrate Communities in the Lower Grand Canyon, Arizona: Rampart and Muav Caves	IAR: 2001, 2004 Publications: Carpenter, M. C. 2003. Late Pleistocene Aves, Chiroptera, Perissodactyla, and Artiodactyla from Rampart Cave, Grand Canyon, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona. Carpenter, M. C., and J. I. Mead. 1999. Late Pleistocene vertebrate communities of the lower Grand Canyon: Rampart and Muav caves. Journal of Vertebrate Paleontology 19(supplement to 3):36A.
GRCA-2001-SCI-0002	2001–2002	Steven Austin	Sedimentary Model for Canyon-Length Mass Kill and Burial of Large Orthocone Nautiloids, Redwall Limestone (Lower Mississippian), Grand Canyon, Arizona: A Proposal for Research	IAR: 2001
GRCA-2003-SCI-0085	2003–2005	Judy Hellmich	Research and Education in Grand Canyon National Park	IAR: 2004, 2005
GRCA-2009-SCI-0005	2009	Karl Karlstrom	The Trail of Time at Grand Canyon: synthesis of Proterozoic research and studies of Quaternary geology of Grand Canyon	IAR: 2009
GRCA-2009-SCI-0032	2009	John Foster	Trilobite Taphonomy of a Quarry in the Middle Cambrian Bright Angel Shale, Grand Canyon National Park	IAR: 2009 Foster, J. R. 2011. Trilobites and other fauna from two quarries in the Bright Angel Shale (Middle Cambrian, Series 3; Delamaran), Grand Canyon National Park, Arizona. Pages 99–120 *in* J. S. Hollingsworth, F. A. Sundberg, and J. R. Foster, editors. Cambrian stratigraphy and paleontology of northern Arizona and southern Nevada. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 67.

Appendix Table 12-A-1 (continued). List of paleontological research permits issued from 1999 to 2019.

Permit Number	Year	Investigator(s)	Study Title	IAR/Report Published
GRCA-2010-SCI-0001	2010–2011	Paul Strother, John H. Beck, Eben Rose	Investigations of the Earliest Land Plants: The Collection of Microfossils from the Bright Angel Shale at Red Canyon	IAR: 2010, 2011
GRCA-2010-SCI-0017	2010–2011	Karl Karlstrom, Owen Shufeldt, Jake McDermott, Ryan Crow, Laura Crossey	The Trail of Time at Grand Canyon: synthesis of Proterozoic research and studies of Quaternary geology of Grand Canyon	IAR: 2010, 2011
GRCA-2010-SCI-0027	2010	Jill Allen	Grand Canyon Rocks	IAR: 2010
GRCA-2010-SCI-0039	2010–2012	Leonard Brand, John Whitmore	Coconino Sandstone fossil trackways: relationship to sedimentological features	IAR: 2010, 2011, 2012
GRCA-2011-SCI-0034	2011–2013	Deanna Greco, Erica Clites, James Super, Jeff Dobbins	Paleontological Inventory of Grand Canyon National Park	IAR: 2011
GRCA-2012-SCI-0021	2012–2013	Steve Dworkin, Stacy Atchley, Lee Nordt	Reconstructing Paleozoic Climates Using Paleosols in the Grand Canyon	IAR: 2012
GRCA-2013-SCI-0001	2013–2014	Jessica Metcalfe, Michael Richards	Late Pleistocene Paleoecology of the Colorado Plateau	IAR: 2013, 2014
GRCA-2013-SCI-0052	2013–2014	Jeffrey Martin, Jim Mead	Holocene and late-Pleistocene Bison of the Grand Canyon and the Colorado Plateau	IAR: 2013, 2014
GRCA-2015-SCI-0011	2015–2016	Jessica Metcalfe, Michael Richards	Late Pleistocene Paleoecology of the Colorado Plateau	IAR: 2015, 2016 Publication: Metcalfe, J. Z. 2018. Pleistocene hairs: microscopic examination prior to destructive analysis. PaleoAmerica, DOI: 10.1080/20555563.2017.1413529
GRCA-2017-SCI-0054	2017–2018	Steve Rowland	Fossil Trackways in the Supai Formation in GRCA	IAR: 2017, 2018
GRCA-2017-SCI-0059	2018–2019	Anne Miller, David Elliott	Ichnology of the Bright Angel Shale Formation, Grand Canyon, AZ: Indicators for Middle Cambrian Paleoecology	IAR: 2018 Ongoing Masters thesis work
GRCA-2018-SCI-0030	2018	Chris Doughty	A metagenomics study of extinct sloth dung	IAR: 2018
GRCA-2019-SCI-0002	2019	Vincent Santucci, Anne Miller, Justin Tweet, Mark Nebel	Grand Canyon National Park Paleontological Resource Inventory and Paleoblitz	In Progress

Appendix 12-B. GRCA Photogrammetry 3D Models

3D photogrammetry models can be viewed on the National Park Service Geologic Resource Division Sketchfab page (https://sketchfab.com/grd_nps/models) or GRCA photogrammetry series website (https://www.nps.gov/articles/series.htm?id=A9E62040-AC6F-A6D7-BE564A036F1D6146).

All photos and files associated with each 3D model have been archived on IRMA and within the GRCA and WASO paleontology archives.

Appendix Table 12-B-1. GRCA specimens which have been documented as 3D photogrammetry models.

Specimen ID	Scientific Name	Common Name
NMNH 25478	*Desmodus stocki*	Stock's Vampire Bat Skull
GRCA 56441	*Oreamnos harringtoni*	Harrington's Goat Skull
GRCA 55908	*Gymnogyps californianus*	California Condor Beak
GRCA 21734	*Miracinonyx trumani*	American Cheetah Maxilla
GRCA 21751	*Nothrotheriops shastensis*	Shasta Ground Sloth Skull
USNM PAL 617525	*Nothrotheriops shastensis*	Shasta Ground Sloth Coprolite
GRCA 20214	*Tainoceras schellbachi*	Coiled Nautiloid Holotype
GRCA 21372	*Productus bassi*	Brachiopod
GRCA 14466	Undetermined	Crinoid Columnal Block
GRCA 33184	*Platysomus*	Phyllodont Fish Tooth Plate
N/A	*Chelichnus*	Vertebrate Trackway Block (Field)
N/A	*Ichniotherium*	Vertebrate Trackway Block (Field)
GRCA 3217	Undetermined	Permian Seed Fern
USNM PAL 38033	*Supaia merriami*	Permian Seed Fern Holotype
GRCA 3090	*Typus whitei*	Griffinfly Wing Holotype
GRCA 17187	*Dolichometopus productus*	Trilobite

Appendix A. Fossil Taxa Named From GRCA

Befitting a park with such a long history of paleontological investigations and diversity of fossiliferous rocks, GRCA has been the source for numerous type specimens. At least 167 fossil species have been named from fossils discovered within 2020 GRCA boundaries (Appendix Table A-1), a number which is exceeded only by Florissant Fossil Beds National Monument, Guadalupe Mountains National Park, John Day Fossil Beds National Monument, and Yellowstone National Park. They are also among the most diverse groups of species named from an NPS unit. An additional 13 potentially based on GRCA fossils (see the following Appendix B). The great majority of these taxa were collected and named during the National Park era (1919–present), although not all of the discovery localities were within GRCA at the time of discovery due to park boundary changes. For example, some type specimens recovered from what is now far western GRCA during the 1930s were then within Lake Mead National Recreation Area (LAKE). Pre-1919 taxa are primarily Walcott's Cambrian invertebrates and a few species from the very early era of exploration. Some land once in southern GRCA was transferred to the Havasupai Indian Reservation in 1975; some fossil sites given in the literature as within GRCA are affected by this, but apparently not any type specimens. Two bibliographies by Spamer (1984, 1988) were invaluable for creating this appendix and Appendix B.

Although GRCA is justly famous for its Paleozoic plants and vertebrate ichnofossils, its type record is dominated by invertebrates, particularly trilobites, brachiopods, and ostracode-like crustaceans. Taxa from nearby but not within GRCA were omitted. Three names that would otherwise fall within this table are omitted: the pseudofossil or dubiofossil *Brooksella canyonensis* (Bassler 1941; see Chapter 4); the controversial "algae" *Rivularites permiensis* (White 1929); and the *nomen nudum* or "naked name" *Agostopus robustus*, accidentally created by Gilmore (1927) as he recognized a year later (Gilmore 1928). The table is divided into plants, invertebrates, vertebrates, invertebrate ichnofossils, vertebrate ichnofossils, other ichnofossils, and other fossils. Within each group, sorting is by general type of fossil, and then alphabetically by taxa. No evaluation of validity or preferred taxonomy is intended; only corrections of preoccupied names are included.

Institutional Abbreviations—**ANSP**, Academy of Natural Sciences of Drexel University, Philadelphia, Pennsylvania, USA; **FMNH**, Field Museum of Natural History, Chicago, Illinois, USA; **GCNPM**, Grand Canyon National Park Museum, Grand Canyon, Arizona, USA; **HUPC**, Paleobotanical Collections of Harvard University Herbaria, Cambridge, Massachusetts, USA; **LACM**, Natural History Museum of Los Angeles County, Los Angeles, California, USA; **LO**, Lund University (Lunds universitet), Lund, Sweden; **MNA**, Museum of Northern Arizona, Flagstaff, Arizona, USA; **NGS**, Nebraska Geological Survey (now at UNSM); **UCLA**, University of California Los Angeles (now at LACM); **UCMP**, University of California Museum of Paleontology, Berkeley, California, USA; **UNSM**, University of Nebraska State Museum, Lincoln, Nebraska, USA; **USNM**, National Museum of Natural History, Washington, D.C., USA; **USNMPC**, U.S. National Museum Parasite Collection, Beltsville, Maryland, USA; **UT**, Eberhard Karls University, Tübingen (Eberhards Karls Universität Tübingen), Tübingen, Germany; **YPM**, Peabody Museum of Natural History, New Haven, Connecticut, USA.

Appendix Table A-1. Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Plants	*Brachyphyllum arizonicum*	White 1929	Permian, Hermit	USNM 38060	Conifer
	Brachyphyllum tenue	White 1929	Permian, Hermit	Syntypes USNM 38061, 312783, and 312784	Conifer
	Pagiophyllum dubium	White 1929	Permian, Hermit	Syntypes USNM 38062 and 312790–312792	Conifer
	Palaeotaxites praecursor	White 1929	Permian, Hermit	Syntypes USNM 38058, 38060, and 324560–324564	Conifer
	Voltzia dentiloba	White 1929	Permian, Hermit	Syntype USNM 38056	Conifer
	Walchia dawsoni	White 1929	Permian, Hermit	Syntypes USNM 38052 and 312796–312798	Conifer
	Walchia gracillima	White 1929	Permian, Hermit	Syntypes USNM 38053 and 312793–312795	Conifer
	Brongniartites? aliena	White 1929	Permian, Hermit	Syntypes USNM 38043 and 312785–312789	Pteridophyte
	Brongniartites? yakiensis	White 1929	Permian, Hermit	Syntypes USNM 38042, 312807–312812 and 324554–324559	Pteridophyte
	Sphenophyllum gilmorei	White 1929	Permian, Hermit	Syntypes USNM 38025 and 324567–324575	Pteridophyte
	Supaia anomala	White 1929	Permian, Hermit	Syntypes USNM 38035 and 324581–324588	Pteridophyte
	Supaia breviloba	White 1929	Permian, Hermit	Syntype USNM 38037	Pteridophyte
	Supaia compacta	White 1929	Permian, Hermit	Syntypes USNM 38034 and 324590–324593	Pteridophyte
	Supaia linearifolia	White 1929	Permian, Hermit	Syntypes USNM 38036 and 324594	Pteridophyte
	Supaia merriami	White 1929	Permian, Hermit	Syntype USNM 38033	Pteridophyte
	Supaia rigida	White 1929	Permian, Hermit	Syntypes USNM 38031, 324595, and 324596	Pteridophyte

Appendix Table A-1 (continued). Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Plants (continued)	*Supaia sturdevantii*	White 1929	Permian, Hermit	Syntypes USNM 38032 and 324597	Pteridophyte
	Supaia subgoepperti	White 1929	Permian, Hermit	USNM 38038	Pteridophyte
	Supaia thinnfeldioides	White 1929	Permian, Hermit	Syntypes USNM 38030 and 324598–324603	Pteridophyte
	Yakia heterophylla	White 1929	Permian, Hermit	Syntypes USNM 38044 and 312799–312804	Pteridophyte
	Taeniopteris angelica	White 1929	Permian, Hermit	Syntypes USNM 38048, 324565, 324566, and 342587	Pteridophyte or gingko
	Callipteris arizonae	White 1929	Permian, Hermit	Syntype USNM 38027	Pteridophyte?
	Cyclocarpon angelicum	White 1929	Permian, Hermit	Syntype USNM 38063	Seed/fruit
	Eltovaria bursiformis	White 1929	Permian, Hermit	Syntype USNM 38066	Seed/fruit
Invertebrates	*Avonia subhorrida newberryi*	McKee 1938	Permian, Kaibab	USNM 102301	Brachiopod
	Billingsella obscura	Walcott 1905	Cambrian, Bright Angel	USNM 52258a	Brachiopod
	Chonetes quadratus	King 1931	Permian, Kaibab	YPM 10830a	Brachiopod
	Composita arizonica	McKee 1938	Permian, Kaibab	Syntypes USNM 102303, 102304, and 102305	Brachiopod
	Derbyia regularis	McKee 1938	Permian, Kaibab	Syntypes USNM 102290 and 102291	Brachiopod; preoccupied, renamed *Derbyia arizonensis* (McKee 1941)
	Dictyonina arizonaensis	Resser 1945	Cambrian, Bright Angel?	USNM 108557	Brachiopod
	Finkelnburgia noblei	Walcott 1924	Cambrian, Muav	USNM 69750–69752	Brachiopod
	Iphidea crenistria	Walcott 1897	Cambrian, Bright Angel	USNM 26431a	Brachiopod

Appendix Table A-1 (continued). Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Invertebrates (continued)	*Iphidea superba*	Walcott 1897	Cambrian, Bright Angel?	USNM 26429a	Brachiopod
	Lingulella kanabensis	Resser 1945	Cambrian, Bright Angel / Muav	USNM 33829a	Brachiopod
	Lingulella mckeei	Resser 1945	Cambrian, Bright Angel	USNM 108561a	Brachiopod
	Nisusia (*Jamesella*) *kanabensis*	Walcott 1908	Cambrian, Bright Angel / Muav	USNM 52300	Brachiopod
	Obolus (*Lingulella*) *chuarensis*	Walcott 1898	Cambrian, Bright Angel?	USNM 57020a	Brachiopod
	Obolus (*Lingulella*) *euglyphus*	Walcott 1898	Cambrian, Bright Angel	USNM 27316a	Brachiopod
	Obolus (*Lingulella*) *lineolatus*	Walcott 1898	Cambrian, Bright Angel?	USNM 27325a	Brachiopod
	Obolus (*Lingulella*) *spatulus*	Walcott 1902	Cambrian, Bright Angel	USNM 35290a	Brachiopod
	Obolus (*Lingulella*) *zetus*	Walcott 1898	Cambrian, Bright Angel	USNM 27347b	Brachiopod
	Obolus (*Westonia*) *themis*	Walcott 1905	Cambrian, Bright Angel	USNM 51732a	Brachiopod
	Bascomella subsphaerica	Condra and Elias 1944	Permian, Kaibab	NGS 449 (not found on May 2016 visit to UNSM)	Bryozoan
	Girtypora maculata	McKinney 1983	Permian, Kaibab	FMNH PE 24301	Bryozoan
	Bicorbula arizonica	Condra and Elias 1945a	Permian, Kaibab	NGS 264 (not found on May 2016 visit to UNSM)	Bryozoan or bryozoan-algal consortium; preoccupied, renamed *Bicorbis arizonica* (Condra and Elias 1945b)

Appendix Table A-1 (continued). Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Invertebrates (continued)	*Conularia kaibabensis*	McKee 1935	Permian, Kaibab	USNM 102289 (originally GCNPM FK211)	Conulariid
	Eocrinus multibrachiatus	Kirk (in Resser 1945)	Cambrian, Bright Angel	USNM 108556a	Eocrinoid
	Oreohelix yavapai fortis	Cockerell 1927	Pleistocene?, “red earth”	ANSP 141875	Gastropod
	Typus gilmorei	Carpenter 1927	Permian, Hermit	USNM 71279	Insect (meganeurid wing); generic name now corrected to *Tupus* (ICZN 1985)
	Typus whitei	Carpenter 1928	Permian, Hermit	USNM 71713	Insect (meganeurid wing); generic name now corrected to *Tupus* (ICZN 1985)
	Agamofilaria oxyura	Schmidt et al. 1992	Pleistocene	USNMPC 82076	Nematode
	Strongyloides shastensis	Schmidt et al. 1992	Pleistocene	USNMPC 82075	Nematode
	Bradoria tontoensis	Ulrich and Bassler 1931	Cambrian, unspecified	USNM 81377	Ostracode-like crustacean
	Dielymella appressa	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56506	Ostracode-like crustacean
	Dielymella dorsalis	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56505	Ostracode-like crustacean
	Dielymella nasuta	Ulrich and Bassler 1931	Cambrian, Bright Angel	Cotypes USNM 56508 and 56509	Ostracode-like crustacean
	Dielymella recticardinalis	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56510	Ostracode-like crustacean
	Dielymella recticardinalis angustata	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56511	Ostracode-like crustacean
	Indiana curta	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56466	Ostracode-like crustacean

Appendix Table A-1 (continued). Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Invertebrates (continued)	*Indiana faba*	Ulrich and Bassler 1931	Cambrian, Bright Angel?	USNM 56458	Ostracode-like crustacean
	Indiana faba intermedia	Ulrich and Bassler 1931	Cambrian, Bright Angel?	USNM 56462	Ostracode-like crustacean
	Indiana impressa	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56463	Ostracode-like crustacean
	Tontoia kwaguntensis	Walcott 1912	Cambrian, Bright Angel?	USNM 57660	Ostracode-like crustacean
	Walcottella apicalis	Ulrich and Bassler 1931	Cambrian, Bright Angel?	Cotypes USNM 56477 and 56478	Ostracode-like crustacean
	Walcottella breviuscula	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56481	Ostracode-like crustacean
	Walcottella concentrica	Ulrich and Bassler 1931	Cambrian, Bright Angel?	Cotypes USNM 56479 and 56480	Ostracode-like crustacean
	Walcottella leperditoides	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56484	Ostracode-like crustacean
	Walcottella limatula	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56488	Ostracode-like crustacean
	Walcottella longula	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56491	Ostracode-like crustacean
	Walcottella nitida	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56485	Ostracode-like crustacean
	Walcottella oblonga	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56486	Ostracode-like crustacean
	Walcottella obsoleta	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56487	Ostracode-like crustacean
	Walcottella pulchella	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56483	Ostracode-like crustacean

Appendix Table A-1 (continued). Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Invertebrates (continued)	*Walcottella scitula*	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56482	Ostracode-like crustacean
	Walcottella subtruncata	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56490	Ostracode-like crustacean
	Walcottella ventrosa	Ulrich and Bassler 1931	Cambrian, Bright Angel	USNM 56489	Ostracode-like crustacean
	Scenella hermitensis	Resser 1945	Cambrian, Muav	USNM 108568a	Scenellid (described as a gastropod)
	Acrocephalops? arizonaensis	Resser 1945	Cambrian, Bright Angel	USNM 108624	Trilobite
	Albertella schenki	Resser 1945	Cambrian, Bright Angel	USNM 108583	Trilobite
	Alokistocare althea	Walcott 1916a	Cambrian, Bright Angel	Lectotype USNM 61574	Trilobite
	Bolaspis aemula	Resser 1945	Cambrian, Muav	USNM 108602a	Trilobite
	Clavaspidella kanabensis	Resser 1945	Cambrian, Bright Angel / Muav	USNM 108578a	Trilobite
	Delaria macclintocki	Cisne 1971	Permian, Kaibab	GCNPM 3949	Trilobite
	Dolichometopus tontoensis	Walcott 1916b	Cambrian, Bright Angel	Lectotype USNM 62685	Trilobite
	Ehmaniella arizonaensis	Resser 1945	Cambrian, Bright Angel	USNM 108603a	Trilobite
	Ehmaniella hebes	Resser 1945	Cambrian, Bright Angel or Muav	USNM 108612a	Trilobite
	Elrathia nitens	Resser 1945	Cambrian, Bright Angel	USNM 108625	Trilobite
	Elrathiella? insueta	Resser 1945	Cambrian, Bright Angel or Muav	USNM 108621a	Trilobite

Appendix Table A-1 (continued). Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Invertebrates (continued)	*Glossopleura mckeei*	Resser 1935	Cambrian, Bright Angel	USNM 62714	Trilobite
	Glyphaspis tecta	Resser 1945	Cambrian, Muav	USNM 108596a	Trilobite
	Glyphaspis vulsa	Resser 1945	Cambrian, Bright Angel	USNM 108595a	Trilobite
	Glyphaspsis kwanguntensis	Resser 1945	Cambrian, Bright Angel or Muav	USNM 108618a	Trilobite
	Kochina? angustata	Resser 1945	Cambrian, Bright Angel / Muav	USNM 108610	Trilobite
	Kootenia mckeei	Resser 1945	Cambrian, Bright Angel / Muav	USNM 108588a	Trilobite
	Kootenia schenki	Resser 1945	Cambrian, Muav	USNM 108586a	Trilobite
	Kootenia simplex	Resser 1945	Cambrian, Bright Angel	USNM 108591a	Trilobite
	Pachyaspis fonticola	Resser 1945	Cambrian, Bright Angel	USNM 108608	Trilobite
	Parehmania kwaguntensis	Resser 1945	Cambrian, Bright Angel or Muav	USNM 108620a	Trilobite
	Parehmania nitida	Resser 1945	Cambrian, Bright Angel or Muav	USNM 108613a	Trilobite
	Parehmania tontoensis	Resser 1945	Cambrian, Bright Angel	USNM 108614	Trilobite
	Solenopleurella diligens	Resser 1945	Cambrian, Bright Angel / Muav	USNM 108627a	Trilobite
	Solenopleurella erosa	Resser 1945	Cambrian, Muav	USNM 108616a	Trilobite
	Solenopleurella porcata	Resser 1945	Cambrian, Muav	USNM 108626a	Trilobite
	Spencia tontoensis	Resser 1945	Cambrian, Bright Angel	USNM 108611a	Trilobite

Appendix Table A-1 (continued). Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Invertebrates (continued)	*Trachycheilus typicale*	Resser 1945	Cambrian, Bright Angel or Muav	USNM 108619	Trilobite
Vertebrates	*Megactenopetalus kaibabanus*	David 1944	Permian, Kaibab	MNA G2.2280	Chondrichthyan
Invertebrate Ichnofossils	*Scoyenia gracilis*	White 1929	Permian, Hermit	USNM 201869	Invertebrate trail
	Walpia hermitensis	White 1929	Permian, Hermit	Syntypes USNM 263675 and 263676	Invertebrate trail
	Unisulcus sinuosus	Gilmore 1927	Permian, Coconino	USNM 11498	Invertebrate trail
	Octopodichnus didactylus	Gilmore 1927	Permian, Coconino	USNM 11501	Invertebrate track
	Paleohelcura tridactyla	Gilmore 1926	Permian, Coconino	USNM 11145	Invertebrate track
	Stipsellus annulatus	Howell 1957	Cambrian, Tapeats	GCNPM 8538	Invertebrate trail
	Cruziana arizonensis	Seilacher 1970	Cambrian, Tapeats	UT lc 1329/9	Invertebrate trace
	Mesichnium benjamini	Gilmore 1926	Permian, Coconino	USNM 11155	Invertebrate track
	Triavestigia niningeri	Gilmore 1927	Permian, Coconino	USNM 11510	Invertebrate track
	Angulichnus alternipes	Elliott and Martin 1987	Cambrian, Bright Angel	MNA N3862	Invertebrate trail
Vertebrate Ichnofossils	*Castrocopros martini*	Hunt and Lucas 2018	Pleistocene	USNM PAL 720155	*Nothrotheriops* coprolite
	Agostopus matheri	Gilmore 1926	Permian, Coconino	USNM 11135	Vertebrate track
	Agostopus medius	Gilmore 1927	Permian, Coconino	USNM 11509	Vertebrate track
	Allopus? *arizonae*	Gilmore 1926	Permian, Coconino	USNM 11132	Vertebrate track
	Amblyopus pachypodus	Gilmore 1927	Permian, Coconino	USNM 11511	Vertebrate track
	Ammobatrachus turbatans	Gilmore 1928	Pennsylvanian, Wescogame	USNM 11691	Vertebrate track
	Anomalopus sturdevanti	Gilmore 1927	Pennsylvanian, Wescogame	USNM 11577	Vertebrate track
	Baropezia eakini	Gilmore 1926	Permian, Coconino	USNM 11137	Vertebrate track
	Baropus coconinoensis	Gilmore 1927	Permian, Coconino	USNM 11514	Vertebrate track

Appendix Table A-1 (continued). Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Vertebrate Ichnofossils (continued)	*Barypodus metszeri*	Gilmore 1927	Permian, Coconino	USNM 11505	Vertebrate track
	Barypodus palmatus	Gilmore 1926	Permian, Coconino	USNM 11134	Vertebrate track
	Barypodus tridactylus	Gilmore 1927	Permian, Coconino	USNM 11502	Vertebrate track
	Batrachichnus obscurus	Gilmore 1927	Permian, Hermit	USNM 11529	Vertebrate track
	Collettosaurus pentadactylus	Gilmore 1927	Permian, Hermit	USNM 11527	Vertebrate track
	Dolichopodus tetradactylus	Gilmore 1926	Permian, Coconino	USNM 11123	Vertebrate track
	Dromillopus parvus	Gilmore 1927	Permian, Hermit	USNM 11537	Vertebrate track
	Exocampe? delicatula	Lull 1918	Permian, Hermit	YPM 2146	Vertebrate track
	Hyloidichnus bifurcatus	Gilmore 1927	Permian, Hermit	USNM 11518	Vertebrate track
	Hyloidichnus whitei	Gilmore 1928	Permian, Hermit	USNM 11692	Vertebrate track
	Hylopus hermitanus	Gilmore 1927	Permian, Hermit	USNM 11517	Vertebrate track
	Ichniotherium gilmorei	Haubold 1971	Permian, Hermit	USNM 11707 (now lost)	Vertebrate track
	Laoporus noblei	Lull 1918	Permian, Coconino	YPM 2144	Vertebrate track
	Laoporus schucherti	Lull 1918	Permian, Coconino	YPM 2143	Vertebrate track
	Megapezia? coloradensis	Lull 1918	Permian, Hermit	YPM 2145	Vertebrate track
	Nanopus maximus	Gilmore 1927	Permian, Coconino	USNM 11506	Vertebrate track
	Nanopus merriami	Gilmore 1926	Permian, Coconino	USNM 11146	Vertebrate track
	Palaeopus regularis	Gilmore 1926	Permian, Coconino	USNM 11143	Vertebrate track
	Stenichnus yakiensis	Gilmore 1927	Pennsylvanian, Wescogame	USNM 11533	Vertebrate track
	Tridentichnus supaiensis	Gilmore 1927	Pennsylvanian, Wescogame	USNM 11534	Vertebrate track
Other Ichnofossils	*Cryptozoon? Occidentale*	Dawson 1897	Neoproterozoic, Kwagunt	USNM 33799 (USNM 60710 and 60711 per Rezak 1957)	Stromatolite
Other Fossils	*Archeococcidia antiquus*	Schmidt et al. 1992	Pleistocene	USNMPC 82073	Coccidian oocyst

Appendix Table A-1 (continued). Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Other Fossils (continued)	*Archeococcidia nothrotheriopsiae*	Schmidt et al. 1992	Pleistocene	USNMPC 82074	Coccidian oocyst
	Endothyra baileyi poloumera	Skipp 1969	Mississippian, Redwall	USNM 641727	Foraminifera
	Septaglomospiranella rossi	Skipp 1969	Mississippian, Redwall	USNM 641578	Foraminifera
	Chuaria circularis	Walcott 1899	Neoproterozoic, Kwagunt	Lectotype USNM 33800	Organic-walled microfossil
	Kaibabia gemmulella	Porter and Riedman 2016	Neoproterozoic, Galeros	UCMP 36082a	Organic-walled microfossil
	Microlepidopalla mira	Porter and Riedman 2016	Neoproterozoic, Galeros	UCMP 36104b	Organic-walled microfossil
	Vandalosphaeridium walcottii	Vidal and Ford 1985	Neoproterozoic, Kwagunt	LO 5661	Organic-walled microfossil
	Volleyballia dehlerae	Porter and Riedman 2016	Neoproterozoic, Galeros	UCMP 36080d	Organic-walled microfossil
	Bombycion micron	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 62988	Vase-shaped microfossil
	Bonniea dacruchares	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 64409	Vase-shaped microfossil
	Bonniea pytinaia	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 64410	Vase-shaped microfossil
	Cycliocyrillium simplex	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 64455	Vase-shaped microfossil
	Cycliocyrillium torquata	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 64453	Vase-shaped microfossil
	Hemisphaeriella ornata	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 62990	Vase-shaped microfossil

Appendix Table A-1 (continued). Fossil taxa named from specimens found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Classification
Other Fossils (continued)	*Melanocyrillium fimbriatum*	Bloeser 1985	Neoproterozoic, Kwagunt	UCLA 58968 (now LACM?)	Vase-shaped microfossil
	Melanocyrillium hexodiadema	Bloeser 1985	Neoproterozoic, Kwagunt	UCLA 58959 (now LACM?)	Vase-shaped microfossil
	Melanocyrillium horodyskii	Bloeser 1985	Neoproterozoic, Kwagunt	UCLA 58976 (now LACM?)	Vase-shaped microfossil
	Melicerion poikilon	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 62990	Vase-shaped microfossil
	Palaeoarcella athanata	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 62988	Vase-shaped microfossil
	Trachycyrillium pudens	Porter et al. 2003	Neoproterozoic, Kwagunt	HUPC 64413	Vase-shaped microfossil

Literature Cited

Bassler, R. S. 1941. A supposed jellyfish from the pre-Cambrian of the Grand Canyon. Proceedings of the United States National Museum 89:519–522.

Bloeser, B. 1985. *Melanocyrillium*, a new genus of structurally complex late Proterozoic microfossils from the Kwagunt Formation (Chuar Group), Grand Canyon, Arizona. Journal of Paleontology 59(3):741–765.

Carpenter, F. M. 1927. A fossil insect from the lower Permian of the Grand Canyon. Proceedings of the United States National Museum 71(23).

Carpenter, F. M. 1928. A new protodonatan from the Grand Canyon. Psyche 35:186–189.

Cisne, J. L. 1971. Paleoecology of trilobites of the Kaibab limestone (Permian) in Arizona, Utah, and Nevada. Journal of Paleontology 45(3):525–533.

Cockerell, T. D. A. 1927. A large form of *Oreohelix yavapai* in the Grand Canyon. The Nautilus 40(3):101.

Condra, G. E. and M. K. Elias. 1944. Carboniferous and Permian ctenostomatous Bryozoa. Geological Society of America Bulletin 55:517–568.

Condra, G. E. and M. K. Elias. 1945a. *Bicorbula*, a new Permian bryozoan, probably a bryozoan-algal consortium. Journal of Paleontology 19(2):116–125.

Condra, G. E. and M. K. Elias. 1945b. *Bicorbis arizonica* Condra and Elias, new name for *B. arizonica*. Journal of Paleontology 19:411.

David, L. R. 1944. A Permian shark from the Grand Canyon. Journal of Paleontology 18(1):90–93.

Dawson, J. W. 1897. Note on *Cryptozoon* and other ancient fossils. Canadian Record of Science 7:203–219.

Elliott, D. K. and D. L. Martin. 1987. A new trace fossil from the Cambrian Bright Angel Shale, Grand Canyon, Arizona. Journal of Paleontology 61(4):641–648.

Gilmore, C. W. 1926. Fossil footprints from the Grand Canyon. Smithsonian Miscellaneous Collections 77(9).

Gilmore, C. W. 1927. Fossil footprints from the Grand Canyon: second contribution. Smithsonian Miscellaneous Collections 80(3).

Gilmore, C. W. 1928. Fossil footprints from the Grand Canyon: third contribution. Smithsonian Miscellaneous Collections 80(8).

Haubold, H. 1971. Ichnia amphibiorum et reptiliorum fossilium. Handbuch der Paleoherpetologie, Teil 18. Stuttgart, Gustav Fischer Verlag.

Howell, B. F. 1957. *Stipsellus annulatus*, a *Skolithos*-like Cambrian fossil from Arizona. Bulletin of the Wagner Free Institute of Sciences of Philadelphia 32(2):17–20.

Hunt, A. P., and S. G. Lucas. 2018. The record of sloth coprolites in North and South America: implications for terminal Pleistocene extinctions. New Mexico Museum of Natural History and Science Bulletin 79:277–298.

International Commission on Zoological Nomenclature (ICZN). 1985. Opinion 1317 *Tupus* Sellards, 1906 (Insecta, Protodonata): Conserved. Bulletin of Zoological Nomenclature 42(2):169–170.

Jackson, R. T. 1912. Phylogeny of the Echini, with a revision of Palaeozoic species. Memoirs of the Boston Society of Natural History 7.

King, R. E. 1931. The geology of the Glass Mountains. Part II, Faunal summary and correlation of the Permian formations with description of Brachiopoda. Texas University Bulletin 3042.

Lull, R. S. 1918. Fossil footprints from the Grand Canyon of the Colorado. American Journal of Science (4th series) 45:337–346.

McKee, E. D. 1935. A *Conularia* from the Permian of Arizona. Journal of Paleontology 9:427–429.

McKee, E. D. 1938. The environment and history of the Toroweap and Kaibab formations of northern Arizona and southern Utah. Carnegie Institution of Washington Publication 492.

McKee, E. D. 1941. *Derbya arizonensis*, new name for *D. regularis* McKee. Journal of Paleontology 15(1):91.

McKinney, F. K. 1983. Ectoprocta (Bryozoa) from the Permian Kaibab Formation, Grand Canyon National Park, Arizona. Fieldiana: Geology (New Series) 13(1341).

Newberry, J. S. 1861. Geological report. *In* Ives, J. C., Report upon the Colorado River of the West, explored in 1857 and 1858. U.S. 36th Congress, 1st Session, House Executive Document 90, Serial 1058, Part 3.

Porter, S. M., and L. A. Riedman. 2016. Systematics of organic-walled microfossils from the ca. 780–740 Ma Chuar Group, Grand Canyon, Arizona. Journal of Paleontology 90(5):815–853.

Porter, S. M., R. Meisterfeld, and A. H. Knoll. 2003. Vase-shaped microfossils from the Neoproterozoic Chuar Group, Grand Canyon: a classification guided by modern testate amoebae. Journal of Paleontology 77(3):409–429.

Resser, C. E. 1935. Nomenclature of some Cambrian trilobites. Smithsonian Miscellaneous Collections 93(5):1–46.

Resser, C. E. 1945. Cambrian fossils of the Grand Canyon. Pages 169–220 *in* E. D. McKee and C. E. Resser. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

Rezak, R. 1957. Stromatolites of the Belt series in Glacier National Park and vicinity, Montana. U.S. Geological Survey, Washington, D.C. Professional Paper 294-D.

Schmidt, G. D., D. W. Duszynski, and P. S. Martin. 1992. Parasites of the extinct Shasta Ground Sloth *Nothrotheriops shastensis*, in Rampart Cave, Arizona. Journal of Parasitology 78(5):811–816.

Seilacher, A. 1970. *Cruziana* stratigraphy of non-fossiliferous Paleozoic sandstones. Pages 447–476 *in* T. P. Crimes and J. C. Harper, editors. Trace Fossils. Geological Journal Special Issue 3.

Skipp, B. 1969. Foraminifera. Pages 173–255 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Spamer, E. E. 1984. Geology of the Grand Canyon: An annotated bibliography, with an annotated catalogue of Grand Canyon type fossils. Volume 2. Supplement to the annotated bibliography (1857–1983), Supplement and revisions to the annotated catalogue. Geological Society of America, Boulder, Colorado. Microform Publication 14.

Spamer, E. E. 1988. Geology of the Grand Canyon: An annotated bibliography, with an annotated catalogue of Grand Canyon type fossils. Volume 3. Second supplement (to 1987) with an annotated bibliography of the world literature on the Grand Canyon type fossil *Chuaria circularis* Walcott, 1899, an index fossil for the late Proterozoic. Geological Society of America, Boulder, Colorado. Microform Publication 17.

Ulrich, E. O. and R. S. Bassler. 1931. Cambrian bivalved Crustacea of the order Conchostraca. U.S. National Museum Proceedings 78 (article 4).

Vidal, G. and T. D. Ford. 1985. Microbiotas from the late Proterozoic Chuar Group (northern Arizona) and Uinta Mountain Group (Utah) and their chronostratigraphic implications. Precambrian Research 28(3–4):349–389.

Walcott, C. D. 1897. Cambrian Brachiopoda; genera *Iphidea* and *Yorkia*; with descriptions of new species of each, and of the genus Acrothele. U.S. National Museum Proceedings 19:707–718.

Walcott, C. D. 1898. Cambrian Brachiopoda: *Obolus* and *Lingulella*, with description of new species. U.S. National Museum Proceedings 21:385–420.

Walcott, C. D. 1899. Pre-Cambrian fossiliferous formations. Geological Society of America Bulletin 10:199–244.

Walcott, C. D. 1902. Cambrian Brachiopoda: *Acrotreta*; *Linnarssonella*; *Obolus*; with descriptions of new species. U.S. National Museum Proceedings 25:577–612.

Walcott, C. D. 1905. Cambrian Brachiopoda; with descriptions of new genera and species. U.S. National Museum Proceedings 28:227–337.

Walcott, C. D. 1908. Cambrian geology and paleontology, No. 3: Cambrian Brachiopoda: Descriptions of new genera and species. Smithsonian Miscellaneous Collections 53:53–165.

Walcott, C. D. 1912. Cambrian geology and paleontology II. Middle Cambrian Branchiopoda, Malacostraca, Trilobita, and Merostomata. Smithsonian Miscellaneous Collections 57(6):145–228.

Walcott, C. D. 1916a. Cambrian geology and paleontology III. Cambrian trilobites. Smithsonian Miscellaneous Collections 64(3):157–258.

Walcott, C. D. 1916b. Cambrian geology and paleontology II. Cambrian trilobites. Smithsonian Miscellaneous Collections 64(5):303–570.

Walcott, C. D. 1924. Cambrian geology and paleontology IV. No. 9. Cambrian and Ozarkian Brachiopoda; Ozarkian Cephalopoda and Notostraca. Smithsonian Miscellaneous Collections 67(9):477–554.

White, C. D. 1929. Flora of the Hermit shale, Grand Canyon, Arizona. Carnegie Institution of Washington Publication 405.

Appendix B. Fossil Taxa Potentially Named From GRCA

A small number of taxa have been named from fossils discovered somewhere in the Grand Canyon area, but with insufficient provenance information to determine the exact location. Some, all, or none of the type specimens may have come from GRCA (Appendix Table B-1).

Institutional Abbreviations—**CU**, Columbia University, New York City, New York; **GCNPM**, Grand Canyon National Park Museum, Grand Canyon, Arizona; **USNM**, National Museum of Natural History, Washington, D.C.

Appendix Table B-1. Fossil taxa named from specimens possibly found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Notes
Invertebrates	*Avonia dorsoconcava*	McKee 1938	Permian, Kaibab	GCNPM 9993	Brachiopod
	Chonetes kaibabensis	McKee 1938	Permian, Kaibab	Syntypes USNM 102292, 102293, and 102294	Brachiopod
	Marginifera meridionalis	McKee 1938	Permian, Kaibab	Syntype USNM 102302	Brachiopod
	Productus (*Dictyoclostus*) *bassi*	McKee 1938	Permian, Kaibab	Lectotype USNM 102295a	Brachiopod
	Productus (*Dictyoclostus*) *paraindicus*	McKee 1938	Permian, Kaibab	Lectotype USNM 102297	Brachiopod
	Productus ivesi	Newberry 1861	Permian, Toroweap	Syntypes USNM 5356a–c	Brachiopod
	Archaeocidaris gracilis	Newberry 1861	Permian, Toroweap / Kaibab	Syntypes CU 604 and USNM 5412	Echinoid
	Archaeocidaris longispinus	Newberry 1861	Permian, Toroweap / Kaibab	CU 6419G	Echinoid
	Archaeocidaris ornatus	Newberry 1861	Permian, Toroweap / Kaibab	Syntypes CU 6000G	Echinoid; preoccupied, renamed *Archaeocidaris coloradensis* (Jackson 1912)
	Kootenia havasuensis	Resser 1945	Cambrian, Muav	USNM 108601a	Trilobite
	Pachyaspis moorei	Resser 1945	Cambrian, Muav	USNM 108606a	Trilobite

Appendix Table B-1 (continued). Fossil taxa named from specimens possibly found within GRCA.

Category	Taxon	Citation	Age, Formation	Type Specimen	Notes
Invertebrate Ichnofossils	*Cruziana linnarsoni*	White 1874	Cambrian, Tonto Group	Syntype USNM 8614	Invertebrate (crustacean?) trace
	Cruziana rustica	White 1874	Cambrian, Tonto Group	Syntype USNM 8615	Invertebrate (crustacean?) trace

Literature Cited

Jackson, R. T. 1912. Phylogeny of the Echini, with a revision of Palaeozoic species. Memoirs of the Boston Society of Natural History 7.

McKee, E. D. 1938. The environment and history of the Toroweap and Kaibab formations of northern Arizona and southern Utah. Carnegie Institution of Washington Publication 492.

Newberry, J. S. 1861. Geological report. *In* Ives, J. C., Report upon the Colorado River of the West, explored in 1857 and 1858. U.S. 36th Congress, 1st Session, House Executive Document 90, Serial 1058, Part 3.

Resser, C. E. 1945. Cambrian fossils of the Grand Canyon. Pages 169–220 *in* E. D. McKee and C. E. Resser. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

White, C. A. 1874. Preliminary report upon invertebrate fossils collected by the expeditions of 1871, 1872, and 1873, with descriptions of new species. *In* U.S. Army Engineer Department, Geographical and geological explorations and surveys west of the one hundredth meridian. U.S. Government Printing Office.

Appendix C. Stratigraphic Tables of GRCA Paleozoic Taxa

The following tables collate known Paleozoic taxa reported from localities verified as within GRCA, based on the literature and Grand Canyon National Park museum records. They provide a synoptic look at the groups of fossils discussed in Chapters 5 through 9 in a stratigraphic format. Taxonomically, it was not possible to critically evaluate every cited occurrence. There are undoubtedly instances where the same taxon has been listed more than once due to differing interpretations or general changes in taxonomic usage. In addition, it was not within the scope of the project to evaluate every potential synonymy and genus-species combination. The most recent name employed in Grand Canyon literature for a given taxon is generally used, with usage in the chapters of this document serving as the ultimate arbitrator. Older combinations are included in a section following the tables to facilitate translation between different references. Some cited taxa have outdated names, but the author did not provide enough information to determine more appropriate names. In these cases, the taxonomy is left as the author gave it.

The tables are organized with the first column providing the taxa and the last column providing references, with the intervening tables showing presence or absence. Groups are separated by darker gray rows; within groups, taxa are sorted alphabetically by genus and species, with reports not assigned to a genus placed after those that are. Taxon presence in a given formation is denoted by "Y" in the appropriate cell. Taxon absence is denoted by "–". If there is some question about an occurrence, a "?" is used. Some taxa are present in multiple formations; in these cases, the listed references may apply to one or all of the formations. "GCM" as a reference refers to GRCA collections; only unique results unreported in the literature are included. Names employed these otherwise unattested records should be taken with some caution. Due to the number of taxa and formations, entries have been split into four tables, incorporating the following formations in ascending order:

Acronyms used for Appendix Table C-1: Cambrian

Cu = Frenchman Mountain Dolostone

Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone)

Ctm = Muav Limestone

Ctba = Bright Angel Shale (* = possibly Muav Limestone instead)

Ctt = Tapeats Sandstone

Acronyms used for Appendix Table C-2: Devonian and Mississippian

Msc = Surprise Canyon Formation

Mr = Redwall Limestone

Dtb = Temple Butte Formation

Acronyms used for Appendix Table C-3: Supai Group

PIPsu = Undivided Supai Group

Pse = Esplanade Sandstone

IPswe = Wescogame Formation

IPsm = Manakacha Formation

IPswa = Watahomigi Formation

Acronyms used for Appendix Table C-4: post-Supai Permian

Pk = Kaibab Formation

Pt = Toroweap Formation

Pc = Coconino Sandstone

Ph = Hermit Formation

Appendix Table C-1. Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Plants	Cryptogam spores, various morphologies	–	Y	–	–	–	Strother and Beck 2000, Taylor and Strother 2008
Invertebrates	**Overall Invertebrates**	**Y**	**Y**	**Y**	**Y**	**–**	**–**
Invertebrates: Porifera (sponges)	Archaeocyatha undetermined?	–	?	Y	–	–	Resser 1945, GCM
	Sponge spicules	–	–	Y	–	–	Resser 1945 (potentially *Chancelloria*)
Invertebrates: Brachiopoda (lamp shells)	**Overall Brachiopoda**	**Y**	**Y**	**Y**	**Y**	**–**	**–**
	Acrothele sp.	–	–	–	Y	–	Walcott 1890
	Acrotreta? sp.	–	–	Y	–	–	Noble 1922
	Clitambonites sp.	–	–	Y	–	–	Noble 1922
	Dictyonina arizonaensis	–	Y	–	–	–	Walcott 1912a, Noble 1922, Resser 1945, Bonde et al. 2018
	Dictyonina sp.	–	–	?	–	–	Resser 1945
	Diraphora impressions?	–	Y	–	–	–	Resser 1945
	Discina sp.	–	–	Y	–	–	Noble 1922
	Finkelnburgia sp.	–	–	Y	–	–	Noble 1922
	Iphidea ornatella	–	–	–	Y	–	Walcott 1890
	Lingula sp.	–	Y	Y	–	–	Noble 1922, Strother and Beck 2000
	Lingula monticula?	–	Y	–	–	–	Frech 1893
	Lingulella acutangula	–	Y	–	–	–	Walcott 1912a, Noble 1922
	Lingulella chuarensis	–	Y	–	–	–	Walcott 1898, 1912a, 1916a, Schuchert 1918a, Noble 1922, Resser 1945, Foster 2011
	Lingulella euglypha	–	Y	–	–	–	Walcott 1898, 1912a, 1916a, Resser 1945
	Lingulella kanabensis	–	–	–	Y	–	Resser 1945
	Lingulella lineolata	–	Y	–	–	–	Walcott 1898, 1912a, 1916a, Noble 1922, Resser 1945
	Lingulella mckeei	–	Y	–	–	–	Resser 1945, Foster 2011, Bonde et al. 2018

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Invertebrates: Brachiopoda (lamp shells) (continued)	*Lingulella spatula*	–	Y	–	–	–	Walcott 1902, 1912a, Noble 1922, Resser 1945
	Lingulella themis	–	Y	–	–	–	Walcott 1905, 1912a, Noble 1922, Resser 1945
	Lingulella winona convexa	–	–	–	Y	–	Walcott 1912a
	Lingulella zetus	–	Y	–	Y	–	Walcott 1898, 1912a, 1916a, Resser 1945
	Lingulella n. sp.	–	Y	–	–	–	Schenk and Wheeler 1942
	Lingulella sp.	–	Y	–	Y	–	Walcott 1890, Resser 1945
	Lingulepis prima?	–	–	Y	–	–	Noble 1922
	Lingulepis sp.	–	–	Y	–	–	Walcott 1883, 1890, Noble 1922
	Micromitra pealei	–	Y	–	–	–	Walcott 1912a, 1916a
	Micromitra (*Paterina*) *crenistria*	–	Y	–	Y	–	Walcott 1897, 1912a, 1916a, Resser 1945
	Micromitra (*Paterina*) *superba*	–	Y	–	–	–	Walcott 1897, 1912a, 1916a, Noble 1922, Resser 1945
	Nisusia kanabensis	–	–	–	Y	–	Walcott 1908, 1912a, Resser 1945
	Nisusia noblei	–	–	Y	–	–	Walcott 1924, Resser 1945
	Nisusia noblei?	–	Y	Y	–	–	Resser 1945, GCM
	Nisusia obscura	–	Y	–	–	–	Walcott 1905, 1912a, 1916a, Resser 1945
	Nisusia sp.	–	Y	Y	–	–	Schenk and Wheeler 1942, Palmer E&R 1963/10/17, Bonde et al. 2018
	Nisusia? sp.	–	Y	Y	–	–	Resser 1945, Palmer E&R 1963/10/17
	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
	Obolella polita?	–	Y	–	–	–	Frech 1893
	Obolella sp.	–	Y	–	–	–	Frech 1893
	Obolella sp.?	–	Y	–	–	–	Frech 1893
	Obolus sp.	–	Y	–	–	–	GCM
	Paterina? sp.	–	Y	–	–	–	Resser 1945

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Invertebrates: Brachiopoda (lamp shells) (continued)	*Protorthis* sp.	–	–	–	Y	–	Walcott 1912a
	Syntrophia sp.	–	–	Y	–	–	Noble 1922
	Trematis sp.	–	–	Y	–	–	Noble 1922
	Wimanella sp.	–	Y	–	–	–	GCM
	Obolidae undetermined	Y	–	–	–	–	Schuchert 1918a
	Brachiopoda undetermined	–	Y	Y	–	–	McKee 1945, GCM
Invertebrates: Hyolitha	*Hyolithes primordialis*?	–	–	Y	–	–	Noble 1922
	Hyolithes sp. (small)	–	Y*	–	–	–	Resser 1945
	Hyolithes sp.	–	Y	Y	Y	–	Walcott 1890, 1916a, Noble 1922, Schenk and Wheeler 1942, Resser 1945, Foster 2011, Bonde et al. 2018
	Hyolithes sp. (two species)	–	Y*	–	–	–	Resser 1945
	Hyolithes? sp.	–	–	Y	–	–	Resser 1945
Invertebrates: Mollusca: Helcionelloida	*Helcionella* sp.	–	–	Y	–	–	Palmer E&R 1963/10/17, Bonde et al. 2018
Invertebrates: Mollusca: Gastropoda (snails)	Gastropoda undetermined	–	Y	–	–	–	GCM
Invertebrates: Arthropoda: Trilobita	**Overall Trilobita**	**?**	**Y**	**Y**	**Y**	**–**	**–**
	Acrocephalops? *arizonaensis*	–	Y	–	–	–	Resser 1945
	Albertella schenki	–	Y	–	–	–	Resser 1945
	Albertella n. sp.	–	Y	–	–	–	Schenk and Wheeler 1942
	Albertella sp.	–	Y	–	–	–	Bonde et al. 2018
	Alokistocare lepida	–	Y	–	–	–	Schenk and Wheeler 1942
	Alokistocare sp.	–	Y	Y	–	–	Schenk and Wheeler 1942, Resser 1945

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Invertebrates: Arthropoda: Trilobita (continued)	*Alokistocare*?	–	Y	–	–	–	Schenk and Wheeler 1942
	Alokistocare sp. or *Ehmaniella* sp.	–	Y	–	–	–	Bonde et al. 2018
	Amecephalus althea	–	Y	–	–	–	Walcott 1916a, 1916b, Schuchert 1918a, Noble 1922, Schenk and Wheeler 1942, Resser 1945, Foster 2011
	Amecephalus cf. *A. althea*	–	Y	–	–	–	Schenk and Wheeler 1942
	Amecephalus sp. cf. *A. packi*	–	Y	–	–	–	Foster 2011
	Anomocarella sp.	–	–	Y	–	–	Noble 1922
	Anoria n. sp.	–	Y	–	–	–	Schenk and Wheeler 1942
	Anoria sp.	–	Y	Y	–	–	Schenk and Wheeler 1942, Resser 1945
	"Anoria" *tontoensis*	–	Y	–	–	–	Walcott 1916a, Schuchert 1918a, Noble 1922, Resser 1945, Foster 2011
	Antagmus arizonaensis	–	Y	–	–	–	Resser 1945, Bonde et al. 2018
	Antagmus sp.	–	Y	–	–	–	GCM
	Athabaskia kanabensis	–	–	–	Y	–	Resser 1945
	Athabaskia sp.	–	Y	Y	–	–	Resser 1945
	Bathyurus? sp.	–	–	Y	–	–	Noble 1922
	Bolaspis aemula	–	–	Y	–	–	Resser 1945
	Bolaspis? sp.	–	–	Y	–	–	Resser 1945
	Dolichometopus sp.	–	–	–	Y	–	Walcott 1890
	Dorypyge sp.	–	–	Y	–	–	Stoyanow 1936
	Ehmaniella arizonaensis	–	Y	–	–	–	Resser 1945
	Ehmaniella basilica	–	Y	–	–	–	Bonde et al. 2018
	Ehmaniella hebes	–	–	–	Y	–	Resser 1945
	Ehmaniella sp.	–	Y	–	–	–	Bonde et al. 2018
	Elrathia nitens	–	Y	–	–	–	Resser 1945

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Invertebrates: Arthropoda: Trilobita (continued)	*Elrathia* n. sp.	–	Y	–	–	–	Schenk and Wheeler 1942
	Elrathia sp.	–	Y	–	–	–	Schenk and Wheeler 1942, Resser 1945
	Elrathiella? *insueta*	–	–	–	Y	–	Resser 1945
	Glossopleura boccar	–	Y	–	Y	–	Walcott 1916a, Schuchert 1918a, Noble 1922, Resser 1935, Schenk and Wheeler 1942, Resser 1945, Palmer E&R 1963/10/17, Foster 2011, Bonde et al. 2018
	Glossopleura meriwitica	–	Y	–	–	–	GCM
	Glossopleura walcotti?	–	Y	–	–	–	Foster 2011
	Glossopleura sp.	–	Y	Y	Y	–	Schenk and Wheeler 1942, Resser 1945, Palmer E&R 1963/10/17, Elliott and Martin 1987a, Bonde et al. 2018
	Glossopleura? sp.	–	Y	–	–	–	GCM
	Glossopleura sp. or *Anoria* sp.	–	–	Y	–	–	Resser 1945
	Glyphaspis kwaguntensis	–	–	Y	Y	–	Resser 1945, GCM
	Glyphaspis tecta	–	–	Y	–	–	Resser 1945
	Glyphaspis vulsa	–	Y	–	–	–	Resser 1945
	Glyphaspis sp.	–	Y	Y	–	–	Schenk and Wheeler 1942, Resser 1945
	Glyphaspis? sp.	–	–	Y	–	–	Resser 1945
	Kochina? *angustata*	–	–	–	Y	–	Resser 1945
	Kootenia havasuensis	–	–	?	–	–	Resser 1945
	Kootenia mckeei	–	–	Y	Y	–	Resser 1945
	Kootenia schenki	–	–	Y	–	–	Schenk and Wheeler 1942, Resser 1945, Palmer E&R 1963/10/17, Bonde et al. 2018
	Kootenia simplex	–	Y	Y	–	–	Resser 1945

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Invertebrates: Arthropoda: Trilobita (continued)	*Kootenia* sp.	–	Y	Y	Y	–	Schenk and Wheeler 1942, Resser 1945, Palmer E&R 1963/10/17, Foster 2011
	Kootenia? sp.	–	–	Y	–	–	Resser 1945
	Neolenus sp.	–	–	Y	–	–	Noble 1922
	Olenellus sp.	?	Y	–	–	–	Schenk and Wheeler 1942, Resser 1945
	Olenoides sp.	–	–	–	Y	–	Walcott 1890
	Pachyaspis fonticola	–	Y	–	–	–	Resser 1945
	Pachyaspis moorei	–	Y	?	–	–	Resser 1945, Bonde et al. 2018
	Pachyaspis sp.	–	Y*	–	–	–	Resser 1945
	Pagodia? sp.	–	–	Y	–	–	Noble 1922
	Parehmania kwaguntensis	–	Y	–	Y	–	Bonde et al. 2018
	Parehmania nitida	–	–	–	Y	–	Resser 1945
	Parehmania tontoensis	–	Y	–	–	–	Resser 1945
	Parehmania sp.	–	Y*	–	–	–	Resser 1945
	Ptarmigania sp.	–	Y	–	Y	–	Resser 1945
	Ptychoparia sp.	–	–	Y	–	–	Walcott 1883, 1890, Noble 1922
	Ptychoparia? sp.	–	–	Y	–	–	Noble 1922
	Saukia sp.	–	–	Y	–	–	Noble 1922
	Spencella diligens	–	–	–	Y	–	Resser 1945
	Spencella erosa	–	–	Y	–	–	Resser 1945
	Spencella porcata	–	–	Y	–	–	Schenk and Wheeler 1942, Resser 1945, Palmer E&R 1963/10/17, Bonde et al. 2018
	Spencella sp.	–	–	Y	Y	–	Resser 1945, GCM
	Spencia tontoensis	–	Y	–	–	–	Resser 1945
	Trachycheilus typicale	–	–	–	Y	–	Resser 1945
	Zacanthoides cf. *walapai*	–	Y	–	–	–	Resser 1945

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Invertebrates: Arthropoda: Trilobita (continued)	*Zacanthoides* sp.	–	Y	Y	–	–	Resser 1945, GCM
	Olenellidae undetermined	–	Y	–	–	–	Bonde et al. 2018
	Trilobita undetermined	–	Y	Y	–	–	Stoyanow 1936, Palmer E&R 1963/10/17, GCM
Invertebrates: Arthropoda: Bradoriida	**Overall Bradoriida**	**–**	**Y**	**–**	**Y**	**–**	**–**
	"*Bradoria tontoensis*"	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Dielymella appressa	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Dielymella dorsalis	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Dielymella nasuta	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Dielymella recticardinalis	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Dielymella recticardinalis angustata	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Indianites curtus	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Indianites faba	–	Y*	–	–	–	Noble 1922, Ulrich and Bassler 1931a, Resser 1945
	Indianites impressus	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Indianites intermedius	–	Y*	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella apicalis	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella breviuscula	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella concentrica	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella leperditoides	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella limatula	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella longula	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella nitida	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella oblonga	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella obsoleta	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella pulchella	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella scitula	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Invertebrates: Arthropoda: Bradoriida (continued)	*Walcottella subtruncata*	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Walcottella ventrosa	–	Y	–	–	–	Ulrich and Bassler 1931a, Resser 1945
	Three species of "*Leperditia*"	–	–	–	Y	–	Walcott 1890
Invertebrates: Echinodermata: Eocrinoidea	*Eocystites*? undetermined sp.	–	Y	–	–	–	Walcott 1916a, Noble 1922
	Gogia longidactylus	–	Y	–	–	–	GCM
	Gogia ?longidactylus	–	Y	–	–	–	Foster 2011
	Gogia multibrachiatus	–	Y	–	–	–	Resser 1945, Foster 2011
	"Cystoid" (interpreted as Eocrinoidea undetermined)	–	–	Y	–	–	GCM
Invertebrates: Other Invertebrates	*Chancelloria* cf. *C. eros*	–	–	Y	–	–	Schenk and Wheeler 1942
	Chancelloria sp.	–	Y	–	–	–	Elliott and Martin 1987a
	"Echinoid imprint?"	–	Y	–	–	–	GCM
	Margaretia sp.	–	Y	–	–	–	GCM
	Margaretia? sp. (misspelled *Margaritia*)	–	Y	–	–	–	Schenk and Wheeler 1942
	"Pelecypod" (bradoriid?)	–	Y	–	–	–	GCM
	Scenella hermitensis	–	–	Y	–	–	Resser 1945
	Tontoia kwaguntensis	–	Y	–	–	–	Walcott 1912b, Resser 1945
Ichnofossils	**Overall Ichnofossils**	**Y**	**Y**	**Y**	**Y**	**Y**	**–**
Ichnofossils: Microbial Trace Fossils	Wrinkle structures	–	Y	–	–	–	Strother and Beck 2000, Baldwin et al. 2004
Ichnofossils: Invertebrate Trace Fossils	**Overall Invertebrate Trace Fossils**	**Y**	**Y**	**Y**	**Y**	**Y**	**–**
	Angulichnus alternipes	–	Y	–	–	–	Martin 1985, Elliott and Martin 1987b
	Arenicolites isp.	Y	Y	–	–	–	Baldwin et al. 2004, Hagadorn et al. 2011

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Ichnofossils: Invertebrate Trace Fossils (continued)	*Belorhaphe* isp.	–	Y	–	–	–	Martin 1985
	Bergaueria aff. *B. perata*	–	Y	–	–	–	Martin 1985
	Bergaueria isp.	–	Y	–	–	–	Chapter 8
	Cruziana arizonensis	Y	–	–	–	–	Seilacher 1970
	Cruziana linnarssoni	–	Y	–	–	–	White 1874
	Cruziana cf. *C. rusiformis*	–	Y	–	–	–	GCM
	Cruziana rustica	–	Y	–	–	–	White 1874
	Cruziana "*grandcanyonensis*/*rusiformis*"	–	Y	–	–	–	Martin 1985, Rose 2003
	Cruziana isp.	Y	Y	–	Y	–	Walcott 1886, Frech 1893, Palmer E&R 1963/10/17, Elliott and Martin 1987b, Baldwin et al. 2004, Hagadorn et al. 2011
	Dimorphichnus isp.	–	Y	–	–	–	Martin 1985, Baldwin et al. 2004
	Diplichnites isp.	Y	Y	–	–	–	Chapter 8, GCM
	Diplocraterion yoyo	–	Y	–	–	–	Martin 1985
	Diplocraterion isp.	Y	Y	Y	–	–	Elliott and Martin 1987a, 1987b, Baldwin et al. 2004, Chapter 8
	Diplocraterion or *Rhizocorallium*	Y	–	–	–	–	Chapter 8
	"*Flectostriatus imporcatus*"	–	Y	–	–	–	Martin 1985
	"*Fodichnites bitumulus*"	–	Y	–	–	–	Martin 1985
	Monocraterion isp.	–	Y	–	–	–	Baldwin et al. 2004
	Monomorphichnus lineatus var. *giganticus*	–	Y	–	–	–	Chapter 8
	Monomorphichnus isp.	Y	Y	–	–	–	Martin 1985, Baldwin et al. 2004, Hagadorn et al. 2011
	Nereites isp.	–	Y	–	–	–	Martin 1985, Chapter 8

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Ichnofossils: Invertebrate Trace Fossils (continued)	*Palaeophycus* aff. *P. striatus*	–	Y	–	–	–	Martin 1985
	Palaeophycus isp.	Y	Y	–	–	–	Schuchert 1918a, Elliott and Martin 1987a, 1987b, Baldwin et al. 2004
	"*Pholetichnus circinatus*"	–	Y	–	–	–	Martin 1985
	Phycodes aff. *P. circinnatum*	–	Y	–	–	–	Martin 1985
	Phycodes isp.	–	Y	Y	–	–	Elliott and Martin 1987b, Baldwin et al. 2004, Chapter 8
	Planolites isp.	Y	Y	–	–	–	Baldwin et al. 2004, Hagadorn et al. 2011
	Rusophycus didymus	Y	Y	–	–	–	Chapter 8
	Rusophycus cf. *R. dispar*	–	Y	–	–	–	GCM
	Rusophycus "*biungis*"	–	Y	–	–	–	Martin 1985
	Rusophycus isp.	Y	Y	–	Y	–	Elliott and Martin 1987b, Baldwin et al. 2004, Bonde et al. 2018, Chapter 8
	Scolicia isp.	–	Y	–	–	–	Martin 1985
	Sinusites isp.	–	Y	–	–	–	Chapter 8
	Skolithos annulatus	Y	–	–	–	–	Howell 1957, Alpert 1974
	Skolithos linearis	–	Y	–	–	–	Martin 1985
	Skolithos cf. *S. linearis*	Y	–	–	–	–	GCM
	Skolithos isp.	Y	Y	–	–	–	Frech 1893, McKee 1945, Elliott and Martin 1987a, Baldwin et al. 2004, Hagadorn et al. 2011
	Spirophycus isp.	–	Y	–	–	–	Martin 1985
	Teichichnus rectus	–	Y	–	–	–	Martin 1985
	Teichichnus isp.	Y	Y	–	–	–	Elliott and Martin 1987a, Baldwin et al. 2004, Hagadorn et al. 2011
	Teichichnus isp. large	–	Y	–	–	–	Chapter 8
	Teichichus isp. small	–	Y	–	–	–	Chapter 8

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Ichnofossils: Invertebrate Trace Fossils (continued)	Cf. *Teichichnus* isp.	–	Y	–	–	–	GCM
	Treptichnus pedum	–	Y	Y	–	–	Martin 1985, Buatois 2018, GCM
	Treptichnus cf. *T. pedum*	–	–	Y	–	–	GCM
	Treptichnus isp.	Y	–	–	–	–	Hagadorn et al. 2011
	"Fucoids" (including "seaweeds")	Y	Y	Y	–	–	Noble 1922, White 1928, 1929, McKee 1932, 1945
	Indeterminate horizontal burrows	–	Y	–	–	–	Martin 1985
	Invertebrate tracks	–	–	–	Y	–	Palmer E&R 1963/10/17
	Reticulate burrow	–	–	Y	–	–	GCM
	Spiral trace fossil	–	Y	–	–	–	Martin 1985
	Trilobite trails	Y	Y	–	Y	–	Schenk and Wheeler 1942, McKee 1945, Palmer E&R 1963/10/17
	Trilobite? crawling and resting traces	–	Y	–	–	–	Beus 1987
	Vertical tubes	–	–	Y	–	–	GCM
	Worm borings and trails	Y	Y	Y	–	Y	Noble 1922, Schenk and Wheeler 1942, McKee 1945
	Unspecified invertebrate trace fossils	–	Y	Y	–	–	Beus 1987, GCM
Other Fossils	**Overall Other Fossils**	**Y**	**Y**	**Y**	**–**	**–**	**–**
	"*Eophyton*"? (a type of tool mark made by organisms dragged by currents)	–	–	Y	–	–	GCM
	"Fossil spine-like structures"	Y	–	–	–	–	GCM (possibly Muav instead)
	Girvanella structures/spheres	–	–	Y	–	–	McKee 1945
	Large metazoan fragments	–	Y	–	–	–	Baldwin et al. 2004

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-1 (continued). Cambrian fossil taxa. Ctt = Tapeats Sandstone, Ctba = Bright Angel Shale (* = possibly Muav Limestone instead), Ctm = Muav Limestone, Ctu = Undivided Tonto Group (presumably mostly the Bright Angel Shale and Muav Limestone), and Cu = Frenchman Mountain Dolostone.

Category	Taxon Observed	Ctt	Ctba	Ctm	Ctu	Cu	References
Other Fossils (continued)	Leiospheres	–	Y	–	–	–	Baldwin et al. 2004
	Mats of filaments resembling *Nematothallus*	–	Y	–	–	–	Baldwin et al. 2004
	Non-marine cryptospores	–	Y	–	–	–	Baldwin et al. 2004
	Organic impression	Y	–	–	–	–	GCM
	Terrestrial algal cell clusters	–	Y	–	–	–	Baldwin et al. 2004
	Undetermined fossil	–	Y	–	–	–	GCM

* = possibly Muav Limestone instead of Bright Angel Shale

Appendix Table C-2. Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Plants	**Overall Plants**	**–**	**–**	**Y**	**–**
Plants: Equisetopsida	*Calamites* (*Mesocalamites*) *cistiformis*	–	–	Y	Tidwell et al. 1992
	Calamites sp.	–	–	Y	Tidwell et al. 1992, Beus 1999
Plants: Isoetopsida	*Lepidodendron* sp.	–	–	Y	Beus 1990a, 1995, Beus 1999, Billingsley and Beus 1999a
	Lepidostrobophyllum sp.	–	–	Y	Tidwell et al. 1992, Beus 1999
Plants: Other Plants	Carbonized wood	–	–	Y	Hodnett and Elliott 2018
	Undetermined wood	–	–	Y	Billingsley and Beus 1999a
	Undetermined plant fossils	–	–	Y	Beus 1995, Billingsley and Beus 1999a, Hodnett and Elliott 2018
Invertebrates	**Overall Invertebrates**	**Y**	**Y**	**Y**	**–**
Invertebrates: Cnidaria: Anthozoa (corals)	**Overall Anthozoa**	**Y**	**Y**	**Y**	**–**

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Invertebrates: Cnidaria: Anthozoa (corals) (continued)	Anthozoa undetermined	Y	Y	Y	Noble 1922, McKee and Gutschick 1969a, Billingsley and Beus 1985, 1999a, Hodnett and Elliott 2018, GCM
	Anthozoa undetermined?	–	Y	–	GCM
Invertebrates: Cnidaria: Anthozoa: Tabulata (tabulate corals)	**Overall Tabulata**	**–**	**Y**	**Y**	**–**
	Michelinia sp.	–	–	Y	Billingsley and McKee 1982, Beus 1999
	Syringopora aculeata	–	Y	–	McKee and Gutschick 1969b, Sando 1969
	Syringopora surcularia	–	Y	–	Sando E&R 1963/10/14, Bonde et al. 2018
	Syringopora aff. *S. surcularia*	–	Y	–	Sando E&R 1963/10/14
	Syringopora cf. *S. surcularia*	–	Y	–	Schuchert 1918b
	Syringopora sp.	–	Y	–	McKee and Gutschick 1969b, Sando and Bamber 1985
	Tabulata undetermined	–	Y	–	GCM
Invertebrates: Cnidaria: Anthozoa: Rugosa (rugose corals)	**Overall Rugosa**	**Y**	**Y**	**Y**	**–**
	Amplexizaphrentis sp.	–	Y	–	Sando and Bamber 1985
	Amplexus sp.	–	–	Y	Beus 1999
	Barytichisma sp.	–	–	Y	Beus 1999
	Clisiophyllum sp.	–	Y	–	Schuchert 1918b
	Diphyphyllum? sp.	–	Y	–	Noble 1922
	Diphyphyllum (*Lithostrotion*?) sp.	–	Y	–	Noble 1922
	Dorlodotia inconstans	–	Y	–	Easton and Gutschick 1953
	Dorlodotia inconstans?	–	Y	–	Easton and Gutschick 1953
	Dorlodotia sp.?	–	?	–	Sando and Bamber 1985
	Homalophyllites paucicinctus	–	?	–	Easton and Gutschick 1953
	Homalophyllites subcrassus	–	Y	–	McKee and Gutschick 1969b, Sando 1969
	Menophyllum excavatum	–	Y	–	Schuchert 1918b
	Streptelasma? sp.	?	–	–	Schenk and Wheeler 1942

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Invertebrates: Cnidaria: Anthozoa: Rugosa (rugose corals) (continued)	*Sychnoelasma* sp.	–	Y	–	Sando and Bamber 1985
	Vesiculophyllum incrassatum	–	Y	–	Easton and Gutschick 1953, Sando E&R 1963/10/14, McKee and Gutschick 1969b, Sando 1969, Bonde et al. 2018
	Vesiculophyllum sp.	–	Y	–	Sando and Bamber 1985
	Zaphrentites persimilis	–	Y	–	Sando E&R 1963/10/14, Bonde et al. 2018
	"Cyathophylloids"	Y	–	–	Walcott 1883
	Probable "cup corals"	Y	–	–	Noble 1922
	Undetermined horn corals	–	Y	Y	McKee and Gutschick 1969b, Billingsley and Beus 1999a
Invertebrates: Bryozoa (moss animals)	**Overall Bryozoa**	**–**	**Y**	**Y**	**–**
	Archimedes sp.	–	–	Y	Beus 1999
	Cladochonus? sp.	–	Y	–	Noble 1922
	Cystodictya sp.	–	Y	–	McKee and Gutschick 1969b
	Dichotrypa n. sp.	–	Y	–	Duncan 1969
	Dichotrypa sp.	–	Y	–	McKee and Gutschick 1969b
	Fenestella sp.	–	Y	–	Schuchert 1918b, Noble 1922
	Fenestella fine mesh	–	Y	–	McKee and Gutschick 1969b
	Fenestella medium mesh	–	Y	–	McKee and Gutschick 1969b
	Fenestella moderately coarse mesh	–	Y	–	McKee and Gutschick 1969b
	Fenestella coarse mesh	–	Y	–	McKee and Gutschick 1969b
	Fenestella fine mesh?	–	Y	–	McKee and Gutschick 1969b
	Fenestella medium mesh?	–	Y	–	McKee and Gutschick 1969b
	Fenestella coarse mesh?	–	Y	–	McKee and Gutschick 1969b
	Fenestralia? sp.	–	Y	–	McKee and Gutschick 1969b
	Penniretepora sp.	–	Y	–	McKee and Gutschick 1969b
	Polypora sp.	–	Y	–	Schuchert 1918b, McKee and Gutschick 1969b

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Invertebrates: Bryozoa (moss animals) (continued)	*Polypora*? sp.	–	Y	–	McKee and Gutschick 1969b
	Fenestellidae undetermined	–	Y	–	McKee and Gutschick 1969b
	Undetermined “rhomboporoid”?	–	Y	–	McKee and Gutschick 1969b
	Encrusting forms	–	–	Y	Beus 1999
	Fenestrate forms	–	–	Y	Beus 1999, Billingsley and Beus 1999a
	Ramose forms	–	–	Y	Beus 1999, Billingsley and Beus 1999a
	Bryozoa undetermined	–	Y	Y	McKee and Gutschick 1969a, Billingsley and Beus 1985, 1999a, Hodnett and Elliott 2018
Invertebrates: Brachiopoda (lamp shells)	**Overall Brachiopoda**	**Y**	**Y**	**Y**	**–**
	Anthracospirifer bifurcatus	–	–	Y	Gordon E&R 1973/1/3
	Anthracospirifer curvilateralis	–	–	Y	Beus 1999
	Anthracospirifer aff. *A. curvilateralis*	–	–	Y	Billingsley and McKee 1982
	Anthracospirifer cf. *A. curvilateralis*	–	–	Y	Gordon E&R 1979/1/23
	Anthracospirifer sp. A	–	–	Y	Gordon E&R 1979/1/23, Billingsley and McKee 1982
	Beecheria cf. *B. arkansanum*	–	–	Y	Beus 1999
	Beecheria sp.	–	–	Y	Billingsley and McKee 1982
	Brachythyris subcardiiformis	–	–	Y	Gordon E&R 1973/1/3
	Camarotoechia sp.	–	Y	–	Noble 1922
	Camarotoechia? n. sp.	–	Y	–	Noble 1922
	Camarotoechia? sp.	–	Y	–	Noble 1922
	Cleiothyridina sp.	–	–	Y	Beus 1999
	Composita gibbosa	–	–	Y	Billingsley and McKee 1982
	Composita laevis	–	–	Y	Beus 1999
	Composita ovata	–	–	Y	Billingsley and McKee 1982
	Composita subquadrata	–	–	Y	Beus 1999
	Composita sp.	–	Y	Y	Gordon E&R 1979/1/23, GCM

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Invertebrates: Brachiopoda (lamp shells) (continued)	*Composita*? sp.	–	Y	–	Noble 1922
	Cranaena sp.	–	–	Y	Gordon E&R 1979/1/23
	Dielasma sp.	–	Y	–	Carter et al. 2014
	Eumetria sp.	–	–	Y	Beus 1999
	Flexaria sp.	–	–	Y	Beus 1999
	Floweria chemungensis	–	Y	–	Noble 1922
	Inflatia aff. *I. clydensis*	–	–	Y	Gordon E&R 1979/1/23, Billingsley and McKee 1982
	Inflatia sp.	–	–	Y	Gordon E&R 1979/1/23
	Inflatia? sp.	–	–	Y	Gordon E&R 1973/1/3
	Inflatia sp. or *Sandia* sp.	–	–	Y	Billingsley and McKee 1982
	Leiorhynchoidea carbonifera	–	–	Y	Beus 1999
	Leiorhynchoidea sp.	–	–	Y	Gordon E&R 1979/1/23
	Leptagonia sp.	–	Y	–	McKee and Gutschick 1969c
	?*Macropotamorhynchus* cf. *M. purduei*	–	–	Y	Beus 1999
	Mirifusella cf. *M. fortunata*	–	Y	–	Carter et al. 2014
	Neospirifer striatus	–	Y	–	Frech 1893
	Orthotetes sp.	–	–	Y	Beus 1999
	Ovatia sp.	–	–	Y	Gordon E&R 1979/1/23, Beus 1999
	Ovatia sp. (small)	–	–	Y	Billingsley and McKee 1982
	Prospira sp. B.	–	Y	–	Carter et al. 2014
	Pugnoides sp.	–	Y	Y	Billingsley and McKee 1982
	Rhipidomella nevadensis	–	–	Y	Beus 1999
	Rhipidomella sp.	–	Y	–	Noble 1922
	Rotaia neogenes	–	–	Y	Beus 1999
	Schizophoria sp.	–	–	Y	Billingsley and McKee 1982

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Invertebrates: Brachiopoda (lamp shells) (continued)	*Spirifer centronatus*	–	Y	–	Schuchert 1918b, Noble 1922
	Spirifer redwallensis	–	?	–	Carter et al. 2014
	Spirifer aff. *S. incertus*	–	Y	–	Noble 1922
	Spirifer cf. *S. occidentalis*	–	Y	–	Grant E&R 1963/10/16, Bonde et al. 2018
	Spirifer sp.	–	Y	–	Grant E&R 1963/10/16, McKee and Gutschick 1969b
	Spirifer sp. smaller	–	Y	–	Grant E&R 1963/10/16
	Finely striated *Spirifer*	–	Y	–	Schuchert 1918b
	Syringothyris? sp.	–	Y	–	Noble 1922, Grant E&R 1963/10/16, Bonde et al. 2018
	Tomiproductus gallatinensis	–	?	–	Carter et al. 2014
	Torynifer setiger?	–	–	Y	Billingsley and McKee 1982
	Torynifer sp.	–	–	Y	Beus 1999
	Unispirifer minnewankensis	–	Y	–	Carter et al. 2014
	Wellerella osagensis	–	Y	–	GCM
	Orthotetoidea undetermined	–	–	Y	Billingsley and McKee 1982
	Small productoids	–	–	Y	Beus 1999
	Productida undetermined	–	Y	Y	McKee and Gutschick 1969b, 1969c, Billingsley and Beus 1999a
	Rhynchonellida undetermined	–	Y	–	McKee and Gutschick 1969c
	Punctate spiriferid	–	–	Y	Beus 1999
	Spiriferida undetermined	–	Y	Y	Grant E&R 1963/10/16, Billingsley and Beus 1999a, Bonde et al. 2018
	Strophomenida undetermined	–	Y	–	McKee and Gutschick 1969a
	Terebratulida undetermined	–	Y	–	McKee and Gutschick 1969b, McKee and Gutschick 1969c
	“Smooth-shelled brachiopod”	–	–	Y	Billingsley and Beus 1999a
	Brachiopoda undetermined	Y	Y	Y	Walcott 1883, Stoyanow 1936, Grant E&R 1963/10/16, McKee and Gutschick 1969b, 1969c, Billingsley and McKee 1982, Billingsley and Beus 1985, 1999a, Hodnett and Elliott 2018

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Invertebrates: Mollusca	**Overall Mollusca**	**Y**	**Y**	**Y**	**–**
	Mollusca undetermined	–	–	?	Billingsley and Beus 1985
	Mollusca undetermined?	–	Y	–	GCM
Invertebrates: Mollusca: Bivalvia (clams, oysters, etc.)	*Aviculopecten* sp.	–	–	Y	Gordon E&R 1979/1/23, Beus 1999
	?*Edmondia* sp.	–	–	Y	Beus 1999
	Septimyalina sp.	–	–	Y	Beus 1999
	Septimyalina? sp.	–	–	Y	Gordon E&R 1979/1/23
	Schizodus sp.	–	–	Y	Billingsley and McKee 1982
	Pectinida undetermined	–	–	Y	Billingsley and McKee 1982
	Bivalvia undetermined	–	–	Y	Billingsley and McKee 1982, Billingsley and Beus 1999a
Invertebrates: Mollusca: Cephalopoda: Nautiloidea	*Rayonnoceras* sp.	–	Y	–	Breed 1969, Billingsley and Breed 1976
	Coiled nautiloid	–	Y	–	Billingsley and Breed 1976
Invertebrates: Mollusca: Gastropoda (snails)	**Overall Gastropoda (snails)**	**Y**	**Y**	**Y**	**–**
	?*Bellazona* sp.	–	–	Y	Beus 1999
	Bellerophon sp.	–	Y	–	Noble 1922
	Bellerophon (*Bellerophon*) sp.	–	–	Y	Gordon E&R 1979/1/23, Billingsley and McKee 1982
	?*Bellerophon* spp.	–	–	Y	Beus 1999
	Euconospira montezuma	–	Y	–	Bonde et al. 2018
	Euomphalus sp.	–	–	Y	Gordon E&R 1979/1/23, Beus 1999
	Euomphalus? sp.	–	Y	–	Noble 1922
	cf. *Euphemites* sp.	–	Y	–	Bonde et al. 2018
	Glabrocingulum sp.	–	–	Y	Beus 1999
	?*Loxonema* sp.	–	–	Y	Beus 1999
	Straparollus (*Straparollus*?) sp.	–	–	Y	Billingsley and McKee 1982

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Invertebrates: Mollusca: Gastropoda (snails) (continued)	High-spired gastropod	–	–	Y	Billingsley and Beus 1999a
	Low-spired gastropod	–	Y	–	McKee and Gutschick 1969b
	Bellerophontoidea undetermined	–	Y	–	Yochelson 1969
	Euomphaloidea undetermined	–	Y	–	Yochelson 1969
	Murchisoniacea undetermined	–	Y	–	Yochelson 1969
	Gastropoda undetermined	Y	–	–	Walcott 1883
Invertebrates: Arthropoda	**Overall Arthropoda**	**–**	**Y**	**Y**	**–**
Invertebrates: Arthropoda: Trilobita	*Aprathia* sp.	–	Y	–	McKee and Gutschick 1969b, 1969d, Brezinski 2017
	Breviphillipsia n. sp.	–	Y	–	Cisne 1971
	Paladin cf. *P. chesterensis*	–	–	Y	Beus 1999
	Paladin sp.	–	–	Y	Gordon E&R 1979/1/23, Beus 1999
	Phillipsia peroccidens	–	Y	–	GCM
	Trilobita undetermined	–	–	Y	Billingsley and Beus 1999a
Invertebrates: Arthropoda: Ostracoda (seed shrimp)	Ostracoda undetermined	–	–	Y	Billingsley and Beus 1999a
Invertebrates: Echinodermata	**Overall Echinodermata**	**–**	**Y**	**Y**	**–**
	Pelmatozoan debris	–	–	Y	Billingsley and McKee 1982
	Echinodermata undetermined	–	–	?	Billingsley and Beus 1985, 1999a
Invertebrates: Echinodermata: Asteroidea (sea stars)	Uractinida undetermined	–	–	Y	Beus 1999
Invertebrates: Echinodermata: Blastoidea (sea buds)	*Pentremites* n. sp.	–	–	Y	Beus 1999
	Pentremites sp.	–	Y	–	McKee and Gutschick 1969b, Macurda 1969
	Blastoidea undetermined	–	–	Y	Billingsley and Beus 1999a

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Invertebrates: Echinodermata: Crinoidea (sea lilies)	*Cymbiocrinus* n. sp.	–	–	Y	Beus 1999
	Long crinoid columnals	–	Y	–	McKee and Gutschick 1969b
	Crinoidea undetermined	–	Y	Y	Frech 1893, Schuchert 1918b, Noble 1922, McKee and Gutschick 1969a, 1969c, McKee E&R 1978/5/25, Beus 1987, Billingsley and Beus 1999a , Hodnett and Elliott 2018
	Crinoidea undetermined?	–	Y	–	GCM
Invertebrates: Echinodermata: Echinoidea (sea urchins)	Echinoidea undetermined	–	–	Y	Billingsley and Beus 1999a
Invertebrates: Conodonta	**Overall Conodonta**	**Y**	**–**	**Y**	**–**
	Adetognathus unicornis	–	–	Y	Martin and Barrick 1999
	Adetognathus sp.	–	–	Y	Martin and Barrick 1999
	Adetognathus/*Cavusgnathus* sp.	–	–	Y	Martin and Barrick 1999
	Cavusgnathus naviculus	–	–	Y	Martin and Barrick 1999
	Cavusgnathus unicornis	–	–	Y	Martin and Barrick 1999
	Gnathodus bilineatus	–	–	Y	Martin and Barrick 1999
	Gnathodus girtyi simplex	–	–	Y	Martin and Barrick 1999
	Gnathodus sp.	–	–	Y	Martin and Barrick 1999
	Icriodus cf. *I. subterminus*	Y	–	–	Beus 1990b
	Kladognathus sp.	–	–	Y	Martin and Barrick 1999
	Pandorinella insita	Y	–	–	Beus 1990b
	Polygnathus pennatus	Y	–	–	Beus 1990b
	Polygnathus xylus	Y	–	–	Beus 1990b
	Polygnathus cf. *P. angustidiscus*	Y	–	–	Beus 1990b
	“Spathagnotus” cf. *S. gradatus*	Y	–	–	Beus 1990b

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Invertebrates: Conodonta (continued)	Ramiforms	–	–	Y	Martin and Barrick 1999
	Conodonta undetermined	–	–	Y	Hodnett and Elliott 2018
Invertebrates: Other Invertebrates	Spicules	–	–	Y	Billingsley and Beus 1999a
	Unspecified invertebrate fossils	–	–	Y	Billingsley and Beus 1985, Beus 1995
Vertebrates	**Overall Vertebrates**	**Y**	**Y**	**Y**	**–**
Vertebrates: Placodermi	*Bothriolepis coloradoensis* or *B. leidyi*	Y	–	–	Schuchert 1918b, Noble 1922, Denison 1951, Elliott and Blakey 2005
Vertebrates: Chondrichthyes	**Overall Chondrichthyes**	**–**	**Y**	**Y**	**–**
	Amaradontus santuccii	–	–	Y	Hodnett and Elliott 2018
	Cladodus cf. *C. marginatus*	–	–	Y	Hodnett and Elliott 2018
	Cochliodus cf. *C. contortus*	–	–	Y	Hodnett and Elliott 2018
	Cooleyella platera	–	–	Y	Hodnett and Elliott 2018
	Deltodus cf. *D. angularis*	–	–	Y	Hodnett and Elliott 2018
	Deltodus cf. *D. cingulatus*	–	–	Y	Hodnett and Elliott 2018
	Deltodus sp.	–	–	Y	Hodnett and Elliott 2018
	Helodus sp.	–	Y	–	McKee and Gutschick 1969b, 1969d, Elliott and Blakey 2005
	Helodus? sp.	–	–	Y	Hodnett and Elliott 2018
	cf. *Helodus* (*Psephodus*) *didymus*	–	Y	–	Chapter 6
	Heteropetalus sp.	–	–	Y	Hodnett and Elliott 2018
	Microklomax carrieae	–	–	Y	Hodnett and Elliott 2018
	Physonemus sp.	–	Y	–	Elliott and Blakey 2005
	Orodontidae indet.	–	–	Y	Hodnett and Elliott 2018
	Undetermined shark teeth	–	–	Y	Billingsley and Beus 1999a
Vertebrates: Sarcopterygii	*Holoptychius* sp.	Y	–	–	Schuchert 1918b, Elliott and Blakey 2005

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Vertebrates: Miscellaneous Fish	"Placoganoid" fish plates	Y	–	–	Walcott 1883
	Indeterminate fish plates and scales	Y	–	–	Noble 1922, Stoyanow 1936
	Fish teeth	–	Y	Y	McKee and Gutschick 1969b, 1969d, Gordon E&R 1979/1/23
Vertebrates: Other Vertebrates	Undetermined vertebrate remains	–	–	Y	Billingsley and Beus 1999a, Hodnett and Elliott 2018
Ichnofossils	**Overall Ichnofossils**	**Y**	**Y**	**Y**	**–**
Ichnofossils: Microbial Trace Fossils	Oncolites	–	–	Y	Beus 1990a, Billingsley and Beus 1999a
	Stromatolites	–	–	Y	Billingsley and Beus 1985, 1999a
Ichnofossils: Invertebrate Trace Fossils	?*Conostichnus* isp.	–	–	Y	Billingsley et al. 1999
	Cruziana isp.	–	–	Y	Billingsley et al. 1999
	"Fucoids" (including "seaweeds")	–	Y	–	Noble 1922
	Invertebrate borings and burrows	Y	–	Y	Beus 1973, Billingsley and Beus 1999a
	Worm borings and trails	Y	Y	Y	Schuchert 1918b, Stoyanow 1936, McKee E&R 1978/5/25, GCM
	Undetermined invertebrate trace fossils	–	Y	–	GCM
Ichnofossils: Other Trace Fossils	"Algal" laminations	–	–	Y	Billingsley and Beus 1999a
	General bioturbation	–	–	Y	Billingsley and Beus 1999a
	Unspecified trace fossils	–	–	Y	Billingsley and Beus 1999a
Other Fossils	**Overall Other Fossils**	**Y**	**Y**	**Y**	**–**
Other Fossils: Foraminifera	**Overall Foraminifera**	**–**	**Y**	**Y**	**–**
	Earlandia sp.	–	Y	–	McKee and Gutschick 1969b
	Endothyra kleina?	–	Y	–	McKee and Gutschick 1969b
	Endothyra tantala	–	Y	–	McKee and Gutschick 1969b
	Endothyra trachida	–	Y	–	McKee and Gutschick 1969b

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Other Fossils: Foraminifera (continued)	*Endothyra trachida*?	–	Y	–	McKee and Gutschick 1969b
	Endothyra sp.	–	Y	–	McKee and Gutschick 1969b
	Endothyra aff. *E. excellens*	–	Y	–	McKee and Gutschick 1969b
	Endothyra aff. *E. gutschicki*	–	Y	–	McKee and Gutschick 1969b
	Endothyra aff. *E. tantala*	–	Y	–	McKee and Gutschick 1969b
	Eoendothyranopsis spiroides	–	Y	–	McKee and Gutschick 1969b
	Eoendothyranopsis spiroides?	–	Y	–	McKee and Gutschick 1969b
	Eoendothyranopsis aff. *E. spiroides*	–	Y	–	McKee and Gutschick 1969b
	Eoendothyranopsis cf. *E. spiroides*	–	Y	–	McKee and Gutschick 1969b
	Globoendothyra baileyi	–	Y	–	McKee and Gutschick 1969b
	Globoendothyra baileyi poloumera	–	Y	–	McKee and Gutschick 1969b, Skipp 1969
	Globoendothyra baileyi?	–	Y	–	McKee and Gutschick 1969b
	Inflatoendothyra eospiroides	–	Y	–	McKee and Gutschick 1969b
	Paracaligella? sp.	–	Y	–	McKee and Gutschick 1969b
	Paramillerella? sp.	–	Y	–	McKee and Gutschick 1969b
	Pohlia henbesti	–	Y	–	Skipp et al. 1966, McKee and Gutschick 1969b
	Pohlia henbesti?	–	Y	–	McKee and Gutschick 1969b
	Septabrunsiina parakrainica	–	Y	–	McKee and Gutschick 1969b
	Septabrunsiina sp.	–	Y	–	McKee and Gutschick 1969b
	Septabrunsiina sp.?	–	Y	–	McKee and Gutschick 1969b
	Septaglomospiranella chernoussovensis	–	Y	–	Skipp et al. 1966, McKee and Gutschick 1969b
	Septaglomospiranella rossi	–	Y	–	McKee and Gutschick 1969b, Skipp 1969
	Septaglomospiranella sp.	–	Y	–	McKee and Gutschick 1969b
	Septatournayella? sp.	–	Y	–	McKee and Gutschick 1969b

Appendix Table C-2 (continued). Devonian and Mississippian fossil taxa. Dtb = Temple Butte Formation, Mr = Redwall Limestone, and Msc = Surprise Canyon Formation.

Category	Taxon Observed	Dtb	Mr	Msc	References
Other Fossils: Foraminifera (continued)	*Spinobrunsiina torquida*	–	Y	–	McKee and Gutschick 1969b
	Spinobrunsiina torquida?	–	Y	–	McKee and Gutschick 1969b
	Spinoendothyra spinosa	–	Y	–	McKee and Gutschick 1969b
	Spinoendothyra spinosa?	–	Y	–	McKee and Gutschick 1969b
	Spiroplectamminoides cf. *S. parva*	–	Y	–	McKee and Gutschick 1969b
	Tournayella sp.	–	Y	–	McKee and Gutschick 1969b
	Tournayella? sp.	–	Y	–	McKee and Gutschick 1969b
	Tuberendothyra paratumula	–	Y	–	McKee and Gutschick 1969b
	Tuberendothyra tuberculata	–	Y	–	McKee and Gutschick 1969b
	Tuberendothyra sp.	–	Y	–	McKee and Gutschick 1969b
	Tuberendothyra sp.?	–	Y	–	McKee and Gutschick 1969b
	Foraminifera undetermined	–	–	Y	Gordon E&R 1979/1/23
Other Fossils: Miscellaneous	**Overall Miscellaneous**	**Y**	**Y**	**Y**	**–**
	“Algae”	–	Y	–	White 1927, 1928
	“Algal” concretions	–	–	Y	Billingsley and Beus 1999a
	Calcisphaera sp.	–	Y	–	McKee and Gutschick 1969b
	Undetermined calcispheres	–	Y	–	McKee and Gutschick 1969b
	Flaring vertical structures	–	–	Y	Billingsley and Beus 1999a
	Reworked Redwall Limestone fossils	–	–	Y	Billingsley and Beus 1999a
	Twig-like fossils	–	–	Y	Billingsley and Beus 1999a
	Trace fossils or stromatoporoid sponges	Y	–	–	Beus 1973, 1990b
	Unspecified or unidentifiable fossils	Y	Y	Y	McKee 1969, McKee and Gutschick 1969a, 1969c, Billingsley and Beus 1999a

Appendix Table C-3. Supai Group fossil taxa. IPswa = Watahomigi Formation, IPsm = Manakacha Formation, IPswe = Wescogame Formation, Pse = Esplanade Sandstone, and PIPsu = Undivided Supai Group.

Category	Taxon Observed	IPswa	IPsm	IPswe	Pse	References
Plants	**Plants**	**Y**	**Y**	**Y**	**Y**	**–**
Plants: Equisetopsida (horsetails)	*Calamites* sp.	Y	–	–	–	White 1929
Plants: Polypodiopsida (ferns)	Polypodiopsida undetermined	–	Y	–	–	McKee 1982a, 1982b
Plants: “Seed Ferns”	*Neuropteris* sp.	Y	–	–	–	White 1929
Plants: Pinopsida (conifers)	*Cordaites* sp.	Y	–	–	–	White 1929
	Walchia sp.	Y	–	–	Y	White 1929, McKee 1982a, 1982b
	?*Walchia* sp. stems	Y	–	–	–	McKee 1982a
Plants: Other Plants	*Taeniopteris* sp.	Y	–	–	–	White 1929
	“Fern or cycadofilice” fragments	–	–	Y	–	Schuchert 1918b, McKee 1982b (record possibly from Esplanade instead)
	Plant stems	–	–	–	Y	McKee 1982a, 1982b
	Undetermined plants	Y	Y	–	Y	McKee 1982a, 1982b
Invertebrates	**Invertebrates**	**Y**	**Y**	**Y**	**Y**	**–**
Invertebrates: Cnidaria: Anthozoa (corals)	Coral bioclasts	–	–	–	Y	McKee 1982b
	Anthozoa undetermined	Y	–	–	–	McKee 1982a
Invertebrates: Cnidaria: Anthozoa: Tabulata (tabulate corals)	*Michelinia* sp.	Y	–	–	–	McKee 1982b, Gordon 1982
Invertebrates: Cnidaria?: Conulariida	Conulariida undetermined	Y	–	–	–	McKee 1982b, Gordon 1982

Appendix Table C-3 (continued). Supai Group fossil taxa. IPswa = Watahomigi Formation, IPsm = Manakacha Formation, IPswe = Wescogame Formation, Pse = Esplanade Sandstone, and PIPsu = Undivided Supai Group.

Category	Taxon Observed	IPswa	IPsm	IPswe	Pse	References
Invertebrates: Bryozoa (moss animals)	**Overall Bryozoa**	**Y**	**Y**	**Y**	**Y**	**–**
	Cystiodictya? sp.	Y	–	–	–	McKee 1982b, Gordon 1982
	Fenestella sp.	Y	–	–	–	McKee 1982b, Gordon 1982
	Encrusting fistuliporoid	Y	–	–	–	McKee 1982b, Gordon 1982
	Encrusting stenoporoid	Y	–	–	–	McKee 1982b, Gordon 1982
	Ramose trepostome	Y	–	–	–	McKee 1982b, Gordon 1982
	Indeterminate rhomboporoid	Y	–	–	–	McKee 1982b, Gordon 1982
	Bryozoan bioclasts	Y	Y	Y	Y	McKee 1982b
	Bryozoa undetermined	Y	–	–	–	McKee 1982a, Billingsley and Beus 1999a
Invertebrates: Brachiopoda (lamp shells)	**Overall Brachiopoda**	**Y**	**–**	**–**	**–**	**–**
	Anthracospirifer newberryi	Y	–	–	–	McKee 1982b, Gordon 1982
	Anthracospirifer tanoensis	Y	–	–	–	McKee 1982b, Gordon 1982
	Composita ovata	Y	–	–	–	McKee 1982b, Gordon 1982
	Composita subtilita	Y	–	–	–	McKee 1938, 1982b, Gordon 1982
	Composita sp.	Y	–	–	–	McKee 1982a, 1982b, Gordon 1982
	Rhynchonelloid cf. *Cupularostrum* sp.	Y	–	–	–	Gordon 1982
	Derbyia sp. cf. *D. robusta*	Y	–	–	–	Gordon 1982
	Derbyia sp.	Y	–	–	–	McKee 1982a, 1982b
	Lingula sp.	Y	–	–	–	McKee 1982b, Gordon 1982
	Lingula sp.?	Y	–	–	–	McKee 1982a
	Orbiculoidea meekana	Y	–	–	–	McKee 1982b, Gordon 1982
	Orbiculoidea meekana?	Y	–	–	–	Gordon 1982
	Orbiculoidea sp.	Y	–	–	–	McKee 1982a
	Orbiculoidea sp.?	Y	–	–	–	GCM
	Orthotetes sp. A	Y	–	–	–	McKee 1982b, Gordon 1982

Appendix Table C-3 (continued). Supai Group fossil taxa. IPswa = Watahomigi Formation, IPsm = Manakacha Formation, IPswe = Wescogame Formation, Pse = Esplanade Sandstone, and PIPsu = Undivided Supai Group.

Category	Taxon Observed	IPswa	IPsm	IPswe	Pse	References
Invertebrates: Brachiopoda (lamp shells) (continued)	*Punctospirifer transversus*	Y	–	–	–	McKee 1982b, Gordon 1982
	Reticulariina gonionota	Y	–	–	–	McKee 1982b
	?*Reticulariina gonionota*	Y	–	–	–	Gordon 1982
	Schizophoria altirostris	Y	–	–	–	McKee 1982b
	Schizophoria altirostris?	Y	–	–	–	Gordon 1982
	?*Schizophoria altirostris*	Y	–	–	–	Gordon 1982
	Spirifer sp.	Y	–	–	–	McKee 1982a
	Spiriferina sp.	Y	–	–	–	McKee 1982a
	Productida undetermined	Y	–	–	–	McKee 1982a, 1982b
	Brachiopoda undetermined	Y	–	–	–	Billingsley and Beus 1999a
Invertebrates: Mollusca	**Overall Mollusca**	**Y**	**–**	**–**	**–**	**–**
Invertebrates: Bivalvia (clams, oysters, etc.)	**Overall Bivalvia**	**Y**	**–**	**–**	**–**	**–**
	Aviculopecten gravidus	Y	–	–	–	McKee 1982b, Gordon 1982
	Aviculopecten gravidus?	Y	–	–	–	Gordon 1982
	Aviculopecten sp. A	Y	–	–	–	Gordon 1982
	Aviculopecten sp.	Y	–	–	–	McKee 1982b
	Leptodesma sp.	Y	–	–	–	McKee 1982b, Gordon 1982
	Myalina cuneiformis	Y	–	–	–	GCM
	Myalina perattenuata	Y	–	–	–	GCM
	Myalina permiana	Y	–	–	–	GCM
	Myalina sp.	Y	–	–	–	McKee 1982b
	Myalina (*Myalina*) sp. B	Y	–	–	–	Gordon 1982
	Oriocrassatella sp.	Y	–	–	–	Gordon 1982
	Permophorus sp.	Y	–	–	–	McKee 1982b

Appendix Table C-3 (continued). Supai Group fossil taxa. IPswa = Watahomigi Formation, IPsm = Manakacha Formation, IPswe = Wescogame Formation, Pse = Esplanade Sandstone, and PIPsu = Undivided Supai Group.

Category	Taxon Observed	IPswa	IPsm	IPswe	Pse	References
Invertebrates: Bivalvia (clams, oysters, etc.) (continued)	*Promytilus* sp.	Y	–	–	–	McKee 1982b, Gordon 1982
	Schizodus sp. A	Y	–	–	–	Gordon 1982
	Myalinidae undetermined	Y	–	–	–	McKee 1982b, Gordon 1982
Invertebrates: Mollusca: Gastropoda (snails)	*Euomphalus* sp.	Y	–	–	–	Gordon 1982
	Straparollus sp.	Y	–	–	–	McKee 1982b
	High spired gastropod	Y	–	–	–	Gordon 1982, Billingsley and Beus 1999a
	Bellerophontoidea undetermined	Y	–	–	–	McKee 1982b, Gordon 1982
	Pleurotomarioidea undetermined	Y	–	–	–	McKee 1982b
	Gastropoda undetermined	Y	–	–	–	McKee 1982a
Invertebrates: Arthropoda: Trilobita	*Paladin* sp.	Y	–	–	–	McKee 1982b, Gordon 1982
	Paladin? sp.	Y	–	–	–	Gordon 1982
	Trilobite free cheeks (librogenae)	Y	–	–	–	McKee 1982b, Gordon 1982
Invertebrates: Arthropoda: Ostracoda (seed shrimps)	Ostracode bioclasts	–	Y	–	–	McKee 1982b
Invertebrates: Echinodermata	Pelmatozoan (stalked echinoderm) bioclasts	Y	Y	Y	Y	McKee 1982b
Invertebrates: Echinodermata: Echinoidea (sea urchins)	Echinoidea undetermined	Y	–	–	–	Gordon 1982
Invertebrates: Conodonta	*Adetognathus lautus*	Y	–	–	–	Martin and Barrick 1999
	Adetognathus spathus	Y	–	–	–	Martin and Barrick 1999
	Adetognathus sp.	Y	–	–	–	Martin and Barrick 1999
	Gnathodus sp.	Y	–	–	–	Martin and Barrick 1999
	Rhachistognathus muricatus	Y	–	–	–	Martin and Barrick 1999

Appendix Table C-3 (continued). Supai Group fossil taxa. IPswa = Watahomigi Formation, IPsm = Manakacha Formation, IPswe = Wescogame Formation, Pse = Esplanade Sandstone, and PIPsu = Undivided Supai Group.

Category	Taxon Observed	IPswa	IPsm	IPswe	Pse	References
Invertebrates: Conodonta (continued)	*Rhachistognathus websteri*	Y	–	–	–	Martin and Barrick 1999
	Ramiforms	Y	–	–	–	Martin and Barrick 1999
Invertebrates: Other Invertebrates	Bivalve or brachiopod bioclasts	Y	Y	Y	Y	McKee 1982b
	Marine fossil clasts	–	–	Y	–	McKee 1982b
	Unspecified invertebrate fossils	Y	Y	Y	Y	McKee 1982a
Vertebrates	**Overall Vertebrates**	**Y**	**–**	**–**	**–**	**–**
Vertebrates: Chondrichthyes	*Deltodus mercurii*	Y	–	–	–	Gordon 1982
	Deltodus sp.	Y	–	–	–	McKee 1982b, Gordon 1982
Vertebrates: Miscellaneous Fish	Fish teeth	Y	–	–	–	McKee 1982a
Ichnofossils	**Overall Ichnofossils**	**Y**	**Y**	**Y**	**Y**	**–**
Ichnofossils: Microbial Trace Fossils	*Rivularites* (as microbially induced sedimentary structure, not taxon)	Y	Y	–	Y	White 1929, MicKee 1982a, 1982b
	Stromatolites	Y	Y	–	–	Rawson and Turner 1974, McKee 1982a, 1982b
	Possible stromatolitic laminations	Y	–	–	–	McKee 1982a
Ichnofossils: Invertebrate Trace Fossils	*Diplichnites* isp.	–	–	Y	–	Gilmore 1928 (identified in Chapter 8)
	Helminthopsis isp.	–	–	Y	–	Gilmore 1928 (identified in Chapter 8)
	Horseshoe-crab-like invertebrate tracks	–	Y	–	–	McKee 1982a
	Unnamed invertebrate tracks (two types)	–	–	Y	–	McKee 1982b
	Worm borings and trails	Y	Y	Y	Y	McKee 1982a, 1982b
Ichnofossils: Vertebrate Trace Fossils	cf. *Amphisauropus* isp.	–	–	Y	–	Gilmore 1928, Haubold 1971, McKee 1982b, Santucci and Hunt 1998, Chapter 9
	Batrachichnus isp.	–	–	Y	–	Chapter 9

Appendix Table C-3 (continued). Supai Group fossil taxa. IPswa = Watahomigi Formation, IPsm = Manakacha Formation, IPswe = Wescogame Formation, Pse = Esplanade Sandstone, and PIPsu = Undivided Supai Group.

Category	Taxon Observed	IPswa	IPsm	IPswe	Pse	References
Ichnofossils: Vertebrate Trace Fossils (continued)	cf. *Limnopus* isp.	–	–	Y	–	Gilmore 1927, Haubold 1971, McKee 1982b, Santucci and Hunt 1998, Santucci et al. 1998, Marchetti et al. 2019, Chapter 9
	Varanopus isp.	–	–	Y	–	Chapter 9
	cf. *Varanopus* isp.	–	–	Y	–	Gilmore 1927, Haubold 1971, McKee 1982b, Santucci and Hunt 1998, Santucci et al. 1998, Chapter 9
	Undetermined tetrapod tracks ("cf. *Chelichnus* isp.")	–	Y	–	–	Rowland 2017, Rowland and Caputo 2018, Chapter 9
	Undetermined vertebrate tracks	–	–	Y	–	Schuchert 1918b, Gilmore 1927, Haubold 1971, McKee 1982a, 1982b, Santucci and Hunt 1998, Santucci et al. 1998, Marchetti et al. 2019, Chapter 9
Ichnofossils: Other Trace Fossils	Unspecified trace fossils	–	–	–	Y	McKee 1982a
Other Fossils	**Overall Other Fossils**	**Y**	**Y**	**Y**	**Y**	**–**
Other Fossils: Foraminifera	**Overall Subphylum Foraminifera**	**Y**	**Y**	**Y**	**Y**	**–**
	Endothyra media	–	Y	–	–	McKee 1982b (record possibly from Wescogame instead)
Other Fossils: Foraminifera (continued	*Endothyra teres*	–	Y	–	–	McKee 1982b (record possibly from Wescogame instead)
	Endothyra sp.	–	Y	–	–	McKee 1982c
	Eoschubertella sp.	–	Y	–	–	McKee 1982a, 1982b
	Fusulinella sp.	Y	Y	–	–	McKee 1982a, 1982b
	Pseudostaffella sp.	Y	Y	–	–	McKee 1982a, 1982b
	Schubertella sp.	–	–	–	Y	McKee 1982b
	Fusulinida undetermined	–	Y	Y	–	McKee 1982a, 1982c
	Foraminifera bioclasts	–	Y	Y	Y	McKee 1982b

Appendix Table C-3 (continued). Supai Group fossil taxa. IPswa = Watahomigi Formation, IPsm = Manakacha Formation, IPswe = Wescogame Formation, Pse = Esplanade Sandstone, and PIPsu = Undivided Supai Group.

Category	Taxon Observed	IPswa	IPsm	IPswe	Pse	References
Other Fossils: Miscellaneous	**Overall Miscellaneous**	**Y**	**Y**	**Y**	**–**	**–**
	"Algal" laminations	Y	–	–	–	Billingsley and Beus 1999a
	Miscellaneous "algae"	Y	–	Y	–	McKee 1982b
	Probable "algal" limestone nodules	–	Y	–	–	McKee 1982a
	Possible "algal" features	–	–	Y	–	White 1927
	Calcisphere bioclasts	–	Y	–	–	McKee 1982b
	Girvanella bioclasts	–	Y	–	–	McKee 1982b
	Possible organic features	–	–	Y	–	McKee 1982a
	Unspecified fossils	Y	–	–	–	McKee 1982a, Billingsley and Beus 1999a

Appendix Table C-4. Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Plants	**Overall Plants**	**Y**	**–**	**–**	**Y**	**–**
Plants: Ulvophyceae	Dasycladaceae undetermined	–	–	–	Y	GCM
Plants: Equisetopsida (horsetails)	*Sphenophyllum gilmorei*	Y	–	–	–	White 1929
	Sphenophyllum gilmorei?	Y	–	–	–	GCM
	Sphenophyllum sp.	Y	–	–	–	GCM
	Equisetopsida undetermined	Y	–	–	–	GCM
Plants: "Seed Ferns"	**Overall "Seed ferns"**	**Y**	**–**	**–**	**–**	**–**
	Auritifolia anomala	Y	–	–	–	White 1929
	Autunia conferta	Y	–	–	–	White 1929
	Brongniartites sp.	Y	–	–	–	GCM

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Plants: "Seed Ferns" (continued)	*Brongniartites*? *aliena*	Y	–	–	–	White 1929
	Brongniartites? *aliena*?	Y	–	–	–	GCM
	Brongniartites? *yakiensis*	Y	–	–	–	White 1929
	Gracilopteris raymondii	Y	–	–	–	White 1929
	Neuropteridium? sp.	Y	–	–	–	White 1929
	Rhachiphyllum sp.	Y	–	–	–	White 1929
	Rhachiphyllum sp.?	Y	–	–	–	GCM
	Rhachiphyllum? sp.	Y	–	–	–	White 1929
	Supaia breviloba	Y	–	–	–	White 1929
	Supaia compacta	Y	–	–	–	White 1929
	Supaia linearifolia	Y	–	–	–	White 1929
	Supaia merriami	Y	–	–	–	White 1929
	Supaia merriami?	Y	–	–	–	GCM
	Supaia rigida	Y	–	–	–	White 1929
	Supaia sturdevantii	Y	–	–	–	White 1929
	Supaia subgoepperti	Y	–	–	–	White 1929
	Supaia thinnfeldioides	Y	–	–	–	White 1929
	Supaia sp.	Y	–	–	–	White 1929
	Supaia? sp.	Y	–	–	–	White 1929
	Yakia heterophylla	Y	–	–	–	White 1929
	Yakia heterophylla?	Y	–	–	–	GCM
	Yakia sp.	Y	–	–	–	GCM
	Yakia sp.?	Y	–	–	–	GCM
	Seed-bearing pteridosperm frond	Y	–	–	–	White 1929
	Undetermined seed ferns	Y	–	–	–	GCM

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Plants: Gingoopsida	*Psygmophyllum*? sp.	Y	–	–	–	White 1929
	Rhipidopsis sp.?	Y	–	–	–	GCM
Plants: Pinopsida (conifers)	**Overall Pinopsida**	**Y**	**–**	**–**	**–**	**–**
	Brachyphyllum arizonicum	Y	–	–	–	White 1929
	Brachyphyllum tenue	Y	–	–	–	White 1929
	Feysia sp.	Y	–	–	–	White 1929
	Hermitia dawsoni	Y	–	–	–	White 1929
	Hermitia sp.	Y	–	–	–	White 1929
	Otovicia hypnoides	Y	–	–	–	White 1929
	Pagiophyllum dubium	Y	–	–	–	White 1929
	Pagiophyllum sp.?	Y	–	–	–	GCM
	Paleotaxites praecursor	Y	–	–	–	White 1929
	Paleotaxites sp.	Y	–	–	–	GCM
	Paleotaxites sp.?	Y	–	–	–	GCM
	Taxites? sp.	Y	–	–	–	White 1929
	Voltzia dentiloba	Y	–	–	–	White 1929
	Voltzia sp.	Y	–	–	–	White 1929
	Walchia piniformis	Y	–	–	–	White 1929
	Walchia sp.	Y	–	–	–	GCM
	Walchia sp.?	Y	–	–	–	GCM
	Walchiostrobus sp.	Y	–	–	–	GCM
	Pinopsida undetermined	Y	–	–	–	GCM
Plants: Other Plants	**Overall Other Plants**	**Y**	**–**	**–**	**–**	**–**
	Carpolithus sp.	Y	–	–	–	White 1929

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Plants: Other Plants (continued)	*Cyclocarpon angelicum*	Y	–	–	–	White 1929
	Cyclocarpon sp.	Y	–	–	–	White 1929
	Eltovaria bursiformis	Y	–	–	–	White 1929
	Taeniopteris angelica	Y	–	–	–	White 1929
	Taeniopteris coriacea	Y	–	–	–	White 1929
	Taeniopteris eckhardti	Y	–	–	–	GCM
	Taeniopteris cf. *T. eckhardti*	Y	–	–	–	White 1929
	“Gymnospermous ament”	Y	–	–	–	White 1929
	Undetermined plants	Y	–	–	–	White 1929
Invertebrates	**Overall Invertebrates**	**Y**	**–**	**Y**	**Y**	**–**
	Invertebrata undetermined	–	–	–	Y	GCM
Invertebrates: Porifera (sponges)	*Actinocoelia maeandrina*	–	–	–	Y	Griffin 1966
	Actinocoelia sp.	–	–	–	Y	Thompson 1995, Thayer 2009
	Hexactinellida undetermined	–	–	–	Y	GCM
	Porifera undetermined	–	–	–	Y	Schuchert 1918b, McKee 1938, Sorauf and Billingsley 1991
	Porifera undetermined?	–	–	–	Y	GCM
Invertebrates: Cnidaria: Anthozoa (corals)	Anthozoa undetermined	–	–	–	Y	GCM
	Anthozoa undetermined?	–	–	–	Y	GCM
Invertebrates: Cnidaria: Anthozoa: Rugosa (rugose corals)	*Lophophyllum* sp.	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Lophophyllum sp.?	–	–	–	Y	GCM
	Undetermined horn corals	–	–	–	Y	McKee 1938
Invertebrates: Cnidaria?: Conulariida	*Conularia kaibabensis*	–	–	–	Y	McKee 1935, Sinclair 1948, Spamer 1984

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Invertebrates: Bryozoa (moss animals)	**Overall Bryozoa**	**–**	**–**	**Y**	**Y**	**–**
	Bascomella subsphaerica	–	–	–	Y	Condra and Elias 1944
	Bicorbis arizonica	–	–	–	Y	Condra and Elias 1945a, 1945b, McKinney 1983
	Chasmatopora sp.	–	–	–	Y	GCM
	Fistulipora n. sp.	–	–	–	Y	McKee 1938
	Fistulipora sp.	–	–	–	Y	McKinney 1983
	Girtypora maculata	–	–	–	Y	McKinney 1983
	Meekopora parilis	–	–	–	Y	McKinney 1983
	Meekopora n. sp.	–	–	–	Y	McKee 1938
	Meekopora sp.	–	–	–	Y	GCM
	Polypora spinulifera	–	–	–	Y	McKee 1938
	Polypora spinulifera?	–	–	–	Y	GCM
	Polypora sp.	–	–	–	Y	GCM
	Rhabdomeson sp.	–	–	–	Y	McKinney 1983
	Rhipidomella hessensis	–	–	–	Y	Condra and Elias 1944
	Rhombopora lepidodendroides	–	–	–	Y	McKee 1938
	Septopora biserialis	–	–	–	Y	McKee 1938
	Septopora n. sp. #2	–	–	–	Y	McKee 1938
	Septopora sp.	–	–	–	Y	GCM
	?*Stenodiscus* sp.	–	–	–	Y	McKinney 1983
	Stenopora n. sp. #1	–	–	–	Y	McKee 1938
	Stenopora n. sp. #2	–	–	–	Y	McKee 1938
	Stenopora sp.	–	–	–	Y	GCM
	Streblotrypa sp.	–	–	–	Y	McKinney 1983

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Invertebrates: Bryozoa (moss animals) (continued)	Undetermined fenestellid	–	–	–	Y	McKinney 1983
	Bryozoa undetermined	–	–	Y	Y	McKee 1938, Sorauf and Billingsley 1991, Thompson 1995
Invertebrates: Brachiopoda (lamp shells)	**Overall Brachiopoda**	**–**	**–**	**Y**	**Y**	**–**
	Anomaloria sp., *Neophricadothyris* sp., or *Squamularia* sp.	–	–	–	Y	GCM
	Avonia sp.	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Bathymyonia nevadensis	–	–	–	Y	McKee 1938
	Camarophoria sp.	–	–	–	Y	GCM
	Camarotoechia sp.	–	–	–	Y	GCM
	Chonetes sp.	–	–	–	Y	McKee 1938
	Chonetes sp.?	–	–	–	Y	GCM
	Composita arizonica	–	–	–	Y	McKee 1938
	Composita arizonica var.	–	–	–	Y	GCM
	Composita cf. *C. ovata*	–	–	–	Y	GCM
	Composita subtilita	–	–	–	Y	McKee 1938, 1982b, Gordon 1982
	Composita sp.	–	–	Y	Y	McKee 1938, Sorauf and Billingsley 1991
	cf. *Composita* sp.	–	–	–	Y	McKee 1938
	Derbyia arizonensis	–	–	–	Y	McKee 1938, 1941
	Derbyia multistriata	–	–	–	Y	GCM
	Derbyia nasuta	–	–	–	Y	McKee 1938
	Derbyia sp.	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991, Thayer 2009
	Derbyia sp.?	–	–	–	Y	GCM
	Dielasma phosphoriensis	–	–	–	Y	McKee 1938
	Dielasma sp.	–	–	–	Y	GCM

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Invertebrates: Brachiopoda (lamp shells) (continued)	*Dyoros* aff. *D. hillanus*	–	–	–	Y	GCM
	Dyoros kaibabensis	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Dyoros subliratus	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Dyoros tetragonus	–	–	–	Y	King 1931, Cooper and Grant 1975
	Echinauris dorsoconcava	–	–	–	Y	McKee 1938
	Echinauris subhorrida?	–	–	–	Y	GCM
	Echinauris newberryi	–	–	–	Y	McKee 1938
	Echinauris sp.	–	–	–	Y	Thompson 1995
	Hustedia meekana	–	–	–	Y	McKee 1938
	Hustedia sp.	–	–	–	Y	Thompson 1995
	Kozlowskia sp.	–	–	–	Y	Thompson 1995
	Kutorginella meridionalis	–	–	–	Y	McKee 1938
	Liosotella popei	–	–	–	Y	McKee 1938
	Marginifera sp.	–	–	–	Y	McKee 1938
	Meekella occidentalis	–	–	–	Y	GCM
	Meekella pyramidalis	–	–	–	Y	Schuchert 1918b, McKee 1938
	Meekella sp.	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991, Thompson 1995, Thayer 2009
	Neophricadothyris sp.	–	–	–	Y	Thompson 1995
	Neophricadothyris sp., or *Rugatia occidentalis*, or *Rugatia paraindicus*	–	–	–	Y	GCM
	Orbiculoidea sp.	–	–	–	Y	GCM
	Orthotetes sp.	–	–	–	Y	GCM
	Peniculauris bassi	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Peniculauris bassi or *Reticulatia* sp.	–	–	–	Y	GCM
	Peniculauris ivesi	–	–	Y	Y	Schuchert 1918b, McKee 1938

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Invertebrates: Brachiopoda (lamp shells) (continued)	*Peniculauris* sp.	–	–	–	Y	Thompson 1995, Thayer 2009
	Phrenophoria pinguis	–	–	–	Y	McKee 1938
	Phricodothyris guadalupensis	–	–	–	Y	McKee 1938
	Productus sp.	–	–	–	Y	GCM
	Pugnax sp.	–	–	–	Y	GCM
	Pugnoides sp.	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Punctospirifer? sp.	–	–	–	Y	McKee 1938
	Rhipidomella transversa	–	–	–	Y	McKee 1938
	Rhynchonella sp.	–	–	–	Y	GCM
	Rugatia occidentalis	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Rugatia paraindicus	–	–	–	Y	McKee 1938
	Rugatia sp.	–	–	–	Y	Thompson 1995
	Spirifer sp.	–	–	–	Y	GCM
	Spiriferellina hilli	–	–	–	Y	McKee 1938
	Spiriferina sp.	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Squamularia sp.	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Waagenoconcha irginae	–	–	–	Y	GCM
	Waagenoconcha sp.	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Wellerella osagensis	–	–	–	Y	GCM
	Productida undetermined	–	–	–	Y	Thompson 1995
	Brachiopoda undetermined	–	–	–	Y	McKee 1938, Thompson 1995
	Brachiopoda undetermined?	–	–	–	Y	GCM
Invertebrates: Mollusca	**Overall Mollusca**	**–**	**–**	**Y**	**Y**	**–**
	Mollusca undetermined	–	–	Y	Y	McKee 1938, Thompson 1995
Invertebrates: Mollusca: Bivalvia (clams, oysters, etc.)	**Overall Bivalvia**	**–**	**–**	**Y**	**Y**	**–**
	Acanthopecten coloradoensis	–	–	–	Y	Newell 1937

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Invertebrates: Mollusca: Bivalvia (clams, oysters, etc.) (continued)	*Allorisma capax*	–	–	–	Y	GCM
	Allorisma sp.	–	–	Y	–	McKee 1938
	Astartella gurleyi	–	–	–	Y	Shimer 1919, McKee 1938
	Astartella sp.	–	–	–	Y	GCM
	Aviculopecten sp.	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991
	Bakewellia parva	–	–	–	Y	Shimer 1919, McKee 1938
	Deltopecten coreyanus	–	–	–	Y	GCM
	Deltopecten coreyanus var.	–	–	–	Y	GCM
	Deltopecten sp.	–	–	Y	Y	GCM
	Edmondia sp.	–	–	Y	–	McKee 1938
	Modiomorpha sp.	–	–	–	Y	GCM
	Nuculana? sp.	–	–	Y	Y	McKee 1938, GCM
	Pecten sp.	–	–	–	Y	GCM
	Permophorus sp.	–	–	Y	Y	McKee 1938, GCM
	Pseudomonotis sp.	–	–	–	Y	GCM
	Pteria sp.	–	–	Y	Y	McKee 1938, GCM
	Pterinopecten sp.	–	–	–	Y	GCM
	Schizodus canalis	–	–	–	Y	GCM
	Schizodus sp.	–	–	Y	Y	McKee 1938, GCM
	Schizodus? sp.	–	–	Y	Y	McKee 1938
	Schizodus sp.?	–	–	–	Y	GCM
	Solenomya naenia	–	–	–	Y	GCM
	Solenomya trapezoides	–	–	–	Y	GCM
	Solenomya trapezoides?	–	–	–	Y	GCM
	Solenomya sp.	–	–	Y	Y	McKee 1938, GCM
	Bivalvia undetermined	–	–	Y	Y	McKee 1938

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Invertebrates: Mollusca: Cephalopoda	Cephalopoda undetermined	–	–	–	Y	GCM
Invertebrates: Mollusca: Cephalopoda: Nautiloidea	**Overall Nautiloidea**	**–**	**–**	**Y**	**Y**	**–**
	Domatoceras bradyi	–	–	–	Y	Miller and Unklesbay 1942, Miller and Youngquist 1949
	Domatoceras simplex	–	–	–	Y	McKee 1938
	Domatoceras? sp.	–	–	Y	–	McKee 1938
	Orthoceras sp.	–	–	–	Y	McKee 1938
	Stearoceras rotundatum?	–	–	–	Y	Miller and Unklesbay 1942, Miller and Youngquist 1949
	Stearoceras sanandreasense?	–	–	–	Y	Miller and Unklesbay 1942, Miller and Youngquist 1949
	Stearoceras sp.	–	–	–	Y	GCM
	Stearoceras sp.?	–	–	–	Y	GCM
Invertebrates: Mollusca: Gastropoda (snails)	**Overall Gastropoda**	**–**	**–**	**Y**	**Y**	**–**
	Aclisina sp.	–	–	Y	–	McKee 1938
	Bellerophon majusculus	–	–	–	Y	McKee 1938
	Bellerophon sp.	–	–	Y	–	McKee 1938
	Euomphalus kaibabensis	–	–	–	Y	GCM
	Euomphalus sp.	–	–	Y	Y	McKee 1938, GCM
	Euomphalus sp.?	–	–	–	Y	GCM
	Euphemites sp.	–	–	Y	Y	McKee 1938
	Euphemites cf. *E. carbonarius*	–	–	–	Y	Shimer 1919, McKee 1938
	Goniospira sp.	–	–	Y	Y	McKee 1938
	Murchisonia? cf. *M terebra*	–	–	–	Y	Shimer 1919, McKee 1938
	Naticopsis sp.	–	–	Y	Y	McKee 1938

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Invertebrates: Mollusca: Gastropoda (snails) (continued)	*Pleurotomaria sp.*	–	–	–	Y	McKee 1938
	Gastropoda undetermined	–	–	Y	Y	McKee 1938
	Gastropoda undetermined?	–	–	–	Y	GCM
Invertebrates: Mollusca: Scaphopoda (tusk shells)	*Dentalium* sp.	–	–	Y	Y	GCM
	Plagioglypta sp.	–	–	–	Y	GCM
	Prodentalium canna	–	–	–	Y	Shimer 1919, McKee 1938
	Scaphopoda undetermined	–	–	–	Y	GCM
Invertebrates: Arthropoda	**Overall Arthropoda**	**Y**	**–**	**Y**	**Y**	**–**
Invertebrates: Arthropoda: Trilobita	*Ameura* sp.	–	–	–	Y	GCM
	Delaria sevilloidia	–	–	–	Y	Cisne 1971
	Delaria snowi	–	–	–	Y	Cisne 1971
	Ditomopyge scitulus	–	–	–	Y	GCM
	Ditomopyge sp.	–	–	–	Y	McKee 1938, Sorauf and Billingsley 1991, Thayer 2009
	Novoameura mckeei	–	–	–	Y	Cisne 1971
	Trilobita undetermined	–	–	–	Y	McKee 1938, Snow 1945, Sorauf and Billingsley 1991
	Trilobita undetermined?	–	–	–	Y	GCM
Invertebrates: Arthropoda: Eurypterida (sea scorpions)	*Hastimima* sp.?	Y	–	–	–	White 1929
Invertebrata: Arthropoda: Ostracoda (seed shrimp)	Ostracoda undetermined	–	–	Y	–	McKee 1938
Invertebrates: Arthropoda: Insecta	*Tupus gilmorei*	Y	–	–	–	Carpenter 1927, White 1929
	Tupus whitei	Y	–	–	–	Carpenter 1928, White 1929

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Invertebrates: Arthropoda: Insecta (continued)	Blattodea undetermined	Y	–	–	–	Carpenter 1928, Spamer 1984
	Odonata undetermined	Y	–	–	–	Carpenter 1928
Invertebrates: Echinodermata	**Overall Echinodermata**	**–**	**–**	**Y**	**Y**	**–**
	Echinodermata undetermined	–	–	–	Y	GCM
	Echinodermata undetermined?	–	–	–	Y	GCM
Invertebrates: Echinodermata: Crinoidea (sea lilies)	Crinoidea undetermined (star-shape plates)	–	–	–	Y	GCM
	Crinoidea undetermined	–	–	Y	Y	Shimer 1919, McKee 1938, Sorauf and Billingsley 1991, Thompson 1995, Thayer 2009
Invertebrates: Echinodermata: Echinoidea (sea urchins)	*Archaeocidaris ornatus*	–	–	–	Y	GCM
	Archaeocidaris sp.	–	–	Y	Y	McKee 1938, Sorauf and Billingsley 1991
	Echinoidea undetermined	–	–	–	Y	GCM
Invertebrates: Conodonta	*Hindeodus excavatus*	–	–	–	Y	Thompson 1995
	Neostreptognathus newelli	–	–	–	Y	Thompson 1995
	Neostreptognathus prayi	–	–	–	Y	Thompson 1995
	Neostreptognathus ruzhencevi	–	–	–	Y	Thompson 1995
	Neostreptognathus cf *N. tschuvashovi*	–	–	–	Y	Thompson 1995
	Sweetia festiva	–	–	–	Y	Thompson 1995
Vertebrates	**Overall Vertebrates**	**–**	**–**	**–**	**Y**	**–**
Vertebrates: Chondrichthyes	**Overall Chondrichthyes**	**–**	**–**	**–**	**Y**	**–**
	Cooleyella typicalis	–	–	–	Y	Thompson 1995
	Cooperella striatula	–	–	–	Y	Thompson 1995
	Deltodus mercurii	–	–	–	?	David 1944, Gass 1963
	Hybodus sp.	–	–	–	Y	Gass 1963, GCM

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Vertebrates: Chondrichthyes (continued)	*Kirkella typicalis*	–	–	–	Y	Thompson 1995
	Megactenopetalus kaibabanus	–	–	–	Y	David 1944, Gass 1963, Ossian 1976
	Mooreyella typicalis	–	–	–	Y	Thompson 1995
	Petrodus sp.	–	–	–	Y	Thompson 1995
	Psephodus sp. A	–	–	–	Y	Gass 1963
	Sturgeonella quinqueloba	–	–	–	Y	Thompson 1995
	Symmorium cf. *S. reniforme*	–	–	–	Y	Thompson 1995
	Symmoriid mucus membrane denticle	–	–	–	Y	Thompson 1995
	Undescribed ctenacanthoid	–	–	–	Y	Thompson 1995
	Undescribed hybodontoid	–	–	–	Y	Thompson 1995
	Hybodontiformes undetermined	–	–	–	Y	Thompson 1995
	Undetermined dermal denticle	–	–	–	Y	Thompson 1995
	Undetermined shark tooth	–	–	–	Y	McKee 1938
Vertebrates: Actinopterygii	Palaeoniscidae undetermined	–	–	–	Y	Thompson 1995
	Undetermined phyllodont	–	–	–	Y	Johnson and Zidek 1981, Thompson 1995
Vertebrates: Miscellaneous Fish	Fish teeth	–	–	–	Y	McKee 1938, Thompson 1995
Ichnofossils	**Overall Ichnofossils**	**Y**	**Y**	**Y**	**Y**	**–**
Ichnofossils: Microbial Trace Fossils	*Rivularites* (as microbially induced sedimentary structure, not taxon)	Y	–	–	–	White 1929
	Stromatolites	–	–	Y	–	Rawson and Turner 1974, McKee 1982a, 1982b
Ichnofossils: Invertebrate Trace Fossils	**Overall Invertebrate Trace Fossils**	**Y**	**Y**	**–**	**Y**	**–**
	Diplopodichnus isp.	–	Y	–	–	Chapter 8
	Octopodichnus didactylus	–	Y	–	–	Gilmore 1927
	Paleohelcura benjamini	–	Y	–	–	Gilmore 1927

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Ichnofossils: Invertebrate Trace Fossils (continued)	*Paleohelcura tridactyla*	–	Y	–	–	Gilmore 1926, 1927
	Scoyenia gracilis	Y	–	–	–	White 1929
	Sphaerapus larvalis	Y	–	–	–	White 1929, Lucas et al. 2013
	Taenidium serpentinum	–	–	–	Y	DeCourten 1978
	Unisulcus sinuosus	–	Y	–	–	Gilmore 1927
	"Fucoids"	–	–	–	Y	McKee 1932, 1938
	Gastropod trail	–	–	–	Y	GCM
	?*Teichichnus*-derived heavy bioturbation	–	–	–	Y	Thompson 1995
	Trilobite trails	–	–	–	Y	McKee 1938, GCM
	Worm borings and trails	Y	Y	–	Y	White 1929, McKee 1933, 1938, 1982a, GCM
	Unspecified invertebrate traces	–	Y	–	Y	McKee 1933, Thompson 1995
Ichnofossils: Vertebrate Trace Fossils	**Overall Vertebrate Trace Fossils**	**Y**	**Y**	**–**	**–**	**–**
	Amphisauropus kablikae	Y	–	–	–	Chapter 9
	cf. *Amphisauropus* isp.	Y	Y	–	–	Chapter 9
	Batrachichnus salamandroides	Y	–	–	–	Chapter 9
	cf. *Batrachichnus* isp.	Y	–	–	–	Chapter 9
	Dimetropus isp.	Y	–	–	–	Chapter 9
	Cf. *Dimetropus* isp.	Y	–	–	–	Chapter 9
	Dromopus lacertoides	Y	–	–	–	Chapter 9
	cf. *Dromopus* isp.	–	Y	–	–	Marchetti et al. 2019
	Erpetopus isp.	Y	Y	–	–	Chapter 9
	cf. *Erpetopus* isp.	Y	Y	–	–	Chapter 9
	Hyloidichnus bifurcatus	Y	–	–	–	Chapter 9
	cf. *Hyloidichnus* isp.	Y	–	–	–	Chapter 9
	Ichniotherium cottae	Y	–	–	–	Chapter 9

Appendix Table C-4 (continued). Post-Supai Permian fossil taxa. Ph = Hermit Formation, Pc = Coconino Sandstone, Pt = Toroweap Formation, and Pk = Kaibab Formation.

Category	Taxon Observed	Ph	Pc	Pt	Pk	References
Ichnofossils: Vertebrate Trace Fossils (continued)	*Ichniotherium gilmorei*	Y	–	–	–	Haubold 1971
	Ichniotherium sphaerodactylum	–	Y	–	–	Chapter 9
	cf. *Ichniotherium* isp.	–	Y	–	–	Chapter 9
	cf. *Tambachichnium* isp.	–	Y	–	–	Chapter 9
	Varanopus curvidactylus	–	Y	–	–	Chapter 9
	cf. *Varanopus* isp.	–	Y	–	–	Chapter 9
	Mud-puppy-like traces	Y	–	–	–	White 1929
	Undescribed vertebrate track 1	–	Y	–	–	Hunt and Santucci 2001
	Undescribed vertebrate track 2	–	Y	–	–	Santucci et al. 2006
	Undetermined vertebrate traces	Y	Y	–	–	Beus and Billingsley 1989, Chapter 9
Ichnofossils: Other Trace Fossils	Undetermined trace fossils	Y	Y	–	–	GCM
Other Fossils	Possible “algal” features	Y	–	–	–	White 1927
	Unspecified fossils	Y	–	Y	Y	Reeside and Bassler 1922, McKee 1938, Rawson and Turner 1974, GCM

Taxonomic Notes

Cambrian

[Genus] sp. and [Genus] sp. undet. as used in McKee and Resser (1945) are consolidated as [Genus] sp.

Stratigraphy for many of Walcott's records follows Resser (1945).

Stratigraphy of Schenk and Wheeler (1942) revised after McKee and Resser (1945). Similarly, Palmer E&R 1963/10/17 and Bonde et al. (2018) describe Schenk's Cambrian collections as all Bright Angel Shale, when they actually include both Bright Angel Shale and Muav Limestone collections.

Alokistocare althea = Amecephalus althea

Billingsella obscura = Nisusia obscura

Clavaspidella enucleata Resser as mentioned in Schenk and Wheeler (1942) and repeated in Bonde et al. (2018) appears to be the only use of this name, suggesting Resser had proposed it but later dropped it; the same site in Resser (1945) has *Clavaspidella* sp. (=*Athabaskia*), and it is assumed that this was Resser's final identification

Clavaspidella kanabensis = Athabaskia kanabensis

Corophioides = Diplocraterion

Dolichometopus productus (in part) = *Dolichometopus tontoensis* (in part) = *Glossopleura mckeei = Glossopleura boccar*

Dolichometopus tontoensis (in part) = *Anoria tontoensis*

Ehmaniella basilica = Proehmaniella basilica

Ehmaniella hebes = Proehmaniella hebes

Eocrinus = Gogia

Finkelnburgia noblei = Nisusia noblei

Glossopleura mckeei = Glossopleura boccar

Species of *Indiana* named in Ulrich and Bassler (1931a) moved to *Indianites* in Ulrich and Bassler (1931b)

Iphidea crenistria = Micromitra (*Paterina*) *crenistria*

Iphidea superba = Micromitra (*Paterina*) *superba*

Isopodichnus = Cruziana

Kootenia n. sp. of Schenk and Wheeler (1942) = *Kootenia schenki*

Lingulella perattenuata ("*attenuata*" of Resser 1945) = *Lingulella spatula*

Lingulepis and *Crepicephalus*/*Ptychoparia* zone of Walcott (1883, 1890) is Muav Limestone per Noble (1922)

Micromitra (*Iphidella*) *pannula* (in part) = *Dictyonina arizonaensis*

Nisusia? (*Jamesella*) *kanabensis = Nisusia kanabensis*

Obolus (*Lingulella*) *chuarensis = Obolus* (*Westonia*) *chuarensis = Lingulella chuarensis*

Obolus (*Lingulella*) *euglyphus = Obolus* (*Westonia*) *euglyphus = Lingulella euglypha*

Obolus (*Lingulella*) *lineolatus = Lingulella lineolata*

Obolus (*Lingulella*) *spatulus = Lingulella attenuata* (in part) = *Lingulella* (*Lingulepis*) *spatula = Lingulepis spatula = Lingulella spatula*

Obolus (*Lingulella*) *zetus = Lingulella zetus*

Obolus (*Westonia*) *themis = Lingulella themis*

Orthisina = Clitambonites

Phycodes pedum = Treptichnus pedum

Scalarituba = Nereites

Solenopleurella diligens = Spencella diligens

Solenopleurella erosa = Spencella erosa

Solenopleurella porcata = Spencella porcata

Solenopleurella n. sp. of Schenk and Wheeler (1941) = *Solenopleurella porcata = Spencella porcata*

Stipsellus annulatus (sometimes misspelled *Stripsellus*) = *Skolithos annulatus*

Devonian–Mississippian

Fossils reported as Chesterian at Bright Angel Trail in McKee and Gutschick (1969a) are actually Meramecian and from the Horseshoe Mesa Member of the Redwall Limestone (Billingsley and Beus 1999b).

Bellerophontaceae = Bellerophontoidea

Bothriolepis nitidens of Schuchert (1918b) and Noble (1922) accepted as *B. coloradoensis* or *B. leidyi* (Denison 1951)

Caninophyllum incrassatum = Vesiculophyllum incrassatum

Endothyra baileyi = Globoendothyra baileyi

Endothyra baileyi poloumera = Globoendothyra baileyi poloumera

Endothyra eospiroides = Inflatoendothyra eospiroides

Endothyra spinosa = Spinoendothyra spinosa

Endothyra torquida = Spinobrunsiina torquida

Eomillerella spiroides = Eoendothyranopsis spiroides

Lithostrotion (*Diphyphyllum*) *inconstans = Dorlodotia inconstans*

Monilipora? sp. = *Cladochonus* sp.

Phillipsia sampsoni = Aprathia sp.

Phillipsia tuberculata = Phillipsia peroccidens

Pugnoides osagensis = Wellerella osagensis

Schuchertella chemungensis = Floweria chemungensis

Septaglomospiranella primaeva = Septaglomospiranella chernoussovensis

Septatournayella henbesti = Pohlia henbesti

Spirifer striatus = Neospirifer striatus

Straparollus (*Euomphalus*) = *Euomphalus*

Triplophyllites (*Homalophyllites*) *paucicinctus = Homalophyllites paucicinctus*

Supai Group

Undivided Supai: the only fossils not otherwise accounted for in Supai units are cup corals and brachiopods in a limestone bed (Schuchert 1918b), thread-like algae in limestone of red shales of middle Supai (White 1929), calcareous algae (Stoyanow 1936), and the gastropod *Euomphalus* cf. *E. hollingsworthi* (Yochelson E&R 1969/6/23).

[Genus] sp. and [Genus] sp. undet. as used in White (1929) are consolidated as [Genus] sp.

Pre-1982 records from the Supai Group have been distributed according to modern stratigraphic definitions if it was possible to determine the proper usage.

Ammobatrachus turbatans = *Limnopus* isp. = undetermined tracks = cf. *Amphisauropus* isp.

Anomalopus sturdevanti = *Limnopus* isp. = undetermined tracks

Bellerophontaceae = Bellerophontoidea

Derbyia may also be spelled *Derbya* in some references

“Horse-hoof-like” features in Gilmore 1926 and McKee 1982b are pseudofossils

Pleurotomariaceae = Pleurotomarioidea

Stenichnus yakiensis = *Batrachichnus delicatulus* = cf. *Varanopus* isp.

Straparollus (*Euomphalus*) = *Euomphalus*

Tridentichnus supaiensis = *Anomalopus* (*Tridentichnus*) *supaiensis* = *Limnopus* isp. = cf. *Limnopus* isp.

Post-Supai Permian

[Genus] sp. and [Genus] sp. undet. as used in White (1929) are consolidated as [Genus] sp.

The taxonomic histories of the vertebrate ichnotaxa of the Hermit Formation and Coconino Sandstone are extremely convoluted, and the names currently in use (Chapter 9) are almost entirely different from the names introduced in the 1910s and 1920s. Because of this, Chapter 9 is listed as the reference for all such ichnotaxa. The histories of the ichnotaxa can be traced through a number of documents (Lull 1918; Noble 1922; Gilmore 1926, 1927, 1928; Haubold 1971; McKeever and Haubold 1996; Hunt and Santucci 1998; Santucci and Hunt 1998; Santucci et al. 1998; Francischini et al. 2018; Marchetti et al. 2019). Chains of synonyms are included below.

Agostopus matheri = *Chelichnus duncani* = cf. *Amphisauropus* isp.

Agostopus medius = *Chelichnus duncani* = undetermined tracks

Allopus? *arizonae* = *Baropezia arizonae* = *Chelichnus duncani* = undetermined tracks

Amblyopus pachypodus = *Chelichnus gigas* = undetermined tracks

Anisopyge mckeei = *Novoameura mckeei*

Avonia dorsoconcava = *Echinauris dorsoconcava*

Avonia subhorrida newberryi = *Echinauris newberryi*

Baropezia eakini = *Chelichnus duncani* = *Ichniotherium sphaerodactylum*

Baropus coconinoensis = *Chelichnus gigas* = *Ichniotherium sphaerodactylum*

Barypodus metszeri = *Chelichnus duncani* = cf. *Tambachichnium* isp.

Barypodus palmatus = *Chelichnus gigas* = undetermined tracks

Barypodus tridactylus = cf. *Tambachichnium* isp.

Batrachichnus obscurus = *Batrachichnus delicatulus* = cf. *Batrachichnus* isp.

Bicorbula arizonica = *Bicorbis arizonica*

Callipteris arizonae = *Rhachiphyllum* sp.

Callipteris conferta = *Autunia conferta*

Callipteris raymondii = *Gracilopteris raymondii*

Callipteris sp. = *Rhachiphyllum* sp.

Chonetes (*Lissochonetes*) *subliratus* = *Dyoros subliratus*

Chonetes hillanus = *Dyoros hillanus*

Chonetes kaibabensis = *Quadrochonetes kaibabensis* = *Dyoros kaibabensis*

Chonetes quadratus = *Dyoros tetragonus*

Coconino Sandstone "fucoidal coprolites" of McKee 1933 = invertebrate burrows (*Scolecocoprus*)

Collettosaurus pentadactylus = *Gilmoreichnus hermitanus* = cf. *Dimetropus* isp.

Cursipes sp. = *Gilmoreichnus hermitanus* = cf. *Hyloidichnus* isp.

Delaria macclintocki = *Delaria sevilloidia*

Dentalium canna = *Plagioglypta canna* = *Prodentalium canna*

Derbyia may also be spelled *Derbya* in some references

Derbyia regularis = *Derbyia arizonensis*

Dolichopodus tetradactylus = *Chelichnus bucklandi* = undetermined tracks

Dromillopus parvus = *Batrachichnus delicatulus* = *Batrachichnus salamandroides*

Echinocrinus = *Archaeocidaris* (suppressed for *Archaeocidaris*)

Endothyra spinosa = *Spinoendothyra spinosa*

Endothyra torquida = *Spinobrunsiina torquida*

Euphemus = *Euphemites*

Exocampe? *delicatula* = *Batrachichnus delicatula* = *Nanipes delicatulus* = *Batrachichnus delicatulus* = *Batrachichnus salamandroides*

Gilmoreichnus hermitanus = cf. *Hyloidichnus* isp.

Griffithides scitulus = *Ditomoyge scitulus*

Hermit Formation “coprolites” in GCM = *Sphaerapus larvalis* (via *Walpia hermitensis*)

Hyloidichnus whitei = *Hyloidichnus bifurcatus*

Hylopus hermitanus = *Palaeosauropus hermitanus* = *Gilmoreichnus hermitanus* = cf. *Hyloidichnus* isp.

Laoporus coloradoensis = *Chelichnus bucklandi* = cf. *Varanopus* isp.

Laoporus noblei of Gilmore 1926 (in part) = *Chelichnus duncani* = cf. *Varanopus* isp.

Laoporus noblei of Lull 1918 = *Chelichnus bucklandi* = cf. *Varanopus* isp.

Laoporus schucherti = *Chelichnus bucklandi* = cf. *Varanopus* isp.

Leda sp. and *Nucula* sp. of McKee 1938 = *Nuculana*? sp.

Marginifera meridionalis = *Kozlowskia meridionalis* = *Kutorginella meridionalis*

Marginifera popei = *Liosotella popei*

Megapezia? *coloradensis* = *Parabaropus coloradensis* = cf. *Amphisauropus* isp. and *Dimetropus* isp.

Mesichnium benjamini = *Paleohelcura benjamini*

Nanopus maximus = *Chelichnus duncani* = undetermined tracks = cf. *Tambachichnium* isp.

Nanopus merriami = *Chelichnus bucklandi* = cf. *Erpetopus* isp.

Palaeopus regularis = *Chelichnus duncani* = undetermined tracks

Pleurophorus = *Permophorus*

Pleurotomariaceae = Pleurotomarioidea

Productus (*Dictyoclostus*) *bassi* = *Peniculauris bassi*

Productus (*Dictyoclostus*) *ivesi* = *Dictyoclostus ivesi* = *Peniculauris ivesi*

Productus (*Dictyoclostus*) *occidentalis* = *Dictyoclostus occidentalis* = *Rugatia occidentalis*

Productus (*Dictyoclostus*) *paraindicus* = *Rugatia paraindicus*

Productus irginae = *Waagenoconcha irginae*

Productus montpelierensis = *Waagenoconcha montpelieriensis* = *Bathymyonia nevadensis*

Productus subhorridus = *Avonia subhorrida* = *Echinauris subhorrida*

Pugnax osagensis = Wellerella osagensis

Pugnoides pinguis = Wellerella pinguis = Phrenophoria pinguis

Retzia meekana = Hustedia meekana

Scolecocoprus cameronensis = Taenidium serpentinum

Spiriferina hilli = Spiriferellina hilli

Squamularia guadalupensis = Phricodothyris guadalupensis

Straparollus (*Euomphalus*) = *Euomphalus*

Titanoceras rotundatum = Stearoceras rotundatum

Titanoceras sanandreasense = Stearoceras sanandreasense

Triavestigia niningeri = Paleohelcura tridactyla

Typus gilmorei = Tupus gilmorei

Typus whitei = Tupus whitei

Ullmannia frumentaria = Feysia sp.

Walchia dawsoni = Hermitia dawsoni

Walchia gracillima = Hermitia sp.

Walchia hypnoides = Otovicia hypnoides

Walpia hermitensis = Sphaerapus larvalis

Literature Cited

Alpert, S. P. 1974. Systematic review of the genus *Skolithos*. Journal of Paleontology 48(4):661–669.

Baldwin, C. T., P. K. Strother, J. H. Beck, and E. Rose. 2004. Palaeoecology of the Bright Angel Shale in the eastern Grand Canyon, Arizona, USA, incorporating sedimentological, ichnological and palynological data. Pages 213–236 *in* D. McIlroy, editor. The application of ichnology to palaeoenvironmental and stratigraphic analysis. Geological Society, London, United Kingdom. Special Publications 228.

Beus, S. S. 1973. Devonian stratigraphy and paleogeography along the western Mogollon Rim, Arizona. Museum of Northern Arizona Bulletin 49.

Beus, S. S. 1987. Geology along the South Kaibab Trail, eastern Grand Canyon, Arizona. Pages 371–378 *in* S. S. Beus, editor. Rocky Mountain section of the Geological Society of America. Geological Society of America, Boulder, Colorado. Centennial field guide 2.

Beus, S. S. 1990a. Redwall Limestone and Surprise Canyon Formation. Pages 119–145 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology. Oxford University Press, New York, New York.

Beus, S. S. 1990b. Temple Butte Formation. Pages 107–117 *in* S. S. Beus and M. Morales, editors. Grand Canyon geology. Oxford University Press, New York, New York.

Beus, S. S. 1995. Paleontology of the Surprise Canyon Formation (Mississippian) in Grand Canyon, Arizona. Pages 25–36 *in* D. Boaz, S. Bolander, P. Dierking, M. Dornan, and B. J. Tegowski, editors. Proceedings of the third annual fossils of Arizona symposium. Proceedings - Fossils of Arizona Symposium 3.

Beus, S. S. 1999. Megafossil paleontology. Pages 69–96 *in* G. H. Billingsley and S. S. Beus, editors. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

Beus, S. S., and G. H. Billingsley. 1989. Paleozoic strata of the Grand Canyon, Arizona. Pages 122–127 *in* D. P. Elston, G. H. Billingsley, and R. A. Young, editors. Geology of Grand Canyon, northern Arizona (with Colorado River guides): Lees Ferry to Pierce Ferry, Arizona. P. M. Hanshaw. Field trips for the 28th international geological congress. American Geophysical Union, Washington, D.C.

Billingsley, G. H., and S. S. Beus. 1985. The Surprise Canyon Formation: an Upper Mississippian and Lower Pennsylvanian(?) rock unit in the Grand Canyon, Arizona. Pages A27–A33 *in* Stratigraphic notes, 1984. U.S. Geological Survey, Washington, D.C. Bulletin 1605-A.

Billingsley, G. H., and S. S. Beus, editors. 1999a. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

Billingsley, G. H., and S. S. Beus. 1999b. Overview of the Surprise Canyon Formation. Pages 1–8 *in* G. H. Billingsley and S. S. Beus, editors. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

Billingsley, G. H., and W. J. Breed. 1976. Mississippian nautiloids of the Grand Canyon. Plateau 48(3-4):67–70.

Billingsley, H., and E. D. McKee. 1982. Pre-Supai buried valleys. Pages 137–153 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

Billingsley, G. H., S. S. Beus, and P. Grover. 1999. Stratigraphy of the Surprise Canyon Formation. Pages 9–16 *in* G. H. Billingsley and S. S. Beus, editors. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

Bonde, A., V. L. Santucci, J. S. Tweet, E. Eichenberg, and B. Moore. 2018. Lake Mead National Recreation Area: Paleontological resources inventory. Natural Resource Report NPS/LAKE/NRR—2018/1618. National Park Service, Fort Collins, Colorado.

Breed, W. J. 1969. The discovery of orthocone nautiloids in the Redwall Limestone—Marble Canyon, Arizona. Page 134 *in* D. L. Baars, F. G. Bowman, Jr., and C. W. Spencer, editors. Geology and natural history of the Grand Canyon region. Four Corners Geological Society, Durango, Colorado. Field conference guidebook 5.

Brezinski, D. K. 2017. Trilobites from the Redwall Limestone (Mississippian) of Arizona. Annals of Carnegie Museum 84(2).

Buatois, L. A. 2018. *Treptichnus pedum* and the Ediacaran–Cambrian boundary: significance and caveats. Geological Magazine 155(1):174–180.

Carpenter, F. M. 1927. A fossil insect from the lower Permian of the Grand Canyon. Proceedings of the United States National Museum 71(23).

Carpenter, F. M. 1928. A new protodonatan from the Grand Canyon. Psyche 35:186–189.

Carter, J. L., D. K. Brezinski, A. D. Kollar, and J. T. Dutro, Jr. 2014. Brachiopoda taxonomy and biostratigraphy of the Redwall Limestone (Lower Mississippian) of Arizona. Annals of Carnegie Museum 82(3):257–290.

Cisne, J. L. 1971. Paleoecology of trilobites of the Kaibab limestone (Permian) in Arizona, Utah, and Nevada. Journal of Paleontology 45(3):525–533.

Condra, G. E., and M. K. Elias. 1944. Carboniferous and Permian ctenostomatous Bryozoa. Geological Society of America Bulletin 55:517–568.

Condra, G. E., and M. K. Elias. 1945a. *Bicorbula*, a new Permian bryozoan, probably a bryozoan-algal consortium. Journal of Paleontology 19(2):116–125.

Condra, G. E., and M. K. Elias. 1945b. *Bicorbis arizonica* Condra and Elias, new name for *B. arizonica*. Journal of Paleontology 19:411.

Cooper, G. A., and R. E. Grant. 1975. Permian brachiopods of west Texas, III. Smithsonian Contributions to Paleobiology 19.

David, L. R. 1944. A Permian shark from the Grand Canyon. Journal of Paleontology 18(1):90–93.

DeCourten, F. L. 1978. *Scolecocoprus cameronensis* Brady (1947) from the Kaibab Limestone of northern Arizona; a re-interpretation. Journal of Paleontology 52(2):491–493.

Denison, R. H. 1951. Late Devonian fresh-water fishes from the western United States. Fieldiana Geology 11(5):221–261.

Duncan, H. 1969. Bryozoans. Pages 345–433 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Easton, W. H., and R. C. Gutschick. 1953. Corals from the Redwall limestone (Mississippian) of Arizona. Bulletin - Southern California Academy of Sciences 52(1):1–27.

Elliott, D. K., and R. C. Blakey. 2005. The pre-Permian vertebrate record in Arizona. New Mexico Museum of Natural History and Science Bulletin 29:1–9.

Elliott, D. K., and D. L. Martin. 1987a. *Chancelloria*, an enigmatic fossil from the Bright Angel Shale (Cambrian) of Grand Canyon, Arizona. Journal of the Arizona-Nevada Academy of Science 21(2):67–72.

Elliott, D. K., and D. L. Martin. 1987b. A new trace fossil from the Cambrian Bright Angel Shale, Grand Canyon, Arizona. Journal of Paleontology 61(4):641–648.

Foster, J. R. 2011. Trilobites and other fauna from two quarries in the Bright Angel Shale (Middle Cambrian, Series 3; Delamaran), Grand Canyon National Park, Arizona. Pages 99–120 *in* J. S. Hollingsworth, F. A. Sundberg, and J. R. Foster, editors. Cambrian stratigraphy and paleontology of northern Arizona and southern Nevada. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 67.

Francischini, H., S. G. Lucas, S. Voigt, L. Marchetti, V. L. Santucci, C. L. Knight, and P. Dentzien-Dias. 2018. The first record of *Ichniotherium* in the Cisuralian Coconino Sandstone of the Grand Canyon National Park (Arizona, USA). Boletim de Resumos, XI Simpósio Brasileiro de Paleontologia de Vertebrados:52.

Frech, F. 1893. Section in Congress Canyon opposite Point Sublime. Pages 476–481 *in* Compte rendu. International Geologic Congress [Congres geologique international], 5th session. Government Printing Office, Washington, D.C.

Gass, H. L. 1963. A review of the Paleozoic fish of Arizona. Thesis. University of Arizona, Tucson, Arizona.

Gilmore, C. W. 1926. Fossil footprints from the Grand Canyon. Smithsonian Miscellaneous Collections 77(9).

Gilmore, C. W. 1927. Fossil footprints from the Grand Canyon: second contribution. Smithsonian Miscellaneous Collections 80(3).

Gilmore, C. W. 1928. Fossil footprints from the Grand Canyon: third contribution. Smithsonian Miscellaneous Collections 80(8).

Gordon, M., Jr. 1973/01/03. OER-72-5. USGS Internal Memo.

Gordon, M., Jr. 1979/01/23. O&G-78-10. USGS Internal Memo.

Gordon, M., Jr. 1982. Biostratigraphy of the Watahomigi Formation. Pages 113–135 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

Grant, R. E. 1963/10/16. O-60-55. USGS Internal Memo.

Griffin, L. R. 1966. *Actinocoelia maeandrina* Finks, from the Kaibab Limestone of northern Arizona. Brigham Young University Geology Studies 13:105–108.

Hagadorn, J. W., J. L. Kirschvink, T. D. Raub, and E. C. Rose. 2011. Above the Great Unconformity: A fresh look at the Tapeats Sandstone, Arizona-Nevada, U.S.A. Museum of Northern Arizona Bulletin 67:63–77.

Haubold, H. 1971. Ichnia amphibiorum et reptiliorum fossilium. Handbuch der Paleoherpetologie, Teil 18. Gustav Fischer Verlag, Stuttgart, Germany.

Hodnett, J.-P. M., and D. K. Elliott. 2018. Carboniferous chondrichthyan assemblages from the Surprise Canyon and Watahomigi formations (latest Mississippian–Early Pennsylvanian) of the western Grand Canyon, northern Arizona. Journal of Paleontology 92, Memoir 77.

Howell, B. F. 1957. *Stipsellus annulatus*, a *Skolithos*-like Cambrian fossil from Arizona. Bulletin of the Wagner Free Institute of Sciences of Philadelphia 32(2):17–20.

Hunt, A. P., and V. L. Santucci. 1998. Taxonomy and ichnofacies of Permian tetrapod tracks from Grand Canyon National Park, Arizona. Pages 94–96 *in* V. L. Santucci, and L. McClelland, editors. National Park Service paleontological research. U.S. National Park Service, Geologic Resources Division Technical Report NPS/NRGRD/GRDTR-98/01.

Hunt, A. P., and V. L. Santucci. 2001. An unusual tetrapod track morphology from the Permian Coconino Sandstone, Grand Canyon National Park, Arizona. Pages 44–47 *in* V. L. Santucci and L. McClelland, editors. Proceedings of the 6th fossil resource conference: 2001 a fossil odyssey. Technical Report - National Park Service, Geological Resource Division, NRS/NRGRD/GRDTR-01/01.

Johnson, G. D., and J. Zidek. 1981. Late Paleozoic phyllodont tooth plates. Journal of Paleontology 55(3):524–536.

King, R. E. 1931. The geology of the Glass Mountains. Part II, Faunal summary and correlation of the Permian formations with description of Brachiopoda. Texas University Bulletin 3042.

Lucas, S. G., S. Voigt, A. J. Lerner, and E. C. Rainforth. 2013. *Sphaerapus*, a poorly known invertebrate trace fossil from nonmarine Permian and Jurassic strata of North America. Ichnos 20(3):142–152.

Lull, R. S. 1918. Fossil footprints from the Grand Canyon of the Colorado. American Journal of Science 45(269):337–346.

Macurda, D. B., Jr. 1969. Blastoids. Pages 457–473 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Marchetti, L., S. Voigt, S. G. Lucas, H. Francischini, P. Dentzien-Dias, R. Sacchi, M. Mangiacotti, S. Scali, A. Gazzola, A. Ronchi, and A. Millhouse. 2019. Tetrapod ichnotaxonomy in eolian paleoenvironments (Coconino and De Chelly formations, Arizona) and late Cisuralian (Permian) sauropsid radiation. Earth Science Reviews 190:148–170.

Martin, D. L. 1985. Depositional systems and ichnology of the Bright Angel Shale (Cambrian), eastern Grand Canyon, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Martin, H., and J. E. Barrick. 1999. Conodont biostratigraphy. Pages 97–115 *in* G. H. Billingsley and S. S. Beus, editors. Geology of the Surprise Canyon Formation of the Grand Canyon, Arizona. Museum of Northern Arizona, Flagstaff, Arizona. Bulletin 61.

McKee, E. D. 1932. Some Fucoides from Grand Canyon. Grand Canyon Nature Notes 7(8):77–81.

McKee, E. D. 1933. The Coconino Sandstone—its history and origin. Pages 77–115 *in* Papers concerning the palaeontology of California, Arizona, and Idaho. Carnegie Institution of Washington, Washington, D.C. Publication 440.

McKee, E. D. 1935. A *Conularia* from the Permian of Arizona. Journal of Paleontology 9:427–429.

McKee, E. D. 1938. The environment and history of the Toroweap and Kaibab formations of northern Arizona and southern Utah. Carnegie Institution of Washington Publication 492.

McKee, E. D. 1941. *Derbya arizonensis*, new name for *D. regularis* McKee. Journal of Paleontology 15(1):91.

McKee, E. D. 1945. Stratigraphy and ecology of the Grand Canyon Cambrian. Pages 5–168 *in* E. D. McKee and C. E. Resser. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

McKee, E. D. 1969. Paleozoic rocks of Grand Canyon. Pages 78–90 *in* D. L. Baars, F. G. Bowman, Jr., and C. W. Spencer, editors. Geology and natural history of the Grand Canyon region. Four Corners Geological Society, Durango, Colorado. Field conference guidebook 5.

McKee, E. D. 1978/05/25. O&G-78-10. USGS Internal Memo.

McKee, E. D. 1982a. Measured stratigraphic sections. Pages 389–496 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D. 1982b. Distribution and age of flora and fauna. Pages 75–112 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C. Professional Paper 1173.

McKee, E. D. 1982c. Environment of deposition of sandstone bodies. Pages 245–262 *in* E. D. McKee. The Supai Group of Grand Canyon. U.S. Geological Survey, Washington, D.C.

McKee, E. D., and W. J. Breed. 1969. The Toroweap Formation and Kaibab Limestone. Pages 12–26 *in* W. K. Summers and F. E. Kottlowski, editors. The San Andres Limestone, a reservoir for oil and water in New Mexico. New Mexico Geological Survey, Socorro, New Mexico. Special Publication 3.

McKee, E. D., and R. C. Gutschick. 1969a. Appendix: Descriptions of stratigraphic sections of the Redwall Limestone. Pages 613–692 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969b. The Redwall faunas. Pages 125–172 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969c. Sequence of sediments and unconformities. Pages 13–95 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and R. C. Gutschick. 1969d. Miscellaneous fossil groups - algae and stromatolites, holothurians, trilobites, ostracodes, and fish. Pages 545–552 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

McKee, E. D., and C. E. Resser. 1945. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

McKeever, P. J., and H. Haubold. 1996. Reclassification of vertebrate trackways from the Permian of Scotland and related forms from Arizona and Germany. Journal of Paleontology 70(6):1011–1022.

McKinney, F. K. 1983. Ectoprocta (Bryozoa) from the Permian Kaibab Formation, Grand Canyon National Park, Arizona. Fieldiana: Geology (New Series) 13(1341).

Miller, A. K., and A. G. Unklesbay. 1942. Permian nautiloids from western United States. Journal of Paleontology 16(6):719–738.

Miller, A. K., and W. Youngquist. 1949. American Permian nautiloids. Geological Society of America Memoir 41.

Newell, N. D. 1937. Late Paleozoic pelecypods: Pectinacea. Kansas Geological Survey, Lawrence, Kansas. Bulletin 10(1).

Noble, L. F. 1922. A section of the Paleozoic formations of the Grand Canyon at Bass Trail. Pages 23–73 *in* Shorter contributions to general geology, 1922. U.S. Geological Survey, Washington, D.C. Professional Paper 131-B.

Ossian, C. R. 1976. Redescription of *Megactenopetalus kaibabanus* David 1944 (Chondrichthyes: Petalodontidae) with comments on its geographic and stratigraphic distribution. Journal of Paleontology 50(3):392–397.

Palmer, A. R. 1963/10/17. O-60-55. USGS Internal Memo.

Rawson, R. R., and C. E. Turner. 1974. The Toroweap Formation: a new look. Pages 155–190 *in* T. N. V. Karlstrom, G. A. Swann, and R. L. Eastwood, editors. Geology of northern Arizona, with notes on archaeology and paleoclimate. Geological Society of America, Rocky Mountain Section, Flagstaff, Arizona. Guidebook 27, Part 1, regional studies.

Reeside, J. B., Jr., and H. Bassler. 1922. Stratigraphic sections in southwestern Utah and northwestern Arizona. U.S. Geological Survey, Washington, D.C. Professional Paper 129-D.

Resser, C. E. 1935. Nomenclature of some Cambrian trilobites. Smithsonian Miscellaneous Collections 93(5):1–46.

Resser, C. E. 1945. Cambrian fossils of the Grand Canyon. Pages 169–220 *in* E. D. McKee and C. E. Resser. Cambrian history of the Grand Canyon region. Carnegie Institution of Washington Publication 563.

Rose, E. C. 2003. Depositional Environment and History of the Cambrian Tonto Group, Grand Canyon, Arizona. Thesis. Northern Arizona University, Flagstaff, Arizona.

Rowland, S. M. 2017. Trackway of a sideways-walking pelycosaur in the Carboniferous Manakacha Formation in Grand Canyon National Park. Abstracts with Programs - Geological Society of America 49(6):abstract 136-12.

Rowland, S. M., and M. V. Caputo. 2018. Trackway of a sideways-walking basal tetrapod in the Pennsylvanian Manakacha Formation of Grand Canyon National Park. Journal of Vertebrate Paleontology, Program and Abstracts, 2018:206.

Sando, W. J. 1963/10/14. O-60-55. USGS Internal Memo.

Sando, W. J. 1969. Corals. Pages 257–343 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Sando, W. J., and E. W. Bamber. 1985. Coral zonation of the Mississippian System in the Western Interior Province of North America. U.S. Geological Survey, Washington, D.C. Professional Paper 1334.

Santucci, V. L., and A. P. Hunt. 1998. Biostratigraphy and paleoecology of late Paleozoic tetrapod tracks from Grand Canyon National Park, Arizona. Journal of Vertebrate Paleontology 18(supplement to 3):75A.

Santucci, V. L., A. P. Hunt, and M. G. Lockley. 1998. Fossil vertebrate tracks in National Park Service areas. Dakoterra 5:107–114.

Santucci, V. L., A. P. Hunt, T. Nyborg, and J. P. Kenworthy. 2006. Additional fossil vertebrate tracks in National Park Service areas. New Mexico Museum of Natural History and Science Bulletin 34:152–158.

Schenk, E. T., and H. E. Wheeler. 1942. Cambrian sequence in western Grand Canyon, Arizona. Journal of Geology 50(7):822–899.

Schuchert, C. 1918a. The Cambrian of the Grand Canyon of Arizona. American Journal of Science, 4th series 45(269):362–369.

Schuchert, C. 1918b. On the Carboniferous of the Grand Canyon of Arizona. American Journal of Science, 4th series 45(269):347–361.

Seilacher, A. 1970. *Cruziana* stratigraphy of non-fossiliferous Paleozoic sandstones. Pages 447–476 *in* T. P. Crimes and J. C. Harper, editors. Trace Fossils. Geological Journal Special Issue 3.

Shimer, H. W. 1919. Permo-Triassic of northwestern Arizona. Geological Society of America Bulletin 30(4):471–498.

Sinclair, G. W. 1948. The biology of the Conularida. PhD Thesis. McGill University, Montreal.

Skipp, B. 1969. Foraminifera. Pages 173–255 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Skipp, B., L. D. Halcomb, and R. C. Gutschick. 1966. Tournayellinae, calcareous foraminifera, in Mississippian rocks of North America; with translations from the original Russian of descriptions of several key genera and species by Ivan Mittin and Betty Skipp. Cushman Foundation for Foraminiferal Research Special Publication 9.

Snow, J. I. 1945. Trilobites of the Middle Permian Kaibab Formation of northern Arizona. Plateau 18(2):17–24.

Sorauf, J. E., and G. H. Billingsley. 1991. Members of the Toroweap and Kaibab formations, Lower Permian, northern Arizona and southwestern Utah. The Mountain Geologist 28(1):9–24.

Spamer, E. E. 1984. Paleontology in the Grand Canyon of Arizona: 125 years of lessons and enigmas from the late Precambrian to the present. The Mosasaur 2:45–128.

Stoyanow, A. A. 1936. Correlation of Arizona Paleozoic formations. Geological Society of America Bulletin 37:459–540.

Strother, P. K., and J. H. Beck. 2000. Spore-like microfossils from Middle Cambrian strata: expanding the meaning of the term cryptospore. Pages 413–424 *in* M. M. Harley, C. M. Morton, and S. Blackmore, editors. Pollen and spores: morphology and biology. Royal Botanic Gardens, Kew, United Kingdom.

Taylor, W. A., and P. K. Strother. 2008. Ultrastructure of some Cambrian palynomorphs from the Bright Angel Shale, Arizona, USA. Review of Palaeobotany and Palynology 151(1–2):41–50.

Thayer, D. 2009. An introduction to Grand Canyon fossils. Grand Canyon Association, Grand Canyon, Arizona.

Tidwell, W. D., J. R. Jennings, and S. S. Beus. 1992. A Carboniferous flora from the Surprise Canyon Formation in the Grand Canyon, Arizona. Journal of Paleontology 66(6):1013–1021.

Ulrich, E. O., and R. S. Bassler. 1931a. Cambrian bivalved Crustacea of the order Conchostraca. U.S. National Museum Proceedings 78 (article 4).

Ulrich, E. O., and R. S. Bassler. 1931b. *Indianites*, new name for the Cambrian crustacean *Indiana* Ulrich and Bassler. Journal of the Washington Academy of Sciences 21:364.

Walcott, C. D. 1883. Pre-Carboniferous strata in the Grand Cañon of the Colorado, Arizona. American Journal of Science, 3rd series 26:437–442.

Walcott, C. D. 1886. Second contribution to the studies on the Cambrian faunas of North America. U.S. Geological Survey, Washington, D.C. Bulletin 30.

Walcott, C. D. 1890. The fauna of the Lower Cambrian or *Olenellus* zone. U.S. Geological Survey, Washington, D.C. Annual Report 10, Part 1:509–760.

Walcott, C. D. 1897. Cambrian Brachiopoda; genera *Iphidea* and *Yorkia*; with descriptions of new species of each, and of the genus Acrothele. U.S. National Museum Proceedings 19:707–718.

Walcott, C. D. 1898. Cambrian Brachiopoda: *Obolus* and *Lingulella*, with description of new species. U.S. National Museum Proceedings 21:385–420.

Walcott, C. D. 1902. Cambrian Brachiopoda: *Acrotreta*; *Linnarssonella*; *Obolus*; with descriptions of new species. U.S. National Museum Proceedings 25:577–612.

Walcott, C. D. 1905. Cambrian Brachiopoda; with descriptions of new genera and species. U.S. National Museum Proceedings 28:227–337.

Walcott, C. D. 1908. Cambrian geology and paleontology, No. 3: Cambrian Brachiopoda: Descriptions of new genera and species. Smithsonian Miscellaneous Collections 53:53–165.

Walcott, C. D. 1912a. Cambrian Brachiopoda. U.S. Geological Survey, Washington, D.C. Monograph 51.

Walcott, C. D. 1912b. Cambrian geology and paleontology II. Middle Cambrian Branchiopoda, Malacostraca, Trilobita, and Merostomata. Smithsonian Miscellaneous Collections 57(6):145–228.

Walcott, C. D. 1916a. Cambrian geology and paleontology III. No. 5. Cambrian trilobites. Smithsonian Miscellaneous Collections 64(5):303–570.

Walcott, C. D. 1916b. Cambrian geology and paleontology III. No. 3. Cambrian trilobites. Smithsonian Miscellaneous Collections 64(3):157–258.

Walcott, C. D. 1924. Cambrian geology and paleontology IV. No. 9. Cambrian and Ozarkian Brachiopoda; Ozarkian Cephalopoda and Notostraca. Smithsonian Miscellaneous Collections 67(9):477–554.

White, C. A. 1874. Preliminary report upon invertebrate fossils collected by the expeditions of 1871, 1872, and 1873, with descriptions of new species. In U.S. Army Engineer Department, Geographical and geological explorations and surveys west of the one hundredth meridian. U.S. Government Printing Office, Washington, D.C.

White, C. D. 1927. Study of the fossil floras in the Grand Canyon, Arizona. Carnegie Institution of Washington Year Book 26:366–369.

White, C. D. 1928. Study of the fossil floras in the Grand Canyon, Arizona. Carnegie Institution of Washington Year Book 27:389–390.

White, C. D. 1929. Flora of the Hermit shale, Grand Canyon, Arizona. Carnegie Institution of Washington Publication 405.

Yochelson, E. L. 1969. Gastropods and pelecypods. Pages 439–451 *in* E. D. McKee and R. C. Gutschick. History of the Redwall Limestone of northern Arizona. Geological Society of America, Boulder, Colorado. Memoir 114.

Appendix D. Outside Repositories of GRCA Fossils

Institutions known to have collections from GRCA are listed below. Addresses, phone numbers, links, and email addresses to departments are included as available. Hyperlinks are subject to change without warning.

Academy of Natural Sciences of Drexel University
1900 Benjamin Franklin Parkway
Philadelphia, PA 19103
(215) 299-1000
https://ansp.org/

American Museum of Natural History
Central Park West at 79th Street
New York, NY 10024
(212) 769-5100
https://www.amnh.org/research/paleontology

Brigham Young University Museum of Paleontology
1683 North Canyon Road
Provo, UT 84602
(801) 422-3680
http://geology.byu.edu/museum/
byumuseumofpaleontology@byu.edu

California Academy of Sciences
55 Music Concourse Drive
San Francisco, CA 94118
(415) 379-8000
https://www.calacademy.org/
https://www.calacademy.org/scientists/department-of-invertebrate-zoology-and-geology-history
Carnegie Museum of Natural History
4400 Forbes Ave.
Pittsburgh, PA 15213
(412) 622-3131
https://carnegiemnh.org/

Cincinnati Museum Center
1301 Western Avenue
Cincinnati, OH 45203
(513) 287-7000
https://www.cincymuseum.org/
information@cincymuseum.org

Columbia University, Lamont-Doherty Earth Observatory
P.O. Box 1000
61 Route 9W
Palisades, NY 10964-1000 USA
(845) 359-2900
https://www.ldeo.columbia.edu/
director@ldeo.columbia.edu

Eberhard Karls University of Tübingen
Department of Geosciences
Hölderlinstr. 12
72074 Tübingen, Germany
+49-(0)7071-29-76862
https://uni-tuebingen.de/en/faculties/faculty-of-science/departments/geosciences/department/

Field Museum of Natural History
1400 S. Lake Shore Drive
Chicago, IL 60605-2496
(312) 922-9410
https://www.fieldmuseum.org/
collections@fieldmuseum.org

Harvard University Herbaria
22 Divinity Avenue
Cambridge, MA 02138
(617) 495-2365
https://huh.harvard.edu/
huh-requests@oeb.harvard.edu

Lake Mead National Recreation Area
601 Nevada Way
Boulder City, NV 89005
(702) 293-8990
https://www.nps.gov/lake/index.htm

Lund University (Lunds Universitet)
Box 117, 221 00 Lund, Sweden
Telephone +46 (0)46 222-0000
https://www.lunduniversity.lu.se/

Museum of Comparative Zoology, Harvard University
26 Oxford Street
Cambridge, MA 02138
(617) 495-2460
https://mcz.harvard.edu/

Museum of Northern Arizona
3101 N Fort Valley Rd
Flagstaff, AZ 86001
(928) 774-5213
https://musnaz.org/

Natural History Museum of Los Angeles County
900 Exposition Blvd
Los Angeles, CA 90007
(213) 763-3466
https://nhm.org/
https://nhm.org/research-collections/departments/invertebrate-paleontology (Invertebrate Paleontology)
invpaleo@nhm.org (Invertebrate Paleontology general contact)

Peabody Museum of Natural History at Yale University
P.O. Box 208118
170 Whitney Ave
New Haven, CT 06520
(203) 432-5050
https://peabody.yale.edu/
peabody.collections@yale.edu

Raymond M. Alf Museum of Paleontology
1175 West Baseline Road
Claremont, CA 91711-2199
(909) 624-2798
https://www.alfmuseum.org/

Smithsonian Institution, National Museum of Natural History
Department of Paleobiology
P.O. Box 37012
NHB MRC 121
Washington, D.C. 20013
https://naturalhistory.si.edu/research/paleobiology
paleodept@si.edu

University of Arizona
Department of Geosciences
1040 E. 4th Street
Tucson, AZ 85721
(520) 621-6000
https://www.geo.arizona.edu/

University of California Museum of Paleontology
1101 Valley Life Sciences Building
Berkeley, CA 94720-4780
(510) 642-1822
https://ucmp.berkeley.edu/
https://ucmp.berkeley.edu/contact-ucmp/

University of Notre Dame
Department of Civil & Environmental Engineering & Earth Sciences
156 Fitzpatrick Hall
Notre Dame, IN 46556
(574) 631-5380
https://ceees.nd.edu/
ceees@nd.edu

Western Archeological and Conservation Center
255 N Commerce Park Loop
Tucson, AZ 85745
(520) 791-6400
https://www.nps.gov/orgs/1260/index.htm

Appendix E. Paleontological Resource Law and Policy

General Resource Management Considerations

Paleontological resources are non-renewable remains of past life preserved in a geologic context. Fossils possess scientific and educational values and are of great interest to the public; therefore, it is exceedingly important that appropriate management attention be placed on protecting, monitoring, collecting, and curating of these paleontological specimens from federal lands. In 2009, the Paleontological Resources Preservation Act (PRPA) was signed into law as part of the Omnibus Public Land Management Act of 2009. The new paleontology-focused legislation includes provisions related to inventory, monitoring, public education, research and collecting permits, curation, and criminal/civil prosecution associated with fossils from designated DOI lands. Paleontological resource protection training is available for NPS staff through the NPS Geologic Resources Division (GRD). GRD is also available to provide support in paleontological resource theft or vandalism investigations.

National Park Service management policies state

> *…management actions will be taken to prevent illegal collecting [of fossil resources] and may be taken to prevent damage from natural processes such as erosion. Protection may include construction of shelters over specimens for interpretation in situ, stabilization in the field [which can include reburial] or collection, preparation, and placement of specimens in museum collections. The locality and geologic data associated with a specimen will be adequately documented at the time of specimen collection. Protection may also include, where necessary, the salvage collection of threatened specimens that are scientifically significant.*

Effective paleontological resource management serves to protect fossil resources by implementing strategies that mitigate, reduce, or eliminate loss of fossilized materials and their relevant data. Because fossils are representatives of adaptation, evolution, and diversity of life through deep time, they have intrinsic scientific value beyond that of the physical objects themselves. Their geological and geospatial contexts provide additional critical data concerning paleoenvironmental, paleogeographic, paleoecologic, and a number of other conditions that together allow for a more complete interpretation of the physical and biological history of the earth. Therefore, paleontological resource management must act to protect not only the fossils themselves, but to collect and maintain the ancillary data as well.

In general, losses of paleontological resources result from naturally occurring physical processes, by direct or indirect human activities, or by a combination of both. The greatest loss of ancillary data occurs when fossils are removed from their original geological context. Thus, when a fossil weathers and erodes from its surrounding sediments and geologic context, it begins to lose significant ancillary data until, at some point, it becomes more a scientific curiosity than a useful piece of scientific data. A piece of loose fossil “float” can still be of scientific value; however, when a fossil has been completely removed from its original context, such as an unlabeled personal souvenir or a specimen with no provenance information in a collection, it is of very limited scientific utility. Similarly,

fossils inadvertently exhumed during roadway construction or a building excavation may result in the loss or impairment of the scientific and educational values associated with those fossils. It is not necessary to list here all of the natural and anthropogenic factors that can lead to the loss of paleontological resources; rather it is sufficient to acknowledge that anything which disturbs native sediment or original bedrock has potential to result in the loss of the paleontological resources that occur there, or the loss of the associated paleontological resource data.

For management purposes, paleontological localities are evaluated for factors that could cause potential loss of paleontological resources. Their overall conditions are reported as good, fair, or poor based on the situations found at each individual locality. Risks and conditions that influence the degree of potential loss are categorized as Disturbance, Fragility, Abundance, and Site Access. "Disturbance" evaluates conditions that promote accelerated erosion or mass wasting resulting from human activities. "Fragility" evaluates natural conditions that may influence the degree to which fossil transportation is occurring, such as inherently soft rapidly eroding sediment or mass wasting on steep hillsides. "Abundance" judges both the natural condition and number of specimens actually preserved in the deposits as well as the risk of being easily recognized as a fossil-rich area which could lead to the possibility of unpermitted collecting. "Site Access" assesses the risk of a locality being visited by large numbers of visitors or the potential for easy removal of large quantities of fossils or fossil-bearing sediments as a result of proximity to public use areas or other access (along trails, at roadcuts, at beach or river access points, and so on). There is no simple formula for convenience, because it is influenced by each visitor's appreciation of factors such as terrain, distance from the access point, the presence and types of potentially harmful plant and animal life, the weather, and the visitor's capabilities, interest, and energy level. Generally speaking, any sedimentary exposures or accumulations of loose rock that are visible from a trail or other means of access and do not require significant climbing will attract geology enthusiasts, the curious, and the energetic.

Each of the factors noted above may be mitigated by management actions. Localities exhibiting a significant degree of disturbance may require either active intervention to slow accelerated erosion, periodic collection and documentation of fossil materials, or both. Localities developed on sediments of high fragility naturally erode at a relatively rapid rate and would require frequent visits to collect and document exposed fossils in order to prevent or reduce losses. Localities with abundant or rare fossils, or high rates of erosion, may be considered for periodic monitoring in order to assess the stability and condition of the locality and resources, in regard to both natural processes and human-related activities. Localities that are easily accessible by road or trail would benefit from the same management strategies as those with abundant fossils and by occasional unscheduled visits by park staff, documentation of in situ specimens, and/or frequent law enforcement patrols.

Cave localities are in a distinct class for management due to the close connection with archeological resources and unique issues affecting cave resources. See Santucci et al. (2001) for additional discussion of paleontological resources in cave settings.

Management strategies to address any of these conditions and factors could also incorporate the assistance of qualified specialists to collect and document resources rather than relying solely on staff

to accomplish such a large task at GRCA. Active recruitment of paleontological research scientists should also be used as a management strategy.

Management Actions

The following material is reproduced and adapted in large part from Henkel et al. (2015):

In March 2009, the Paleontological Resources Preservation Act (PRPA) (16 USC 460aaa) was signed into law (Public Law 111–11). This act defines paleontological resources as

> *…any fossilized remains, traces, or imprints of organisms, preserved in or on the earth's crust, that are of paleontological interest and that provide information about the history of life on earth.*

The law stipulates that the Secretary of the Interior should manage and protect paleontological resources using scientific principles. The Secretary should also develop plans for

> *…inventory, monitoring, and the scientific and educational use of paleontological resources.*

Paleontological resources are considered park resources and values that are subject to the "no impairment" standard in the National Park Service Organic Act (1916). In addition to the Organic Act, PRPA will serve as a primary authority for the management, protection and interpretation of paleontological resources. The proper management and preservation of these non-renewable resources should be considered by park resource managers whether or not fossil resources are specifically identified in the park's enabling legislation.

The Paleontological Resources Management section of NPS Reference Manual 77 provides guidance on the implementation and continuation of paleontological resource management programs. Administrative options include those listed below and a park management program will probably incorporate multiple options depending on specific circumstances:

No action—no action would be taken to collect the fossils as they erode from the strata. The fossils would be left to erode naturally and over time crumble away, or possibly be vandalized by visitors, either intentionally or unintentionally.

Surveys—will be set up to document potential fossil localities. All sites will be documented with the use of GPS and will be entered into the park GIS database. Associated stratigraphic and depositional environment information will be collected for each locality. A preliminary faunal list will be developed. Any evidence of poaching activity will be recorded. Rates of erosion will be estimated for the site and a monitoring schedule will be developed based upon this information. An NPS Paleontological Locality Database Form will also be completed for each locality. A standard version of this form will be provided by the Paleontology Program of the Geologic Resources Division upon request and can be modified to account for local conditions and needs.

Monitoring—fossil-rich areas would be examined periodically to determine if conditions have changed to such an extent that additional management actions are warranted. Photographic records should be kept so that changes can be more easily ascertained.

Cyclic prospecting—areas of high erosion which also have a high potential for producing significant specimens would be examined periodically for new sites. The periodicity of such cyclic prospecting will depend on locality-specific characteristics such as rates of sediment erosion, abundance or rarity of fossils, and proximity to visitor use areas.

Stabilization and reburial—significant specimens which cannot be immediately collected may be stabilized using appropriate consolidants and reburied. Reburial slows down but does not stop the destruction of a fossil by erosion. Therefore, this method would be used only as an interim and temporary stop-gap measure. In some situations, stabilization of a locality may require the consideration of vegetation. For example, roots can destroy in situ fossils, but can also protect against slope erosion, while plant growth can effectively obscure localities, which can be positive or negative depending on how park staff want to manage a locality.

Shelter construction—it may be appropriate to exhibit certain fossil sites or specimens in situ, which would require the construction of protective shelters to protect them from the natural forces of erosion. The use of shelters draws attention to the fossils and increases the risk of vandalism or theft, but also provides opportunities for interpretation and education.

Excavation—partial or complete removal of any or all fossils present on the surface and potentially the removal of specimens still beneath the surface which have not been exposed by erosion.

Closure—the area containing fossils may be temporarily or permanently closed to the public to protect the fossil resources. Fossil-rich areas may be closed to the public unless accompanied by an interpretive ranger on a guided hike.

Patrols—may be increased in areas of known fossil resources. Patrols can prevent and/or reduce theft and vandalism. The scientific community and the public expect the NPS to protect its paleontological resources from vandalism and theft. In some situations a volunteer site stewardship program may be appropriate (for example the “Paleo Protectors” at Chesapeake & Ohio Canal National Historical Park).

Alarm systems/electronic surveillance—seismic monitoring systems can be installed to alert rangers of disturbances to sensitive paleontological sites. Once the alarm is engaged, a ranger can be dispatched to investigate. Motion-activated cameras may also be mounted to visually document human activity in areas of vulnerable paleontological sites.

National Park Service Management Policies (2006; Section 4.8.2.1) also require that paleontological resources, including both organic and mineralized remains in body or trace form, will be protected, preserved, and managed for public education, interpretation, and scientific research. In 2010, the National Park Service established National Fossil Day as a celebration and partnership organized to promote public awareness and stewardship of fossils, as well as to foster a greater appreciation of their scientific and educational value (https://www.nps.gov/subjects/fossilday/index.htm). National Fossil Day occurs annually on Wednesday of the second full week in each October in conjunction with Earth Science Week.

Related Laws, Legislation, and Management Guidelines

National Park Service Organic Act

The NPS Organic Act directs the NPS to manage units

> *…to conserve the scenery and the natural and historic objects and the wildlife therein and to provide for the enjoyment of the same in such a manner as will leave them unimpaired for the enjoyment of future generations.* (16 U.S.C. § 1).

Congress reiterated this mandate in the Redwood National Park Expansion Act of 1978 by stating that the NPS must conduct its actions in a manner that will ensure no

> *…derogation of the values and purposes for which these various areas have been established, except as may have been or shall be directly and specifically provided by Congress.* (16 U.S.C. § 1 a-1).

The Organic Act prohibits actions that permanently impair park resources unless a law directly and specifically allows for the acts. An action constitutes an impairment when its impacts

> *…harm the integrity of park resources or values, including the opportunities that otherwise would be present for the enjoyment of those resources and values.* (Management Policies 2006 1.4.3).

Paleontological Resources Protection Act (P.L. 111-011, Omnibus Public Land Management Act of 2009, Subtitle D)

Section 6302 states

> *The Secretary (of the Interior) shall manage and protect paleontological resources on Federal land using scientific principles and expertise. The Secretary shall develop appropriate plans for inventory, monitoring, and the scientific and educational use of paleontological resources, in accordance with applicable agency laws, regulations, and policies. These plans shall emphasize interagency coordination and collaborative efforts where possible with non-Federal partners, the scientific community, and the general public.*

The Federal Cave Resources Protection Act of 1988 (16 USC 4301)

The law provides a legal authority for the protection of all cave resources on NPS and other federal lands. The definition for "Cave Resource" in Section 4302 states

> *Cave resources include any material or substance occurring naturally in caves on Federal lands, such as animal life, plant life, paleontological deposits, sediments, minerals, speleogens, and speleothems.*

NPS Management Policies 2006

NPS Management Policies 2006 include direction for preserving and protecting cultural resources, natural resources, processes, systems, and values (NPS 2006). It is the goal of the NPS to avoid or minimize potential impacts to resources to the greatest extent practicable consistent with the management policies.

NPS Director's Order-77, Paleontological Resources Management

DO-77 describes fossils as non-renewable resources and identifies the two major types, body fossils and trace fossils. It describes the need for managers to identify potential paleontological resources using literature and collection surveys, identify areas with potential for significant paleontological resources, and conduct paleontological surveys (inventory). It also describes appropriate actions for managing paleontological resources including: no action, monitoring, cyclic prospecting, stabilization and reburial, construction of protective structures, excavation, area closures, patrols, and the need to maintain confidentiality of sensitive location information.

Excerpt from Clites and Santucci (2012):

Monitoring

An important aspect of paleontological resource management is establishing a long-term paleontological resource monitoring program. National Park Service paleontological resource monitoring strategies were developed by Santucci et al. (2009). The park's monitoring program should incorporate the measurement and evaluation of the factors stated below.

Climatological Data Assessments

These assessments include measurements of factors such as annual and storm precipitation, freeze/thaw index (number of 24-hour periods per year where temperature fluctuates above and below 32 degrees Fahrenheit), relative humidity, and peak hourly wind speeds.

Rates of Erosion Studies

These studies require evaluation of lithology, slope degree, percent vegetation cover, and rates of denudation around established benchmarks. If a park does not have this information, there may be opportunities to set up joint projects, because erosion affects more than just paleontological resources.

Assessment of Human Activities, Behaviors, and Other Variables

These assessments involve determining access/proximity of paleontological resources to visitor use areas, annual visitor use, documented cases of theft/vandalism, commercial market value of the fossils, and amount of published material on the fossils.

Condition Assessment and Cyclic Prospecting

These monitoring methods entail visits to the locality to observe physical changes in the rocks and fossils, including the number of specimens lost and gained at the surface exposure. Paleontological prospecting would be especially beneficial during construction projects or road repair.

Periodic Photographic Monitoring

Maintaining photographic archives and continuing to photo-document fossil localities from established photo-points enables visual comparison of long-term changes in site variables.

Literature Cited

Clites, E. C., and V. L. Santucci. 2012. Protocols for paleontological resource site monitoring at Zion National Park. Natural Resource Report NPS/ZION/NRR—2012/595. National Park Service, Fort Collins, Colorado.

Henkel, C. J., W. P. Elder, V. L. Santucci, and E. C. Clites. 2015. Golden Gate National Recreation Area: Paleontological Resource Inventory. Natural Resource Report NPS/GOGA/NRR—2015/915. National Park Service, Fort Collins, Colorado.


National Park Service. 2004. Paleontological resource management. *In* Natural Resource Management Reference Manual 77. U.S. National Park Service, Washington, D.C.

National Park Service. 2006. Management Policies 2006. U.S. Government Printing Office, Washington, D.C.

Santucci, V. L., J. Kenworthy, and R. Kerbo. 2001. An inventory of paleontological resources associated with National Park Service caves. NPS Geological Resources Division, Denver, Colorado. Technical Report NPS/NRGRD/GRDTR-01/02.

Santucci, V. L., J. P. Kenworthy, and A. L. Mims. 2009. Monitoring in situ paleontological resources. Pages 189–204 *in* R. Young and L. Norby, editors. Geological monitoring. Geological Society of America, Boulder, Colorado.

Appendix F. Geologic Time Scale



Ma=Millions of year old. **Bndy Age**=Boundary Age. Colors are standard USGS colors for geologic maps. Modified from 1999 Geological Society of America Timescale (https://www.geosociety.org/documents/gsa/timescale/timescl-1999.pdf). Dates and additional information from International Commission on Stratigraphy update 2019/05 (http://www.stratigraphy.org/index.php/ics-chart-timescale) and USGS Fact Sheet 2007-3015 (https://pubs.usgs.gov/fs/2007/3015/).

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS 113/167727, March 2020



Natural Resource Stewardship and Science
1201 Oakridge Drive, Suite 150
Fort Collins, CO 80525