



Shoreline Change at Fort Matanzas National Monument

2019–2020 Data Summary

Natural Resource Data Series NPS/SECN/NRDS—2021/1327



ON THE COVER

Photo taken of the beach at Fort Matanzas National Monument displaying the high tide line that was mapped during the fall 2019 data collection. The photo is facing south. NPS photo / SECN staff.

Shoreline Change at Fort Matanzas National Monument

2019–2020 Data Summary

Natural Resource Data Series NPS/SECN/NRDS—2021/1327

Lisa Cowart Baron

National Park Service
Southeast Coast Inventory and Monitoring Network
13165 Mount Pleasant Road
Jacksonville, FL 32225

May 2021

U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science
Fort Collins, Colorado

The National Park Service, Natural Resource Stewardship and Science office in Fort Collins, Colorado, publishes a range of reports that address natural resource topics. These reports are of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Data Series is intended for the timely release of basic data sets and data summaries. Care has been taken to assure accuracy of raw data values, but a thorough analysis and interpretation of the data has not been completed. Consequently, the initial analyses of data in this report are provisional and subject to change.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner. Data in this report were collected and analyzed using methods based on established, peer-reviewed protocols and were analyzed and interpreted within the guidelines of the protocols.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available from the [Southeast Coast Network website](#) and the [Natural Resource Publications Management](#) website. If you have difficulty accessing information in this publication, particularly if using assistive technology, please email irma@nps.gov.

Please cite this publication as:

Baron, L. C. 2021. Shoreline change at Fort Matanzas National Monument: 2019–2020 data summary. Natural Resource Data Series NPS/SECN/NRDS—2021/1327. National Park Service, Fort Collins, Colorado.

Contents

	Page
Figures	iv
Tables	iv
Summary and Key Findings	v
Introduction	1
Study Area	3
Methods	5
Standard Operating Procedure	5
Shoreline Data Collection	5
Shoreline Change Analysis	5
Water Levels	6
Protocol Changes	6
Results	7
Shoreline Data	7
Annual Shoreline Change Analysis	7
Seasonal Shoreline Change Analysis	9
Spring 2019 to Fall 2019	9
Fall 2019 to Spring 2020	11
Summary	15
Literature Cited	17
Appendix A. Transect End Point Rates	19

Figures

	Page
Figure 1. Map showing the location where long-term shoreline monitoring data is collected at Fort Matanzas NM, highlighted by the red oval (study area).	2
Figure 2. Map of the shoreline surveys collected from 2019 to 2020 and the spring 2019 to spring 2020 shoreline change at Fort Matanzas National Seashore.	8
Figure 3. Histogram of transect shoreline change distance for spring 2019 to spring 2020.	9
Figure 4. Map of the shoreline surveys collected from 2019 to 2020 and the spring 2019 to fall 2019 shoreline change at Fort Matanzas National Seashore.).	10
Figure 5. Histogram of transect shoreline change distance for spring 2019 to fall 2019.	11
Figure 6. Map of the shoreline surveys collected from 2019 to 2020 and the fall 2019 to spring 2020 shoreline change at Fort Matanzas National Seashore.	12
Figure 7. Histogram of transect shoreline change distance for fall 2019 to spring 2020.	13

Tables

	Page
Table 1. Standard operating procedures used for shoreline position data collection, processing, and reporting.	4
Table 2. High tide levels surveyed during the spring FOMA shoreline monitoring.	5
Table 3. Table listing the length and positional error of the shorelines collected during 2019 and 2020 with the mean length and horizontal precision calculated by averaging across the entire shoreline feature.	6
Table A-1. Table of the shoreline change distances (net shoreline movement), in meters, for the study area analyzed.	15

Summary and Key Findings

In 2019 and 2020 the Southeast Coast Network collected shoreline data at Fort Matanzas National Monument as a part of the NPS Vital Signs Monitoring Program. Monitoring was conducted following methods developed by the National Park Service Northeast Barrier Coast Network and consisted of mapping the high tide swash line using a global positioning system (GPS) unit in the spring of each year (Psuty et al. 2010). Shoreline change was calculated using the Digital Shoreline Analysis System (DSAS) developed by USGS (Theiler et al. 2008).

Key findings from this effort are as follows:

- A mean of 2,226.56 meters (7,321.7 feet [ft]) of shoreline were mapped from 2019 to 2020 with a mean horizontal precision of 9.90 centimeters (3.9 inches [in]) at Fort Matanzas National Monument during 2019 to 2020.
- In the annual shoreline change analysis, the mean shoreline distance change from spring 2019 to spring 2020 was -5.70 meters (-18.7 ft) with a standard deviation of 26.25. The shoreline change distance ranged from -48.38 to 119.56 meters (-158.7 to 392.3 ft). Two erosion and one accretion areas were identified in the study area beyond the uncertainty of the data (± 10 meters [32.8 ft]).
- In the spring 2019 to fall 2019 seasonal shoreline change analysis, the mean shoreline distance change from was -8.79 meters (28.8 ft) with a standard deviation of 22.73. The shoreline change distance ranged from -55.75 to 75.20 meters (-182.9 to 246.7 ft). Four erosion and two accretion areas were identified in the study area beyond the uncertainty of the data (± 10 meters [32.8 ft]).
- In the fall 2019 to spring 2020 seasonal shoreline change analysis, the mean shoreline distance change from was 3.23 meters (10.6 ft) with a standard deviation of 19.36. The shoreline change distance ranged from -25.82 to 104.93 meters (-84.7 to 344.3 ft). One and four erosion areas were identified in the study area beyond the uncertainty of the data (± 10 meters [32.8 ft]).
- The annual shoreline change from 2019 to 2020 determined erosion has taken place on the east side of A1A where the Matanzas Inlet is located. Some accretion occurred on the west side of A1A along the Matanzas River, where a spit has formed.
- The seasonal shoreline change saw that on average, during the spring 2019 to fall 2019 season, almost half the shoreline eroded, with some accretion west of A1A and along the oceanfront beach, southeast of the east parking area. The fall 2019 to spring 2020 seasonal shoreline change analysis determined almost a quarter of the shoreline accreted and the area that eroded during the spring 2019 to fall 2019 season accreted during the fall 2019 to spring 2020 season.
- Overall, the most dynamic area of shoreline change within Fort Matanzas National Monument appears to be on the east and west side of A1A, along the Matanzas River inlet.

Introduction

The problem of land loss and gain and landscape alteration at the marine edge is fundamental to the many issues facing coastal park resource stewards. Shoreline change is a prime geoindicator of coastal environmental resource threats within parks. Change in shoreline position drives the alteration and replacement of established natural habitats, and shoreline retreat may destroy cultural resources, facilities, and other infrastructure. Geomorphologic change is a basic concern because it also influences change in other natural resource areas of interest within the National Park Service such as estuarine and subsurface water quality, species and habitats of concern, recreational visitor use, and even resource extraction. Spatial variability is inherent in shoreline change. Early identification of changes in past trends, along with an understanding of normal variability, is key to recognizing ecological problems in coastal parks. For managers, an understanding of the spatial and temporal patterns of geomorphologic change is important for optimal management of any coastal park because: (1) the interface of marine and land systems is very dynamic and is driven by multiple forcing mechanisms, (2) it results in alterations to resource patterns and dynamics at habitat and ecosystem level, and (3) it will eventually result in the loss of static resources. The establishment of local long-term monitoring programs will enhance understanding of these effects.

To effectively monitor diverse and dynamic coastal resources, the Southeast Coast Network has implemented a monitoring

approach which consists of mapping the high tide swash line along the oceanfront beaches using GPS equipment. The data collected is post-processed to attain higher accuracy data and analyzed using the Digital Shoreline Analysis System (DSAS) created by USGS (Theiler et al. 2008). This approach is modeled after the *Northeast Coastal Barrier Network Geomorphological Protocol: Part 1—Ocean Shoreline Position* (Psuty et al. 2010).

The monitoring objectives for which data are collected and analyzed as part of this effort are (Baron 2018a):

1. Determine inter-annual changes in mean high-water shoreline position (average and standard deviation of change in position at fixed transects, and cumulatively within management units).
2. Determine the magnitude, direction, and spatial distribution of observed shoreline change including the identification of areas of highly dynamic shorelines, and areas of accretion versus erosion.

This report provides a summary of shoreline monitoring data collected by the network to address these objectives at Fort Matanzas National Monument (FOMA) from 2019 through 2020 (Figure 1). Data used in this report are available at <https://irma.nps.gov/DataStore/Reference/Profile/2267045>.

Fort Matanzas National Monument Shoreline Study Area

Southeast Coast Network
National Park Service
U.S. Department of the Interior




Figure 1. Map showing the location where long-term shoreline monitoring data is collected at Fort Matanzas NM, highlighted by the red polygon (study area). The area monitored includes the entire ocean shoreline of the monument and a portion of the shoreline along the Matanzas River.

Study Area

Fort Matanzas National Monument is located 22 kilometers (14 miles [mi]) south of St. Augustine on the northeast Atlantic coast of Florida, embedded within the boundaries of the Guana Tolomato Matanzas National Estuarine Reserve (GTMNERR). It encompasses a total of 298 acres (120.6 hectares [ha]) divided between the southern tip of Anastasia Island and the northern end of Rattlesnake Island. Both are barrier islands separated from the Florida mainland by the Matanzas River and the Intracoastal Waterway (DeVivo et. al. 2015).

The site's namesake, Fort Matanzas, is an historic structure, prominent to this area as it was integral in protecting the southern waterway access to St. Augustine during the colonial wars. The sediment along the banks of the historic Fort is continually changing based on the dynamics of shifting sediment along the ocean shoreline as well as the Matanzas Inlet spit. Also, the area is an important stopover spot for migratory and nesting birds as well as nesting sea

turtles. Mapping efforts have been focused on the ocean shoreline of the monument, which is a 1.3-kilometer (0.8-mi) long stretch of beach beginning at the northern boundary line and ending at Matanzas Inlet. The section of back barrier beach stretching from Matanzas Inlet to the pedestrian walkway along the Matanzas River was added to the study area at the request of the park. Documenting long-term trends in shoreline change will greatly benefit future wildlife and cultural management decisions for this site.

The purpose of this report is to deliver the most recently collected data from within the park as part of an ongoing long-term shoreline-change monitoring program.

Methods

Standard Operating Procedure

Table 1 lists the standard operating procedures (SOPs) utilized in the long-term shoreline monitoring data collection, processing, and reporting. These SOPs are critical to maintaining continuity between data sets across many surveys to ensure comparable results. The version numbers in the table apply to both 2019 and 2020 data sets.

The shoreline was mapped along the oceanfront sandy shoreline of Fort Matanzas National Monument following the methods developed by the National Park Service Northeast Coast Barrier Network (Psuty et al. 2010). The Digital Shoreline Analysis System (DSAS; Theiler et al. 2008) was used to analyze the shoreline change. Annual shoreline change distances were calculated by determining the distance between two shorelines. Three shoreline surveys are included in this report, including a spring 2019, fall 2019, and spring 2020.

As stated in SOP 2.3.11 *Coastal Shoreline Monitoring Annual Change Analysis—Version 1.0* (Baron 2018b), the uncertainty of the individual shoreline survey is ± 5 meters (16.4 ft). This value is derived from the: (1) GPS accuracy, (2) differential tide position, (3) meteorological effects—wave dimension and run-up, and (4) interpretation of the line (Psuty et al. 2010). The annual comparisons deduced using two individual shoreline surveys will have an uncertainty of ± 10 meters (32.8 ft; Psuty et al. 2010). Refer to Psuty et al. (2010) for additional information regarding shoreline survey uncertainty.

Shoreline Data Collection

Shoreline data is collected according to SOP 1.3.22 *Conducting the Shoreline Survey—Version 1.0* (Baron 2018c).

The high tide swash line is mapped using a Trimble® GPS unit. The data collected is post-processed according to SOP 2.3.10 *Post survey GPS data processing—Version 1.0* (Baron 2018d) using Pathfinder Office software to obtain higher accuracy data. The corrected data is exported as an ESRI shapefile. The ESRI shapefile is then edited, formatted, and uploaded to a personal geodatabase to analyze using DSAS.

Shoreline Change Analysis

The three shoreline surveys collected during the 2019–2020 time period were analyzed using DSAS. The annual shoreline change was performed by analyzing the spring 2019 and spring 2020 shoreline data collected. Seasonal shoreline change was determined from the spring 2019 to fall 2019 and fall 2019 to spring 2020. The spring 2019 shoreline data was collected on April 8, 2019; the fall 2019 shoreline data was collected on November 5, 2019; and the spring 2020 shoreline data was collected on March 19, 2020. Three shoreline analyses were performed, including an annual analysis and two seasonal analyses. The annual analysis is from spring 2019 to spring 2020. The two seasonal analyses are from spring 2019 to fall 2019 and fall 2019 to spring 2020. All analyses were performed by calculating the distance the high tide line moved between the two shorelines analyzed, along cross-shore transects spaced 10 meters (32.8 ft) apart. This analysis is termed "net shoreline movement" in DSAS.

Erosion areas were determined as areas having negative shoreline change distance less than the uncertainty range of the data (± 10 meters [32.8 ft]). Accretion areas were determined as locations with positive shoreline change distances greater than the uncertainty range (± 10 meters [32.8 ft]).

Table 1. Standard operating procedures used for shoreline position data collection, processing, and reporting.

SOP number	SOP title	SOP version
1.3.21	Survey timing and GPS mission planning	1.0
1.3.22	Conducting the GPS shoreline survey	1.0
2.3.10	Post survey GPS data processing	1.0
2.3.11	Coastal shoreline monitoring annual change analysis	1.0
2.3.09	Coastal shoreline monitoring annual reporting	1.0

Water Levels

Three shoreline surveys of the high tide swash line included in this report were conducted during neap tide windows. The survey window was chosen based on the predicted tides using the closest NOAA tide gage with predicted tides (Mayport Bar Pilot Dock, FL—Station ID: 8720218; NOAA 2020). The actual tide heights that occurred before the surveys included in this annual report are listed in Table 2. The high tides prior to the shoreline surveys conducted have similar water levels.

Protocol Changes

No changes have occurred to the protocol. The data collected for this report was collected using SOP 1.3.22 *Conducting the Shoreline Survey*—Version 1.0 (Baron 2018c).

Table 2. High tide levels surveyed during the spring FOMA shoreline monitoring (m—meters).

Season	Survey collection date	Previous high tide height* (m)	Previous high tide time
Spring 2019	4/8/2019	1.50	1113
Fall 2019	11/5/2019	1.60	0248
Spring 2020	3/19/2020	1.53	0612

* Tide heights are relative to Mean Lower Low Water, Mayport Bar Pilot Dock, FL - Station ID: 8720218

Results

Shoreline Data

Three shoreline surveys were collected in in the 2019 and 2020 time period. On average, 2,232.56 meters (7,324.7 ft) of shoreline were mapped during the time period analyzed with a mean horizontal error of 9.90 centimeters (3.9 in; Table 3).

Annual Shoreline Change Analysis

Figure 2 displays a map of the shorelines collected in 2019 and 2020 and the annual shoreline change that occurred from spring 2019 to spring 2020. The annual shoreline change distance is displayed by the transects between the two shorelines, which are clipped to the extent of shoreline change and colored according to the shoreline change distance. A negative shoreline change distance denotes erosion, shown in warmer colors. Positive shoreline change distances are depicting accretion, and is shown in cooler colors. The lowest shoreline change distance was -48.38 meters (-158.7 ft) and the highest shoreline change distance was 119.56 meters (392.3 ft; Appendix A). The mean shoreline distance change for the area mapped was -5.90 meters (-18.7ft) with a standard deviation of 26.25.

Two erosion areas were present in the study area, represented as red, orange, and yellow transects in Figure 2. Erosion was identified for 29.50 % of the transects (59 of 200) in the analysis (Figure 3). Because the transects are located every 10 meters (32.8 ft) along the baseline, it can be inferred that erosion occurred along a cumulative 590 meters (1,935.7 ft) of the study area surveyed.

One accretion area was present in the spring 2019 to spring 2020 shoreline survey analysis and is represented as green and blue transects in Figure 2. Accretion was identified for 4.50% of the transects (9 of 200) in the analysis (Figure 3). Because transects are located every 10 meters (32.8 ft) along the baseline, it can be inferred that accretion occurred along a cumulative 90 meters (295.3 ft) of the study area surveyed.

Data used in this report are available at <https://irma.nps.gov/DataStore/Reference/Profile/2267045>.

Table 3. Table listing the length and positional error of the shorelines collected during 2019 and 2020 with the mean length and horizontal precision calculated by averaging across the entire shoreline feature [cm—centimeters; m—meters].

Collection Date	Mean Horizontal Accuracy (cm)	Length (m)
4/8/2019	10.00	2,147.23
11/5/2019	9.79	2,204.45
3/19/2020	9.87	2,345.99

Shoreline Data

Spring 2019 to Spring 2020




Figure 2. Map of the shoreline surveys collected from 2019 to 2020 and the spring 2019 to spring 2020 shoreline change at Fort Matanzas National Monument. The shoreline survey collected in spring 2019 is shown as a solid black line, the fall 2019 shoreline is a black hatched line, and the shoreline survey collected in spring 2020 is shown as a dotted black line. The more negative shoreline change denotes more erosion and are shown in warmer colors (yellow, orange, and red). The positive shoreline change or accretion are displayed as cooler colors (greens and blues). The grey transects denote shoreline change distance within the uncertainty range of the data (± 10 meters [32.8 ft]).




Figure 3. Histogram of transect shoreline change distance for spring 2019 to spring 2020. The shoreline change distance is binned by 2-meter (6.56 ft) increments, except for the bins less than -48 and greater than 48. The value of the x-axis represents the bin range. The value on the y-axis represents the frequency. The accretion transects are shown in blue and the erosion transects are shown in red. The transects within the uncertainty range (± 10 meters [32.8 ft]) are displayed as grey.

Seasonal Shoreline Change Analysis

In addition to the annual shoreline change, the seasonal shoreline change can be analyzed with the 2019–2020 shoreline data because the fall 2019 shoreline survey was collected. To look at seasonal variations in shoreline change, the spring 2019 to fall 2019 and fall 2019 to spring 2020 shoreline change was calculated.

Spring 2019 to Fall 2019

Figure 4 displays a map of the shorelines collected in 2019 and 2020 and the seasonal shoreline change that occurred from spring 2019 to fall 2019. The seasonal shoreline change distance is displayed by the transects between the two shorelines, which are clipped to the extent of shoreline change and colored according to the shoreline change distance. A negative shoreline change distance denotes erosion, shown in warmer colors. Positive shoreline change distances are depicting accretion and are shown in cooler colors. The lowest shoreline change distance was -55.75 meters (-182.9 ft) and the highest shoreline change distance was 75.20 meters (246.7 ft; Appendix A). The mean shoreline distance change for the area mapped was -8.79 meters (28.8 ft) with a standard deviation of 22.73.

Four erosion areas were present in the study area, represented as red, orange, and yellow transects in Figure 4. Erosion was identified for 41.41% of the transects (82 of 198) in the analysis (Figure 5). Because the transects are located every 10 meters (32.8 ft) along the baseline, it can be inferred that erosion occurred along a cumulative 820 meters (2,690.3 ft) of the study area surveyed.

Two accretion areas were present in the spring 2019 to fall 2019 shoreline survey analysis and are represented as green and blue transects in Figure 4. Accretion was identified for 17.17% of the transects (34 of 198) in the analysis (Figure 5). Because transects are located every 10 meters (32.8 ft) along the baseline, it can be inferred that accretion occurred along a cumulative 340 meters (1,115.5 ft) of the study area surveyed.

Data used in this report are available at <https://irma.nps.gov/DataStore/Reference/Profile/2267045>.

Shoreline Data

Spring 2019 to Fall 2019




Figure 4. Map of the shoreline surveys collected from 2019 to 2020 and the spring 2019 to fall 2019 shoreline change at Fort Matanzas National Monument. The shoreline survey collected in spring 2019 is shown as a solid black line, the fall 2019 shoreline is a black hatched line, and the shoreline survey collected in spring 2020 is shown as a dotted black line. The more negative shoreline change denotes more erosion and are shown in warmer colors (yellow, orange, and red). The positive shoreline change or accretion are displayed as cooler colors (greens and blues). The grey transects denote shoreline change distance within the uncertainty range of the data (± 10 meters [32.8 ft]).




Figure 5. Histogram of transect shoreline change distance for spring 2019 to fall 2019. The shoreline change distance is binned by 2 meters (6.56 ft) increments, except for the bins less than -50 and greater than 48. The value of the x-axis represents the bin range. The value on the y-axis represents the frequency. The accretion transects are shown in blue and the erosion transects are shown in red. The transects within the uncertainty range (± 10 meters [32.8 ft]) are displayed as grey.

Fall 2019 to Spring 2020

Figure 6 displays a map of the shorelines collected in 2019 and 2020 and the annual seasonal change that occurred from fall 2019 to spring 2020. The annual shoreline change distance is displayed by the transects between the two shorelines, which are clipped to the extent of shoreline change and colored according to the shoreline change distance. A negative shoreline change distance denotes erosion, shown in warmer colors. Positive shoreline change distances are depicting accretion and are shown in cooler colors. The lowest shoreline change distance was -25.82 meters (-84.7 ft) and the highest shoreline change distance was 104.93 meters (344.3 ft; Appendix A). The mean shoreline distance change for the area mapped was 3.23 meters (10.6 ft) with a standard deviation of 19.36.

One erosion area was present in the study area, represented as red, orange, and yellow transects in Figure 6. Erosion

was identified for 18.00% of the transects (36 of 200) in the analysis (Figure 7). Because the transects are located every 10 meters (32.8 ft) along the baseline, it can be inferred that erosion occurred along a cumulative 360 meters (1,181.1 ft) of the study area surveyed.

Four accretion areas were present in the fall 2019 to spring 2020 shoreline survey analysis and are represented as green and blue transects in Figure 6. Accretion was identified for 23.50% of the transects (47 of 200) in the analysis (Figure 7). Because transects are located every 10 meters (32.8 ft) along the baseline, it can be inferred that accretion occurred along a cumulative 470 meters (1,542 ft) of the study area surveyed.

Data used in this report are available at <https://irma.nps.gov/DataStore/Reference/Profile/2267045>.




Figure 6. Map of the shoreline surveys collected from 2019 to 2020 and the fall 2019 to spring 2020 shoreline change at Fort Matanzas National Seashore. The shoreline survey collected in spring 2019 is shown as a solid black line, the fall 2019 shoreline is a black hatched line, and the shoreline survey collected in spring 2020 is shown as a dotted black line. The more negative shoreline change denotes more erosion and are shown in warmer colors (yellow, orange, and reds). The positive shoreline change or accretion are displayed as cooler colors (greens and blues). The grey transects denote shoreline change distance within the uncertainty range of the data (± 10 meters [32.8 ft]).




Figure 7. Histogram of transect shoreline change distance for fall 2019 to spring 2020. The shoreline change distance is binned by 2 meters (6.56 ft) increments, except for the bins less than -24 and greater than 48. The value of the x-axis represents the bin range. The value on the y-axis represents the frequency. The accretion transects are shown in blue and the erosion transects are shown in red. The transects within the uncertainty range (± 10 meters [32.8 ft]) are displayed as grey.

Summary

The annual shoreline change from 2019 to 2020 determined erosion occurred on the east side of A1A, along the Matanzas Inlet. Some accretion occurred on the west side of A1A along the Matanzas River where a spit formed; however, erosion was also determined on the west side of A1A, north of the accretion area. The shoreline change along the oceanfront beach within the park, east of A1A, was within the uncertainty range of the data.

The seasonal shoreline change analysis determined that on average during the spring 2019 to fall 2019 season, almost half the shoreline eroded, with erosion occurring northeast of the east parking area access, on the east side of A1A approaching the inlet, on the west side of A1A just north of the accreting spit along the Matanzas River, and also on the west side of A1A where the island narrows just south of the west parking area access. The two areas of accretion denoted in this seasonal analysis were west of A1A along the inlet where a spit is forming and southeast of the east parking area access. Additionally, this time period included the passing of Hurricane Dorian, which may have contributed to some of the erosion.

The fall 2019 to spring 2020 seasonal change determined almost a quarter of the shoreline accreted, less erosion occurred than determined for the spring 2019 to fall 2019 seasonal shoreline change, and the area that accreted during the spring 2019 to fall 2019 season eroded during this time span. Accretion was determined for a stretch of oceanfront beach northeast of the east parking area access, east of A1A approaching the Matanzas Inlet, east and west of A1A along the Matanzas Inlet, and a pocket of accretion occurred where the island narrows, west of A1A. Notably, the areas of accretion were areas of erosion in the spring 2019 to fall 2019 analysis.

Overall, the seasonal fluctuations from the spring 2019 to fall 2019 and fall 2019 to spring 2020 analysis flip-flopped erosion and accretion areas. However, the most dynamic area of shoreline change is evident in the annual analysis, which appears to be on the east and west side of A1A, along the Matanzas River Inlet. In this area, erosion on the ocean side of A1A occurs and accretion occurs on the west side of A1A, where a spit has formed.



Photo taken of the beach at Fort Matanzas National Monument displaying the high tide line that was mapped during the spring 2020 data collection. The photo is facing south. NPS photo / SECN staff.

Literature Cited

- Baron, L. C. 2018a. Monitoring coastal shoreline change in Southeast Coast Network parks: Protocol implementation plan. Natural Resource Report NPS/SECN/NRR—2018/1744. National Park Service, Fort Collins, Colorado. Available at <https://irma.nps.gov/DataStore/Reference/Profile/2256509> (last accessed October 2020).
- Baron, L. C. 2018b. Coastal shoreline monitoring annual change analysis—Version 1.0. Southeast Coast Network Standard Operating Procedure NPS/SECN/SOP—2.3.11. Southeast Coast Network, Athens, Georgia. Available at: <https://irma.nps.gov/DataStore/Reference/Profile/2253897> (last accessed October 2020).
- Baron, L. C. 2018c. Conducting the GPS Shoreline Survey—Version 1.0. Southeast Coast Network Standard Operating Procedure NPS/SECN/SOP—1.3.22. Southeast Coast Network, Athens, Georgia. Available at: <https://irma.nps.gov/DataStore/Reference/Profile/2253893> (last accessed October 2020).
- DeVivo, J. C., J. Asper, S. Eastman, N. M. Rankin, A. C. Curtis, C. J. Wright, E. Thompson, and M. W. Byrne. 2015. Protocol for monitoring coastal salt marsh elevation and vegetation communities in Southeast Coast Network parks. Natural Resource Report NPS/SECN/NRR—2015/985. National Park Service, Fort Collins, Colorado. Available at <https://irma.nps.gov/DataStore/Reference/Profile/2222936> (last accessed October 2020).
- National Oceanic and Atmospheric Administration (NOAA). 2020. The Center for Operational Oceanographic Products and Services. Available at: tidesandcurrents.noaa.gov (last accessed October 2020).
- Psuty, N. P., T. M. Silveira, M. Duffy, J. P. Pace, and D. E. Skidds. 2010. Northeast Coastal and Barrier Network geomorphological monitoring protocol: Part I—ocean shoreline position. National Park Service, Fort Collins, Colorado. Available at: <https://irma.nps.gov/DataStore/Reference/Profile/664308> (last accessed October 2020).
- Thieler, E. R., E. A. Himmelstoss, J. L. Zichichi, and A. Ergul. 2008. The digital shoreline analysis system (DSAS) version 4.0—an ArcGIS extension for calculating shoreline change. Open-File Report 2008-1278. U.S. Geological Survey. Available at: <https://cmgds.marine.usgs.gov/publications/DSAS/of2008-1278/> (last accessed October 2020).

Appendix A. Transect End Point Rates

Table A-1. Table of the shoreline change distances (net shoreline movement), in meters, for the study area analyzed. Only the transects that intersected both shorelines in one of the three analyses are included in the table. Transects that did not intersect both shorelines in the analysis time period are shown as NA. Negative net shoreline movement denotes erosion and positive net shoreline movement denotes accretion.

Transect ID	Spring 2019 to Fall 2019	Spring 2019 to Spring 2020	Fall 2019 to Spring 2020
9	0.33	0.24	-0.09
10	1.31	-1.85	-3.17
11	1.54	0.74	-0.80
12	0.62	1.73	1.11
13	-0.95	-0.05	0.90
14	-2.76	-3.65	-0.89
15	-0.96	-4.11	-3.15
16	0.67	-3.32	-3.99
17	-2.62	-4.03	-1.41
18	0.49	-3.69	-4.18
19	-1.41	-1.79	-0.37
20	-1.58	-2.73	-1.14
21	-3.54	-3.34	0.20
22	-2.83	-2.09	0.74
23	-2.03	-3.05	-1.02
24	-2.69	-3.24	-0.55
25	-1.12	-3.33	-2.20
26	-1.25	-1.42	-0.17
27	-4.08	-2.40	1.68
28	-2.35	-2.81	-0.46
29	-1.97	-1.69	0.29
30	-4.04	-1.18	2.86
31	-2.64	0.31	2.95
32	-1.79	-0.29	1.50
33	-1.40	-1.34	0.06
34	-5.47	0.13	5.60
35	-3.10	0.24	3.33
36	-5.18	-0.45	4.74
37	-7.36	-1.16	6.20
38	-5.93	-0.95	4.98
39	-7.11	-1.12	5.99
40	-7.25	-1.31	5.94
41	-7.76	-1.24	6.52
42	-10.38	-2.64	7.75
43	-12.05	-2.32	9.73
44	-14.29	-3.32	10.98
45	-16.25	-3.82	12.43
46	-19.15	-5.66	13.49
47	-18.47	-4.65	13.82
48	-20.61	-5.68	14.93
49	-22.43	-5.18	17.25

Transect ID	Spring 2019 to Fall 2019	Spring 2019 to Spring 2020	Fall 2019 to Spring 2020
50	-21.55	-4.74	16.80
51	-22.52	-4.52	18.00
52	-22.93	-6.58	16.34
53	-19.54	-4.84	14.70
54	-19.37	-4.88	14.49
55	-20.72	-4.72	16.00
56	-18.50	-3.99	14.52
57	-17.99	-5.25	12.74
58	-20.33	-4.94	15.39
59	-19.30	-5.31	13.99
60	-21.38	-7.22	14.15
61	-21.72	-9.71	12.01
62	-19.33	-8.03	11.29
63	-19.25	-9.65	9.59
64	-17.52	-8.18	9.34
65	-15.27	-7.88	7.39
66	-14.86	-8.88	5.98
67	-12.31	-9.68	2.63
68	-9.59	-8.29	1.30
69	-2.87	-6.54	-3.67
70	-0.53	-7.75	-7.23
71	2.51	-7.78	-10.29
72	5.19	-6.36	-11.56
73	7.45	-4.94	-12.39
74	8.45	-6.13	-14.58
75	13.13	-6.26	-19.38
76	12.42	-6.97	-19.39
77	12.03	-4.40	-16.44
78	14.41	-4.46	-18.87
79	15.26	-3.15	-18.42
80	15.87	-2.38	-18.24
81	18.25	-0.71	-18.96
82	16.25	-3.29	-19.54
83	18.68	-2.30	-20.98
84	17.31	-2.82	-20.12
85	19.49	-3.49	-22.98
86	22.25	-3.57	-25.82
87	22.94	-1.26	-24.20
88	24.15	-0.47	-24.62
89	23.23	-0.29	-23.52
90	25.24	2.75	-22.49
91	25.58	1.64	-23.95
92	23.28	3.33	-19.95
93	24.08	3.64	-20.44
94	25.01	4.59	-20.42
95	25.47	5.01	-20.45

Transect ID	Spring 2019 to Fall 2019	Spring 2019 to Spring 2020	Fall 2019 to Spring 2020
96	27.07	7.22	-19.85
97	20.31	2.94	-17.37
98	20.59	1.88	-18.71
99	17.31	-2.08	-19.39
100	16.55	0.75	-15.79
101	12.40	-1.19	-13.59
102	12.57	-0.99	-13.56
103	8.93	-4.83	-13.77
104	7.52	-3.90	-11.42
105	6.20	-4.34	-10.54
106	5.13	-5.95	-11.07
107	3.75	-6.15	-9.90
108	1.58	-6.48	-8.07
109	-0.90	-10.21	-9.30
110	-2.24	-9.49	-7.25
111	-7.01	-10.49	-3.47
112	-14.36	-17.36	-3.00
113	-20.31	-20.68	-0.36
114	-24.89	-21.39	3.50
115	-27.99	-19.94	8.06
116	-30.48	-22.04	8.44
117	-33.94	-23.47	10.46
118	-39.19	-25.16	14.02
119	-42.91	-27.91	15.00
120	-45.40	-28.54	16.86
121	-49.97	-31.05	18.92
122	-50.64	-30.86	19.78
123	-50.79	-30.78	20.01
124	-52.10	-32.48	19.62
125	-51.39	-30.81	20.59
126	-53.10	-33.70	19.39
127	-53.68	-36.03	17.65
128	-54.40	-37.48	16.92
129	-54.87	-38.21	16.65
130	-53.48	-37.58	15.90
131	-51.68	-37.61	14.08
132	-50.50	-37.31	13.19
133	-47.75	-36.52	11.23
134	-48.41	-38.52	9.89
135	-44.92	-38.28	6.64
136	-37.43	-40.09	-2.66
137	-40.11	-41.11	-1.00
138	-39.92	-42.07	-2.15
139	-42.13	-43.59	-1.46
140	-42.10	-45.05	-2.95
141	-44.30	-48.05	-3.75

Transect ID	Spring 2019 to Fall 2019	Spring 2019 to Spring 2020	Fall 2019 to Spring 2020
142	-46.20	-48.23	-2.03
143	-49.05	-48.38	0.67
144	-53.30	-48.14	5.15
145	-55.75	-45.97	9.78
146	-48.91	-38.83	10.07
147	-47.05	-31.49	15.56
150	47.03	67.62	20.58
151	68.38	93.26	24.87
152	75.20	110.31	35.10
153	43.19	117.59	74.40
154	28.13	119.56	91.43
155	12.66	113.67	101.01
156	-1.75	103.18	104.93
157	-12.48	76.76	89.24
158	-17.21	68.84	86.05
159	-20.91	-26.08	-5.17
160	-20.57	-22.49	-1.92
161	-19.26	-18.35	0.90
162	-18.29	-16.41	1.88
163	-18.23	-17.73	0.49
164	-19.42	-19.30	0.12
165	-19.18	-20.28	-1.10
166	-19.22	-21.45	-2.23
167	-19.30	-21.93	-2.63
168	-18.52	-21.38	-2.86
169	-16.08	-18.92	-2.83
170	-13.12	-15.60	-2.48
171	-9.72	-14.71	-4.99
172	-8.70	-15.21	-6.51
173	-8.19	-14.79	-6.60
174	-7.85	-12.95	-5.10
175	-6.01	-9.07	-3.07
176	-3.73	-5.76	-2.03
177	-2.04	-3.50	-1.46
178	-0.74	-2.29	-1.54
179	0.56	-0.78	-1.34
180	0.23	-0.78	-1.01
181	-9.15	-7.90	1.25
182	-4.59	-5.84	-1.26
183	-8.41	-8.98	-0.57
184	-11.74	-10.96	0.78
185	-14.01	-10.61	3.41
186	-12.11	-8.56	3.56
187	-8.10	-5.35	2.75
188	-3.58	-1.07	2.51
189	1.10	2.54	1.44

Transect ID	Spring 2019 to Fall 2019	Spring 2019 to Spring 2020	Fall 2019 to Spring 2020
190	3.42	4.21	0.79
191	NA	NA	30.78
192	NA	NA	15.41
193	NA	NA	4.78
194	NA	NA	1.28
195	-18.43	-18.00	0.43
196	-15.17	-14.46	0.71
197	-11.45	-10.40	1.05
198	-6.74	-5.58	1.16
199	-3.34	-2.96	0.38
200	-1.43	-0.95	0.48
201	-1.08	-0.63	0.45
202	-1.37	-0.69	0.68
203	-1.69	-0.77	0.93
204	-1.25	-0.54	0.70
205	-1.15	-0.11	1.04
206	-1.51	-0.33	1.18
207	-0.63	0.25	0.88
208	-0.22	0.31	0.53
209	0.03	0.59	0.56
210	-0.02	0.27	0.29
211	-0.06	0.66	0.72
212	-0.14	1.00	1.13

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

National Park Service
U.S. Department of the Interior



Natural Resource Stewardship and Science

1201 Oak Ridge Drive, Suite 150
Fort Collins, Colorado 80525