

Cumberland Piedmont Network

National Park Service
Inventory and Monitoring Division



Standard Operating Procedure 7

Species Diversity and Tree Regeneration Measurements— Version 1.40

Please cite this as:

Cumberland Piedmont Network. 2023. Species diversity and tree regeneration measurements—Version 1.40. Cumberland Piedmont Network Standard Operating Procedure NPS/CUPN/SOP—7. Cumberland Piedmont Network, Mammoth Cave, Kentucky.

Overview

This standard operating procedure (SOP) documents data collection for overall plant species diversity, cover and tree seedling/sapling counts in nested subplots within the 20×20 -meter (65.6×65.6 feet [ft]) plot. Quantitative data are collected in eight sets of nested subplots, the 10×10 -meter (32.8×32.8 ft) module, and the entire 20×20 -meter plot following a modified Carolina Vegetation Survey approach (Lee et al. 2008). While the tree data will help us understand long term patterns, the understory data, comprised of herbaceous plants, seedlings and saplings, will potentially alert us to short term patterns and changes in the vegetation. As stressors affect the vegetation, seedling and herbaceous species composition and structure will respond more quickly than will large trees, coarse woody debris, or other measures we are actively monitoring. The data will help to discern trends including invasive exotic plant species numbers and seedling regeneration rates for key canopy tree species.

Most of the monitoring SOPs can be completed with basic plant identification skills. However, this SOP requires at least one team member to have demonstrated expertise and experience with the full diversity of vascular plant species in the park and to be able to quickly identify each plant species as data are collected. This protocol will be applied in late spring–late summer (late April to late September). Efforts will be made to revisit plots within two weeks of the time they were initially sampled (i.e., plots initially sampled in mid-June in 2011 were revisited in mid-June in 2016). To avoid over-trampling of the vegetation in the subplots, all crew members must be aware of the corner nested subplot locations and, to the extent practical, avoid walking in these areas.

Contents

	Page
Definitions.....	3
Step by Step Instructions	3
1.0 × 1.0 meter (Depth 3)	5
3.16 × 3.16 meters Corner (Depth 2).....	7
10 × 10 meters (Depth 1).....	9
Sample Remaining Modules.....	9
Assign Cover Classes	9
Notes on Plot Revisits	10
Changing or Amending the Previous Sampling Event's Data.....	11
Notes on Collecting.....	12
Equipment List.....	13
Completing Field Data Forms.....	14
Literature Cited	19
Revision Log.....	19

Definitions

Cover is defined as the percentage of the ground that is covered by the vertical projection of the above ground material (e.g., leaves, branches) for a particular species (Lee et al. 2008). For this protocol, we record canopy cover. In canopy cover, unlike foliar cover, the tiny gaps between leaves and limbs are considered filled when estimating.

Height of a seedling or sapling is the distance from the soil surface to the top of the terminal bud (the large bud at the top of the stem). Do not include leaves or needles in this measurement. Seedling or sapling height is determined from stems as they are naturally growing. Height is measured vertically from the ground to the tip of the terminal bud, not along the axis of the stem. Stems should **not** be straightened for measurement.

Seedlings are live juvenile **trees** with at least two true leaves and are less than 1.37 meters (4.5 ft) tall. A seedling is counted only if it is rooted in the nested subplot. Seedlings less than 5 centimeters (2 inches [in]) in height are not counted in the tally. Tree seedlings that originate at a common root collar but branch under the leaf litter are independent seedlings. If a tree seedling forks or branches above ground, measure the height of the tallest terminal bud and count the seedling as one. Do not count stump sprouts from a live tree as seedlings. Tally stump sprouts from standing dead trees, counting each sprout as a separate seedling. Refer to the park-specific plant species list which is printed from the protocol database each field season. This list indicates which species are considered trees.

Saplings are live juvenile **trees** more than 1.37 meters in height but less than 10.0 centimeters (3.9 in) DBH. If a sapling forks or branches above 0.5 meters (1.6 ft), measure the height of the tallest terminal bud and count the sapling as one; if a sapling forks or branches below 0.5 meters, measure each fork or branch as a separate sapling. Saplings are alive if there are any living parts (i.e., leaves, buds, cambium) at or above 1.37 meters (breast height). Saplings that have been temporarily defoliated are considered alive. A sapling is counted only if it is rooted in the nested subplot. Refer to the park-specific plant species list to determine which species are considered trees.

Vines are not tallied as a seedling or sapling. Presence of vine species is recorded in the nested subplots (Depths 3 and 2), and 10 × 10-meter (32.8 × 32.8 ft) modules (Depth 1) along with other plant species.

Step by Step Instructions

Each of the 10 × 10-meter modules (1–4) contains four numbered corners with teams sampling two of the corners within each of the modules for a total of eight corners sampled (Figure 1). Corners sampled are 1–2, 1–4, 2–2, 2–3, 3–2, 3–4, 4–2, and 4–3 (Figure 1), where the first number refers to the module number and the second refers to the specific corner within that module. In other words: corners two and four are sampled in module one; corners two and three in module two; corners two and four in module three; and corners two and three in module four. It is important for the field crew

to keep themselves oriented, so the correct corners are completed and documented. When in doubt, check with other crew members or the plot diagram included in the field datasheets.

Following the Carolina Vegetation Survey protocols (Lee et al. 2008), nested subplots in eight corners are sampled in a particular pattern. This means each corner sampled is not adjacent to another sampled corner and there is a permanent post anchoring each nested subplot. The corners are indicated at the top of the columns on the datasheet, as pairs, module number and corner number (e.g., 1–2, 1–4). Once at the nested corner location, take two 1-meter (3.3 ft) sticks and use them along with the previously laid out tapes to form a square 1 square meter (10.8 square feet [ft²]) nested subplot, which corresponds to Depth 3 for that corner (Figure 7.2). If for some reason you cannot sample a corner, mark through the corner and module numbers at the top of the column on each page of the field datasheet for this SOP and replace with 0–0. Also, make a note of why the corner was not sampled in the Corner Notes (e.g., very active yellowjacket nest in corner, concerned for crew safety).

In recent operational reviews concern has been expressed from multiple sources about the use of minimally trained or inexperienced field crew members as data recorders for the botanist. This role requires a great deal of experience or oversight from the botanist and whenever possible should be performed by someone with substantial and demonstrated ability. This may mean that if the individual responsible for completing tree work is the only experienced data recorder on the plot than they will need to alternate between recording for the botanist and collecting tree data. One way to partition this workload with two-person teams is to have the tree data collector record for the botanist when completing the two intensive corners within the module. When the two corners are complete the tree data recorder can hand the clipboard back to the botanist and switch to tree work, while the botanist collects the residual data for the module (i.e., level 1). This approach of putting the clipboard in the hands of the botanist in each module also allows an opportunity for the botanist to review the module data before moving on to the next module (i.e., it potentially benefits data quality). Note it is also a good time for the botanist to check with the tree recorder to make sure all trees are accounted for and there is agreement on identification. This increase in training and experience level requirements for data recorders also means that some thought needs to be given to engaging and utilizing the skills of “temporary” or minimally experienced field crew members. This shouldn’t be forgotten! It is important to engage park staff and others in this process and utilize their help. Some potential tasks have been identified within this and other SOPs.




Figure 1. 20 × 20-meter (65.6 × 65.6 ft) plot diagram delineating location of eight sampled subplots (shaded dark).




Figure 2. Diagram showing placement of sampled corners in module 1 (Lee et al. 2008).

1.0 × 1.0 meter (Depth 3)

Once the 1 square meter (10.8 ft²) corner nested subplot is laid out, species presence data are collected. First, identify and record all plant species in the 1 × 1 meter (3.3 × 3.3 ft) nested subplot for module 1 (refer again to Figure 2). Skip over any plants that are very small (less than 5

centimeters [2 in]) and cannot be attributed to a species. In the box for the nested subplot on the datasheet, mark a “3” to indicate that the species is present in module one, corner/subplot two at the 1 × 1 meter level. For plot revisits, the datasheets for the plot will be filled with a preprinted species list from the last survey, with species codes listed in alphabetical order and extra blank rows to add species not encountered during the last survey. Refer to SOP 14 *Recording Species Nomenclature—Version 1.37* (CUPN 2023a) for detailed information on species nomenclature. Certain species have duplicative seven letter codes, so eight letter codes are provided for these species on the park-specific species list. They include species such as *Acer saccharum* and *Acer saccharinum*, *Carya ovata* and *Carya ovalis*, *Polygonum hydropiper* and *Polygonum hydropiperoides*, and *Polygonum virginianum* and *Polypodium virginianum*. Record your confidence in the identification value under the field of **ID conf** (Table 1).

If there is any doubt about species identification, an ID confidence of **Probable** or **Uncertain** is recorded instead of **Certain** (Table 1). If needed, utilize the **Note** number column to reference a more detailed note about the species ID at the end of the species list. If a plant cannot be identified to species quickly and confidently, then identify the species to either family or genus and assign it with a descriptive name for later reference (for instance, “*Carex* slender rhizome” or “*Symphyotrichum* large purple flower” and utilize the **Note** number column to reference a more detailed note about the species ID at the end of the species list. If the species cannot be determined in the field, collect a sample with foliage, buds, cones, or flowers, and/or a photograph to make a positive identification later. Put a check in the **Collected** column on the data sheet to indicate that a specimen was collected. For more information on species collecting and identification, refer to the “Notes on Collecting” section at the end of this SOP.

Table 1. Species confidence level codes.

Code	Description	Symbol
Certain	The person(s) making the taxon interpretation has a high confidence in the accuracy of the field or herbarium identification.	C
Probable	The person(s) making the taxon interpretation has a moderately high confidence in the accuracy of the field or herbarium identification, but uncertainly is typically recorded as “cf” or “?” (e.g., taxon is most likely the species named; occasionally another taxa may be listed as a possibility—it can be named in a comments field).	P
Uncertain	The person(s) making the taxon interpretation has only moderate confidence in the accuracy of the field or herbarium identification, as shown by “??” or other comments about uncertainty. (For example, taxon may be one of two or more species and although one name is chosen, it could as easily be another.)	U

Once you have collected species data at the 1 × 1 meter level (Depth 3), then move on to the tree seedling tallies at the 1 × 1 meter level. Blank datasheets will be provided for seedlings and saplings for both plot establishment and plot revisit surveys. Identify tree species and tally (by species) all individual seedlings in the 1 × 1 meter nested subplot that are 5–15, 15–30, 30–50, or 50–137

centimeters (2–6, 6–12, 12–20, or 20–54 in) tall. Skip over any plants that are very small (less than 5 centimeters) and cannot be attributed to a species. Use the dots and square method described at the top of the seedling and sapling datasheet to document numbers of seedlings and saplings (Figure 3).

Based on reviews of QA/QC plot data, it appears that crews are sometimes forgetting to go back and conduct a seedling count for all seedling species identified in the subplot. Therefore, crews can and are encouraged to ID and tally seedlings before turning their attention to the herb species.




Figure 3. Tally method for recording (counting) seedling and saplings occurring within sampled corners.

3.16 × 3.16 meters Corner (Depth 2)

At this point, you have completed the 1 × 1 meter (3.3 × 3.3 ft) nested subplot but still must complete the 3.16 × 3.16 meters (10 square meters [107.6 ft²]) nested subplot in the same corner. The 3.16 × 3.16 meters (Depth 2) nested subplot includes the 1 × 1 meter (Depth 3) nested subplot that was already examined, so you simply need to expand the size to 3.16 meters (10.4 ft) on each side using the set of collapsible fiberglass tent poles and module boundary tapes. This protocol relies on repeated measures (i.e., consistency). Thus, it is important to invest some effort in getting an accurate 10 square meter subplot and doing it in such a way that the next team, whether that be a QA/QC team or another monitoring effort in five years, lays their subplot out in such a way as to be sampling the same 10 square meter area.

If the tape measures delineating the module boundary are anchored in the corner conduit, then two of the sides of the 10 square meter subplot are already established. Remember the “permanent marker” for the subplot is the conduit, not where your field crew happened to anchor the chaining pin when laying out the plot. You may move the chaining pin into the conduit at this time or loop the tape around the conduit. Because the tapes have already been extended and crew members are no longer pulling on the tapes, there should no longer be a risk of inadvertently pulling up a conduit. So, beginning at the conduit measure 3.16 meters along each tape from the conduit. Place a flag at each of these points along with a meter stick extending into the module along the two currently unmarked sides of the 10 square meter subplot. Once the meter sticks are adjusted to right angles from the tapes a crew member can visually extend these lines to where they would meet at the opposite corner from the conduit. Place a flag at this point, which should be 4.47 meters (14.7 ft) at a 45° angle from the conduit. However, this approach is discouraged as the Cumberland Piedmont Network seeks to improve repeatability of sampled corners.

The preferred method for delineating the 10 square meter subplot is to use the pair of collapsible fiberglass tent poles which have been added to each of the equipment boxes. Each of the two poles have been precut to 3.16 meters, making them the exact length needed to form one side of the 10

square meter subplot. If the tape measures delineating the module boundary are anchored in the corner conduit, then two of the sides of the 10 square meter subplot are already established. If it is not in the conduit, move it or make sure the tape is looped around the outside of the conduit. Next, place one end of the fiberglass pole against the conduit and extend the pole along one of the 2-meter (6.6 ft) tapes. At the end of the fiberglass pole opposite the conduit place a survey flag on the meter tape. Retrieve the fiberglass pole and extend it into the module perpendicular to the tape measure with the end butted against the survey flag. If a second crew member is assisting (which is the recommended approach), they can be following the same procedure along the other subplot tape boundary. Each of the two crew members can adjust their poles so the ends of the two fiberglass poles meet, delineating one of the plots eight 10-square meter subplots. At this time, the fiberglass poles can be picked up and moved to the next corner if a minimum of three survey flags (one on each end and one in the middle) have been placed along the pole's line. Additional survey flags are placed inside or outside of plants where it is not obvious as to whether the species is inside or outside of the subplot boundary. Placement of these flags is completed before picking up the fiberglass poles and moving on to the next corner. If there are questions about whether large size class saplings are within the subplot, the previous sampling event's datasheets can be consulted. Using the fiberglass poles delineates the full subplot boundary without relying on the meter sticks used by the botanist. This is an excellent task for extra or minimally trained field crew members to perform independently and helps the botanist.

Two types of data are collected in the 10 square meter subplot. This includes any new species not already documented in the corner's 1 square meter (10.8 ft²; Depth 3) subplot. The other data type recorded in the 10 square meter subplot is a tally of tree saplings by species and size class. Because crews can sometimes get in a hurry and forget to collect tree sapling data before moving to the next corner, it is recommended that crews count tree saplings prior to recording the new species for the corner. To do this, tally each tree species by size class (0 to < 1.0 centimeter DBH, 1.0 to < 2.5 centimeter DBH, 2.5 to < 5 centimeters DBH, and 5 to < 10 centimeters DBH) within the 10 square meter subplot (including the 1 square meter subplot area as it is within the Depth 2 area). It can be helpful to carry a small DBH tape to measure borderline saplings that are close to 10 centimeters DBH, and it is a good idea to communicate with the team member completing tree measurements whether the suspect tree is below or above 10cm DBH, so that is not missed or double-counted. **Do not** collect any data on **seedlings** (woody plants below 1.37 meters tall) for the 10 square meter subplot, as these are only recorded at Depth 3 (1 square meter subplot) in each corner.

Now move on to recording Depth 2 species for the corner. Depth 2 species include any species present that did not receive a presence value of 3 for the corner (i.e., present in the 1 square meter subplot). Instead of writing a "3" in the box for the corner to indicate presence, write a "2" to indicate the new "Depth" (modified from Lee et al. 2008). Skip over any plants that are very small (less than 5 centimeters) and cannot be attributed to a species.

If after sampling a subplot at both Depth 3 and 2, and you do not encounter any seedlings or saplings, write a 0 (zero) on the datasheet for that module and corner in the seedling and sapling data section so it is documented the field team surveyed that corner for seedlings and saplings and it was not

missed. It is also imperative the field team tallies all seedlings and saplings at the appropriate Depth in each subplot because these data are used for regeneration analysis. A quick check of tree species written on the species list for that subplot will let you know if there might be any seedlings or saplings missed in the seedling and sapling tallies.

Take up all nested subplot markers and repeat the previous steps in the second nested subplot for the module. Start over with recording of species in the new corner (i.e., if a species occurs in both sampled corners within the module then it should be recorded in both corners). If you are recording data and unsure how to code a species presence on the datasheet, stop and ask questions of other field team members. It is important to get it right as there are no second chances to record the data.

10 × 10 meters (Depth 1)

After completing both nested subplots within the module, move on to the whole 10 × 10 meter (32.8 × 32.8 ft) module, which is referred to as Depth “1” (3.16 × 3.16 meters was Depth “2” and 1 × 1 meter was Depth “3”). Now identify and write down any new species not already documented in either corner of the module. Skip over any plants that are very small (less than 5 centimeters [2 in]) and cannot be attributed to a species. This time, instead of writing a “2” in the box for the corner to indicate presence at that Depth, write a “1” to indicate Depth of 1 (modified from Lee et al. 2008). Do not forget to look up at the trees in the module and consult with the team member surveying trees to make sure that you agree on species ID and have accounted for species that may not be present as seedlings or saplings.

While there is no strict time limit established on searching for residual species at Depth 1, the goal is to be thorough but at the same time recognize that invariably some species will be missed. As a guideline, the botanist will track time spent since they added a species to the module. If it is two or more minutes of active searching without a new species being found, it is probably a good indicator that it is time to move to the next module. When collecting data at the module or Depth 1 level, it is a good idea for the botanist to have the data sheets and record the data at the module level. This serves two purposes: the botanist can review the species list for completeness and legibility before moving on to the next module, and it allows the data recorder to take a short break or move on to another task, such as laying out the corners in the next module.

Sample Remaining Modules

Move on to the next 10 × 10 meter (32.8 × 32.8 ft) module and repeat the steps above until you have completed all four modules of data collection.

Assign Cover Classes

After the four modules have been sampled, a cover value (see definitions) will be assigned for each plant species recorded in the plot. One way of estimating cover is to ignore all but one species and estimate the percent of the ground that would be shaded if the sun were directly overhead. When assigning cover **do not** assign separate cover classes for the canopy, subcanopy, shrub and herbaceous strata. Also, **do not** attempt to assign a separate cover value for each module. When finished you should have a single cover value for each species for the entirety of the 20 × 20 meter (65.6 × 65.6 ft) plot. The cover classes are assigned rapidly (completed in 10 minutes or less) so

teams should not spend extensive time equivocating between two cover classes for a particular species. Remember if reasonably continuous in outline, the canopy of a plant is considered continuous. However, if there are significant gaps within an individual canopy, it is not considered continuous. Cover ignores the vertical distribution of leaves and includes all leaves from those near ground level to those in the canopy (Lee et al. 2008).

Data are recorded in cover classes which represent ranges of percent cover assigned to 10 categories (Table 2; Lee et al. 2008). When assigning cover classes, the botanist should consult with the team member that surveyed the trees (SOP 6 *Tree Measurements—Version 1.48* [CUPN 2023b]) in order to make sure all tree species are accounted for correctly in the cover values and all trees are accounted for in each module (i.e., a presence value exists in the herb data for each tree species or module combination from the tree data). Do not forget about assigning cover values for species that overhang the plot but not rooted in the plot. For those species a presence value of 0 is added to the module where it is overhanging.

Table 2. Canopy Cover Categories, Range and Notes on Cut-offs between Categories [m²=square meters; m=meters].

Category	Range	Cut-off Descriptors
1	<0.1%	0.1% = 0.4 m ² or ~ 0.6 x 0.6 m
2	0.1–1%	1% = 4 m ² or 2 x 2 m
3	1–2%	2% = 8 m ² or 2 x 4 m
4	2–5%	5% = 20 m ² or 4 x 5 m
5	5–10%	10% = 40 m ² or 5 x 8 m
6	10–25%	25% = 100 m ² or 1 module
7	25–50%	50% = 200 m ² or 2 modules
8	50–75%	75% = 300 m ² or 3 modules
9	75–95%	95% = 380 m ² or nearly covered
10	95%–100%	—

Notes on Plot Revisits

For plot revisits, there are several notes to keep in mind:

1. When identifying additional plant species at Depth 1, do not add additional search time to residual level searching for species found during the previous sampling event but not in the current sampling event at Depths 2 or 3 (i.e., don't go back just to locate species found in previous sampling events). Consistency in effort and independence between sampling events is important.

2. If the previous sample excluded a corner for a temporary reason (such as the presence of a yellowjacket nest) that is not applicable during the revisit, sample the corner during the revisit.
3. During plot revisits, there will be an established species list for the plot from the previous sampling event. This list will be in alphabetical order on the datasheet. If you notice obvious species ID errors or incorrect species code usage, cross out the original code on the datasheet and write in the correct code. If helpful, utilize the “Note#” column to explain why the previous ID was incorrect. It can be helpful to identify a previously identified genus to species if possible. Refer to “Changing or Amending Previous Sampling Event’s Data” section below for more information.
4. For those species on the pre-populated species list detected during previous sampling event but not detected during the current sampling event, it is good practice to draw a horizontal line through the entire row of the datasheet to indicate this species was not observed for the current sampling event. Some species found during previous surveys will not be seen during current surveys and new species will be seen during the current survey. Add new species to the bottom of the species list, spelling out the full scientific name.

Changing or Amending the Previous Sampling Event’s Data

While changes to previous sampling event data should be rare, there are circumstances where changes are valid. For example, an obvious misidentification occurred, or a data entry error was discovered in the field. Other examples include:

- Based on the presence of fertile material that was not present in the plot during the previous survey, the botanist changes an identification from *Lonicera standishii*-Probable to *Lonicera maackii*-Confident.
- The botanist realizes they made an obvious mistake on the previous plot visit, such as using an incorrect code-an example is writing CHAMMAC instead of CHIMMAC.

One thing that may help whether to make a taxonomic change is to consult the presence and cover values recorded during the previous sampling event. For instance, if a species was found in only one corner during both sampling events, or if a species was widely dispersed throughout the plot with a relatively high cover value during both sampling events, this would add greater evidence that both botanists were looking at the same taxon. Note the botanist should avoid consulting presence and cover values from the previous sampling event until after data collection for the current event is concluded, if possible. If the botanist concludes that a species name should be changed, they should indicate this by marking through the species code on the field form in pencil and writing the new “corrected” species code next to it. However, before such a change the botanist must consider the following questions:

- How do you know the proposed species change is not just a second species that was either missed or not present during the previous sampling event? This is where you may want or need to consult the previous presence values.
- If the previous field crew were present, would they quickly and easily agree that a data error occurred and should be corrected?
- Does the change improve the quality of the data, both in terms of accuracy and consistency?

In determining whether to make a species change, the botanist must also keep in mind there are some taxonomic concepts where accurate identification from vegetative material alone is difficult at best and there is going to be some disagreement even among trained, experienced botanists. A couple of examples of ambiguity in species calls include *Acer barbatum* vs. *Acer saccharum* and just about anything in the genus *Carex*. These are examples of “errors” that not every botanist may quickly agree upon. Also, because there is some ambiguity with these taxonomic concepts, the data are likely to be lumped for most if not all analyses. So, a proposed change may not necessarily improve the quality of the data. What we’re trying to avoid are situations where taxonomic concepts are constantly changing and in the grand scheme these changes would have little impact on the overall quality of the data and subsequent analyses. Instead of making the change, another possible solution for consideration is to lower the **ID conf** from **Certain** to **Probable** or **Uncertain** and adding a note that the “YYYY botanist believes species X may be species Y.” The updated **ID conf** and note would be on the field form for the next sampling event. The future botanist would be alerted to the fact there was some uncertainty regarding that taxonomic concept.

Notes on Collecting

It is very important that all data collectors for this SOP attempt to thoroughly catalog the species within each of the nested subplots and the plot. By doing this, we hope to minimize the error associated with differences in data collectors’ abilities to find the same specimens and maximize the power to detect change over time. A secondary goal of monitoring efforts should be to enhance plant species lists for each park, so a species new to the park should be collected **if** a good specimen is available and per guidelines below. All relevant specimens or photographs collected will be sent to Curatorial Specialist Brenda Wells, Cumberland Piedmont Network, P.O. Box 8 Mammoth Cave, KY 42259. Following are notes on specimen collection and subsequent identification:

- For any Early Detection invasive species observed in the plot, make sure the Incidental Observation form is filled out and a specimen or photo has been collected to accompany the form. Early detection plant species found on the plot will not be targeted for removal because a management action that specifically targets a plot will compromise the integrity of the plot. Instead, try to locate additional populations of the species outside of the plot so those populations can be targeted, or if the species is very widespread, the plot will be part of a larger management action that doesn’t specifically target the plot.

- Do not collect any plants represented by three or fewer individuals (i.e., locally sparse) in the general vicinity of the plot. If a plant is considered locally sparse and there are individuals outside of the plot, consider taking your collection from outside of the plot.
- Do not remove entire plants (including roots) from within the plot. If the plant occupies greater than 2% cover and is sufficiently robust, a portion of the plant may be removed from within the plot so the overall cover of the plant within the plot is not significantly affected.
- If it is not possible to collect a specimen, take a photograph. First take a photo of the plot sheet so we can easily identify which plot the specimen photo is from. Then place a ruler or an object of known size like a pencil or pen in the photograph for scale and allow the camera to automatically select the appropriate aperture for the given light conditions. Use the Macro setting or Super Macro setting to take detailed close-up photographs of smaller plants.

If a specimen is the same species as one already collected in a previous plot, then mark which plot it was from in the **Collected** column and rely on that for specimen identification.

- For a specimen that isn't identifiable to species (particularly vegetative *Carex* sp. and *Dichanthelium* sp. specimens), it is ok to collect, but mark it clearly as not identifiable and continue to look for a viable specimen in other plots in the park that you can link this specimen to. Do not spend a lot of time on species identification, particularly if spikes or other distinguishing characters are absent. The recommended approach is to use **P** or **U** for **ID confidence** or write the genera, with notes on characters used to distinguish it from other species in the plot. Spending less time on these genera will increase efficiency on the plot and during data validation and reduce turnover between plot samples.
- Keep only quality specimens for vouchers. Voucher specimens must have clearly distinguishable features, such as flowers or fruit. Do not voucher specimens that were given an **ID confidence** of **P** or **U**.
- If you collect a specimen and check the **Collected** column, but later realize you cannot use the specimen and discard it, ensure that you cross out the checkmark in the **Collected** column so others do not mistakenly think that a specimen has been lost.

Equipment List

- Two collapsible fiberglass tent poles, pre-cut to 3.16 meters in length
- Pin flags (20–30)
- 3 meter sticks
- DBH Tape (in centimeters), pocket-sized if available
- Small binoculars

- Hand lens
- Plant press and newspapers (should not be carried in the field)
- Zip seal plastic bags in case you need to collect plant specimens DBH Tape (in centimeters)
- Camera
- Tree identification references
- Vascular plant identification references

Completing Field Data Forms

A detailed list of field descriptions for each field on the field data form associated with this SOP (Figure 5 and 6) can be found below in Table 3.

SOP 7

COWP001

Start Date: / /21
(mm/dd/yy)

Seedling and Sapling

		1	2	3	4	5	6	7	8	9	10	11	12	etc.
Module- Corner	Species Code	Seedlings Ht Classes (cm)				Tree Saplings by DBH (cm)								
	line #	5-15	15-30	30-50	50-137	0-1	1-2.5	2.5-5	5-10					
	1													
	2													
	3													
	4													
	5													
	6													
	7													
	8													
	9													
	10													
	11													
	12													
	13													
	14													
	15													
	16													
	17													
	18													
	19													
	20													
	21													
	22													
	23													

Notes line # _____

_____ p. 10 (____) of 12 (____)

Figure 6. Example CUPN Forest Vegetation pre-printed data form (revisit form) for recording seedling and sapling data.

Table 3. Field data form field names and descriptions for this SOP.

Field Name	Field Description
Botanist Initials	Initials of botanist assigned to plot and designated as final authority for all species identifications.
Reviewed by	Initials of individual who reviewed completed data sheet for this SOP elements. In most situations this should be someone other than individual who recorded data on this portion of datasheet. Botanist can conduct this review while recording cover values.
Module or Corners Sampled	With few exceptions, eight corners are sampled within each plot and the module and corner designations are pre-populated on revisit forms. If for some reason (e.g., crew safety), a corner is not sampled the corner and module designation should be crossed out and a line drawn down through the two columns of the first page.
Corner Notes	If less than eight corners sampled or if different from default corners sampled, record information in this block. Be sure to include the identification of all corners that were sampled (e.g., 1–3, 2–4).
Species Code	Seven or in some cases eight letter species code, typically comprised of first four letters of genus and first three letters of species. Refer to park species list to determine which species require the eight letter tie breaker codes. In instances where a species does not occur on the species list (i.e., no known species code), write out the entire scientific name. Field crew leader should confirm spelling. For revisit plots, a pre-populated species list will be provided. Update/correct as necessary.
Line #	Refer to species-specific note written out at the end of the species list.
ID Confidence	In cases where species identification is not certain record confidence level as either C=certain, P=probable, or U=uncertain.
Collect	Place a check mark in this column if field crew leader collects plant specimen for later identification and/or to make a species voucher.
Module 1-2, 1-4, ... 4-3	For each corner, record the level at which a species is first recorded where 3 = 1 × 1 meter (3.3 × 3.3 ft) subplot and 2 = 3.16 × 3.16 meter (10.4 × 10.4) subplot. If a species is not recorded in either corner but is detected in the module, record it as a 1 in either the first or second corner for that module (second preferred).
Cover Class (20 × 20 meter [65.6 × 65.6 ft])	Botanist should record a cover class for each species for the overall plot where 1 = trace, 2 = 0.1–1%, 3 = 1–2%, 4 = 2–5%, 5 = 5–10%, 6 = 1–25%, 7 = 25–50%, 8 = 50–75%, 9 = 75–95%, and 10 = greater than 95%.
Notes (optional):	Write out note for a specific species mentioned in Note # column.
Seedling/sapling only: Module and Corner	Record the module and corner combination where the seedling or sapling species is found.
Seedling/sapling only: Species Code	Seven or in some cases eight letter species code, typically comprised of first four letters of genus and first three letters of species. Refer to park species list to determine which species require the eight letter tie breaker codes. In instances where a species does not occur on the species list (i.e., no known species code), write out the entire scientific name. Field crew leader should confirm spelling.

Field Name	Field Description
Seedling/sapling only: Seedling Height Classes centimeters (cm): 5–15 cm 15–30 cm 30–50 cm 50–137 cm	Within the 1m x1m nested subplot (i.e., level 3), using tally method (outlined on top or bottom of datasheet) record the number of individuals within each height category for the species in question within the specified corner.
Seedling/sapling only: Tree Sapling by diameter breast height (DBH; centimeters): 0–1 cm DBH 1–2.5cm DBH 2.5–5cm DBH 5–10cm DBH	Within the 3m x 3m nested subplot (i.e., level 2, which includes the level 3 nested subplot), using tally method (outlined on top or bottom of datasheet) record the DBH of tree species which are at least 137 cm (i.e., DBH height) tall and less than 10cm DBH into the appropriate size category. Note, all trees 10cm DBH or above will be recorded on the tree data sheet (SOP 6 [CUPN 2023b]).

Literature Cited

- Cumberland Piedmont Network (CUPN) 2023a. Recording species nomenclature—Version 1.37. Cumberland Piedmont Network Standard Operating Procedure NPS/CUPN/SOP—14. Cumberland Piedmont Network, Mammoth Cave, Kentucky. Available at: <https://irma.nps.gov/DataStore/Reference/Profile/2299613> (last accessed October 2023).
- Cumberland Piedmont Network. 2023b. Tree measurements—Version 1.48. Cumberland Piedmont Network Standard Operating Procedure NPS/CUPN/SOP—6. Cumberland Piedmont Network, Mammoth Cave, Kentucky. Available at: <https://irma.nps.gov/DataStore/Reference/Profile/2299607> (last accessed October 2023).
- Lee, M. T., R. K. Peet, S. D. Roberts, and T. R. Wentworth. 2008. CVS-EEP Protocol for Recording Vegetation, All Levels of Plot Sampling. Version 4.2. Carolina Vegetation Survey. Available at: <http://cvs.bio.unc.edu/protocol/cvs-EEP-protocol-v4.2-lev1-5.pdf>

Revision Log

Revision History Log.

Version #	Date	Revised by	Changes
1.1	01/15/12	B. Moore and R. White	Time allotted was changed from 45 to 55 minutes (Note: unsure of the significance of this change as teams have never been on a time limit when recording data – bmoore 2/24/2020). Clarified that sampling window extends late April to late September with plot (park) revisits occurring same time each year within two weeks of initial sampling. Noted a large number of collections from 2011 were Carex and Dichanthelium lacking identifiable inflorescences. Pointed out it is ok to not spend large amount of time attempting to identify as the species within these genera will likely be lumped for analysis. Additional tips to streamline collections process were provided. Depth 1 (10 x 10 m) time constraint changed from 5 to 10 minutes (As noted, previously botanists have never been on a time limit when recording data; in other words, this “limit” has been ignored – bmoore 2/24/2020).
1.2	09/27/12	B. Moore and R. White	Added step that if a collection for later identification is not useable/identifiable the checkmark in the collected column on the field data form should be changed to an X to indicate no further identification is planned or possible. Emphasized the process for assigning species cover classes should be a relatively quick process in the field.

Version #	Date	Revised by	Changes
1.3	4/4/2018	C. Bledsoe and B. Moore	Note some edits were made prior to 2017 field season. However, these edits remained draft(?) and were rolled into an additional round of edits made prior to the 2018 season. In short, this version contains edits made prior to two seasons but only implemented (i.e. utilized by field crews) beginning in the 2018 season. "Time allotted: 55 minutes" changed to Time estimate: 120 minutes" (As noted earlier in this revision log, we are unsure of the significance of this change as teams have never been on a time limit when recording data – bmoore 2/24/2020). Opening paragraph references soils data. A note was added that as of 2018 soils data collection had not been initiated. Added note that "if a sapling forks below 0.5 m measure each fork as a separate sapling." Clarified that vines are not counted as a seedling or sapling. Added note that if an intensive corner is not sampled a note should be made in Plot Layout Notes. Emphasized importance of asking questions of more experienced crew members if any confusion about how data are to be collected or recorded. Removed statement that Depth 1 "must be completed in 10 minutes or less" (As noted, previously botanists have never been on a time limit when recording data – bmoore 2/25/2020). Added text that there is no strict time limit for searching of residual species (Depth 1). If it has been 2+ minutes since a species has been added than this might be an indicator that it is time to move on to the next module. Removed sentence that cover class should be assigned quickly (completed in 10 minutes or less). Added reminder to record cover classes for species overhanging, but not actually rooted in plot. Added several notes/reminders regarding revists including: Avoid extra time spent going back to previous modules in an attempt to relocate all species from a previous sampling event; If a corner was not sampled in a previous sampling event for whatever reason (e.g., yellow jacket nest), it should still be sampled in the current sampling event; What to do if find obvious ID errors (from previous sampling event) on the preprinted species list for the current event. This included questions to consider when making this determination as to whether previous sampling event data should be modified; Made note on what to do if early detection species observed in the plot; Re-emphasized that it is OK to not identify every taxon to species level as it is not always possible particularly for genera such as Carex and Dichanthelium; Only quality specimens should be kept for vouchers.
1.33	3/29/2020	B. Moore	Added definition of (canopy) cover. Improved directions for documenting when one of the eight corners is not sampled. Added note discouraging the use of minimally experienced crew members as data recorders. Improved directions for establishing 10 m ² subplots including adding a more robust method using incremented light duty chains as an alternative. Identified potential tasks that could be performed by extra or minimally trained field crew members. Added cover table with category ranges and some descriptors (Table 2) to aid in making determinations between classes. Moved Revision History to end of SOP and added more detailed list of changes made to date.
1.34	8/10/2020	B. Moore	Address grammatical changes and clarifying comments made by IMD regional program manager.

Version #	Date	Revised by	Changes
1.35	2/19/2021	B. Moore	Added image of field data forms and table of field data form descriptions (moved from SOP 14). Added reminders on the importance of making sure all tree species/module combinations from tree data have a corresponding presence value in the herb data forms.
1.37	1/14/2022	B. Moore	Added instructions on delineating the 10 m ² subplots using precut tent poles and deleted instructions using a chain or additional tape measures, since this is no longer a preferred approach. Added text encouraging field crews to collect seedling (Depth 3) and sapling (Depth 2) count data prior to turning attention to herb species. The reason for this is to reduce possibility that some or all of the seedling/sapling data might be overlooked. Added text recommending the clipboard and datasheets be handed off to the botanist at the module (Depth 1) level. Justification for this is also included in the added text. Removed SOP specific equipment list as the plot equipment checklist is regularly reviewed and maintained in SOP 1.
1.40	8/7/2023	B. Moore	Minor editorial changes including a new title resulting from regional IMD manager review. Formatted by editor to regional I&M publication standards.