



Sierra Nevada Network White Pine Monitoring

2022 Annual Report

Natural Resource Data Series NPS/SIEN/NRDS—2023/1402



ON THE COVER

Beetle-killed whitebark pine (*Pinus albicaulis*), near Plot 52, South Fork of Kings River drainage, Kings Canyon National Park, California.

NPS / SEAN AUCLAIR

Sierra Nevada Network White Pine Monitoring

2022 Annual Report

Natural Resource Data Series NPS/SIEN/NRDS—2023/1402

Kimiora Ward

National Park Service
Yosemite National Park
9736 Chapel Lane
El Portal, CA 95318

December 2023

U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science
Fort Collins, Colorado

The National Park Service, Natural Resource Stewardship and Science office in Fort Collins, Colorado, publishes a range of reports that address natural resource topics. These reports are of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Data Series is intended for the timely release of basic data sets and data summaries. Care has been taken to assure accuracy of raw data values, but a thorough analysis and interpretation of the data has not been completed. Consequently, the initial analyses of data in this report are provisional and subject to change.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible and technically accurate.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available in digital format from the [Sierra Nevada Network Inventory & Monitoring Program website](#) and the [Natural Resource Publications Management website](#). If you have difficulty accessing information in this publication, particularly if using assistive technology, please email irma@nps.gov.

Please cite this publication as:

Ward, K.L. 2023. Sierra Nevada Network white pine monitoring: 2022 annual report. Natural Resource Data Series NPS/SIEN/NRDS—2023/1402. National Park Service, Fort Collins, Colorado. <https://doi.org/10.36967/2301003>

Contents

	Page
Figures.....	iv
Tables	v
Executive Summary	vi
Acknowledgments.....	viii
Background and Objectives	1
Whitebark Pine	2
Foxtail Pine.....	4
Objectives	4
Methods.....	6
Sampling Frame.....	6
Frequency and Timing of Sampling	10
Plot Layout	10
Plot Measurements	11
2022 Sampling Logistics	14
Results.....	15
Yosemite	15
Sequoia and Kings Canyon	17
Whitebark Pine	19
Foxtail Pine.....	20
Summary	22
Literature Cited	24
Appendix 1. Notes on Field Methods	27
Appendix 2. Plot Status and Location.....	28

Figures

	Page
Figure 1. Distribution of whitebark pine, limber pine, and foxtail pine (from Little 1971) and locations of three Pacific West Region networks and associated parks.	2
Figure 2. Whitebark pine in Upper Kern, Sequoia National Park.....	3
Figure 3. Foxtail pine in the northern Kern River drainage, Sequoia National Park.	4
Figure 4. Whitebark pine sampling frame (blue shading) and GRTS-based plot locations for Yosemite National Park (colored dots).....	7
Figure 5. Whitebark pine sampling frame (blue shading) and GRTS-based plot locations for Sequoia and Kings Canyon National Parks (colored dots).	8
Figure 6. Foxtail pine sampling frame (green shading) and GRTS-based plot locations for Sequoia and Kings Canyon National Parks (colored dots).	9
Figure 7. 50 x 50 m permanent plot layout used in SIEN white pine monitoring.	11
Figure 8. Location of plots sampled in Yosemite during the 2022 field season.....	16
Figure 9. Location of 24 plots sampled in Sequoia and Kings Canyon during the 2022 field season.....	18
Figure 10. 2022 SIEN forest monitoring crew near South Guard Lake in Kings Canyon National Park..	23

Tables

	Page
Table 1. Revisit design for monitoring white pine species in the Sierra Nevada Network.	10
Table 2. Relationship among measured variables, data, and objectives for long-term monitoring of white pine communities in the Pacific West Region.....	12
Table 3. Summary statistics of Panel 3 whitebark pine plots measured at YOSE in 2022 (n = 12).....	17
Table 4. Summary statistics on whitebark pine plots sampled in SEKI in 2022 (n = 12).	19
Table 5. Summary statistics of Panel 3 foxtail pine plots visited at SEKI in 2022 (n = 12).	20

Executive Summary

Five-needle white pines (Family Pinaceae, Genus *Pinus*, Subgenus *Strobus*), and in particular whitebark pine (*Pinus albicaulis*), limber pine (*P. flexilis*), and foxtail pine (*P. balfouriana*) are foundation species in upper subalpine and treeline forests of several National Park Service Pacific West Region parks, including Sequoia and Kings Canyon National Parks (SEKI) and Yosemite National Park (YOSE). The Sierra Nevada Network Inventory & Monitoring Program, in collaboration with the Klamath Network, Upper Columbia Basin Network, and Mojave Desert Network have implemented a joint long-term monitoring protocol to assess the current status and future trends in high elevation white pine communities. Key demographic parameters within white pine forest communities will be estimated by monitoring individual trees within permanent plots through time.

This report documents the results of the 2022 field season, which was the ninth year of monitoring in SEKI and YOSE. The 2022 goal was to complete the first full measure of the third of three rotating panels (Panel 3) for each species-park population: YOSE-whitebark pine, SEKI-whitebark pine, and SEKI-foxtail pine. Each panel consists of 12 permanent 50 x 50 m (2,500 m²) plots that were randomly selected for each of the three populations. The full sampling array thus includes a total of 36 whitebark pine plots in YOSE, 36 whitebark pine plots in SEKI, and 36 foxtail pine plots in SEKI. Data from plot surveys will be used to characterize white pine forest community dynamics in SEKI and YOSE, including changes in tree species composition, forest structure, forest health, and demographics.

Partial measures of Panel 3 were completed in 2017 (11 plots) in Yosemite whitebark pine, in 2017 (9 plots) in SEKI whitebark pine, and in 2014 (7 plots) and 2017-2018 (8, 1 plots) in foxtail pine. In 2022, the first full measure of all Panel 3 plots (and 2nd or 3rd remeasure of most plots) was successfully completed, and installation was completed on four of these plots in SEKI whitebark pine and two in foxtail pine.

In total, the crew visited 36 sites during the 2022 field season, all from Panel 3. Within the 36 completed Panel 1 plots, a total of 6,398 trees were measured. Species composition, forest structure, and factors affecting tree health and reproduction including incidence and severity of white pine blister rust (*Cronartium ribicola*) infection, mountain pine beetle (*Dendroctonus ponderosae*) infestation, dwarf mistletoe (*Arceuthobium spp.*) infection, canopy kill, and female cone production were recorded. During the 2022 field season crews continued to count the total number of mature cones per tree for whitebark and foxtail pine, use crown condition codes to assess crown health, and tag individual seedlings to be tracked through time. All three of these procedures started in 2017 and are to be evaluated by each of the three participating networks over several years, to determine whether they should become permanent changes to the monitoring protocol.

In YOSE, all 12 Panel 3 whitebark pine plots were measured. A total of 2,720 trees were sampled, which included 977 live whitebark pine trees and 1,605 other live conifers. An additional 135 trees (including 26 whitebark) were recorded as dead. The average number of live whitebark pine trees per plot was 81 (SD = 94). White pine blister rust (WPBR) aecia were observed on five whitebark pine in

one plot in YOSE in 2022, and no trees in any plot had inactive cankers showing three or more indicators of WPBR. WPBR had previously been documented in this plot, so the number of plots where rust has ever been observed in Yosemite remains unchanged at six. However, an infection documented in plot 42 in 2021 was not observed again when the plot was resampled in 2022, so it is possible this number should be five. Mountain pine beetle activity was observed on one live whitebark pine and three live and one dead lodgepole pine in YOSE in 2022. Despite documentation of many stands impacted by beetle attack in the field crew notes, the quantified rate of MPB attack was lower than in 2021. Twenty-one percent of live whitebark pine trees produced female cones. Cone-bearing trees averaged seven ($SD = 10$) cones/tree. Whitebark pine seedling density averaged 80 ($SD = 152$) seedlings per hectare. The largest number of whitebark pine seedlings found in a plot was 51 and five of the twelve plots contained whitebark seedlings.

All 12 Panel 3 SEKI whitebark pine plots were measured in 2022, and installation was completed on four of these, so this Panel is now fully installed. Within these plots, 2,179 live whitebark pine, 10 live foxtail pine, and 297 other live conifers were sampled (including five live western white pine). The average number of live whitebark pine trees per plot was 181 ($SD = 125$). Although the crew observed white pine blister rust in seven SEKI whitebark Panel 3 plots, no active cankers (aecia) were observed, and no trees displayed three of five indicators, so no infections were quantified. Mountain pine beetle activity was observed in 18 live and 23 dead whitebark pine and one live and one dead lodgepole pine within three plots in SEKI. Dwarf mistletoe was not encountered. Seven percent of live whitebark pine trees produced female cones. Cone-bearing trees averaged 3.7 ($SD = 3.6$) cones/tree. Whitebark seedling regeneration averaged 700 ($SD = 752$) seedlings per hectare. The largest number of whitebark seedlings found in a plot was 19, and two of the 12 plots did not contain any whitebark seedlings.

In the foxtail pine Panel 3, all 12 plots were measured in 2022, and installation was completed on two of these, so installation of the panel is now complete. Within these plots we measured 309 live foxtail pine, 302 live whitebark pine, and 380 other live conifers, including four live western white pine. An additional 112 dead or recently dead trees and 22 unidentified snags were also measured, 19 of which were foxtail pine. The average number of foxtail pine trees per plot was 26 ($SD = 26$). No signs of blister rust infection or mistletoe were observed on foxtail pine. Mountain pine beetle activity was observed on one dead foxtail pine, one live whitebark pine, and seven live and one dead lodgepole pines within four plots. Sixty-two percent of the foxtail pine trees produced female cones. Cone-bearing trees averaged 33 ($SD = 53$) cones/tree. Seven foxtail pine seedlings were recorded within five plots, resulting in an estimated 72 ($SD = 98$) seedlings per hectare. Eight whitebark pine seedlings and three lodgepole pine seedlings were also found within three additional plots.

Acknowledgments

Funding for this project was provided through the National Park Service Inventory and Monitoring Program. Monica Buhler, Devils Postpile National Monument, provided valuable logistical support to field operations in 2022 and supervised the crew for the first half of the season. The initial high-elevation white pine monitoring annual report (Stucki et al. 2012) on which this document is based was developed by Devin Stucki, Tom Rodhouse, and Shawn McKinney. This work would not have been possible without the amazing contributions of our dedicated field crew: Sean Auclair (lead), Emma Kahle, and Ana Tobio.

Background and Objectives

Many western North American coniferous forests are currently facing unprecedented health challenges, including upsurges of native pests and pathogens, invasive exotic species, and altered disturbance regimes. Increased atmospheric warming, carbon dioxide concentration, and nitrogen deposition, as well as changes in precipitation patterns (i.e., timing, magnitude, and type) portend additional short- and long-term changes in high elevation forest ecosystem processes. Each factor alone can alter forest structure, function, and species composition, and additive or synergistic effects are likely if multiple agents act jointly. How forest ecosystems will respond to modern perturbations is uncertain. However, the magnitude of change in structure, composition, and key ecological processes will likely be exceptional. Indeed, increased tree mortality rates over the last several decades have recently been documented across a broad range of latitude and forest types in western North America (van Mantgem et al. 2009), which may have important consequences for forest stand dynamics and ecosystem functions.

Five-needle white pines (Family Pinaceae, Genus *Pinus*, Subgenus *Strobus*), and in particular whitebark pine (*Pinus albicaulis*), limber pine (*P. flexilis*), and foxtail pine (*P. balfouriana*) are foundational species (Tomback and Achuff 2010) in upper subalpine and treeline forests of several National Park Service (NPS) Pacific West Region (PWR) parks, including Sequoia and Kings Canyon National Parks (SEKI) and Yosemite National Park (YOSE) (Figure 1). Ongoing declines of foundation tree species pose an especially compelling problem because these species provide fundamental structure to ecosystems and are therefore irreplaceable (Ellison et al. 2005). If a foundation tree species is lost from these systems, it will likely lead to a cascade of secondary losses, shifts in biological diversity, and ultimately affect the functioning and stability of the community (Ebenman and Jonsson 2005).




Figure 1. Distribution of whitebark pine, limber pine, and foxtail pine (from Little 1971) and locations of three Pacific West Region networks and associated parks. Sampling also occurs at Great Basin National Park, within the Mojave Desert Network, which is not shown in this figure.

Whitebark Pine

Whitebark pine occurs across a broad geographic range, occurring on both the west and the more arid east side of the Sierra Nevada crest and reaching its southern limit in central California in the Mount Whitney vicinity. Throughout its range, whitebark pine can occur in the montane, upper subalpine, and treeline zones (Arno and Hoff 1990). It often occurs as the only tree species on the coldest and driest sites near treeline (Figure 2) and as a seral species on protected, slightly lower sites more favorable to its shade-tolerant competitors (Arno and Weaver 1990).



Figure 2. Whitebark pine in Upper Kern, Sequoia National Park. NPS/ EMMA KAHLE

In the Pacific West Region (PWR), whitebark pine is scattered across tens of thousands of hectares in the high elevations of SEKI and YOSE (Figure 1). White pine blister rust (*Cronartium ribicola*) infections on whitebark pine decrease from north to south in the PWR, resembling the trend seen in the Rocky Mountains. White pine blister rust (WPBR) is relatively rare in high elevation forests of SEKI and YOSE when compared to northern portions of the PWR (e.g., North Cascades). Mountain pine beetles (*Dendroctonus ponderosae*) are currently abundant in the northern Cascades, but also decrease with latitude (Gibson et al. 2008) in the PWR.

Whitebark pine acts as a foundation species in high-elevation forest communities by regulating ecosystem processes, community composition and dynamics, and by influencing regional biodiversity (Ellison et al. 2005, Tomback and Kendall 2001). Whitebark pine plays a role in initiating community development after fire, influencing snowmelt and stream flow, and preventing soil erosion at high elevations (Tomback et al. 2001, Farnes 1990). The large, wingless seeds of whitebark pine are high in fats and carbohydrates, and provide an important food source for many granivorous birds and mammals (Tomback and Kendall 2001). Whitebark pine is a coevolved mutualist with Clark's nutcracker (*Nucifraga columbiana*) and is dependent upon nutcrackers for dispersal of its seeds (Tomback 1982, McKinney et al. 2009).

Foxtail Pine

Foxtail pine is endemic to two distinct areas in California, the Klamath Mountains in the northwest part of the state and the southern Sierra Nevada (Figure 1). Research on community and population dynamics is lacking for foxtail pine compared to whitebark pine. Foxtail pine occurs in four different forest types: 1) stands dominated by foxtail pine, 2) stands with foxtail pine and whitebark pine, 3) stands with foxtail pine and red fir (*Abies magnifica*), and 4) stands with foxtail pine, red fir, and western white pine (*P. monticola*) (Eckert and Sawyer 2002). Foxtail and whitebark pine overlap in some portions of their southern Sierra Nevada distribution, however, south of the Kings-Kern divide, foxtail pine is the major (sometimes exclusive) subalpine and treeline tree species (e.g., >3,000 m). Foxtail pine provides important habitat and food resources for birds and mammals, and influences snowmelt and soil erosion (Figure 3).



Figure 3. Foxtail pine in the northern Kern River drainage, Sequoia National Park. NPS /EMMA KAHLE

The southern population of foxtail, subspecies *austrina*, provides important data for dendrochronological research on paleoclimate (Lloyd 1997) as a consequence of its great longevity (> 2,000 years) and slow growth. In fact, five-needle pines, in general, have proven valuable in enhancing our understanding of past climates through dendrochronological investigations (e.g., Kipfmüller and Salzer 2010, Woodhouse et al. 2011).

Objectives

At least four networks in the Pacific West Region identified white pine species as targets for long-term monitoring (Sarr et al. 2007, Mutch et al. 2008, Garrett et al. 2007, Hupp et al. 2018). Ecologists in three of these networks (Klamath Network [KLMN], SIEN, and Upper Columbia Basin Network [UCBN]) collaborated to devise a common set of monitoring objectives and procedures which are documented in our multi-network white pine monitoring protocol (McKinney et al. 2012a, 2012b). This protocol was later implemented in Great Basin National Park (GRBA) by the Mojave

Desert Network (MOJN) in 2018 (Hupp et al. 2018). The anticipated impacts from WPBR, dwarf mistletoe, mountain pine beetle, and climate change on high-elevation pines were primary factors considered in developing the monitoring objectives. Key demographic parameters within white pine forest communities will be estimated by monitoring individual trees within permanent plots. Specific objectives of white pine monitoring are to detect status and trend in:

- 1) Trees species composition and structure
- 2) Tree species birth, death, and growth rates
- 3) Incidence of white pine blister rust and level of crown kill
- 4) Incidence of bark beetles
- 5) Incidence of dwarf mistletoe
- 6) Cone production of white pine species

Information gathered from this white pine monitoring project will be integral to providing a more comprehensive understanding of the populations within SIEN parks as well as providing for comparisons across broader geographic areas. It will also allow early detection of important changes in populations that may trigger management intervention.

Methods

This section summarizes the methods used for white pine monitoring in SIEN in 2022. A full description of methods and standard operating procedures may be found in the multi-network white pine monitoring protocol (McKinney et al. 2012a, 2012b). Specific deviations from the published methods or clarifications on how the methods were interpreted in the field during the 2022 field season are documented in Appendix 1.

Sampling Frame

The sample frames for each species-park population (SEKI-whitebark pine, SEKI-foxtail pine, and YOSE-whitebark pine) were based on the distribution of whitebark and foxtail pine as identified in the YOSE and SEKI vegetation maps (Figures 4 through 6). An ordered list of plot locations was generated using a randomized, spatially-balanced sampling design via the Generalized Random Tessellation Stratified (GRTS) algorithm (Stevens and Olsen 2004). Plots in the ordered list were assigned to one of three panels and each panel will be sampled on a rotating basis every three years (Appendix 2). Extra plot locations were drawn in the event that plots from the original sample draw were rejected (the oversample). Common reasons for rejection included 1) plots were on slopes too steep to be safely sampled (slopes ≥ 35 degrees), 2) sites were not accessible in a safe manner, and 3) sampling frame errors (i.e., plot did not contain the target vegetation type). Based on this sampling design, our scope of inference extends broadly across mapped stands of whitebark and foxtail pine on < 35 -degree slopes within YOSE and SEKI.




Figure 4. Whitebark pine sampling frame (blue shading) and GRTS-based plot locations for Yosemite National Park (colored dots). Plots are assigned to one of three panels (12 plots each), which are sampled every three years.




Figure 5. Whitebark pine sampling frame (blue shading) and GRTS-based plot locations for Sequoia and Kings Canyon National Parks (colored dots). Plots are assigned to one of three panels (12 plots each), which are sampled every three years, or the oversample.




Figure 6. Foxtail pine sampling frame (green shading) and GRTS-based plot locations for Sequoia and Kings Canyon National Parks (colored dots). Plots are assigned to one of three panels (12 plots each), which are sampled every three years.

Frequency and Timing of Sampling

We adopted a three-year rotating panel design for re-surveying permanent plots in SEKI and YOSE (McDonald 2003). Sampling typically occurs between June and October and each plot is surveyed once per three-year rotation (Table 1). A total of 36 plots are monitored in each park (YOSE and SEKI) for each species, resulting in an overall total sampling effort of 72 plots in SEKI (36 whitebark and 36 foxtail) and 36 plots in YOSE (whitebark only).

Table 1. Revisit design for monitoring white pine species in the Sierra Nevada Network. This panel design is followed for each of the 3 species-park populations (YOSE-whitebark pine, SEKI-whitebark pine, and SEKI-foxtail pine) for a total SIEN $n = 108$ plots.

Panel	Year ^a												
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
1 (n = 12)	x	–	–	–	x	–	–	– ^b	x	–	–	–	x
2 (n = 12)	–	– ^b	x	–	–	x	–	–	–	– ^b	x	–	–
3 (n = 12)	–	–	–	x	–	–	x	–	–	–	–	x	–

^a An “x” indicates the years that the corresponding panel of monitoring plots was sampled. An “–” (or endash) indicates that the corresponding panel of monitoring plots was not sampled that year.

^b Signifies the panels of monitoring plots originally planned to be sampled but were not sampled those years.

Plot Layout

Quarter-hectare (50 x 50 m) macroplots consisting of five subplots are used to measure and track forest demographic parameters, disease, and insect occurrence, and the magnitude of their impact (Figure 7). The response design for this protocol is compatible with the *Interagency Whitebark Pine Monitoring Protocol for the Greater Yellowstone Ecosystem* (GYWPMWG 2007) but differs in some respects, most notably, plot size. The 10 x 50 m plot size from the Yellowstone protocol has been increased to accommodate the often sparse or patchy distribution of white pines in our PWR parks and to adequately address forest demographic objectives. This design effectively represents five parallel 10 x 50 m subplots as used in the GYWPMWG and as proposed by the Whitebark Pine Ecosystem Foundation (Tomback et al. 2005).

A total of nine square regeneration plots (3 x 3 m) are established within each macroplot to measure seedling regeneration (Figure 7). Regeneration plots are located at each corner (4), at each midpoint between corners (4), and in the middle (1) of the macroplot. The current design was chosen because it provides a reasonable balance among sampling time constraints, observer accuracy and precision, and total area sampled.




Figure 7. 50 x 50 m permanent plot layout used in SIEN white pine monitoring. Smaller regeneration plots are where seedlings are measured.

Plot Measurements

Table 2 describes variables measured, raw data collected, summarized values, and the monitoring objectives addressed. Detailed instructions on response design measurements are provided by McKinney et al. (2012a, 2012b) and only a general overview is presented here.

Table 2. Relationship among measured variables, data, and objectives for long-term monitoring of white pine communities in the Pacific West Region. p/a indicates presence/absence.

Variable	Raw Data	Summarized Data	Objectives Addressed
Species	Tree (nominal)	Trees per hectare (TPH); all spp., each spp., proportion of total by spp.	1. composition & structure
Diameter	Tree (cm)	Basal area (m ² /ha); all spp., each spp., proportion of total by spp. Mean diameter (cm) by spp. Diameter classes (5 cm); proportion and TPH by spp.	1. composition & structure 2. growth rate
Height	Tree (m)	Mean ht. (m); all spp. and by each spp. Height classes (3 m); proportion and TPH by spp.	1. composition & structure 2. growth rate
Status	Tree (live or dead)	Proportion live and dead; all spp and by each sp. TPH and proportion by 5 cm diameter classes in each condition; all spp and by each sp.	2. birth and death rates
Crown kill	Each of three parts of a tree (%)	Mean (%) individual white pine trees.	3. level of crown kill
Active canker	Each of three parts of a tree (p/a)	Proportion and TPH with active cankers by each white pine sp.	3. rust infection incidence
Inactive canker	Each of three parts of a tree (p/a)	Proportion and TPH with inactive cankers by each white pine sp.	3. rust infection incidence
Rust infection	Tree (p/a of active <u>or</u> inactive canker)	Proportion and TPH infected and healthy by each white pine sp. TPH by 5 cm diameter classes in each condition by each white pine sp.	3. rust infection incidence
Bark beetle	Tree (p/a)	Proportion and TPH with beetle sign; all spp and each sp.	4. incidence of bark beetle
Dwarf mistletoe	Tree (p/a)	Proportion and TPH with mistletoe sign; all spp and each sp.	5. incidence of dwarf mistletoe

Table 2 (continued). Relationship among measured variables, data, and objectives for long-term monitoring of white pine communities in the Pacific West Region. p/a indicates presence/absence.

Variable	Raw Data	Summarized Data	Objectives Addressed
Female cones	Tree (p/a)	Proportion and TPH with cones by each white pine sp.	6. cone production
Seedlings	9 m ² plot; number of each of three size classes by species	Mean (number per m ²); all spp and each sp for each size class.	1. composition & structure 2. birth rates

Each live tree at least 1.37 m tall is affixed with a uniquely numbered metal tag; its species is identified, and diameter at 1.37 m (breast height, DBH) and tree height are measured. Standing dead trees ≥ 5 cm DBH are also affixed with a metal tag and are recorded as either recently dead or dead. Recently dead trees have needles present (but no green needles) and dead trees have no needles present. For plots that are re-measured, recently dead trees are defined as those that were alive during the previous visit but are now dead and dead trees are those that were already dead. White pine blister rust infection is assessed for all living white pine trees. The bole and branches of white pine trees are each vertically divided into thirds (upper, middle, and bottom) and each third is assigned one of three rust condition classes: 1) absent—no sign of rust infection, 2) active cankers (aeciospores present), or 3) no active cankers, but with the presence of at least three of the following five indicators of infection: rodent chewing, flagging, swelling, roughened bark, and oozing sap. Mountain pine beetle occurrence is recorded for all pine trees using three indicators of beetle activity: pitch tubes, frass, and J-shaped galleries. The presence of galleries is only determined for recently dead and dead trees because bark has to be removed for this assessment (unless bark is already missing and galleries can be observed on the live tree). Current dwarf mistletoe infection is recorded for all living white pine trees by noting presence or absence of mistletoe for each third of a tree using the Hawksworth (1977) rating system. The level of canopy kill in live trees is determined by dividing the tree's canopy (all the main branches, encompassing all foliage and supporting twigs and side branches) into thirds and obtaining an ocular estimate of the percentage of each third of the canopy that is dead. Cone production is recorded based on whether female cones are present or absent on each live whitebark or foxtail pine tree. Starting in 2017 we also counted mature female cones on target species to assess cone abundance in addition to presence/absence. This was continued in 2022. Live seedlings are recorded by species and assigned to a height class in regeneration plots. Height classes are: 1) 20 to <50 cm, 2) 50 to <100 cm, and 3) 100 to <137 cm. Seedlings <20 cm are not measured. Beginning in 2017 all seedlings in the regeneration plots were tagged by using wire to attach a tag around the base of the stem. These seedlings will be individually tracked in the same way as the mature trees going forward.

2022 Sampling Logistics

A three-person crew was hired to establish or re-measure up to 36 plots within YOSE and SEKI during the 2022 field season. Training occurred over several weeks in June at SEKI and included information on forest pathology by the SIEN Ecologist and USFS Region 5 Entomologist and Pathologist. Training on safety, backcountry communication, and project-specific sampling methods was provided by SIEN. Sylvia Haultain, Sierra Nevada Inventory and Monitoring Network Program Manager, and Monica Buhler, Devils Postpile, Natural Resources Program Manager, supervised the crew until the new SIEN ecologist, Erika Blomdahl, started in early July. Forest monitoring fieldwork occurred in SEKI in June – July and the first half of September, and in YOSE in August. Detailed notes for each trip and route descriptions were recorded by the crews and will be used in future years to help guide planning. These documents are stored in the forest project folder on the SIEN network drive. For a complete list of plot status and locations see Appendix 2.

Results

Yosemite

On June 6, and from August 2 to 23, 12 plots were sampled throughout whitebark pine stands in Yosemite National Park (Figure 8). All sampled plots were from Panel 3.

Summary statistics for the YOSE whitebark pine plots are provided in Table 3. The 12 completed plots contained a total of 2,720 trees: 977 live whitebark pine (14 recently dead, 12 dead), 1,512 live lodgepole pine (41 recently dead, 58 dead), 93 live mountain hemlock (*Tsuga mertensiana*) (two recently dead, eight dead), 3 dead trees of unidentified species. Of the 977 live whitebark pine trees, 24% displayed a krummholz growth form and approximately 30% of stems grew in clumps, with the average stems per clump being 3.1 (SD = 1.8). The average number of live whitebark pine trees per plot (2,500 m²) was 81 (SD = 94) with a range of 3 to 256 trees.

The forest crew noted five YOSE_PINALB Panel 3 plots with signs and symptoms of white pine blister rust in their field notes. However, only five trees in a single plot, YOSE_PINALB_56 near Merced Peak, met the criteria for infection. All five trees had active cankers, and for four of them, these cankers were on the bole, suggesting the infections were not new. None of these trees, nor any other trees in the panel, had inactive cankers showing 3 of 5 signs of infection, so no additional infections were documented. WPBR had been previously documented in plot YOSE_PINALB_56, so the number of plots where rust has ever been observed in Yosemite remained unchanged at six. However, the number of trees infected in plot 56 may have changed. In 2016, 12 trees met criteria for infection in this plot. All of these trees were still alive in 2022, but only one still met criteria for infection (3 out of 5 indicators), while the other 11 displayed just 2 of 3 indicators of rust this year. The four other infected trees in 2022 were new infections as of this year. In addition, Plot YOSE_PINALB_42 had one infected whitebark pine tree in 2021 when this plot was included in a Panel 2 sample round, but this tree had no indicators of rust in 2022. The five infected trees in YOSE_PINALB_56 led to an incidence of white pine blister rust infection across all 977 live whitebark pine trees of 0.5%, and an average per plot infection rate of 0.2% (SD = 0.5) of live whitebark trees. Mountain pine beetle activity was observed in four plots on four trees (1 live whitebark pine in YOSE_PINALB_56 and 3 live and 1 dead lodgepole pine in other plots). Dwarf mistletoe was encountered on a single whitebark pine tree in YOSE_PINALB_56.

Twenty-one percent of the 977 live whitebark pine trees produced female cones this year, with the cone crop varying among stands between 0 and 30% of live whitebark trees bearing cones. The number of cones per tree among all 204 cone-bearing trees averaged 7 (SD = 10). Whitebark pine seedling regeneration averaged 82 (SD = 152) seedlings per hectare. Lodgepole pine (14 seedlings) and mountain hemlock (15 seedlings) were also encountered in the regeneration plots.





Figure 8. Location of plots sampled in Yosemite during the 2022 field season.

Table 3. Summary statistics of Panel 3 whitebark pine plots measured at YOSE in 2022 (n = 12). Mountain pine beetle infestation rate is calculated using only live and recently dead host trees.

Plot Metrics	Average (SD)	Range
<i>P. albicaulis</i> density (trees/ha)	326 (376)	12 – 1024
Other species density (trees/ha)	535 (542)	0 – 1752
Snag density (dead trees/ha)	46 (44)	0 – 156
<i>P. albicaulis</i> DBH (cm)	10.1 (7.5)	2.4 – 30.7
Other species DBH (cm)	17.4 (8.3)	4.2 – 35.3
<i>P. albicaulis</i> Basal Area (m ² /ha)	4.2 (6.1)	0 – 20.1
Other species Basal Area (m ² /ha)	27.5 (25.0)	0 – 69.7
<i>P. albicaulis</i> blister rust infection rate (# of infected trees/ha)	1.7 (5.8)	0 – 20
Dwarf mistletoe infection rate (# of infected trees/ha)	0.3 (1.2)	0 – 4
Mountain pine beetle infestation rate (# of infested trees/ha)	1.3 (2.0)	0 – 4
<i>P. albicaulis</i> seedling regeneration 20-136 cm (seedlings/ha)	864 (1801)	0 – 6296
Other species seedling regeneration 20-136 cm (seedlings/ha)	298 (409)	0 – 1358
<i>P. albicaulis</i> female cone production (# of trees with cones/ha)	68 (107)	0 – 280
<i>P. albicaulis</i> # of cones/tree for cone-bearing trees	4.3 (3.2)	1 – 9

Sequoia and Kings Canyon

From June 26 to September 17, 2022, the forest crew sampled 12 plots in Panel 3 of the SEKI-whitebark pine population (Figure 9). They completed installation on three of these (SEKI PINALB plots 58, 67, and 79). From June 10 to September 6, 2022, they sampled 12 SEKI-foxtail pine plots in Panel 3. Installation was completed on SEKI PINBAL plot 46.



Whitebark Pine

Twelve whitebark plots were sampled in 2022 in SEKI Panel 3. Summary statistics are provided in Table 4. The 12 sampled whitebark plots contained a total of 2,598 trees: 2,179 live whitebark (58 dead, 38 recently dead), 252 live lodgepole (1 dead, 8 recently dead), 40 live mountain hemlock (1 recently dead), 10 live foxtail pine, 5 live western white pine (1 dead), and 5 dead snags of unidentified species. The average number of live whitebark pines per plot was 181 (SD = 125) with a range of 16 to 395. Less than 1% (0.5%) of the 2,179 sampled live whitebark pine trees displayed a krummholz growth form, and 21% grew in clumps, with the average clump size 2.5 (SD = 0.8) stems.

Table 4. Summary statistics on whitebark pine plots sampled in SEKI in 2022 (n = 12). Mountain pine beetle infestation rate is calculated using only live and recently dead host trees.

Plot Metrics	Average (SD)	Range
<i>P. albicaulis</i> density (trees/ha)	726 (498)	64 – 1580
Other species density (trees/ha)	102 (125)	4 – 352
Snag density (dead trees/ha)	37 (52)	0 – 180
<i>P. albicaulis</i> DBH (cm)	11.1 (4.2)	5.2 – 19.6
Other species DBH (cm)	18.0 (12.1)	3.1 – 44.8
<i>P. albicaulis</i> Basal Area (m ² /ha)	13.5 (12.2)	0.7 – 35.0
Other species Basal Area (m ² /ha)	5.5 (7.3)	0 – 24.5
<i>P. albicaulis</i> blister rust infection rate (# of infected trees/ha)	0	0
Dwarf mistletoe infection rate (# of infected trees/ha)	0	0
Mountain pine beetle infestation rate (# of infested trees/ha)	14.3 (27.2)	0 – 72
<i>P. albicaulis</i> seedling regeneration 20-136 cm (seedlings/ha)	700 (752)	0 – 2,346
Other species seedling regeneration 20-136 cm (seedlings/ha)	123 (247)	0 – 864
<i>P. albicaulis</i> female cone production (# of trees with cones/ha)	52 (61)	0 – 212
<i>P. albicaulis</i> # of cones/tree per plot, for cone-bearing trees	4.9 (5.3)	2.5 – 20.0

The field crew noted white pine blister rust in seven SEKI_PINALB Panel 3 plots (plots 51, 52, 59, 60, 66, 67, and 73), however no active cankers (aecia) were observed, and no trees had 3 out of 5 indicators, so the criteria for documenting an infection were not met for any of the three five needle pine species present in the plots. WPBR has been observed in a total of five plots in the SEKI_PINALB sample frame in the past. Two of these were resampled this year and no WPBR was found. Three infected whitebark trees were observed in 2016 with inactive bole cankers in plot 51; these trees had zero indicators of rust when the site was resampled in 2021, and in 2022 they had 0, 1, and 2 indicators respectively (3 of 5 indicators are required to qualify as a WPBR infection). Plot 59 had one infected whitebark in 2016; this tree had an active bole canker (with aecia visible) and an

inactive branch canker in 2016, while in 2022 it displayed just 2 of 5 indicators at the bole canker. Mountain pine beetle activity was found in three plots, at relatively high rates: in plot 52 on 3 live and 14 recently dead whitebark pine and 1 recently dead lodgepole pine; in plot 60 on 6 live and 9 recently dead whitebark pine and 1 live lodgepole pine; and in plot 66 on 9 live whitebark pine. No dwarf mistletoe was encountered.

Cone production was moderate in 2022, with approximately 7% of the 2,179 live whitebark pine trees producing female cones, and the per-plot percentage ranging from 0 to 22%. Among the 157 cone-bearing whitebark pine trees observed in 2022, the average number of cones per tree was 3.7 (SE=3.6) with a maximum load of 20 cones. Cone production in foxtail pines within the whitebark plots was higher, with 20% of the ten live trees observed producing female cones, at an average of 28.5 cones per tree (SD = 23.3) and maximum of 45 cones per tree. There were 68 whitebark pine seedlings recorded in the plots, resulting in an estimated 700 (SD = 752) seedlings per ha. The largest number of seedlings found in a plot was nineteen, and these were whitebark pine seedlings. Ten of the 12 plots contained at least one whitebark seedling. One foxtail pine, one Jeffrey pine, and ten lodgepole pine seedlings were found.

Foxtail Pine

Twelve foxtail pine plots from Panel 3 were re-measured in 2022 (Figure 9). Summary statistics are provided in Table 5. The 12 foxtail pine plots contained a total of 1,080 trees: 309 live foxtail pine (18 dead, 1 recently dead), 302 live whitebark pine (11 dead, 19 recently dead), 375 live lodgepole pine (10 dead, 8 recently dead), four live western white pine, 1 Jeffrey pine and 22 dead snags of unidentified species. The average number of foxtail pine trees per plot was 26 (SD = 26). Symptoms of WPBR and mistletoe were not observed on any trees within the foxtail pine panel. Mountain pine beetle activity was documented in four plots, on one recently dead foxtail pine in Plot 34, six live and one recently dead lodgepole pine in Plot 40, one live whitebark pine in Plot 44, and one live lodgepole pine in Plot 45.

Table 5. Summary statistics of Panel 3 foxtail pine plots visited at SEKI in 2022 (n = 12). Mountain pine beetle infestation rate is calculated using only live and recently dead host trees.

Plot Metrics	Average (SD)	Range
<i>P. balfouriana</i> density (trees/ha)	103 (105)	4 – 348
<i>P. albicaulis</i> density (trees/ha)	101 (169)	0 – 452
Other <i>species</i> density (trees/ha)	127 (169)	0 – 400
Snag density (dead trees/ha)	37 (52)	0 – 180
<i>P. balfouriana</i> DBH (cm)	45.6 (28.0)	3.1 – 111.2
<i>P. albicaulis</i> DBH (cm)	9.1 (4.9)	4.2 – 16.3
Other species DBH (cm)	18.7 (12.1)	3.1 – 44.9
<i>P. balfouriana</i> Basal Area (m ² /ha)	21.0 (21.1)	0 – 73.5

Table 5 (continued). Summary statistics of Panel 3 foxtail pine plots visited at SEKI in 2022 (n = 12). Mountain pine beetle infestation rate is calculated using only live and recently dead host trees.

Plot Metrics	Average (SD)	Range
<i>P. albicaulis</i> Basal Area (m ² /ha)	1.7 (2.9)	0 – 8.1
Other species Basal Area (m ² /ha)	9.9 (14.1)	0 – 36.5
<i>P. balfouriana</i> blister rust infection rate (# of infected trees/ha)	0	0
<i>P. albicaulis</i> blister rust infection rate (# of infected trees/ha)	0	0
Dwarf mistletoe infection rate (# of infected trees/ha)	0	0
Mountain pine beetle infestation rate (# of infested trees/ha)	3.3 (8.0)	0 – 28
<i>P. balfouriana</i> seedling regeneration 20-136 cm (seedlings/ha)	72 (98)	0 – 247
<i>P. albicaulis</i> seedling regeneration 20-136 cm (seedlings/ha)	82 (219)	0 – 741
Other species seedling regeneration 20-136 cm (seedlings/ha)	31 (77)	0 – 247
<i>P. balfouriana</i> female cone production (# of trees w/ cones/ha)	64 (66)	0 – 188
<i>P. balfouriana</i> # cones/tree per plot, for cone bearing trees	36 (28)	5 – 84
<i>P. albicaulis</i> female cone production (# of trees w/ cones/ha)	4 (10)	0 – 36
<i>P. albicaulis</i> # cones/tree per plot, for cone bearing trees	8 (10)	1 – 20

Sixty-two percent (62%) of the 309 live foxtail pine trees produced female cones in 2022, with the per-plot percentage varying between 0% and 100%. These 191 trees produced 33 cones per tree on average (max 450), with average production rates per plot ranging from 5 to 84 cones per tree. Approximately four percent of the 302 whitebark pine trees in all PINALB plots produced female cones, averaging 4 cones per tree (max 20), with average production rates per plot ranging from 1-20 cones per cone-bearing tree. There were seven foxtail pine seedlings recorded in five plots, resulting in an estimated 72 (SD = 98) seedlings per hectare. Eight whitebark pine seedlings were also observed in two plots, along with three lodgepole pine seedlings in two plots.

Summary

Our data suggest that the whitebark pine and foxtail pine populations in areas sampled during the 2022 field season continue to have low incidence of white pine blister rust and mountain pine beetle, although anecdotal observations made by the forest crew suggested that WPBR spread to four new whitebark plots in YOSE, and to five new whitebark and three new foxtail pine plots in Sequoia and Kings Canyon, which would amount to substantial expansion since the last Panel 3 sample in 2017. Five whitebark pine trees were found within a single plot in Yosemite that displayed active cankers (aecia) of WPBR. This plot previously had 12 infected trees, and just one of those still displays enough indicators to be diagnosed as infected. An additional seven trees in six plots that were previously infected were not documented as infected in 2022 (one whitebark pine in Yosemite, four whitebark pines in two SEKI whitebark plots, and one foxtail in one foxtail plot, which was the only infection ever documented in foxtail pine in this data set).

This was the first full measure of all Panel 3 plots in SIEN, the second measure of most whitebark pine plots and the third measure of most foxtail plots in the panel. Tracking changes in forest composition, structure, and health through time will be particularly powerful in subsequent re-measures if incorporated into an adaptive management framework. Figure 10 shows the 2022 SIEN forest monitoring crew posing near South Guard Lake in Kings Canyon National Park.



Figure 10. 2022 SIEN forest monitoring crew near South Guard Lake in Kings Canyon National Park. Left to right, Ana Tobio, Emma Kahle, and Sean Auclair. NPS/ SEAN AUCLAIR.

Literature Cited

- Arno, S. F., and T. Weaver. 1990. Whitebark pine community types and their patterns on the landscape. Pages 97-105 in W. C. Schmidt and K. J. McDonald, editors. Proceedings of the Symposium on whitebark pine ecosystems: ecology and management of a high-elevation resource. USDA Forest Service General Technical Report INT-GTR-270. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, Colorado.
- Arno, S. F., and R. J. Hoff. 1990. *Pinus albicaulis* Engelm. Whitebark pine. Pages 268-279 in R. P. Burns and B. H. Honkala, editors. Silvics of North America, Volume 1, Conifers. Agriculture Handbook 654. USDA Forest Service, Washington, D.C.
- Ebenman, B., and T. Jonsson. 2005. Using community viability analysis to identify fragile systems and keystone species. *Trends in Ecology and Evolution* 20:568–575.
- Eckert, A. J., and J. O. Sawyer. 2002. Foxtail pine importance and conifer diversity in the Klamath Mountains and southern Sierra Nevada, California. *Madroño* 49:33-45.
- Ellison, A. E., M. S. Bank, B. D. Clinton, E. A. Colburn, K. Elliott, C. R. Ford, D. R. Foster, B. D. Kloeppel, J. D. Knoepp, G. M. Lovett, and others. 2005. Loss of foundation species: consequences for the structure and dynamics of forested ecosystems. *Frontiers in Ecology and the Environment* 3:479–486.
- Farnes, P. E. 1990. SNOTEL and snow course data describing the hydrology of whitebark pine ecosystems. Pages 302–304 in W. C. Schmidt and K. J. McDonald, editors. Proceedings of the Symposium on whitebark pine ecosystems: ecology and management of a high-elevation resource, 29–31 March 1989, Bozeman, Montana. USDA Forest Service General Technical Report INT-GTR-270. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, Colorado.
- Garrett, L. K., T. J. Rodhouse, G. H. Dicus, C. C. Caudill, and M. R. Shardlow. 2007. Upper Columbia Basin Network vital signs monitoring plan. Natural Resource Report NPS/UCBN/NRR—2007/002. National Park Service, Fort Collins, Colorado.
- Gibson, K., K. Skov, S. Kegley, C. Jorgensen, S. Smith, and J. Witcosky. 2008. Mountain pine beetle impacts in high-elevation five-needle pines: Current trends and challenges. U.S. Department of Agriculture, Forest Service, Forest Health Protection Report R1-08-020.
- Greater Yellowstone Whitebark Pine Monitoring Working Group (GYWPMWG). 2007. Interagency whitebark pine monitoring protocol for the Greater Yellowstone Ecosystem, v 1.0. Greater Yellowstone Coordinating Committee, Bozeman, Montana. Unpublished.
- Hawksworth, F. G. 1977. The 6-class dwarf mistletoe rating system. General Technical Report RM-48. USDA Forest Service Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 7p.

- Hupp, N., D. S. Stucki, J. Lonneker, M. Lehman, S. Wright, and J. Brackin. 2018. Protocol Implementation Plan for Monitoring White Pine in Great Basin National Park: Standard Operating Procedures, version 1.0. Mojave Desert Network, National Park Service, Boulder City, Nevada.
- Kipfmueeller, K. F., and M. W. Salzer. 2010. Linear trend and climate response of five-needle pines in the western United States related to treeline proximity. *Canadian Journal of Forest Research* 40: 134-142.
- Little, Elbert L., Jr. 1971. Atlas of United States trees. Volume 1. Conifers and important hardwoods. Misc. Publ. 1146. Washington, DC: U.S. Department of Agriculture, Forest Service. 320 p.
- Lloyd, A. H. 1997. Response of tree-line populations of foxtail pine (*Pinus balfouriana*) to climate variation over the last 1000 years. *Canadian Journal of Forest Research* 27:936-942.
- McDonald, T. L. 2003. Review of environmental monitoring methods: Survey designs. *Environmental Monitoring and Assessment* 85:277-292.
- McKinney, S.T., C.E. Fiedler, and D.F. Tomback. 2009. Invasive pathogen threatens bird-pine mutualism: implications for sustaining a high-elevation ecosystem. *Ecological Applications*, 19:597-607.
- McKinney, S. T., T. Rodhouse, L. Chow, A. Chung-MacCoubrey, G. Dicus, L. Garrett, K. Irvine, S. Mohren, D. Odion, D. Sarr, and L. A. Starcevich. 2012a. Monitoring white pine (*Pinus albicaulis*, *P. balfouriana*, *P. flexilis*) community dynamics in the Pacific West Region - Klamath, Sierra Nevada, and Upper Columbia Basin Networks: Narrative version 1.0. Natural Resource Report NPS/PWR/ NRR—2012/532. National Park Service, Fort Collins, Colorado.
- McKinney, S. T., T. Rodhouse, L. Chow, A. Chung-MacCoubrey, G. Dicus, L. Garrett, K. Irvine, S. Mohren, D. Odion, D. Sarr, and L. A. Starcevich. 2012b. Monitoring white pine (*Pinus albicaulis*, *P. balfouriana*, *P. flexilis*) community dynamics in the Pacific West Region - Klamath, Sierra Nevada, and Upper Columbia Basin Networks: Standard operating procedures version 1.0 (Appendix A to Narrative Version 1.0). Natural Resource Report NPS/PWR/NRR—2012/533. National Park Service, Fort Collins, Colorado.
- Mutch, L. S., M. G. Rose, A. M. Heard, R. R. Cook, and G. L. Entsminger. 2008. Sierra Nevada Network vital signs monitoring plan. Natural Resource Report NPS/ SIEN/NRR—2008/072. National Park Service, Fort Collins, Colorado.
- Sarr, D. A., D. C. Odion, S. R. Mohren, E. E. Perry, R. L. Hoffman, L. K. Bridy, and A. A. Merton. 2007. Vital signs monitoring plan for the Klamath Network: Phase III report. Natural Resource Technical Report NPS/KLMN/NRR—2007/016, National Park Service, Fort Collins, Colorado.
- Stevens, D. L., and A. R. Olsen. 2004. Spatially balanced sampling of natural resources. *Journal of the American Statistical Association* 99:262-278.

- Tomback D. F. 1982. Dispersal of whitebark pine seeds by Clark's nutcracker: A mutualism hypothesis. *Journal of Animal Ecology* 51:451–467.
- Tomback, D. F., and P. Achuff. 2010. Blister rust and western forest biodiversity: ecology, values, and outlook for white pines. *Forest Pathology* 40:186-225.
- Tomback, D. F., S. F. Arno, and R. E. Keane. 2001. The compelling case for management intervention. Pages 3–25 *in* D. F. Tomback, S. F. Arno, and R. E. Keane, editors. *Whitebark pine communities: Ecology and restoration*. Island Press, Washington, D.C.
- Tomback, D. F., and K. C. Kendall. 2001. Biodiversity losses: the downward spiral. Pages 243–262 *in* D. F. Tomback, S. F. Arno, and R. E. Keane, editors. *Whitebark pine communities: Ecology and restoration*. Island Press, Washington, D.C.
- Tomback, D. F., A. W. Schoettle, K. E. Chevalier, and C. A. Jones. 2005. Life on the Edge for limber pine: Seed dispersal within peripheral population. *Ecoscience* 12:519-529.
- van Mantgem, P. J., N. L. Stephenson, J. C. Byrne, L. D. Daniels, J. F. Franklin, P. Z. Fulé, M. E. Harmon, A. J. Larson, J. M. Smith, A. H. Taylor, and others. 2009. Widespread increase of tree mortality rates in the western United States. *Science* 323:521–523.
- Woodhouse, C. A., G. T. Pederson, and S. T. Gray. 2011. An 1800-yr record of decadal-scale hydroclimatic variability in the upper Arkansas River basin from bristlecone pine. *Quaternary Research* 75: 483-490.

Appendix 1. Notes on Field Methods

From year to year there can be procedures in the protocol sampling methods that are interpreted differently or are ambiguous (i.e., plot and tree level measurements), and crews have to resolve them while conducting field sampling. This appendix attempts to document these known issues and interpretations for the 2022 field season.

We continued to collect electronic data using Juniper Systems MESA 2 Rugged Tablet. We continued to implement some of the changes made in 2017 including recording a crown condition rating, counting cones, and tagging seedlings.

Appendix 2. Plot Status and Location

Table A-1. List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
YOSE-PIAL	01	272730	4211038	Panel_1x	Rejected	–	>35_Slope
YOSE-PIAL	02	288803	4213209	Panel_1x	Rejected	–	No trees
YOSE-PIAL	03	296088	4182250	Panel_1	2011	2019	–
YOSE-PIAL	04	290567	4185341	Panel_1	2011	2019	–
YOSE-PIAL	05	285267	4218542	Panel_1x	Rejected	–	No trees
YOSE-PIAL	06	279029	4195055	Panel_1	2015	2019	–
YOSE-PIAL	07	286332	4191015	Panel_1	2011	2019	Plot boundaries moved in 2015
YOSE-PIAL	08	295120	4177650	Panel_1	2011	2019	–
YOSE-PIAL	09	289361	4208460	Panel_1x	Rejected	–	>35_Slope
YOSE-PIAL	10	296034	4197576	Panel_1	2011	2019	–
YOSE-PIAL	11	295423	4193440	Panel_1x	Rejected	–	No trees
YOSE-PIAL	12	265845	4218338	Panel_1x	Rejected	–	No trees
YOSE-PIAL	13	290527	4215554	Panel_1x	Rejected	–	>35_Slope
YOSE-PIAL	14	298862	4191734	Panel_1	2011	2019	–
YOSE-PIAL	15	294711	4185917	Panel_1	2011	2019	–

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
YOSE-PIAL	16	268730	4223684	Panel_1	2015	2019	–
YOSE-PIAL	17	272205	4217495	Panel_1x	Rejected	–	>35_Slope
YOSE-PIAL	18	295937	4212170	Panel_1	2013	2019	–
YOSE-PIAL	19	303605	4189886	Panel_1	2014	2019	–
YOSE-PIAL	20	286082	4173212	Panel_1	2013	2015	–
YOSE-PIAL	21	284547	4213602	Panel_2	2014	2021	–
YOSE-PIAL	22	303861	4191215	Panel_2	2013	2021	–
YOSE-PIAL	23	290671	4198498	Panel_2x	Rejected	–	No Trees
YOSE-PIAL	24	296891	4175231	Panel_2	2013	2021	–
YOSE-PIAL	25	293052	4208800	Panel_2	2013	2021	–
YOSE-PIAL	26	300265	4196210	Panel_2	2013	2021	–
YOSE-PIAL	27	292618	4193551	Panel_2x	Rejected	–	No Trees
YOSE-PIAL	28	287655	4220242	Panel_2x	Rejected	–	No Route
YOSE-PIAL	29	288002	4215672	Panel_2x	Rejected	–	>35_Slope
YOSE-PIAL	30	297811	4192202	Panel_2x	Rejected	–	No Trees
YOSE-PIAL	31	292132	4186649	Panel_2x	2014	2021	Plot installed in 2014, read in 2016, then uninstalled in 2021 due to safety concerns

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
YOSE-PIAL	32	282831	4220099	Panel_2	2014	2021	–
YOSE-PIAL	33	281485	4206089	Panel_2x	Rejected	–	>35_Slope
YOSE-PIAL	34	295896	4203615	Panel_2x	Rejected	–	>35_Slope
YOSE-PIAL	35	300685	4188869	Panel_2	2014	2021	–
YOSE-PIAL	36	286581	4174238	Panel_2	2014	2016	–
YOSE-PIAL	37	282114	4211493	Panel_2x	Rejected	–	>35_Slope
YOSE-PIAL	38	302534	4193310	Panel_2	2014	2021	–
YOSE-PIAL	39	291731	4196540	Panel_2x	Rejected	–	No Trees
YOSE-PIAL	40	296396	4178608	Panel_2	2014	2021	–
YOSE-PIAL	41	292441	4201020	Panel_2	2016	2021	–
YOSE-PIAL	42	301791	4196147	Panel_3	2014	2022	–
YOSE-PIAL	43	294453	4183421	Panel_3	2014	2022	–
YOSE-PIAL	44	276329	4225930	Panel_3	2017	2022	–
YOSE-PIAL	45	294473	4210376	Panel_3x	Rejected	–	>35_Slope
YOSE-PIAL	46	298760	4187833	Panel_3	2017	2022	–
YOSE-PIAL	47	288739	4186425	Panel_3x	Rejected	–	–
YOSE-PIAL	48	278555	4224311	Panel_3	Rejected	–	Unsafe terrain

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
YOSE-PIAL	49	285782	4214700	Panel_3	Rejected	–	Plot was installed in 2011 and rejected in 2017 due to unsafe terrain
YOSE-PIAL	50	279271	4196291	Panel_3x	Rejected	–	No Trees
YOSE-PIAL	51	302369	4185092	Panel_3	2015	2022	–
YOSE-PIAL	52	294465	4174505	Panel_3	2011	2022	–
YOSE-PIAL	53	280552	4217599	Panel_3	2011	2022	Plot boundaries moved in 2017 due to unsafe terrain.
YOSE-PIAL	54	297249	4199689	Panel_3	2011	2022	–
YOSE-PIAL	55	294244	4198696	Panel_3	2017	2022	–
YOSE-PIAL	56	287430	4166358	Panel_3	2017	2022	–
YOSE-PIAL	57	293520	4200811	Panel_3x	Rejected	–	>35_Slope
YOSE-PIAL	58	297189	4193810	Panel_3x	Rejected	–	No Trees
YOSE-PIAL	59	293338	4182489	Panel_3x	Rejected	–	Trail within plot boundaries
YOSE-PIAL	60	267368	4220379	Panel_3	2017	2022	
YOSE-PIAL	61	291778	4214161	Panel_3	2022	2022	–
YOSE-PIAL	62	297642	4188103	Panel_2	Scouted	2021	To be installed in next panel rotation
SEKI-PIAL	01	367290	4027141	Panel_1x	Rejected	–	No Trees

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
SEKI-PIAL	02	345651	4118343	Panel_1	2014	2019	–
SEKI-PIAL	03	374530	4075808	Panel_1	2015	2019	–
SEKI-PIAL	04	375909	4084188	Panel_1	2015	2019	–
SEKI-PIAL	05	372907	4093446	Panel_1	2015	2019	–
SEKI-PIAL	06	343590	4112006	Panel_1x	Rejected	–	>35_Slope
SEKI-PIAL	07	368793	4067831	Panel_1x	Rejected	–	>35_Slope
SEKI-PIAL	08	357644	4085977	Panel_1	2019	2019	–
SEKI-PIAL	09	366236	4101569	Panel_1x	Rejected	–	>35_Slope
SEKI-PIAL	10	364469	4059015	Panel_1	2019	2019	–
SEKI-PIAL	11	372573	4075346	Panel_1	2015	2019	–
SEKI-PIAL	12	357557	4083750	Panel_1	2012	2019	First full read in 2019
SEKI-PIAL	13	360381	4105785	Panel_1	2015	2019	–
SEKI-PIAL	14	376492	4057576	Panel_1x	Rejected	–	No Trees
SEKI-PIAL	15	368275	4091574	Panel_1	2015	2019	–
SEKI-PIAL	16	356989	4077572	Panel_1x	Rejected	–	>35_Slope
SEKI-PIAL	17	362761	4033663	Panel_1x	Rejected	–	No Trees
SEKI-PIAL	18	343283	4112827	Panel_1	Incomplete	2019	Subplots 4 – 5 not installed in 2019

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
SEKI-PIAL	19	375619	4074608	Panel_1x	Rejected	–	>35_Slope
SEKI-PIAL	20	361155	4084009	Panel_1	2013	2019	–
SEKI-PIAL	21	371958	4097865	Panel_2x	Rejected	–	>35_Slope
SEKI-PIAL	22	351408	4095437	Panel_2x	Rejected	–	No Route
SEKI-PIAL	23	372112	4065836	Panel_2x	Rejected	–	>35_Slope
SEKI-PIAL	24	358026	4089119	Panel_2	2013	2021	–
SEKI-PIAL	25	365621	4094633	Panel_2	2013	2021	–
SEKI-PIAL	26	355753	4057139	Panel_2x	Rejected	–	No Trees
SEKI-PIAL	27	367661	4072341	Panel_2	2013	2021	–
SEKI-PIAL	28	361949	4091581	Panel_2	2013	2021	–
SEKI-PIAL	29	357722	4109427	Panel_2x	Rejected	–	>35_Slope
SEKI-PIAL	30	367938	4039738	Panel_2x	Rejected	–	No Trees
SEKI-PIAL	31	368793	4083219	Panel_2	2013	2021	–
SEKI-PIAL	32	365839	4066495	Panel_2x	Rejected	–	>35_Slope
SEKI-PIAL	33	339897	4121834	Panel_2	2014	2021	–
SEKI-PIAL	34	345544	4113490	Panel_2x	Rejected	–	>35_Slope
SEKI-PIAL	35	376672	4068290	Panel_2x	Rejected	–	>35_Slope

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
SEKI-PIAL	36	362322	4084241	Panel_2x	Rejected	–	>35_Slope
SEKI-PIAL	37	361596	4104498	Panel_2x	Rejected	–	No Route
SEKI-PIAL	38	346720	4094780	Panel_2	2014	2016	–
SEKI-PIAL	39	372095	4070727	Panel_2x	Rejected	–	>35_Slope
SEKI-PIAL	40	354109	4085347	Panel_2x	Rejected	–	No Trees
SEKI-PIAL	41	363987	4095543	Panel_2x	Rejected	–	>35_Slope
SEKI-PIAL	42	361901	4057273	Panel_2x	Rejected	–	No Route
SEKI-PIAL	43	372447	4072712	Panel_2	2014	2021	Subplots 4 - 5 not completed in 2016. Uninstalled after reading in 2021 due to safety concerns.
SEKI-PIAL	44	360112	4088754	Panel_2	2014	2021	–
SEKI-PIAL	45	347366	4115429	Panel_2	2014	2021	–
SEKI-PIAL	46	370788	4038875	Panel_2x	Rejected	–	No Trees
SEKI-PIAL	47	372218	4084481	Panel_2x	Rejected	–	>35_Slope
SEKI-PIAL	48	347886	4081797	Panel_2	2014	2016	–
SEKI-PIAL	49	367097	4094696	Panel_2	2014	2021	–
SEKI-PIAL	50	342764	4117298	Panel_3x	Rejected	–	>35_Slope
SEKI-PIAL	51	376111	4070297	Panel_3	2014	2022	–

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
SEKI-PIAL	52	364852	4084028	Panel_3	2017	2022	–
SEKI-PIAL	53	363263	4099360	Panel_3	Rejected	–	No Route
SEKI-PIAL	54	338892	4118855	Panel_3	Rejected	–	Unsafe terrain
SEKI-PIAL	55	368946	4075659	Panel_3x	Rejected	–	>35_Slope
SEKI-PIAL	56	359700	4080686	Panel_3x	Rejected	–	>35_Slope
SEKI-PIAL	57	359051	4104460	Panel_3x	Rejected	–	No Route
SEKI-PIAL	58	372709	4063450	Panel_3	2017	2022	Subplots 1-3 completed in 2017; installation completed in 2022
SEKI-PIAL	59	372495	4091307	Panel_3	2017	2022	2017
SEKI-PIAL	60	361751	4085403	Panel_3	2019	2022	Plot tagged but not read in 2019
SEKI-PIAL	61	348673	4115246	Panel_3	Rejected	–	Unsafe terrain
SEKI-PIAL	62	376844	4044787	Panel_3	Rejected	–	No trees
SEKI-PIAL	63	371096	4080482	Panel_3x	Rejected	–	>35_Slope
SEKI-PIAL	64	349766	4082680	Panel_3x	Rejected	–	>35_Slope
SEKI-PIAL	65	348874	4112395	Panel_3	Rejected	–	Unsafe terrain
SEKI-PIAL	66	358734	4091091	Panel_3	2017	2022	
SEKI-PIAL	67	364688	4062127	Panel_3	2022	2022	–
SEKI-PIAL	68	358068	4082392	Panel_3	2017	2022	–

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
SEKI-PIAL	69	367417	4051647	Panel_3x	Rejected	–	No trees
SEKI-PIAL	70	363423	4076489	Panel_3x	Rejected	–	>35_Slope
SEKI-PIAL	71	359118	4093582	Panel_3x	Rejected	–	>35_Slope
SEKI-PIAL	72	372846	4069794	Panel_3x	Rejected	–	>35_Slope
SEKI-PIAL	73	360249	4093039	Panel_3	2017	2022	Subplots 1-3 completed in 2017; fully read in 2022
SEKI-PIAL	74	369176	4073598	Panel_3	2017	2022	–
SEKI-PIAL	75	356689	4106954	Panel_3	Rejected	–	No route
SEKI-PIAL	76	367611	4079353	Panel_3x	Rejected	–	No Route
SEKI-PIAL	77	346084	4099017	Panel_3x	Rejected	–	>35_Slope
SEKI-PIAL	78	355533	4082622	Panel_3	2017	2022	
SEKI-PIAL	79	373490	4059513	Panel_3	2017	2022	Plot previously installed in 2017; first full read in 2022
SEKI-PIAL	86	372470	4090216	Panel_3x	Rejected	2021	
SEKI-PIAL	82	348004	4084609	Panel_2	–	–	To be installed in next panel rotation
SEKI-PIBA	01	355122	4029380	Panel_1	2015	2019	–
SEKI-PIBA	02	375721	4039069	Panel_1	2011	2019	–

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
SEKI-PIBA	03	363289	4043329	Panel_1	2015	2015	–
SEKI-PIBA	04	362613	4067179	Panel_1x	Rejected	–	No trees
SEKI-PIBA	05	370617	4026165	Panel_1	2015	2019	–
SEKI-PIBA	06	376988	4042810	Panel_1	2011	2019	–
SEKI-PIBA	07	374789	4056572	Panel_1	2011	2019	–
SEKI-PIBA	08	362507	4062625	Panel_1	2019	2019	–
SEKI-PIBA	09	377826	4033520	Panel_1	2011	2019	–
SEKI-PIBA	10	380915	4038349	Panel_1	2011	2019	–
SEKI-PIBA	11	375675	4050116	Panel_1	2011	2019	–
SEKI-PIBA	12	367913	4076907	Panel_1x	Rejected	–	No Trees
SEKI-PIBA	13	363692	4033498	Panel_1x	Rejected	–	>35_Slope
SEKI-PIBA	14	366283	4038676	Panel_1x	Rejected	–	>35_Slope
SEKI-PIBA	15	386345	4040899	Panel_1x	Rejected	–	Plot was installed in 2011 and rejected in 2015 due to unsafe terrain
SEKI-PIBA	16	356739	4061068	Panel_1	2013	2019	–
SEKI-PIBA	17	355948	4031543	Panel_1	2013	2019	–
SEKI-PIBA	18	375967	4045554	Panel_2	2013	2021	–

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
SEKI-PIBA	19	379846	4047488	Panel_2	2013	2021	–
SEKI-PIBA	20	367540	4067377	Panel_2x	Rejected	–	>35_Slope
SEKI-PIBA	21	375557	4032343	Panel_2	2013	2021	–
SEKI-PIBA	22	380284	4042853	Panel_2	2013	2021	–
SEKI-PIBA	23	374094	4056105	Panel_2	2013	2021	–
SEKI-PIBA	24	361953	4059325	Panel_2x	2016	2019	Plot read and uninstalled in 2019 due to safety concerns
SEKI-PIBA	25	369054	4032210	Panel_2	2013	2021	–
SEKI-PIBA	26	371357	4046768	Panel_2	2016	2016	–
SEKI-PIBA	27	372176	4051432	Panel_2	2013	2016	–
SEKI-PIBA	28	373060	4084548	Panel_2x	Rejected	–	>35_Slope
SEKI-PIBA	29	364439	4034149	Panel_2	2013	2016	–
SEKI-PIBA	30	371006	4042667	Panel_2	2014	2021	–
SEKI-PIBA	31	386287	4038225	Panel_2	2014	2021	–
SEKI-PIBA	32	360362	4059458	Panel_3x	Rejected	–	>35_Slope
SEKI-PIBA	33	384714	4036309	Panel_2	2014	2021	–
SEKI-PIBA	34	373202	4045338	Panel_3	2016	2022	Rebar & tags installed in 2016, first read 2022

Table A-1 (continued). List of sampling locations for white pine monitoring plots in the SIEN. The column EvalStatus indicates whether a site was established, dropped because it was non-target, or if it was not established (not visited) in the field. For sites that were visited, they are listed as either “Incomplete” indicating that not all subplots have been completed or “Scouted”, indicating that the plot has been visited and confirmed as target habitat, but not installed. If the plot was completed, the year it was fully installed is listed. Note that UTM X and UTM Y coordinates for established plots are the plot corner 1 (SW corner) coordinates as established in the field, and no longer match exactly the coordinates produced by the GRTS algorithm used to navigate to the plot during initial set-up.

Park-Species	Plot ID	UTM X	UTM Y	Panel	Installed	LastVisit	EvalNotes
SEKI-PIBA	35	377887	4048446	Panel_3	2014	2017	–
SEKI-PIBA	36	373289	4057861	Panel_3	2014	2017	–
SEKI-PIBA	37	376739	4029729	Panel_3	2014	2022	–
SEKI-PIBA	38	376735	4036915	Panel_3	2014	2022	–
SEKI-PIBA	39	375604	4053148	Panel_3	2014	2017	–
SEKI-PIBA	40	371733	4071478	Panel_3	2014	2022	–
SEKI-PIBA	41	367280	4034622	Panel_3	2014	2014	–
SEKI-PIBA	42	369440	4050366	Panel_3x	Rejected	–	No trees
SEKI-PIBA	43	382685	4040573	Panel_3	2017	2022	–
SEKI-PIBA	44	373659	4077444	Panel_3	2017	2022	–
SEKI-PIBA	45	363766	4029223	Panel_3	2018	2022	–
SEKI-PIBA	46	369487	4038685	Panel_3	2022	2022	–

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS 963/191788, December 2023

National Park Service
U.S. Department of the Interior



[Natural Resource Stewardship and Science](#)

1201 Oakridge Drive, Suite 150
Fort Collins, CO 80525