

U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science Directorate
Geologic Resources Division



Yucca House National Monument

GRI Ancillary Map Information Document

Produced to accompany the Geologic Resources Inventory (GRI) Digital Geologic
Data for Yucca House National Monument

yuho_geology.pdf

Version: 2/11/2025

Geologic Resources Inventory Map Document for Yucca House National Monument

Table of Contents

Geologic Resources Inventory Map Document.....	1
About the NPS Geologic Resources Inventory Program.....	3
GRI Digital Maps and Source Map Citations.....	5
Index Map	6
Map Unit List	7
Map Unit Descriptions.....	8
Qal - Alluvium (Recent and Pleistocene).....	8
Qs - Eolian sand and silt (Recent and Pleistocene).....	8
Qt - Talus (Recent and Pleistocene).....	8
Ql - Landslide deposits (Recent and Pleistocene).....	8
Qb - Block rubble (Recent and Pleistocene).....	8
Qfg - Fanglomerate (Recent and Pleistocene).....	8
Qpg - Pediment deposits (Recent and Pleistocene).....	8
Qsw - Slopewash (Holocene and Pleistocene).....	9
TKdp - Diorite porphyry (Tertiary or Upper Cretaceous).....	9
Kch - Cliff House Sandstone (Upper Cretaceous).....	9
Kme - Menefee Formation (Upper Cretaceous).....	9
Kpl - Point Lookout Formation (Upper Cretaceous).....	9
Km - Mancos Formation, undifferentiated (Upper Cretaceous).....	9
Kmc - Mancos Formation, Cortez Member (Upper Cretaceous).....	10
Kms - Mancos Formation, Smoky Hill Member (Upper Cretaceous).....	10
Kmv - Mancos Formation, Montezuma Valley Member (Upper Cretaceous).....	10
Kmj - Mancos Formation, Juana Lopez Member (Upper Cretaceous).....	10
Kmlm - Mancos Formation, Lower Mancos (Upper Cretaceous).....	10
Kd - Dakota Sandstone (Upper Cretaceous).....	10
Kbc - Burro Canyon Formation (Lower Cretaceous).....	10
Jmb - Morrison Formation, Brushy Basin Shale Member (Upper Jurassic).....	11
Jmw - Morrison Formation, Westwater Canyon Sandstone Member (Upper Jurassic).....	11
Jms - Morrison Formation, Salt Wash Sandstone Member (Upper Jurassic).....	11
Jjc - Junction Creek Sandstone (Upper Jurassic).....	11
Js - Summerville Formation (Upper Jurassic).....	11
Je - Entrada Sandstone (Upper Jurassic).....	11
Jn - Navajo Sandstone (Jurassic).....	11
Geologic Cross Section.....	12
Cross Section A-A'.....	12
Ancillary Source Map Information.....	13
Carrara, 2009.....	13
Cortez SW Quadrangle.....	13
Index Map.....	13
Legend	14
Explanation.....	15
References.....	15
Griffitts, 1999.....	16
Haynes et. al., 1972.....	16

Yucca House National Monument Area (Bedrock Geology and Paleontology).....	16
Report	16
References.....	27
Appendix	29
GRI Digital Data Credits.....	35

Geologic Resources Inventory Map Document



Yucca House National Monument, Colorado

Document to Accompany Digital Geologic-GIS Data

[yuho_geology.pdf](#)

Version: 2/11/2025

This document has been developed to accompany the digital geologic-GIS data developed by the Geologic Resources Inventory (GRI) program for Yucca House National Monument, Colorado (YUHO).

Attempts have been made to reproduce all aspects of the original source products, including the geologic units and their descriptions, geologic cross sections, the geologic report, references and all other pertinent images and information contained in the original publication.

This document contains the following information:

- 1) **About the NPS Geologic Resources Inventory Program** – A brief summary of the Geologic Resources Inventory (GRI) Program and its products. Included are web links to the GRI GIS data model, and to the GRI products page where digital geologic-GIS datasets, scoping reports and geology reports are available for download. In addition, web links to the NPS Data Store and GRI program home page, as well as contact information for the GRI coordinator, are also present.
- 2) **GRI Digital Maps and Source Map Citations** – A listing of all GRI digital geologic-GIS maps produced for this project along with sources used in their completion. In addition, a brief explanation of how each source map was used is provided.
- 3) **Map Unit List** – A listing of all geologic map units present on maps for this project, generally listed from youngest to oldest.
- 4) **Map Unit Descriptions** – Descriptions for all geologic map units. If a unit is present on multiple source maps the unit is listed with its source geologic unit symbol, unit name and unit age followed by the unit's description for each source map.
- 5) **Geologic Cross Sections** – Geologic cross section graphics with source geologic cross section abbreviations.
- 6) **Ancillary Source Map Information** – Additional source map information presented by source map.
- 7) **GRI Digital Data Credits** – GRI digital geologic-GIS data and ancillary map information document production credits.

For information about using GRI digital geologic-GIS data contact:

Stephanie O'Meara
Geologist/GIS Specialist/Data Manager
Colorado State University Research Associate, Cooperator to the National Park Service
Fort Collins, CO 80523
email: stephanie_o'meara@partner.nps.gov

About the NPS Geologic Resources Inventory Program

Background

The Geologic Resources Inventory (GRI) provides geologic map data and pertinent geologic information to support resource management and science-informed decision making in more than 270 natural resource parks throughout the National Park System. Geologic resources for management consideration include both the processes that act upon the Earth and the features formed as a result of these processes. Geologic processes include: erosion and sedimentation; seismic, volcanic, and geothermal activity; glaciation, rockfalls, landslides, and shoreline change. Geologic features include mountains, canyons, natural arches and bridges, minerals, rocks, fossils, cave and karst systems, beaches, dunes, glaciers, volcanoes, and faults.

The GRI is one of 12 inventories funded by the National Park Service (NPS) Inventory and Monitoring Program. The Geologic Resources Division of the NPS Natural Resource Stewardship and Science Directorate administers the GRI. The NPS Geologic Resources Division partners with the Colorado State University Department of Geosciences to produce GRI products. Many additional partners participate in the GRI process by contributing source maps or reviewing products.

The GRI team undertakes three tasks for each park in the Inventory and Monitoring program: (1) conduct a scoping meeting and provide a summary document, (2) provide digital geologic map data in a geographic information system (GIS) format, and (3) provide a GRI report. These products are designed and written for nongeoscientists.

Products

Scoping Meetings: These park-specific meetings bring together local geologic experts and park staff to inventory and review available geologic data and discuss geologic resource management issues. A summary document is prepared for each meeting that identifies a plan to provide digital map data for the park.

Digital Geologic Maps: Digital geologic maps reproduce all aspects of traditional paper maps, including notes, legend, and cross sections. Bedrock, surficial, and special purpose maps such as coastal or geologic hazard maps may be used by the GRI to create digital Geographic Information Systems (GIS) data and meet park needs. These digital GIS data allow geologic information to be easily viewed and analyzed in conjunction with a wide range of other resource management information data.

For detailed information regarding GIS parameters such as data attribute field definitions, attribute field codes, value definitions, and rules that govern relationships found in the data, refer to the NPS Geology-GIS Data Model document available at: <https://www.nps.gov/articles/gri-geodatabase-model.htm>

Geologic Reports: GRI reports synthesize discussions from the original scoping meeting, follow up conference call(s), and subsequent research. Chapters of each report discuss the geologic setting of the park, distinctive geologic features and processes within the park, highlight geologic issues facing resource managers, and describe the geologic history leading to the present-day landscape. Each report also includes a poster illustrating these GRI digital geologic-GIS data.

For a complete listing of GRI products visit the GRI publications webpage: <https://go.nps.gov/gripubs>. GRI digital geologic-GIS data is also available online at the NPS Data Store: <https://irma.nps.gov/DataStore/Search/Quick>. To find GRI data for a specific park or parks select the appropriate park(s), enter "GRI" as a Search Text term, and then select the Search button.

For more information about the Geologic Resources Inventory Program visit the GRI webpage: <https://>

www.nps.gov/subjects/geology/gri.htm. At the bottom of that webpage is a “Contact Us” link if you need additional information. You may also directly contact the program coordinator:

Jason Kenworthy
Inventory Coordinator
National Park Service Geologic Resources Division
P.O. Box 25287
Denver, CO 80225-0287
phone: (303) 987-6923
fax: (303) 987-6792
email: Jason_Kenworthy@nps.gov

The Geologic Resources Inventory (GRI) program is funded by the National Park Service (NPS) Inventory and Monitoring (I&M) Division. Learn more about I&M and the 12 baseline inventories at the I&M webpage: <https://www.nps.gov/im/inventories.htm>.

GRI Digital Maps and Source Map Citations

The GRI digital geologic-GIS maps for Yucca House National Monument, Colorado (YUHO) are listed below. With each map is the source map used in its completion, as well as how the source map was used.

Digital Geologic-GIS Map of the Mud Creek Quadrangle, Colorado (*GRI MapCode MUCR*)

Ekren, E.B., and Houser, F.N., 1959, Preliminary Geologic Map of the Cortez SW Quadrangle Montezuma County, Colorado: U.S. Geological Survey, Mineral Investigations Field Studies Map MF-217, scale 1:24,000 ([Cortez SW Quadrangle](#)). (*GRI Source Map ID 94*).

The GRI used the full extent of the above source map, and all geologic features within its extent were captured.

Digital Bedrock Geologic-GIS Map of Yucca House National Monument and Vicinity, Colorado (*GRI MapCode YUHO_bedrock*)

Griffitts, Mary O., 2001, Bedrock Geology and Paleontology of Yucca House National Monument Area, Colorado, unpublished map, scale 1:62,500 ([Yucca House National Monument Area \(Bedrock Geology and Paleontology\)](#)). (*GRI Source Map ID 1341*).

The GRI used the full extent of the above source map, and all geologic features within its extent were captured.

Digital Geologic-GIS Map of a Portion of Yucca House National Monument, Colorado (*GRI MapCode YUHO*)


Carrara, Paul, 2009, Yucca House Geologic Map: U.S. Geological Survey, unpublished map, scale 1:12,000 ([Yucca House \(Geologic Map\)](#)). (*GRI Source Map ID 75483*).

The GRI used the full extent of the above source map, and all geologic features within its extent were captured.

Additional information pertaining to each source map is also presented in the GRI Source Map Information (YUHO MAP) table included with the GRI digital geologic-GIS data.

Index Map

The following index map displays the extents of the GRI digital geologic-GIS maps produced for Yucca House National Monument (YUHO). The boundary for Yucca House National Monument (as of February, 2025) is outlined in green. The outline for the Digital Geologic-GIS Map of the Mud Creek Quadrangle is outlined in magenta, the outline for the Digital Geologic-GIS Map of a Portion of Yucca House National Monument is outlined in orange, and the outline for the Digital Bedrock Geologic-GIS Map of Yucca House National Monument and Vicinity is outlined in blue.



Index map by James Winter (Colorado State University).

Map Unit List

The geologic units present in the digital geologic-GIS data produced for Yucca House National Monument, Colorado (YUHO) are listed below. Units are listed with their assigned unit symbol and unit name (e.g., Qal - Alluvium). Units are listed from youngest to oldest. No description for water is provided. Information about each geologic unit is also presented in the GRI Geologic Unit Information (YUHOUNIT) table included with the GRI digital geologic-GIS data. Some source unit symbols, names and/or ages may have been changed in this document and in the GRI digital geologic-GIS data. This was done if a unit was considered to be the same unit as one or more units on other source maps used for this project, and these unit symbols, names and/or ages differed. In this case a single unit symbol and name, and the unit's now recognized age, was adopted. Unit symbols, names and/or ages in unit descriptions, correlation figures or other source map figures were not edited. If a unit symbol, name or age was changed by the GRI the unit's source map symbol, name and/or age appears with the unit's source map description.

Cenozoic Era

Quaternary Period

[Qal](#) - Alluvium

[Qs](#) - Eolian sand and silt

[Qt](#) - Talus

[Ql](#) - Landslide deposits

[Qb](#) - Block rubble

[Qfg](#) - Fanglomerate

[Qpg](#) - Pediment deposits

[Qsw](#) - Slopewash

Cenozoic or Mesozoic Eras

Tertiary or Cretaceous Periods

[TKdp](#) - Diorite porphyry

Mesozoic Era

Cretaceous Period

[Kch](#) - Cliff House Sandstone

[Kme](#) - Menefee Formation

[Kpl](#) - Point Lookout Formation

[Km](#) - Mancos Shale, undifferentiated

[Kmc](#) - Mancos Shale, Cortez Member

[Kms](#) - Mancos Shale, Smoky Hill Member

[Kmv](#) - Mancos Shale, Montezuma Valley Member

[Kmj](#) - Mancos Shale, Juana Lopez Member

[Kmlm](#) - Mancos Shale, lower Mancos

[Kd](#) - Dakota Sandstone

[Kbc](#) - Burro Canyon Formation

Jurassic Period

[Jmb](#) - Morrison Formation, Bushy Basin Shale Member

[Jmw](#) - Morrison Formation, Westwater Canyon Sandstone Member

[Jms](#) - Morrison Formation, Salt Wash Sandstone Member

[Jjc](#) - Junction Creek Sandstone

[Js](#) - Summerville Formation

[Je](#) - Entrada Sandstone

[Jn](#) - Navajo Sandstone

Map Unit Descriptions

Descriptions of all geologic map units, generally listed from youngest to oldest, are presented below.

Qal - Alluvium (Recent and Pleistocene)

Qal - Alluvium (Pleistocene and Recent)

No unit description provided on source map: [Cortez SW Quadrangle](#).

Qal - Alluvium (Pleistocene and Recent)

Unconsolidated sands, silts, and gravels deposited mainly in stream beds and flood plains.

Description from source map: [Griffitts, 1999](#).

Qa - Alluvial Deposits (Pleistocene and Holocene)

Silt, sand, and gravel in stream valleys and flood plains; includes soil and locally some colluvial and eolian deposits. Description from source map: [I-629](#).

Qs - Eolian sand and silt (Recent and Pleistocene)

No unit description provided on source map: [Cortez SW Quadrangle](#).

Qt - Talus (Recent and Pleistocene)

No unit description provided on source map: [Cortez SW Quadrangle](#).

Ql - Landslide deposits (Recent and Pleistocene)

No unit description provided on source map: [Cortez SW Quadrangle](#).

Qb - Block rubble (Recent and Pleistocene)

Chiefly porphyry blocks lying on unevenly eroded surfaces; forms hummocky topography. Description from source map: [Cortez SW Quadrangle](#).

Qfg - Fanglomerate (Recent and Pleistocene)

Mainly cobbles and boulders from the Point Lookout sandstone of the Mesaverde group of Cretaceous age. Description from source map: [Cortez SW Quadrangle](#).

Qpg - Pediment deposits (Recent and Pleistocene)

Qpg - Pediment deposits (Pleistocene and Recent)

Predominantly pebbles, cobbles, and boulders of igneous rocks lying on erosion surfaces that extend radially from the mountain area. Description from source map: [Cortez SW Quadrangle](#).

Qap - Pediment Gravels (Pleistocene and Holocene)

Poorly sorted angular to well-rounded sand, pebbles, and boulders; mostly diorite porphyry; in thin partly dissected deposits near the Abajo and Sleeping Ute Mountains. Description from source map: [I-629](#).

Qsw - Slopewash (Holocene and Pleistocene)

The following units describe Qsw - slopewash.

Qc - Colluvial deposits (Pleistocene and Holocene)

Talus, slope wash, block rubble rock glaciers, and, in some high cirques in the San Miguel Mountains, young glacial till. Description from source map: [I-629](#).

Qcl - Landslide deposits (Pleistocene and Holocene)

Unsorted material, commonly in large displaced blocks with characteristic arcuate hummocky surfaces. Description from source map: [I-629](#).

Qct - Talus deposits (Pleistocene and Holocene)

Accumulations of angular blocks fallen from cliffs and outcrops of resistant rock. Description from source map: [I-629](#).

TKdp - Diorite porphyry (Tertiary or Upper Cretaceous)

Medium gray diorite porphyry; phenocrysts of hornblende and andesine; forms laccoliths in extreme southwest part of quadrangle. Description from source map: [Cortez SW Quadrangle](#).

Kch - Cliff House Sandstone (Upper Cretaceous)

White to red-brown, fine to medium grained marine sandstones interbedded with sandy shales. Upper and lower units of massive sandstones separated by a unit of thinner bedded sandy shales. Thickness 200-300 feet. Part of "Mesaverde Group", with [Menefee](#) and [Point Lookout](#) Formations. Description from source map: [Griffitts, 1999](#).

Kme - Menefee Formation (Upper Cretaceous)

Dark gray and brown carbonaceous non-marine shales, thin siltstones and thin coal beds in upper and lower units separated by a middle sandy unit of poorly sorted, irregular bedded sandstones, sandy shales and bentonite beds. Thickness 400-800 feet. Part of "Mesaverde Group", with [Cliff House](#) and [Point Lookout](#) Formations. Description from source map: [Griffitts, 1999](#).

Kpl - Point Lookout Formation (Upper Cretaceous)

White to yellow fine to medium grained marine sandstone with shaly sandstone breaks, highly cross-bedded. Thickness 360 feet. Part of "Mesaverde Group", with [Cliff House](#) and [Menefee](#) Formations. Description from source map: [Griffitts, 1999](#).

Km - Mancos Formation, undifferentiated (Upper Cretaceous)

Predominantly gray to black shaly mudstone. A medium- to coarse-grained glauconitic sandstone and sandy fossiliferous limestone (10-50 feet thick), about 475 feet above base, is mapped separately as Kms. This unit may be equivalent in part to the Juana Lopez sandstone member (of the Carlile shale) of Rankin (1944). Description from source map: [Cortez SW Quadrangle](#).

Marine light to dark gray thin bedded shales and siltstones with well developed faunal zones. 2000 feet (at Mesa Verde National Park). Description from source map: [Griffitts, 1999](#).

Gray to dark-gray soft fissile sparsely fossiliferous marine clay shale; a few thin distinctive calcareous sandstone and sandy clayey limestone ledges in lower 500 feet. Grades vertically into and intertongues with formations of the Mesaverde Group. 2,000 to 3,000 feet thick; thickens northeastward. Description from source map: [I-629](#).

Kmc - Mancos Formation, Cortez Member (Upper Cretaceous)

No unit description available (see Mancos Shale ([Km](#)) for undifferentiated unit description). Description from source map: [Griffitts, 1999](#).

Kms - Mancos Formation, Smoky Hill Member (Upper Cretaceous)

Predominantly gray to black shaly mudstone. A medium- to coarse-grained glauconitic sandstone and sandy fossiliferous limestone (10-50 feet thick), about 475 feet above base, is mapped separately as Kms. This unit may be equivalent in part to the Juana Lopez sandstone member (of the Carlile shale) of Rankin (1944). Description from source map: [Cortez SW Quadrangle](#).

A prominent oyster (*Pseudoperna congesta*) bench about 900 feet above base of [Mancos Formation](#) (also referred to as 'Niobara' on Mesa Verde National Park map). Description from source map: [Griffitts, 1999](#).

Kmv - Mancos Formation, Montezuma Valley Member (Upper Cretaceous)

No unit description available (see Mancos Shale ([Km](#)) for undifferentiated unit description). Description from source map: [Griffitts, 1999](#).

Kmj - Mancos Formation, Juana Lopez Member (Upper Cretaceous)

About 500 feet above base of [Mancos](#), 140 feet of highly fossiliferous dark silty shale with numerous beds of orange weathering calcarenite and thin bentonites (at Mesa Verde National Park). Description from source map: [Griffitts, 1999](#).

Kmlm - Mancos Formation, Lower Mancos (Upper Cretaceous)

No unit description available (see Mancos Shale ([Km](#)) for undifferentiated unit description). Description from source map: [Griffitts, 1999](#).

Kd - Dakota Sandstone (Upper Cretaceous)

Yellowish lenticular sandstone and conglomerate with interbedded carbonaceous shale and impure coal. Average thickness about 125 feet. Description from source map: [Cortez SW Quadrangle](#).

Kbc - Burro Canyon Formation (Lower Cretaceous)

Sandstone and conglomerate with interbedded green and red shale; ranges from 30 to 200 feet in thickness; intertongues with mudstone of the Brushy Basin Shale Member of the Morrison Formation. Description from source map: [Cortez SW Quadrangle](#).

Jmb - Morrison Formation, Brushy Basin Shale Member (Upper Jurassic)

150 to 300 feet thick, consists of varicolored bentonitic mudstone with a few conglomeratic sandstone lenses. Adjacent members intertongue and inter-grade to such an extent that in many places the contacts are arbitrary. Description from source map: [Cortez SW Quadrangle](#).

Jmw - Morrison Formation, Westwater Canyon Sandstone Member (Upper Jurassic)

75 to 100 feet thick, consists of pale yellow-brown fine-to medium-grained sandstone interbedded with green bentonitic mudstone. Adjacent members intertongue and inter-grade to such an extent that in many places the contacts are arbitrary. Description from source map: [Cortez SW Quadrangle](#).

Jms - Morrison Formation, Salt Wash Sandstone Member (Upper Jurassic)

150 to 250 feet thick, consists of tan-gray fine- to medium-grained sandstone interbedded with red-brown and purple mudstone. The Salt Wash sandstone member is uranium bearing in many areas on the Colorado Plateau but no deposits are known in this quadrangle. Adjacent members intertongue and inter-grade to such an extent that in many places the contacts are arbitrary. Description from source map: [Cortez SW Quadrangle](#).

Jjc - Junction Creek Sandstone (Upper Jurassic)

Consists of three gradational units. The upper unit, 20 to 50 feet thick, is argillaceous fine-grained reddish sandstone with obscure flat stratification. The middle unit, 150 to 200 feet thick, is pink to orange-red poorly sorted fine- to coarse-grained quartz sandstone with high-angle cross-stratification; weathers to a "slick rim". The lower unit, 30 to 50 feet thick, has the same general lithology as the middle unit, but with low-angle cross-stratification and numerous horizontal truncations. Thickness is 230 to 280 feet; contact with underlying Summerville formation is gradational. Correlates with the Bluff sandstone of Utah and Arizona. Description from source map: [Cortez SW Quadrangle](#).

Js - Summerville Formation (Upper Jurassic)

Brick-red mudstone and pink to red-brown argillaceous fine-grained well-sorted sandstone, 120 to 130 feet thick. Description from source map: [Cortez SW Quadrangle](#).

Je - Entrada Sandstone (Upper Jurassic)

Consists of two units. Upper unit is sandstone, 70 to 80 feet thick, pale brown and light pink at top grading to white and finally to orange red in lower part; weathers to a "slick rim". The lower unit, 25 to 35 feet thick, is argillaceous very fine grained brick-red sandstone. Description from source map: [Cortez SW Quadrangle](#).


Jn - Navajo Sandstone (Jurassic)

Orange fine-grained eolian sandstone containing numerous prominent horizontal truncations. Description from source map: [Cortez SW Quadrangle](#).

Geologic Cross Section

The geologic cross section present in the GRI digital geologic-GIS data produced for Yucca House National Monument, Colorado (YUHO) is presented below. The cross section graphic was scanned at a high resolution and can be viewed in more detail by zooming in (if viewing the digital format of this document).

Cross Section A-A'



Graphic from source map: [Cortez SW Quadrangle](#).

Ancillary Source Map Information

The following sections present ancillary source map information associated with sources used for this project.

Carrara, 2009

The formal citation for this source.

Carrara, Paul, 2009, Yucca House Geologic Map: U.S. Geological Survey, unpublished map, scale 1:12,000 (*GRI Source Map ID 75483*).

This source only consisted of digital GIS data. No additional information from this source is provided.


Cortez SW Quadrangle

The formal citation for this source.

Ekren, E.B., and Houser, F.N., 1959, Preliminary Geologic Map of the Cortez SW Quadrangle Montezuma County, Colorado: U.S. Geological Survey, Mineral Investigations Field Studies Map MF-217, scale 1:24,000 (*GRI Source Map ID 94*).

Prominent graphics and text associated with this source:

Index Map



Graphic from source map: [Cortez SW Quadrangle](#)

Legend



Contact
Dashed where approximately located; dotted where concealed



Fault, showing displacement
Dashed where approximately located; dotted where concealed; U, upthrown side, D, downthrown side; showing displacement in feet



Anticline
Showing trace of axial plane and direction of plunge; dashed where approximately located



Strike and dip of beds



Lineament
Prominent joints observed in Mancos shale in southern part of quadrangle; believed to indicate faulting at depth; dotted where projected under surficial deposits



Bearing and plunge of linear flow structure



Structure contours
Drawn on base of Mancos shale; long dashes where approximately located; short dashes where projected above surface; dots where contour on a phantom horizon is projected through a sill or laccolith. Laccoliths were intruded at or near the contact of the Dakota sandstone and Mancos shale; locally portions of upper beds of the Dakota sandstone were domed upward by the intrusives (section A-A'); datum is projected through laccoliths in order to define underlying structure. Contour interval is 100 feet. Datum is mean sea level




Gravel pit



Dry hole

Graphic from source map: [Cortez SW Quadrangle](#)

Explanation



Graphic from source map: [Cortez SW Quadrangle](#)

References

Rankin, C.H., 1944, Stratigraphy of the Colorado Group, Upper Cretaceous in northern New Mexico: N. Mex. School of Mines Bull. 20, 27 p.

References from source map: [Cortez SW Quadrangle](#)

Griffitts, 1999

The formal citation for this source.

Griffitts, Mary O., 1999, Mesa Verde National Park Geology, National Park Service, scale 1:24,000. (*GRI Source Map ID 978*).

Only unit descriptions were taken from this source map. No additional information from this source is provided.

Haynes et. al., 1972

The formal citation for this source.

Haynes, D.D., Vogel, J.D., and Wyant, D.G., 1972, Geology, Structure, and Uranium Deposits of the Cortez Quadrangle, Colorado and Utah, U.S. Geological Survey, Miscellaneous Geologic Investigations Map I-629, scale 1:250,000. (*GRI Source Map ID 105*).

Only unit descriptions were taken from this source map. No additional information from this source is provided.

Yucca House National Monument Area (Bedrock Geology and Paleontology)

The formal citation for this source.

Griffitts, Mary O., 2001, Bedrock Geology and Paleontology of Yucca House National Monument Area, Colorado, unpublished map, scale 1:62,500 (*GRI Source Map ID 1341*).

Prominent graphics and text associated with this source:

Report

INTRODUCTION

A detailed study of the bed rock geology in the area presently owned by Mrs Hallie Ismay, surrounding the established Yucca House National Monument, Montezuma County, Colorado was initiated by Mesa Verde National Park in May, 2000. The area of study (figure 1) includes portions of Montezuma Valley in the Mud Creek and Towaoc Quadrangles, extending approximately from the Aztec Divide on the north to the "Old Southern Ute Indian Reservation Boundary" on the south, and from about 1/2 mile west of the Ute Mountain Ute Reservation Boundary to the west edge of the Mesa Verde on the east.




Figure 1. Location of Yucca House National Monument study area (diagonal ruling) within Towaoc and Mud Creek Quadrangles, in southwest Colorado. Ute Mountain Ute Reservation boundary south of shaded bar.

Existing geologic maps of the area were compiled, aerial photographs reviewed and a field reconnaissance was made to map the faunal horizons and surface outcrops. A composite aerial photographic base map was provided by Allan Loy of the Mesa Verde staff. Although much of this area has been mapped in some detail by E.B. Ekren and F.N. Hauser (1965), paleontologic zones were not studied at that time. Mud Creek and Towaoc Quadrangles were remapped to determine the distribution of possible sources of building material used in the Yucca House site. The paleontology is of major significance in the Yucca House archeology. The maps generated in this study include portions on the west edge of the Mesa Verde which were taken from A.A. Wanek's 1959 map of the Mesa Verde area.

SETTING

Montezuma Valley lies to the east of the Ute Mountains and west of the Mesa Verde scarp. A broad, relatively flat area, it is largely covered by alluvial deposits and has been heavily cultivated for many years. There are few bed rock outcrops. Much of the surface is covered by Quaternary deposits, the upper layers of which have been plowed and reworked for generations. Along the east border, and in places along the west side of the study area, bed rock crops out. Numerous early surveys of the Four Corners area include some discussion of the Yucca House vicinity. The earliest recorded is the description by W.H. Holmes during his 1875-76 field excursions to the Aztec Springs Ruins, as the locality was then known. He made a quite detailed sketch map of the mound as it then appeared or as he envisioned it, and noted the "Stone used is chiefly the fossiliferous limestone that outcrops along the base of the Mesa Verde a mile or more away" (Holmes, 1878, p399-400). The most

comprehensive report is the "Geology and Petrology of the Ute Mountains Area, Colorado" by Ekren and Hauser (1965) which details the laccolithic intrusives of the Ute Mountains and includes the northern part of the present field study. Detailed discussions of the stratigraphy of the Four Corners region were published in 1997 in the report of the New Mexico Geological Society Forty Eighth Field Conference.

SEDIMENTARY HISTORY

The Yucca House National Monument and the Ismay property surveyed is underlain by sedimentary rocks, deposited in the great Late Cretaceous inland seaway. Older rocks crop out to the north in the McElmo Valley and to the west along the Ute Mountain front. Pre Cretaceous rocks, Jurassic Morrison and older, are well exposed in the McElmo Canyon. Although they are not found in place on the surface in the area of study, material from these older rocks may be found in the archeological record and in the superficial Cenozoic deposits. The Burro Canyon Formation (Buckhorn), of controversial age, either very late Jurassic or early Cretaceous, which lies between the Morrison and the Dakota Formations, has many silicic pebbles which were used by the early Puebloan people for tool making. The Jurassic Morrison Formation has numerous beds of chert, quartzite and other highly silicic material which are commonly found in artifacts throughout the archeological sites.

The Cretaceous history begins with the deposition of the Dakota Formation. This very widespread sandstone formation is recognized throughout the Western Interior. It represents the first invasion of the Late Cretaceous seas this far inland, and generally consists of light brown fine- to coarse-grained sandstone. In this area, many of the sand grains are subround to angular and mostly poorly sorted. There is much carbonaceous material present and mineable coal beds occur a short distance west of Towaoc. These mines are no longer worked, but some of the seams were two to four feet thick where observed. The brown sandstones and muddy sandstones were deposited in shallow water, in lagoons and near the shore of the transgressing sea. The Dakota is conformably overlain by the marine Mancos Formation.

Only the very thick Mancos Formation is exposed within most of the Ismay holdings and the existing Yucca House National Monument and in the Towaoc and Mud Creek Quadrangles. The closest exposures of the underlying Dakota Sandstone are about 2 1/2 miles to the west in Ute land and about 2 1/2 miles north in the Mud Spring area. Although the Dakota Sandstone is solid enough to make good building material, no blocks in the exposed rubble mounds of the prehistoric structures were recognizably Dakota material. It is very possible that future excavation of the site will reveal Dakota building blocks.

As the sea slowly encroached, this area became more and more distant from the shore line. During a period of 13 million years the fluctuating transgressions of the inland sea deposited more than 2000 feet of Mancos Formation. A measured section at the type locality near Mesa Verde National Park (Leckie, et al 1997) has been divided into eight members. The members as established by Leckie are from the oldest: the Graneros shale, the Bridge Creek Limestone, Fairport shale, Blue Hill shale, the Juana Lopez shale and calcarenite, Montezuma Valley shale, Smoky Hill shale and limestone, and the Cortez sandy shale. The lowest unit, the Graneros, is dated by its invertebrate fauna as upper Cenomanian age (approximately 94 Ma) and the top of the Mancos Cortez unit is dated Campanian in age (approximately 81 Ma). A detailed discussion of regional correlation is available in the 1997 paper by Leckie, et al. Only the upper four members including the Juana Lopez, Montezuma Valley, Smoky Hill and Cortez crop out in the expanded Yucca House study area. The Graneros Shale and Bridge Creek Limestone are recognizable in T35N, R17W Sec.20, about 2 1/2 miles west of the proposed monument boundary. The Graneros Shale is a soft, very dark gray shale and silty shale with a characteristic and easily recognizable fauna including small oysters *Pycnodonte newberryi* (figure 12A-B) which are quite resistant to weathering and accumulate in stream washes. The Bridge Creek Limestone (Greenhorn Limestone) has several relatively massive, solid limestone layers up to 3 feet in thickness which make well defined outcrops with sparse vegetation. The limestone is solid enough that it might have been used for building stone, but no blocks were seen in the presently exposed prehistoric mounds. Even if it had been used, the limestone blocks tend to split

along relatively thin bedding planes after being exposed to weathering, and would probably not have been preserved on the surface to the present. Again, future excavation may reveal the use of this rock in some parts of the site. The Fairport and Blue Hill Members are without doubt present also between the Bridge Creek and the prominent Juana Lopez, but both are soft shaly deposits, poorly resistant to erosion and are not easily recognized in small outcrops and are generally covered in gullies.

The Juana Lopez member of the Mancos Formation is exposed on a dip slope (figure 2) over much of the western part of the Mud Creek and Towaoc Quadrangles. The very low angle of dip of the bedrock allows a relatively thin lithologic unit to be exposed over a very large surface area. The dip is so low, 3 to 5 degrees where measurable, that bedding plane irregularities mask minor structural anomalies. The Juana Lopez is a distinctive and widespread member of the Mancos Formation. With a very abundant late Turonian invertebrate fauna, it is correlated with widely distributed rocks throughout the Western Interior of North America. Originally described by Dane and others (1966) on the east side of the San Juan Basin, the type section was measured "on the Mesita Juana Lopez Grant, six miles northwest of Cerrillos, Santa Fe County, New Mexico." (Rankin, 1944 p. 19). Later studies by Kaufman (in Dane and others 1966), and Hook and Cobban (1980) revised details of location and provide a more comprehensive type section measuring 106 foot thickness. At Mesa Verde a detailed section (Leckie, et al 1997) of the measured thickness of the Juana Lopez totaled approximately 140 feet. Due to the poor exposures it is not practical to measure a detailed section at Yucca House, but it is unlikely to differ seriously from the Mesa Verde section 15 miles to the northeast.



Figure 2. Thin bedded calcarenite of the Juana Lopez (zone 3) exposed on the dip slope west of Yucca House.

The Juana Lopez member of the Mancos Formation is relatively resistant to weathering and frequently forms low terraces and irregular hills because of the numerous layers of solid calcarenite. Calcarenite is a granular rock similar in appearance to sandstone, but composed almost entirely of calcium carbonate. Solution of pieces of this calcarenite in hydrochloric acid left only a very small residue of silt and fine clay. Much of the calcarenite is thin bedded, (figure 2) and could not be used as building stone. But some of the beds are locally more than six inches in thickness (figure 3). The calcarenite is often a dense, dark colored solid rock which weathers to a rusty brown color. It naturally breaks into building blocks along bedding planes and joints (figure 4). Most of the slabs show at least fragmental fossils, and many excellent ammonite and bivalve fossils are found. In the exposed prehistoric walls of the present mounds, many of the building stones are made of the brown,

fossiliferous calcarenite of the Juana Lopez Member of the Mancos Formation (figure 5). The fossils are diagnostic and many are easily recognized even when fragmental. Although the Juana Lopez Member of the Mancos underlies the area, Quaternary terrace gravels, pediment deposits, alluvium, and loess soil cover much of the surface, and some of the boulders from these deposits were also used as building stone. Outcrops of the Juana Lopez typical building stone are not found nearer the archeologic sites of Yucca House than about 1/2 mile (map distance) to the west.



Figure 3. Thick solid beds of calcarenite in gully about ½ mile southwest of Yucca House mound.



Figure 4. Calcarenite beds in gully southwest of Yucca House mound, showing ready made building blocks shaped by bedding planes and joints.



Figure 5. Building block in Yucca House mound, surface covered with *Lopha lugubris* shells from the Juana Lopez zone 2 calcarenite.

Although much of the Juana Lopez is silty shale, there are several consistent calcarenite beds of varying thicknesses. The paleontologic zones which can be recognized in these calcarenite beds should be useful to archeologists during any future excavation. Although alluvial and other Quaternary sediments cover most of the broad flats of the area, in some places deep gullies, ditches and roadcuts expose the bed rock. Leckie (1997) recognized and described four calcarenite zones in the Mesa Verde section, each with a distinctive fauna. Hook and Cobban (1980) recognized the same faunal sequence in the type section in New Mexico. In the Yucca House area, the two uppermost paleontologic zones of the Juana Lopez are recognizable over quite a large area, mainly because the calcarenites are thicker, more resistant to weathering and form long, low ridges trending northeast/southwest. The calcarenite layers are quite solid, frequently very dark and dense. Although some of them look very sandy, most have very little quartz sand, and will dissolve in hydrochloric acid leaving only a residue of fine silt. Both of these zones have distinct enough faunas to be recognizable not only when they occur in place but also when found as building stones. Many of the blocks are made up of almost solid molluscan fossils, both fragmental and entire. The thickest calcarenite beds in the Mesa Verde section occur near the top of the Juana Lopez Member, where they total about 9 feet, though the individual beds are usually less than a foot thick and most are one to six inches in thickness. Some of the material used for building stone comes from this zone (referred to hereafter in this report as "zone 4"). About 30-35 feet lower in the section is another group of calcarenites with a somewhat different fauna. This will be referred to as "zone 3" and is the most widespread zone exposed in the Yucca House environs. The most diagnostic fossils are the ammonite assemblages. The upper (zone 4) calcarenites contain *Prionocyclus novimexicanus* (Marcou) (figures 11 C-D) and *Scaphites whitfieldi* Cobban (figure 10A). Zone 3 contains *Prionocyclus wyomingensis* Meek (figures 11A-B, 18) and *Scaphites warreni* Meek and Hayden (figure 10A) in addition to *Inoceramus perplexus* Whitfield (figure 15). About 30 feet lower a calcarenite unit (unit 2) contains a much more limited fauna mostly of *Lopha lugubris* (Conrad) (figure 13) and *Inoceramus dimidiatus* White (figures 14 A-B). A still lower calcarenite unit has been recognized in the Mesa Verde section, characterized by the presence of *Prionocyclus macombi* Meek, but was not found at Yucca House, probably due to poor exposures.



Figure 6. *Prionocyclus wyomingensis* Meek (zone 3) in place in gully bed rock.

Outcrops of the more solid calcarenite which would make good building stone, similar to building blocks in the mounds, are found in the gullies mainly to the west and southwest of the present Yucca House National Monument. The same faunal beds are present to the north, but mostly in thin-bedded layers. Good solid building stone was not found capping the tops of the mesas. Although Quaternary deposits and vegetation cover much of the area, the three distinctive faunal units were found and recorded as well as possible from the limited exposures along a reasonably continuous trend (figure 7).




Figure 7. Approximate distribution of the faunal zones of the Mancos Formation in the vicinity of Yucca House. (Not outcrop pattern)

Identification of the zone 3 and zone 4 ammonites by field archeologists may pose some problems. An illustrated, simplified description of each of the more important guide fossils, with its diagnostic characteristics is appended to this report. However, many other macrofossils are present in the Juana Lopez. Illustrations of some of these may be found in Kauffman, 1977 and Leckie, et al., 1997.

In fragments, it is often difficult to separate *Prionocyclus novimexicanus* from *P. wyomingensis*, but the association with the two *Scaphites* species is helpful. While the *Lopha lugubris* can occur in a much longer range, the large accumulations are characteristic of the lower paleontologic zone. *Inoceramus dimidiatus* also has a longer range, but the faunal assemblage again will help with recognizing a zone. Many other fossils are found in the Juana Lopez, but are not as diagnostic or easily recognized as the above. Small *Baculites* are frequently found. *Baculites undulatus* (figures 10A, 18) and *B. yokoyami* occur in zones 3 and 4 respectively as fragments and may be useful as

guide fossils to the zones if identifiable. Inoceramids are very common, both as fragments and entire. They are frequently difficult zones.

The Juana Lopez represents a period of quiet deposition far from shore, with little clastic material being brought into the area. Although the calcarenite is largely composed of bioclastic debris such as broken shell material, some probably represents a chemical recrystallization of calcium carbonate derived from the molluscan shells.

Overlying the Juana Lopez Member are the Montezuma Valley and the Smoky Hill Members of the Mancos Formation. The Montezuma Valley Member consists primarily of soft calcareous shale and bentonitic shale and was not recognized as a mappable unit in outcrops. The overlying Smoky Hill Member of calcareous shale and fossiliferous limestone is slightly more resistant and erosion remnants were found with the characteristic *Pseudoperna*-*Inoceramus* fauna (figures 16,17). The biostromes of oysters are more resistant to erosion than the shales above and below them and are frequently preserved in isolated low hills or even as broken fragments in cultivated fields. This member does not actually crop out in the proposed monument area, but has been mapped for reference. The best outcrop of the Smoky Hill is on a small butte T34N, R17W, sec 1 (figure 8). Just to the west and south of the butte is a very deep draw with soft dark gray shale in the bottom, overlain by about 40 feet of alluvium and soil. The gray shale is probably the Montezuma Valley Member.




Figure 8. Smoky Hill erosion remnant capped with yellow Point Lookout Formation colluvium. *Inoceramus* fragments with *Pseudoperna* encrusting them in lower slopes of butte.

Between the Smoky Hill Member and the Point Lookout Formation, lies the very thick unit of the Cortez member, formerly referred to as the Transitional zone. Deposited at the beginning of a temporary withdrawal of the sea from this area, this soft unit consists of nearly 1200 feet of interbedded soft shale, silty to sandy shale, shaly sandstone, siltstone, and thin bedded sandstone (figure 9). The unit coarsens upwards to grade into the Point Lookout Sandstone Formation. The soft gray slopes below the Mesa Verde are all part of the Cortez Member of the Mancos Formation. Scattered fossils are present, but not common. Although scattered fragments of *Inoceramids*, *baculites*, and trace fossils are found, fossil zones in this unit are scarce and difficult to recognize. Several beds of dense dolomitic siltstone form resistant ledges locally protecting soft shales in gullies (figure 9). Most of the deeply gullied slopes are covered with Point Lookout colluvium.




Figure 9. Soft light gray to tan sandy shales of the Cortez Member of the Mancos Formation, overlain by resistant Point Lookout Sandstone. Dolomitic siltstone layers in right gully locally protect soft shales to form small scarps.

The Point Lookout, Menefee and Cliff House Formations border the east side of the study area of the Montezuma Valley along the Mesa Verde cliffs (figure 9). The Point Lookout Formation is the basal member of the Mesaverde Group and conformably and gradationally overlies the Mancos Formation. The outcrops on the West Rim of the Mesa Verde are within Ute lands, and it was not possible to get a permit to land there with a helicopter. Presumably, the Point Lookout, Menefee and Cliff House Formations, which all crop out along the West Rim in Ute land are similar in character to those studied within Mesa Verde National Park. They do not crop out on Ismay land. The Point Lookout Formation influences the Yucca House Monument because it is found in colluvial deposits capping the top of the erosion remnants near the new eastern boundary. The yellow tops of the small hills and buttes are remnants of Point Lookout colluvium that was deposited before the hills were isolated by erosion.

The Point Lookout Formation includes about 300 feet of light colored, resistant sandstone and shaly sandstone. In this area two massive units of sandstone make up the bulk of the formation thickness, with relatively thin shaly sandstone breaks. The contact between the Mancos Formation Cortez member and the Point Lookout Sandstone is gradational, and difficult to mark, and the base of the Point Lookout Formation is generally placed where the sandstone layers become non argillaceous and are more numerous than the muddy layers. The Point Lookout Formation represents a regressive interval in the sedimentary history and was deposited as shore face and strand plain sand. Recent studies by Wright--Dunbar, Zech, and others have shown that some of the deposits are deltaic and lagoonal as well. Here the well cemented sandstones form high cliffs protecting the soft Mancos shales. Although the solid sandstone blocks would be good building stone, none is visible in the unexcavated mounds of Yucca House. Presumably it would have been impractical to transport so far when there was good building material much closer. If there are prehistoric dwelling sites on the eastside of the Montezuma Valley, they will probably include material from the Point Lookout Sandstone, rather than from the Juana Lopez calcarenites.

Overlying the Point Lookout Formation is the nonmarine Menefee Formation. This unit of sediments is composed mostly of relatively soft shales, woody shales, coal and irregular soft sandstones. It forms the slopes above the Point Lookout cliffs and below the overlying Cliff House Sandstone cliffs high

above the Montezuma Valley. The Menefee Formation was deposited on land on the low relief former sea bottom. Low interstream areas developed soil and vegetation, giving rise to carbonaceous shales. The stream sandstones within the Menefee Formation are discontinuous and not well enough indurated in most places to be utilized for building material. In the Yucca House archeological collection are pendants made of red burnt shale. Well above the Point Lookout Sandstone, bright red burnt shale is visible, presumably the result of the burning of a coal bed in the Menefee Formation. The overlying shales are baked and the iron content gives a red color. These burned beds are found in many locations along the mesa edges and are very common along the Mancos River to the south, so the exact source of the carved burnt shale artifacts cannot be determined. Coal beds within the Menefee have been mined in this area in historic times, but prehistoric use of coal has not been documented at this locality.

Overlying the Menefee Formation is the top member of the Mesaverde Group, the Cliff House Formation deposited in the readvancing seaway. The Cliff House Formation consists of two massive sandstone units separated by shaly sandstones in Mesa Verde National Park. The formation was not examined at this locality. It is the source for much of the prehistoric building stone in Mesa Verde National Park and in the Ute Mountain Tribal Park to the south, but no building blocks of Cliff House Sandstone were recognized at the Yucca House mounds. If there are prehistoric building sites on the east side of the valley, nearer the Mesa Verde, they will very likely be found to be built using the local Point Lookout and Cliff House sandstones. .

Because most of the valley has been cultivated over the years, and alluvial soils and crops cover much of the surface, bed rocks crop out only in the deepest gullies, along ditches and other excavations and in a few low hills on the extreme west and on the east below the Mesa Verde cliffs. Sediments of Quaternary terraces, pediments, and alluvium bury the bed rock to at least 40 feet depth in some places. Therefore in many parts of the map the contacts of underlying sediments have had to be approximated.

STRUCTURE

The Cretaceous sedimentary beds underlying the Yucca House Monument area are along the northern part of the San Juan structural basin. The gentle southeast dip toward the basin is interrupted by the structural dome resulting from the laccolithic intrusives of the Ute Mountains, but there is very little effect of this doming visible within the study area. The tectonic setting of the Four Corners region has been recently summarized by Woodward, Anderson and Lucas (1997). Most of the sediments in the study area are very gently dipping to the southeast generally at less than 5 degrees, with a strike about N60E. Dips are gentle and many bedding planes are so irregular due to burrowing and reworking of surfaces during deposition that an accurate dip is difficult to measure. Exposures are poor, mainly in the gully bottoms, where the gully has been eroded through the Quaternary overburden to bed rock and follows down along the gentle dip slope. No faults are visible in the bed rock of the valley, but small faults are visible along the mesa rim, though not within the area covered by a detailed map. The faults shown on the west edge of the Mesa Verde, were located from aerial photographs and Wanek's (1959) map. There is so much soil and overburden on top of the bedrock in the Montezuma Valley, that minor structures are hidden. No stratigraphic anomalies were noted which would require structural explanation.

IGNEOUS HISTORY

Although no igneous rocks crop out in the Ismay-Yucca House holdings, igneous rocks are very common, both as loose cobbles and pebbles, and in building blocks in the prehistoric structures. The Ute Mountain laccoliths and other intrusive bodies crop out less than three miles to the west. These igneous rocks are dated as of Laramide age - very late Cretaceous or very early Tertiary. They were intruded through and between layers of old sedimentary rocks as stocks, laccolithic bodies, dikes and sills. Although there is variation from one intrusive to another, most are light to dark gray, fine- to medium-grained, porphyritic, plagioclase rich rocks. Most of the fine grained gabbro is very dark gray to almost black. The crystalline structure of the ground mass is very fine, but visible to the naked eye.

Mineralogically the ground mass is mostly composed of plagioclase and chlorite with phenocrysts of augite or hornblende (Ekren and Houser, p. 28). More common than the fine grained gabbro, are diorite porphyry, granodiorite porphyry and monzonite porphyry. All of the igneous intrusives examined were porphyritic, and the phenocrysts are usually plagioclase, hornblende or augite. These rocks are found as hammer stones and other artifacts, as well as building stones.

SUMMARY

The bed rocks of the Izmay holdings surveyed are entirely of Upper Cretaceous age. Most of the area is underlain by Mancos Formation sediments. The Point Lookout, Menefee and Cliff House Formations crop out only along the Mesa Verde scarp on the east border of the surveyed area. Igneous rocks from the Ute Mountain intrusives do not crop out in the area, but are abundantly present as pebbles to large boulders in stream beds, and as alluvial and terrace deposits. Yucca House building blocks as seen in the present rubble mounds were derived from two main sources. Many of the blocks are debris from the laccolithic masses to the west which were brought down to the valley in streams and landslides. What appears at this time to be the majority of the building material was derived from the upper two units of calcarenite from the Juana Lopez member of the Mancos Formation. These crop out in several gullies about 1/2 mile west of the Yucca House mounds. When this site is eventually excavated, it may disclose use of some of the earlier Juana Lopez layers or even Dakota Sandstone. Holmes (1878, p. 399-400) assumption that the "stone used is chiefly the fossiliferous limestone that outcrops along the base of the Mesa Verde" was erroneous. There is no limestone cropping out at the base of the Mesa Verde. The diagnostic fossiliferous limestone (calcarenite) crops out to the west of the building site. No sandstone blocks of the Point Lookout or Cliff House Formations were seen in the mounds, and it certainly is unlikely that the early Pueblos would have carried in heavy building blocks from the east side of the valley, when the local stone was so readily available.

ACKNOWLEDGMENTS

This work was financed by the Mesa Verde National Park. The author received much assistance in the field from Marilyn Colyer and direction and encouragement from George San Miguel both of the Natural Resources Division of Mesa Verde National Park. Base maps were furnished by Allan Loy and digitizing of the final geologic map was done in his office. William S. Cobban of the United States Geological Survey (retired) advised on fossil identification and stratigraphy. We wish to thank the Ute Mountain Ute Tribal Council for permission to access part of their land. Don Whyte very helpfully drove us across some of the back country of the Ute land.

Report from source map: [Yucca House National Monument Area \(Bedrock Geology and Paleontology\)](#)

References

- Anderson, O.J. and Lucas, S.G., 1997, The Upper Jurassic Morrison Formation in the Four Corners region: in Mesozoic Geology and Paleontology of the Four Corners region, New Mexico Geological Society, 48th Annual Field Conference Guidebook, p.139-155.
- Cobban, W.A., 1951, Scaphitoid cephalopods of the Colorado group: U.S. Geological Survey Prof. Paper 239, p. 21-25, plates 3-5.
- Cobban, W.A. and Hook, S.C., 1983, Mid-Cretaceous (Turonian) ammonite fauna from Fence Lake area of west-central New Mexico: New Mexico Bureau of Mines and Mineral Resources, Memoir 41, 50 p.
- Cobban, W.A. and Hook, S.C., 1989, Mid-Cretaceous molluscan record from west-central New

- Mexico: New Mexico Geological Society Guidebook, 40th Field Conference, Southeastern Colorado Plateau, p. 247-264.
- Condon, S.M., 1991, Geologic and structure contour map of the Ute Mountain Ute Indian Reservation and adjacent areas, southwest Colorado and northwest New Mexico: U.S. Geological Survey Misc. Invest. Series Map 1-2083.
- Condon, S.M. and Huffman, C., 1984, Stratigraphy and depositional environments of Jurassic rocks, San Juan Basin, New Mexico, with emphasis on the south and west sides: Rocky Mountain Section of Geological Society of America Field Trip Guidebook, 37th Annual Meeting, Durango, Colorado, p.93-107.
- Dane, C.H., Cobban, W.A. and Kauffman, E.G., 1966, Stratigraphy and regional relationships of a reference section for the Juana Lopez Member, Mancos Shale, in the San Juan Basin, New Mexico: U.S. Geological Survey, Bulletin 1224-H, 15 p.
- Ekren, E.B. and Houser, F.N., 1965, Geology and petrology of the Ute Mountains area Colorado: U.S. Geological Survey Professional Paper 481, 74 p., map.
- Haas, Otto, 1946. Intraspecific variation in, and ontogeny of, *Prionotropis woollgari* and *Prionocyclus wyomingensis*: American Museum of Natural History. Bull., v.86, art. 4, p. 141-224.
- Holmes, W.H., 1875-76. Report on the Ancient Ruins of Southwestern Colorado, examined during the summers of 1875 and 1876, in F.V. Hayden, 1878, Tenth Annual Report of the United States Geological and Geographical Survey of the Territories, p. 399-400, Plate XL.
- Hook, S.C. and Cobban, W.A., 1977, *Pycnodonte newberryi* (Stanton) - common guide fossil in Upper Cretaceous of New Mexico: New Mexico Bureau of Mines and Mineral Resources, Annual Report 1976-1977, p. 48-54.
- Hook, S.C. and Cobban, W.A., 1979, *Prionocyclus novimexicanus* (Marcou)- common. Upper Cretaceous guide fossil in New Mexico: New Mexico Bureau of Mines and Mineral Resources, Annual Report 1977 -1978, p.34-42.
- Hook, S.C. and Cobban, W.A., 1980, Some guide fossils in the Upper Cretaceous Juana Lopez Member of Mancos and Carlile Shales, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Annual Report 1978-1979, p. 38-49.
- Hook, S.C. and Cobban, W.A., 1980, Reinterpretation of type section of Juana Lopez Member of Mancos Shale: New Mexico Geology, v. 2, No. 2, 6p.
- Irwin, James H., 1966, Geology and availability of ground water on the Ute Mountain Indian Reservation, Colorado and New Mexico: U.S. Geological Survey Water Supply Paper 1576-G, p. G35-46, Plate I.
- Kauffman, E.G., 1977, Illustrated guide to biostratigraphically important Cretaceous macrofossils, Western Interior basin, U.S.A.: in *The Mountain Geologist*, Cretaceous facies, faunas, and paleoenvironments across the Western Interior basin. V. 14, no. 3-4, p. 225-274.
- Leckie, R.M., Kirkland, J.I., and Elder, W. P., 1997, Stratigraphic framework and correlation of a principle reference section of the Mancos Shale (Upper Cretaceous), Mesa Verde, Colorado: in *Mesozoic geology and paleontology of the Four Corners Region*, New Mexico Geological Society Guidebook, 48th Field Conference, p. 163-216.
- Rankin, C.H., 1944, Stratigraphy of the Colorado Group, Upper Cretaceous, in northern New Mexico: New Mexico Bureau of Mines and Mineral Resources, Bull. 20, p.19.

Scott, G.R., Cobban, W.A., and Merewether, E.A., 1986, Stratigraphy of the Upper Cretaceous Niobrara Formation in the Raton basin, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Bulletin 115, 34p.

Wanek, A.A., 1959, Geology and fuel resources of the Mesa Verde area, Montezuma and La Plata Counties, Colorado: U.S. Geological Survey, Bulletin 10722-M, p. 667-717, plate 49.

Woodward, L.A., Anderson, O.J., and Lucas, S.G., 1997, Tectonics of the Four Corners region of the Colorado Plateau: in Mesozoic geology and paleontology of the Four Corners Region, New Mexico Geological Society Guidebook, 48th Field Conference, p.57-64.

Wright, C.W., 1957, Family Collignoniceratidae, in Treatise on Invertebrate Paleontology, R.C. Moore ed.: Geological Society of America and University of Kansas, part L, p. L426-427, Fig. 547.

References from source map: [Yucca House National Monument Area \(Bedrock Geology and Paleontology\)](#)

Appendix

The fossils collected during this study are catalogued and stored in the Mesa Verde National Park museum collection.

DESCRIPTIVE PALEONTOLOGY OF GUIDE FOSSILS

AMMONITES

Scaphites warreni Meek and Hayden (see Figure 10A)

Scaphites are a group of ammonites which have involute early coils, followed by a living chamber that is loosely coiled or almost straight. *Scaphites warreni* averages 35 mm length, subovate in cross section, stout with small umbilicus. The living chamber is distinctly free from coiled portion. The early whorls are smooth. Adult forms show primary ribs which bend slightly where crossing the flank and at the ventrolateral margin swell into rounded nodes and divide into secondary ribs which extend straight across the venter. A second form of this species (dimorphic) is more slender, less ovate in cross section and smaller, but has similar ornamentation. This species of Scaphite is found in unit 3 of the Juana Lopez Member of the Mancos Formation, and is a very good guide fossil for this horizon.

Scaphites whitfieldi Cobban (see Figure 10B)

Scaphite similar in size to *S. warreni*, but more slender and distinctly quadrangular in cross section. Ornamentation on adult form is more dense and has more evenly spaced ribs on the venter, and this ornamentation combined with the slender, compressed, quadrangular cross section easily separates *S. whitfieldi* from *S. warreni*. This species also has two forms- a macroconch or female which is more robust and has a characteristic umbilical swelling on the living chamber, and a slender smaller male form.

This species of Scaphite is found in unit 4 of the Juana Lopez Member of the Mancos Formation, and is a very good guide fossil for this horizon.

Prionocyclus wyomingensis Meek (see Figures 11A-B)

A moderately large evolute ammonite with a somewhat rectangular whorl cross section, a broad venter and a continuous serrate keel. The inner whorls are more compressed and have finer ribs, much closer together, and with less prominent nodes. On the outer whorls of larger specimens, 7 cm in diameter and larger, the whorl section begins to show slight convexity between the umbilical and

ventrolateral nodes. The ribs are strong with the primary ribs usually bearing umbilical nodes. Secondary ribs in some cases branch from the primaries and in other cases are intercalated between the primaries. As the ribs approach the ventrolateral shoulder, another node is formed on the primaries. At that point, the rib bends abruptly forward toward the keel, disappearing just before contact, so that a slight depression exists on either side of the strong median keel. The keel shows moderately strong serrations.

The diagnostic characteristics of this species is the ovoid rectangular whorl section, the strong clear ribs, and the double row of ventrolateral nodes.

The faunal association with *Scaphites warreni* and *Inoceramus dimidiatus* with *Prionocyclus wyomingensis* helps separate the two common *Prionocyclids*. This faunal group is found in unit 3 of the Juana Lopez Member of the Mancos Formation.

Prionocyclus novimexicanus (Marcou) (see Figures 11C-D)

Similar in many characteristics to *P. wyomingensis*, in fact was originally described as *P. wyomingensis* var. *elegans*. *P. novimexicanus* has only a single row of ventrolateral tubercles instead of the double row of *P. wyomingensis* and has a low ventral ridge bordering a smooth ventral furrow. Haas (1946, p. 200-201) observed that it also has a more compressed whorl section and more distinctly ribbed juvenile whorls. In the field, the most obvious difference is the more compressed, rectangular cross section of the whorl and lack of serration on the keel.

Again the faunal association is useful in identifying this species. It is slightly younger than *P. wyomingensis* and occurs with *Scaphites whitfieldi* and characterizes the upper unit 4 calcarenite zone of the Juana Lopez Member of the Mancos Formation.




Figure 10A. *Scaphites warreni* Meek and Hayden, *Baculites undulatus* d'Orbigny, with juvenile *Inoceramus* and fragment of *Prionocyclus* on slab from Yucca House with zone 3 faunal assemblage. (YUHO # 17). X2.




Figure 10B. *Scaphites whitfieldi* Cobban (YUHO #98) X1




Figure 11.
Prionocyclus
wyomingensis Meek
(YUHO #19)

Prionocyclus
novimexicanus Cobban
(MEVE 51231)

Baculites undulatus d'Orbigny (see Figure 10A, 18)

The Baculites are a group of ammonites with only a minute coil of one to two whorls before growing out into a straight or very slightly bent shaft. Associated with the *Scaphites warreni*-*Prionocyclus wyomingensis* fauna are fragments of small, straight Baculites. Most fragmental sections are less than 5 cm in length and usually only about 5 to 7 mm in diameter, although some specimens measure up to 12 mm in diameter. All show a rounded oval cross-section and distinctive, though rather weak, arcuate swellings along the flanks, dorsum and venter, with a slight taper visible even on short fragments.

Baculites yokoyami Tokunaga and Shimizu

Associated with the *Scaphites whitfieldi*-*Prionocyclus novimexicanus* fauna are fragments of another small, straight Baculite. These also are preserved only as fragmental sections 5 cm or less in length, but though the venter may show closely spaced ribs, the flanks are usually relatively smooth. The cross section is oval. This Baculite is found in a longer vertical time range.

BIVALVES




Figure 12. *Pycnodonte newberryi* (Stanton)
 A. Convex left valve with small concave right valve in place. (MEVE #11185)
 B. Right valve interior showing muscle scar (MEVE #11185)




Figure 13. *Lophalugubris* (Conrad) Typical slab of multiple individuals.
 (MEVE 7286)

Pycnodonte newberryi (Stanton) (see Figures 12A-B)

Pycnodonte newberryi (Stanton) is a small, highly inequivalve, strongly inrolled species of oyster which is very diagnostic of Graneros shale of the Mancos Formation. It has a subovate to subtriangular overall outline. The two valves are very unequal, with the larger left valve strongly inrolled and usually with a conspicuous radial sulcus. Ornamentation consists of irregular growth lines. The right valve is usually flat to slightly concave and ornamentation consists of concentric growth lines. The interior of each valve shows a conspicuous single muscle scar. The attachment area is usually at the apex of the left valve umbone.

It is a relatively unique, easily recognized fossil, and also is quite resistant to weathering. With its almost spherical shape, it rolls out of outcrops and often accumulates in large quantities in stream beds.

Lophalugubris (Conrad) (see Figure 13)

Lophalugubris (Conrad) is a small, inequivalve oyster whose presence in quantities is very diagnostic of the lower calcarenite unit 2 of the Juana Lopez. This oyster has a longer vertical range, found from unit 1 through unit 3 of the Juana Lopez Member of the Mancos Formation, but only in unit 2 is it the dominant species and found covering most of the slabs of calcarenite.

Lophalugubris is easily recognized by its crenulate, subovate outline. The left valve is usually more

convex than the right, and has a large attachment area. The shells of this species are attached to *Inoceramus* shells, and often have attachment areas covering half of the left valve. The right valve is flat to slightly convex. Both valves show a single conspicuous adductor muscle scar. Both valves have strongly plicate outlines which are quickly diagnostic in the field.



Figure 14. *Inoceramus dimidius* White (YUHO #151)
A. Cast of interior of left valve showing sharp change in slope of shell.
B. Profile of same specimen. X1

***Inoceramus dimidius* White (see Figures 14A-B)**

Inoceramus dimidius White is a small, inequivalve inoceramid that occurs with *Lopha lugubris* in unit 2 and also in unit 3 with the *Scaphites warreni-Prionocyclus wyomingensis* faunas. It is easily recognized because of the major break in ornamentation between the earlier stages of growth where there are strong rugae and the later part of the shell which is almost smooth. At the juncture of the two types of ornamentation, a prominent angular change in the slope of the shell, makes an easy diagnostic characteristic.





Figure 15. *Inoceramus perplexus* Whitfield (YUHO 62) A. Cast of exterior left valve, showing regular ornamentation over entire shell.
B. Profile view of same specimen.

Figure 16. *Pseudoperna congesta* (Conrad) (YUHO 158) oysters encrusting large *Inoceramus* X2

Figure 17. Side view of prismatic shell layer of large *Inoceramus* Upper surface encrusted with *Pseudoperna congesta*. (YUHO 158) X2



***Inoceramus perplexus* Whitfield (see Figures 15A-B)**

Inoceramus perplexus Whitfield is a small inoceramid with regular, sharp, concentric growth lines


throughout the shell. This species occurs in Juana Lopez unit 4 with the *Scaphites whitfieldi*-*Prionocyclus novimexicanus* fauna

Pseudoperna congesta (Conrad) (see Figure 16)

A small irregular to ovate oyster, usually found in masses encrusting the prismatic layer of large Inoceramid such as *Inoceramus grandis*. Where crowded, develop a very irregular shape due to impingement by neighboring individuals. Where sparse population, may be nearly oval. Valves very unequal with attachment area on left (larger) valve very large. Right valve often crushed into left valve by weight of overlying sediments. The solid oyster beds are quite resistant to erosion. This is a very widespread, diagnostic species found throughout the Western Interior in Niobrara age sediments. In the Mancos section it is referred to the Smoky Hill Member. The Smoky Hill is the upper unit of the Niobrara elsewhere. The lower unit is missing in the Four Corners area.

Inoceramus (Volviceramus) grandis Conrad (see Figure 17)

A very large, irregular Inoceramid, frequently found only as cross sections of the prismatic layer of the shell. Occasionally these cross sections are up to a meter in width. More often, they are seen only as fragments, densely encrusted by *Pseudoperna congesta*. Individually, it is rare to see a complete *I. grandis*.



Appendix from source map: [Yucca House National Monument Area \(Bedrock Geology and Paleontology\)](#)

GRI Digital Data Credits

This document was developed and completed by Georgia Hybels and James Winter (Colorado State University) for the NPS Geologic Resources Division (GRD) Geologic Resources Inventory (GRI) Program. Quality control of this document by Stephanie O'Meara (Colorado State University).

The information in this document was compiled from GRI source maps, and is intended to accompany the digital geologic-GIS maps and other digital data for Yucca House National Monument, Colorado (YUHO) developed by Georgia Hybels, James Winter, and Stephanie O'Meara (Colorado State University) (see the [GRI Digital Maps and Source Map Citations](#) section of this document for all sources used by the GRI in the completion of this document and related GRI digital geologic-GIS maps).

GRI finalization by Stephanie O'Meara and James Winter (Colorado State University).

GRI program coordination and scoping provided by Bruce Heise, Jason Kenworthy and Tim Connors (NPS GRD, Lakewood, Colorado).