

THE SOLITARY RUGOSE CORAL *ACANTHOPHYLLUM* IN THE SHELLABARGER LIMESTONE (LATE EMSIAN, LATE EARLY DEVONIAN) AT DENALI NATIONAL PARK AND PRESERVE, SOUTH-CENTRAL ALASKA

ROBERT B. BLODGETT¹, VINCENT L. SANTUCCI², ALAN E.H. PEDDER³ and VALERYI V. BARANOV⁴

¹Consulting Geologist, 2821 Kingfisher Drive, Anchorage, Alaska 99502, email: robertbbloodgett@gmail.com; ²Senior Paleontologist, National Park Service – Geologic Resources Division, 1849 C Street, N.W., Room 2644, Washington, D.C. 20240, email: vincent_santucci@nps.gov;

³deceased, 8859 Park Pacific Terrace, Sidney, British Columbia V8L 4SL, Canada; ⁴Diamond and Precious Metals Geology Institute, Siberian Division, Russian Academy of Sciences, pr. Lenina 39, Yakutsk, 677000, Russia, email: baranowvalera@yandex.ru

Abstract—In this short note we provide documentation of the first rugose coral (*Acanthophyllum* sp.) to be formally illustrated from the Shellabarger Limestone of late Early Devonian (Emsian) age in Shellabarger Pass, Talkeetna C-6 quadrangle, south-central Alaska. This also comprises the first rugose coral to ever be illustrated from Denali National Park and Preserve. In addition, we provide information on its stratigraphic occurrence as well as of other reported occurrences of *Acanthophyllum* noted in Alaska.

INTRODUCTION

In this note we provide the first illustrations of Devonian rugose corals from Shellabarger Pass (see Fig. 1 for general location), Denali National Park, Alaska. Devonian brachiopods are now prominently known from this richly fossiliferous area. We are currently trying to document most of the Devonian megafauna, and hope soon to finish research on our richly diverse Emsian brachiopod megafauna. In the near future we will emphasize the Frasnian age fauna of the region. Detailed studies of the megafauna did not begin here until the later part of the 1990s, and these and subsequent studies indicate a strong Eurasian alliance, especially with Northeast Russia, rather with adjacent cratonic North American Devonian rocks of east-central Alaska, suggesting the host Farewell terrane originated as a rift block derived from NE Russia.

The Emsian fauna is from the Shellabarger Limestone, which was established by Blodgett et al. (2022). See Figure 2 for its outcrop belt in the Shellabarger Pass region in south-central Alaska where it primarily crops out in the Talkeetna C-6 quadrangle of south-central Alaska. The type section of the Shellabarger Limestone has a maximum measured thickness of 38.7 m (127 ft) (see Fig. 3).

GEOLOGIC SETTING

Blodgett et al. (2022, p. 73-75) provided the following discussion of the geologic setting of the Devonian of the Shellabarger Pass. The specimens described here are from the base of the Mystic sequence of the Farewell terrane of southwestern and west-central Alaska. The Mystic terrane was originally named by Jones and Silberling (1979) and described further by Jones et al. (1981) as a separate tectonostratigraphic entity of full terrane rank. More recently, Decker et al. (1994) recognized that the Mystic terrane, as well as the Nixon Fork and Dillinger terranes, were genetically related, and all were reduced in rank to subterrane of a larger terrane, termed the Farewell terrane. Gilbert and Bundtzen (1984) considered the Dillinger and Mystic terranes, each of whose type sections are close to one another, to represent a single stratigraphic succession of Paleozoic to Triassic age, assigning the term “sequence” to each. They considered the underlying Dillinger sequence to be a Cambrian to Lower Devonian deep-water succession overlain by the Mystic sequence, consisting of laterally variable Devonian to Triassic? shallow-water to nonmarine sedimentary rocks with intrusive and extrusive mafic and ultramafic rocks. The close stratigraphic relationship between the Dillinger and Mystic sequences has been studied by Blodgett and Gilbert (1992) to the southwest in the Lime Hills D-4 and D-5, 1:63,360 scale quadrangles.

The Nixon Fork subterrane, defined originally as a full-rank terrane by Patton (1978), represents a predominately shallow-water carbonate platform whose lower part is mostly laterally equivalent to deeper-water Cambrian to Lower Devonian rocks of the Dillinger subterrane (Blodgett, 1983; Blodgett and Gilbert, 1983; Bundtzen and Gilbert, 1983; Gilbert and Bundtzen, 1983). Until the late 1990s, most interpretations of the gross lithostratigraphy and faunal affinities of the best-studied subterrane of the Farewell terrane, the Nixon Fork subterrane, had indicated either fragmentation from or lithic continuity with northwestern Canada (Blodgett, 1983; Churkin et al., 1984; Blodgett and Clough, 1985; Rohr and Blodgett, 1985; Abbott, 1995). Newly acquired biogeographic data from Lower Devonian strata of the Farewell terrane, together with reconsideration of the affinities of the faunas and floras of other time intervals, indicate that this continental margin sequence originated by rifting away from the Siberian continent, rather than an origin from northwestern North America (Blodgett and Brease, 1997; Blodgett, 1998; Blodgett et al., 2002). Much of this rifting appears to have occurred during Late Devonian to early Carboniferous time (Blodgett et al., 2010).

Richly fossiliferous Devonian rocks of the basal part of the Mystic sequence are well exposed in the Shellabarger Pass area, Talkeetna C-5 & C-6 quadrangles of south-central Alaska. The



FIGURE 1. Index map of Alaska showing location of Shellabarger Pass.




FIGURE 2. Map showing location of Late Early Devonian (Emsian) limestone outcrop in Shellabarger Pass, Denali National Park and Preserve, Talkeetna C-6 quadrangle, south-central Alaska. Outcrop shown in yellow color surrounding the measured type section. Line indicates measured section. Base of section located at its north end.

pass is situated in the remote western part of the Preserve portion of Denali National Park & Preserve (Fig. 4). Field work was first seriously attempted there in the mid-1970's and was followed by our 1996 summer field work with financial support from the National Geographic Society to study an unnamed limestone unit that forms the ultimate local base of the Mystic sequence. Two separate limestone packages are separated by a siliciclastic/chert unit. The lower limestone interval is of Emsian (late Early Devonian) age and is known from only one outcrop. The upper limestone interval, representing the vast majority of Devonian limestone outcrops in Shellabarger Pass, is of Frasnian (early Late Devonian) age. The fauna from the Emsian limestone includes sponges (Rigby et al., 2009), rugose and tabulate corals, approximately 20 or more species of brachiopods, and trilobites. Common brachiopod genera observed from the Emsian outcrop include *Carinagypa*, *Iydelinia*, *Clorinda*, *Myriospirifer*, *Sibirirhynchia*, *Spinatrypa*, *Spinatrypina*, *Variatrypa*, *Carinatina*, *Opsiconidion*, and *Teichertina*. Previous work (Blodgett and Brease, 1997, Blodgett, 1998, Blodgett and Boucot, 1999, Garcia-Alcalde and Blodgett, 2001, and Blodgett et al., 2002) indicates a much stronger Eurasian component to the Farewell terrane, as opposed to the Cordilleran fauna known from east-central Alaska and Yukon Territory of westernmost Laurentia. This overwhelmingly Eurasian aspect suggests that the host terrane, the Farewell terrane, was a continental fragment of Siberian origin.

The Frasnian limestone fauna has not yet been thoroughly reinvestigated taxonomically, but it is a proposed target for our future study. Preliminary identifications indicated the presence of the brachiopod genera *Schizophoria*, *Productella*, *Atrypa*, *Spinatrypa*, *Allanaria*?, *Tenticospirifer*, *Cyrtospirifer*, *Crurithyris*, *Eleutherokomma*, *Gypidula*, "*Camarotoechia*,"




FIGURE 3. Measured type section of the Shellabarger Limestone. This section forms a single isolated limestone outcrop and constitutes the oldest Devonian interval within the D1 unit of Reed and Nelson (1977, 1980) and forms the oldest stratigraphic interval within the Mystic sequence of Gilbert and Bundtzen (1984). This outcrop is situated on the eastern side of a small hillock situated near the center of the NE $\frac{1}{4}$, SW $\frac{1}{4}$, NE $\frac{1}{4}$, sec. 15, T28N, R18W, Talkeetna C-6 1:63,360 scale quadrangle, latitude 62°31'21"N., longitude 152°35'32" W. The measured section type section of the outcrop indicates a minimum thickness of 38.7 m (127 ft) of lime mudstone and wackestone for the unit (Blodgett et al., 2022, see Fig. 3 and 4). The discussed corals here were found in talus rubble derived from 14.0-21.3 m (46-70 ft.) above the base of the section.



FIGURE 4. Outcrop of the upper part of the type section of the Shellabarger Limestone (Emsian, Late Early Devonian) in Shellabarger Pass (view looking east-northeast). The two people in the photo are Pam Sousanes (NPS) (left) and one of the authors, Robert B. Blodgett (on right).

Ladogioides?, *Theodossia*, *Emanuella?*, and *Tylothyris*, as well as rugose and tabulate corals, the ammonoid *Manticoceras*, and the trilobite *Scutellum*.

CORALS FROM THE DEVONIAN OF SHELLABARGER PASS

The report of *Acanthophyllum* here is from richly fossiliferous Devonian rocks of the basal part of the Mystic sequence well exposed in the Shellabarger Pass area, Talkeetna C-6 quadrangle of south-central Alaska. These basal Emsian age beds were named the Shellabarger Limestone by Blodgett et al. (2022) and recognized to be minimally 38.7 m thick (see Fig. 3). The pass is situated in the remote western part of the Preserve portion of Denali National Park & Preserve. Fossils from this unit (megafossils and conodonts) indicate a late Emsian age.

Corals were listed previously in Reed and Nelson (1977, 1978) from this then unnamed stratigraphic unit, but no detailed taxonomic information was provided. J.T. Dutro, Jr. in an unpublished USGS E&R report dated February 2, 1976 (Shipment number A-75-18) on collection 75AR67 collection from the unit noted the rugose coral *Dohmophyllum* sp. and horn coral, indet. Unfortunately no illustrations of either form are available, but a concerted effort should be able to relocate them in the former USGS collections, which have been turned over to the National Museum of Natural History (Smithsonian Institution).

Overlying Devonian strata in Shellabarger Pass appear to belong to an unnamed carbonate unit of Frasnian (Early Late Devonian) age. William A. Oliver, Jr. of the USGS provided




FIGURE 5. Five specimens of the solitary rugose coral *Acanthophyllum* sp. found in talus adjacent to 14.0-21.3 m (46-70 ft.) above the base of the section. All are transverse sections and are deposited in the Alaska Geologic Materials Center (prefix AKGMC) in Anchorage, Alaska. Specimen numbers are AKGMC 62-66. A – AKGMC-62, B – AKGMC-63, C – AKGMC-64, D – AKGMC-65, and E – AKGMC-66.

identifications on fossils in a series of USGS E&R reports (available online through the Alaska Paleontological Database (www.alaskafossil.org)). Both rugose and tabulate corals were listed, including rugosans: *Tabulophyllum* sp., *Phillipsastraea* sp. (2 or 3 spp.), *Phillipsastrea?* sp., *Macgeea* sp., *Disphyllum* sp., *Disphyllum?* sp., *Syringaxon* sp., *Smithophyllum* sp., *Peneckiella?* sp., *Thamnophyllum* sp., *Paracanthus?* sp. and tabulates: *Alveolites* sp., *Thamnopora* sp., *Cladopora* sp., *Hexagonaria* sp., and *Syringopora* sp. None of these forms have previously been illustrated. In light of the relatively high diversity of the Frasnian coral collections, they would be an excellent target for future taxonomic study.

SYSTEMATIC PALEONTOLOGY

Repository: The rugose corals specimens illustrated in this paper are deposited at the Alaska Geologic Materials Center (prefix AKGMC), Anchorage, Alaska. These consist of about 9 coralla, the best of which are illustrated in Figure 5.

Order Rugosa Milne-Edward & Haime, 1850
Suborder Columnariina Rominger, 1876
Family Ptenophyllidae Wedekind, 1923
Genus *Acanthophyllum* Dybowski, 1873

Type species: *Cyathophyllum heterophyllum* Milne-Edwards & Haime, 1851

Acanthophyllum sp.
 Fig. 5

Discussion: The genus *Acanthophyllum* is a well known and widely distributed solitary rugose coral genus. It is known from the Devonian (Pragian-Frasnian, see Bayer et al., 1956), but is most common in Emsian-Middle Devonian strata. Occurrences are widespread throughout the tropical Old World Realm. Reported occurrences of *Acanthophyllum* in Alaska can be found in Oliver et al. (1975) as well as The Alaska Paleontological Database website (www.alaskafossil.org). The genus is notably common in slightly younger strata (early Eifelian) of the Cheeneetnuk Limestone, McGrath A-5 quadrangle, west-central Alaska.

Material: About 9 coralla are present in the talus collection gathered at 46-70 feet (14.0-21.3 m) above the base of the Shellabarger Limestone. The five best coralla, represented by transverse thin sections, are illustrated here. The associated specimen numbers are: AKGMC-62, AKGMC-63, AKGMC-64, AKGMC-65, and AKGMC-66. These are deposited at the Alaska Geologic Materials Center (prefix AKGMC) in Anchorage, Alaska.

ACKNOWLEDGMENTS

Blodgett expresses his thanks to the Committee for Research and Exploration of the National Geographic Society for providing funds in 1996, which permitted five days of fieldwork in the Shellabarger Pass area of south-central Alaska in late July of the same year. He also thanks Phil F. Brease (deceased) and Pam Sousanes of the U.S. National Park Service, Denali National Park & Preserve, for their able assistance with field work. Carl Bentley, helicopter pilot for Era Helicopters, capably got us into and out of the remote and beautiful area of the central Alaska Range. The Alaska Division of Geological & Geophysical Surveys of Alaska is also acknowledged for providing several brief reconnaissance helicopter trips into the Shellabarger Pass area to Blodgett in the early 1980s.

This work greatly benefited from the recent (post-1995) collaborative efforts on the part of other paleontologists who have studied or made observations of our fossil collections from the Shellabarger Pass area. These include the following individuals: Jenaro L. Garcia-Alcalde (Oviedo, Spain, Devonian brachiopods), Norman M. Savage (University of Oregon, Eugene, Devonian conodonts), and J. Keith Rigby, Sr. (Provo,

Utah, deceased, Devonian sponges). Reviews of a preliminary draft of this paper were graciously provided by Howard R. Feldman, Biology Department, Touro University, New York, New York and Tim Connors, U.S. National Park Service, Lakewood, Colorado.

Finally, we wish to honor one of our co-authors, Alan E.H. Pedder (1932-2023), prominent Devonian coral worker (recently deceased) of Sidney, British Columbia (Fig. 6). He was formerly with the Canadian Geological Survey in Calgary, Alberta, Canada, and was the leading student of Devonian corals from western North America. He was truly a great colleague and friend. We will miss you!



FIGURE 6. Alan Pedder at work in his office at the Geological Survey of Canada in Calgary studying Devonian corals.

REFERENCES

- Abbott, G., 1995, Does Middle Cambrian rifting explain the origin of the Nixon Fork terrane? [abst.]: Geological Society of America Abstracts with Programs, v. 27, no. 5, p. 1.
- Bayer, F.H., Boschma, Hilbrand, Harrington, H.H., Hill, Dorothy, Hyman, Libbie H., Lecompte, Marius, Montanaro-Gallitelli, Eugenia, Moore, R.C., Stumm, E.C., and Wells, J.W., 1956, Treatise on Invertebrate Paleontology, Pt. F., Coelenterata. Geological Society of America and University of Kansas Press, ed., R.C. Moore.
- Blodgett, R.B., 1983, Paleobiogeographic implications of Devonian fossils from the Nixon Fork terrane, southwestern Alaska; in Stevens, C.H., ed., Pre-Jurassic Rocks in Western North America Suspect terranes: Pacific Section, Society of Economic Paleontologists and Mineralogists, Los Angeles, California, p. 125-130.
- Blodgett, R.B., 1998, Emsian (Late Early Devonian) fossils indicate a Siberian origin for the Farewell terrane; in Clough, J.G., and Larson, F., eds., Short Notes on Alaskan Geology 1997: Alaska Division of Geological & Geophysical Surveys Professional Report 118, p. 53-61.
- Blodgett, R.B., Baranov, V.V., and Santucci, V.L., 2002, Two new late Emsian (latest Early Devonian) brachiopods from the Shellabarger Limestone (new formation), Shellabarger Pass, Denali Pass, Denali National Park & Preserve, South-Central Alaska: New Mexico Museum of Natural History and Science, Bulletin 90, p. 73-83.
- Blodgett, R.B., and Boucot, A.J., 1999, Late Early Devonian (late Emsian) eospiriferinid brachiopods from Shellabarger Pass, Talkeetna C-6 quadrangle, south-central Alaska and their biogeographic importance; further evidence for a Siberian origin of the Farewell and allied Alaskan accreted terranes: Senckenbergiana Lethaea, v. 72, no. 1, p. 209-221.
- Blodgett, R.B., Boucot, A.J., Rohr, D.M., and Pedder, A.E.H., 2010, The Alexander terrane – a displaced fragment of Northeast Russia? Evidence from Silurian-Middle Devonian megafossils

- and stratigraphy: Memoirs of the Association of Australasian Paleontologists, v. 39, p. 323-339.
- Blodgett, R.B., and Brease, P.F., 1997, Emsian (late Early Devonian) brachiopods from Shellabarger Pass, Talkeetna C-6 quadrangle, Denali National Park, Alaska indicate Siberian origin for Farewell terrane: Geological Society of America Abstracts with Programs, v. 29, no. 5, p. 5.
- Blodgett, R.B., and Clough, J.G., 1985, The Nixon Fork terrane – part of an in-situ peninsular extension of the Paleozoic North American continent: Geological Society of America Abstracts with Programs, v. 17, no. 6, p. 342.
- Blodgett, R.B., and Gilbert, W.G., 1983, The Cheeneetnuk Limestone; A new Early(?) to Middle Devonian formation in the McGrath A-4 and A-5 Quadrangles, west-central Alaska: Alaska Division of Geological & Geophysical Surveys, Professional Report 85, 6 p.
- Blodgett, R.B., and Gilbert, W.G., 1992, Upper Devonian shallow-water marine siliciclastic strata and associated fauna and flora, Lime Hills D-4 quadrangle, southwest Alaska; in Bradley, D.C., and Dusel-Bacon, C., eds., Geologic studies in Alaska by the U.S. Geological Survey, 1991. U.S. Geological Survey Bulletin 2041, p. 106-115.
- Blodgett, R.B., Rohr, D.M., and Boucot, A.J., 2002, Paleozoic links among some Alaskan accreted terranes and Siberia based on megafossils; in Miller, E.L., Grantz, Art, and Klemperer, S.L., eds., Tectonic Evolution of the Bering Shelf-Chukchi Sea-Arctic Margin and Adjacent Landmasses: Geological Society of America Special Paper 360, p. 273-290.
- Blodgett, R.B., Santucci, V.L., Baranov, V.V., and Hodges, M.S., 2021, The gypidulid brachiopod genus *Carinagypa* in late Emsian (latest Early Devonian) strata of the Shellabarger Pass area (Farewell terrane), Denali Park & Preserve, south-central Alaska: New Mexico Museum of Natural History and Science, Bulletin 82, p. 19-28.
- Bundtzen, T.K., and Gilbert, W., 1983, Outline of geology and mineral resources of the upper Kuskokwim region, Alaska: Journal of the Alaska Geological Society, v. 3, p. 101-119.
- Churkin, M., Jr., Wallace, W.C., Bundtzen, T.K., and Gilbert, W.G., 1984, Nixon Fork-Dillinger terranes: A dismembered Paleozoic craton margin in Alaska displaced from Yukon Territory: Geological Society of America Abstracts with Programs, v. 16, no. 5, p. 275.
- Decker, J., Bergman, S.C., Blodgett, R.B., Box, S.E., Bundtzen, T.K., Clough, J.G., Coonrad, W.L., Gilbert, W.G., Miller, M.L., Murphy, J.M., Robinson, M.S., and Wallace, W.K., 1994, Geology of southwestern Alaska; in Plafker, G., and Berg, H.C., eds., The Geology of Alaska. The Geology of North America, volume G-1. Geological Society of America, Boulder, Colorado, p. 285-310.
- Dybowski, W.N., 1873, Monographie de Zoantharia Sclerodermata Rugosa aus der Silurformation Estlands, Nord-Livlands und der Insel Gotland. Archiv für die Naturkunde. Liv-, Est- und Kurlands. v. 5, p. 257-414, pl. 1-5.
- Garcia-Alcalde, J., and Blodgett, R.B., 2001, New Lower Devonian (Upper Emsian) *Myriospirifer* (Brachiopoda, Eospiriferinae) species from Alaska and northern Spain and the paleogeographic distribution of the genus *Myriospirifer*; in Frýda, J., Blodgett, R.B., and Mergl, M., eds., Havlíček Volume, Journal of the Czech Geological Society, v. 46, no. 3/4, p. 145-154.
- Gilbert, W.G., and Bundtzen, T.K., 1983, Paleozoic stratigraphy of Farewell area, southwest Alaska Range, Alaska [abst.]: Alaska Geological Society Symposium, New Developments in the Paleozoic Geology of Alaska and the Yukon, Anchorage, Anchorage, 1983, Program and Abstracts, p. 10-11.
- Gilbert, W.G., and Bundtzen, T.K., 1984, Stratigraphic relation between Dillinger and Mystic terranes, western Alaska Range, Alaska [abst.]: Geological Society of America Abstracts with Programs, v. 16, no. 5, p. 286.
- Johnson, J.G., and Lane, N.G., 1969, Two new crinoids from central Nevada: Journal of Paleontology, v. 43, no. 1, p. 69-73.
- Jones, D.L., and Silberling, N.J., 1979, Mesozoic stratigraphy—The key to tectonic analysis of southern and central Alaska: U.S. Geological Survey Open-File Report 79-1200, 37 p.
- Jones, D.L., Silberling, N.J., Berg, H.C., and Plafker, George, 1981, Map showing tectonostratigraphic terranes of Alaska: U.S. Geological Survey Open-File Report 81-792, 20 p., 2 sheets, scale 1:2,500,000.
- Milne-Edwards, H., and Haime, J., 1850, A Monograph of the British Fossil Corals: The Palaeontographical Society of London, London, p. i-lxxv + 1-72.
- Milne-Edwards, H., and Haime, J., 1851, Monographie des Polypiers fossiles des terrains paléozoïques: Archives du Muséum d'Histoire Naturelle, v. 5, p. 1-502.
- Oliver, W.A., Merriam, C.W., and Churkin, M., 1975, Ordovician, Silurian, and Devonian corals of Alaska: U.S. Geological Survey Professional Paper 823-B, p. B13-B44.
- Patton, W.W., Jr., 1978, Juxtaposed continental and ocean-island arc terranes in the Medfra quadrangle, west-central Alaska ; in Johnson, K.M., ed., The United States Geological Survey in Alaska: Accomplishment during 1977: U.S. Geological Survey Circular 772-B, p. B38-B39.
- Reed, B.L., and Nelson, S.W., 1977, Geologic map of the Talkeetna Quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map 870-A, 1 sheet, scale 1:250,000.
- Reed, B.L. and Nelson, S.W., 1980, Geologic map of the Talkeetna quadrangle, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map I-1174, 15 p., scale 1:250,000.
- Rigby, J.K., and Blodgett, R.B., 1983, Early Middle Devonian sponges from the McGrath Quadrangle of west-central Alaska: Journal of Paleontology, v. 57, no. 4, p. 773-786.
- Rigby, J.K., Blodgett, R.B., and Anderson, N.K., 2009, Emsian (Late Early Devonian) sponges from west-central and south-central Alaska: Journal of Paleontology, v. 83, no. 2, p. 293-298.
- Rominger, C.L., 1876, Paleontology. Fossil Corals. Geological Survey of Michigan, v. III, pt. II, Lower Peninsula. New York, J. Bien.
- Rohr, D.M., and Blodgett, R.B., 1985, Upper Ordovician Gastropoda from west-central Alaska: Journal of Paleontology, v. 59, no. 3, p. 667-673.
- Wedekind, R., 1923, Die Gliederung des Mitteldevons auf Grund von Korallen. S.B. Ges. Naturw, Marburg (1922), p. 24-35.

