

Nature Bib ID# 46701

Final Report

**FIRE HISTORY DETERMINATION IN THE MIXED CONIFER/ASPEN
COMMUNITY OF BRYCE CANYON NATIONAL PARK**

**Michael J. Jenkins
Department of Forest Resources
Utah State University
Logan, Utah 84322-5215**

Project UW #5/34080

For period ending March 31, 1995

ABSTRACT

A fire history conducted in the mixed conifer community types of Bryce Canyon National Park found the mean fire return interval lengthened from 7.5 to 45 years since 1900. Dendrochronology, species, diameter and age class analysis showed a change in stand composition in favor of white fir and Douglas-fir and a two fold increase in fuel load. Recommendations for prescribed burning and mechanical fuels reductions were made in keeping with National Park Service fire management planning.

A comparison of today's landscape at Bryce Canyon National Park (BCNP) with the landscape shown in historic photographs indicates that a major change has occurred in the Park's vegetative mosaic (Roberts et al. 1993). The density of woody vegetation and down and dead woody fuels appears to have increased significantly in the present century. Other studies have concluded that these effects occurred in part from changes in the frequency and intensity of wildfires (Fisher et al. 1986; Wadleigh and Jenkins 1994). In these studies results indicated that frequent, low intensity surface fires were much more common in presettlement times. It was concluded that grazing and fire suppression practices begun in about 1900 had a significant impact on the vegetational changes observed. In such cases the abundance of woody vegetation generally increases over the short term, but when fires finally do occur they may be of greater intensity (Kilgore and Taylor 1979). Prescribed fires can be used as a means of reducing hazardous fuel loads and reestablishing presettlement fire regimes.

Techniques analyzing tree rings are one way of constructing fire histories for areas supporting trees of sufficient age (Arno and Sneek 1977). Other methods including vegetative mapping, species, size and age class distributions and fuel load inventories are also useful in preparing site specific fire histories. These data are necessary to determine the extent to which present day fires deviate from presettlement patterns regarding size, intensity and location.

Previous fire history studies in BCNP have documented fire return intervals using fire scar analysis of ponderosa pine (*Pinus ponderosa*) in the park (Bucannan and Tolman 1983; Wight 1989) and for the Paunsaugunt Plateau (Stein 1988). Numerous other studies have similarly documented the fire regime in pre-European settlement ponderosa pine forests in

western North America. For reviews of these previous studies see Wright and Bailey (1982) and Agee (1993).

This study was specifically document vegetative changes resulting from a lengthening the fire return interval by suppression of frequent low intensity surface fires and grazing.

STUDY AREA

BCNP occupies approximately 14,250 hectares on the east face of the Paunsaugunt Plateau in south central Utah. This study was conducted in the mesic mixed conifer/aspen communities located in the south and west portions of the park (Figure 1).

METHODS

Three areas in the mixed conifer/aspen community type were selected for this study based on the description by Roberts et al. (1992). These were Whiteman Bench, Bridge Hollow and Yovimpa Pass. Using maps and aerial photos a transect was established to represent the variety of elevation, aspect and other features of each area. Ten .01 hectare plots were selected on each of the three transects for sampling.

All trees in each plot were tallied by species and diameter was measured with a diameter tape or ruler for small trees. A subsample from each diameter class was collected using an increment core for larger trees and by cutting a cross section of trees less than 4 inches in diameter. While traversing the transects any trees possessing visible fire scars were marked and later sampled using the chain saw wedge technique described by Arno and Sneek (1977). Each sample tree was permanently labeled with a metal tag for maintenance in BCNP records. All

samples were labeled and taken to the fire ecology laboratory at Utah State University for analysis.

RESULTS AND DISCUSSION

The analysis of the data yielded the results shown in Figures 2-4. The graphs compare numbers of trees by diameter class, species and location. Superimposed is the average age of each species in each diameter class. The results show that large numbers of the shade tolerant white fir (*Abies concolor*) and Douglas-fir (*Pseudotsuga menziesii*) have become established in the last 100 years. The majority of older, larger trees are ponderosa pine which were the dominant species historically and are now being replaced.

Table 1 shows the mean presettlement fire return interval found from fire scar specimens collected. For purposes of this study, 1900 was considered as the end of the presettlement era. The return interval was constructed by counting the number of fires documented by scars and dividing the number into the total number of years from the oldest fire document to 1950 (Agee 1993). This value represents the number of years between fires sufficiently intense to scar a tree in the study area. Clearly each fire varied in its ecological significance depending on its size and intensity. Conversely there were surely other fires during the time period that did not scar trees or where fire scars were not found in the present study. The main purpose of presenting these data is to illustrate the frequency with which ignitions occurred even in the mesic mixed conifer type. The return interval can also be evaluated in comparison to the fact that only 2 fires were documented by fire scars since 1900 representing a mean fire return interval of about 45 years. These new data support previous fire history studies by Buchannan and Tolman (1983)

who recorded 87 fires in the mixed conifer type from 1616 to 1900 and 6 fires since 1900. Their data suggested that a fire sufficient to scar a tree occurred in the mixed conifer type once every 3-5 years in the period before 1900. Since that time the interval has been lengthened to 16 years with one fire in 1959 being the only recorded since 1928. Stein's (1988) fire history of the mixed conifer forests of the Paunsaugunt Plateau on the Powell Ranger District of the Dixie National Forest, found a fire return interval of 15.2-18.4 years. The last recorded fires in the study occurred in 1892, 1902 and 1911.

The abundance of white fir and Douglas-fir could not have occurred over such a wide area under a pattern of frequent low intensity surface fires. There were areas where these species became established and survived to produce bark thick enough to withstand surface fires. This is indicated in Figures 2-4 by the presence of older individuals of white fir and Douglas-fir. It is the abundance of small, young trees of these species present that reflect the consequences of an altered fire regime. Ponderosa pine is the most fire tolerant species in the mixed conifer type by producing a 0.12-0.25 inch thick dead bark layer in trees as small as 2 inches in diameter. White fir bark on the other hand remains thin, green and photosynthetically active and therefore very fire susceptible in trees up to 4 inches in diameter. White fir bark is so sensitive to heat that mortality can occur from sunscald after an overstory removal. Even low intensity surface fires will kill white fir and Douglas-fir seedlings and saplings (Hall 1976). In addition the low branching habit of white fir when compared to ponderosa pine increases the potential for a surface fire to move into the crown and more likely kill the tree. Douglas-fir is intermediate to white fir and ponderosa in fire tolerance.

Comparison of these data from Hall (1976) with those presented in Figures 2-4 shows that 4 inch diameter white fir are on average 75 years old and Douglas-fir over 50 years old. By the time these species reach 6 inches in diameter they are nearly 100 years old. Under the presettlement fire return interval of 7.5 years it is clear that a majority of white fir and Douglas-fir seedlings and saplings would have been thinned out due to fire sensitive bark. The decrease in fire frequency since 1900 reported by Buchannan and Tolman (1983) and further documented in this study support the notion of an altered stand composition in favor of white fir and Douglas-fir. These data support the conclusions of Roberts et al. (1992) that reduction in surface fire has altered community type structure reducing the area occupied by quaking aspen and ponderosa pine. Specifically this study demonstrated an increase in trees/acre and density and an overall change in ecosystem diversity.

The effect that an altered fire regime has had from a fuels perspective is more important in terms of management implications. BCNP has an explicit objective of restoration of the park ecosystem to a condition typical of pre-European settlement and prior to the establishment of fire exclusion policies (Bryant 1994). The preferred alternative for vegetative restoration is to use scheduled prescribed fires combined with mechanical fuel pretreatments. This is an appropriate decision since the altered fire regime has significantly impacted the fuels complex. From a fire behavior perspective the presettlement fuels complex is best described by Fire Behavior Fuel Model 2 represented by a timber overstory with a grass or herbaceous surface stratum (Anderson 1982). With increasing density and mortality associated with older trees the fuel model tends to shift toward Fuel Model 8 or 10 represented by closed timber litter and understory. Based on fuel loading alone a three fold increase in tons/acre may occur (Anderson 1982). Roberts

et al. (1993) found 14 tons/acre in the ponderosa pine community (Fuel Model 2) versus 31 tons/acre in the mixed conifer/aspen community type (Fuel Model 10). In addition the horizontal and vertical continuity increase from model 2 to model 10 increasing the potential for torching and crowning. Clearly fire behavior under the present fuels regime will be quite different than what might have been expected prior to fire exclusion. Roberts et al. (1993) concluded that the presettlement fuels complex was better represented by Fuel Model 2 due to more frequent fires. Fire suppression of all wildfires is the policy presently in place at BCNP and will result in increasingly hazardous fuels complexes and more damaging fires in the future. Recent wildfires in the western United States continue to demonstrate the relationship between hazardous fuel complexes and costly and dangerous suppression efforts.

The only solution for BCNP and many other NPS and National Forest Lands is to enter into an aggressive, proactive effort to reduce hazardous fuels by mechanical means and application of prescribed fire using scheduled ignitions. The same resources of equipment, person-power and dollars need to be made available to accomplish the task as are made available to fight wildfires. BCNP has instituted a prescribed fire program in the ponderosa pine type near the visitor center and staff housing. The fire management plan recognizes the need for prescribed fires in the mixed conifer type to reduce hazardous fuels and restore stands to presettlement condition. Prescribed fires in this type may occur as early as the fall of 1995 (Bryant, personal communication).

This study and many others have demonstrated the negative impacts of fire suppression on vegetation, fuels and other resources including wildlife, soil and aesthetics. Attempts to

reestablish pre settlement fire regimes will be difficult, dangerous and expensive, but not as great as attempts to further exclude fire from fire dependent ecosystems.

Table 1. Fire frequency calculations for fire scarred trees from approximately 1500-1900.

No. of sample trees	No. years	Total fires	Fire freq.
20	399	53	7.5 YEARS




Literature Cited

- Agee, J.K. 1993. Fire ecology of Pacific Northwest forests. Island Press, Washington D.C. 493 pp.
- Anderson, H.E. 1982. Aids to determining fuel models for estimating fire behavior. USDA Forest Service, General Technical Report INT-122.
- Arno S.F. and K.M. Sneek. A method for determining fire history in coniferous forests of the Mountain West. U.S. For. Serv. Gen. Tech. Rep. INT-42. 28p.
- Bryant, R. 1994. Draft environmental assessment for fire management plan Bryce Canyon National Park, Utah. U.S. Department of Interior, National Park Service. 28 pp.
- Bucannan, H. and J.M. Tolman. 1983. Fire history of the forest of Bryce Canyon National Park, Utah. Final Report USDI Contract No. PX 1330-0-0161. 40 pp.
- Fisher, R.F., M.J. Jenkins and W.F. Fisher. 1986. Fire and the prairie-forest mosaic of Devils Tower National monument. Amer. Midland Natur. 117(2):250-257.
- Kilgore, B.M. and D. Taylor. 1979. Fire history of a sequoia-mixed conifer forest. Ecology 60(1): 129-142.
- Hall, F.C. 1976. Fire and vegetation in the Blue Mountains--implications of land managers. Pro. Tall Timbers Fire Ecol. conf. 15:155-170.
- Roberts, D.W., D. Wight and G.P. Hallsten. 1992. Plant community distribution and dynamics in Bryce Canyon National Park. Final Report USDI Contract No. PX 1200-7-0966. 203 pp.
- Roberts, D.W., M.J. Jenkins, D.W. Wight. 1993. Historical vegetation, fuel loads, and integrated resource information system for Bryce Canyon National Park. Final Report USDI Contract No. 88-264. 249 pp.
- Stein, S.J. 1988. Fire history of the Paunsaugunt Plateau in southern Utah. Great Basin Naturalist 48 (1):58-63.
- Wadleigh, L.L. and M.J. Jenkins. Fire frequency and the vegetative mosaic of a spruce-fir forest in northern Utah. Great Basin Naturalist, In review.
- Wight, D.W. 1989. Fire history, residential study area, Bryce Canyon National Park, Resource Management Office Report. 33pp.

Wright, H.A. and B.A.W. Bailey. 1982. Fire ecology, United States and Southern Canada.
Wiley Interscience, New York. 501 pp.

FIGURE CAPTIONS

- Figure 1. Map of Bryce Canyon National Park, Utah showing location of mixed conifer community type.
- Figures 2-4. Number, age, and diameter distribution of trees on the Bridge Hollow, Whiteman Bench and Yovimpa Pass transects respectively.



scale (km)

Figure 1

Bridge Hollow Transect




Figure 2

Whiteman Bench Transect




Figure 3

Yovimpa Pass Transect




Figure 4