

132-J, ANIAKCHAK CRATER,
ALASKA PENINSULA, SMITH, W. R.
1925

Please do not destroy or throw away this publication. If you have no further use
for it write to the Geological Survey at Washington and ask for a frank to return it

DEPARTMENT OF THE INTERIOR

Hubert Work, Secretary

U. S. GEOLOGICAL SURVEY

George Otis Smith, Director

Property of the
University of Alaska

Professional Paper 132—J

ANIAKCHAK CRATER, ALASKA PENINSULA

BY

WALTER R. SMITH

Published May 15, 1925

Shorter contributions to general geology, 1923-1924

(Pages 139-145)

Property of the
University of Alaska



WASHINGTON
GOVERNMENT PRINTING OFFICE
1925

18142

ANIACHAK CRATER, ALASKA PENINSULA.

By WALTER R. SMITH.

The discovery of a gigantic crater northwest of Aniakhak Bay (see fig. 11) closes what had been thought to be a wide gap in the extensive series of volcanoes occurring at irregular intervals for nearly 600 miles along the axial line of the Alaska Peninsula and the Aleutian Islands. In this belt there are more active and recently active volcanoes than in all the rest of North America. Exclusive of those on the west side of Cook Inlet, which, however, belong to the same group, this belt contains at least 42 active or well-preserved volcanoes and about half as many mountains suspected or reported to be volcanoes. The locations of some of these mountains and the hot springs on the Alaska Peninsula and the Aleutian Islands are shown on a map prepared by G. A. Waring.¹ Attention has been called to these volcanoes for nearly two centuries, but a record of their activity since the discovery of Alaska is far from being complete, and an adequate description of them as a group has never been written. Owing to their recent activity or unusual scenic beauty, some of the best known of the group are Mounts Katmai, Bogoslof, and Shishaldin, but there are many other beautiful and interesting cones and craters.

Aniakhak Crater (Pls. XLI, XLII) was discovered in August, 1922,² by a United States Geological Survey party, in charge of R. H. Sargent, topographic engineer, in the course of a reconnaissance topographic and geologic survey of the country west of the Aleutian Mountains between Wide and Aniakhak bays. The first evidence of an ancient volcanic eruption in the district was observed 30 miles northeast of the crater, in the form of thin deposits of fine ash concentrated in small depressions on the hill-sides. As the work progressed southwestward

the deposits of volcanic ash became more numerous and the material coarser. Along Ray Creek, 18½ miles from the crater, pieces of scoria and fine ash had consolidated and subsequently been cut by the stream until vertical walls stand 6 to 12 feet high on both banks. (See Pl. XLIII, A, B.) The floor of the broad valley of Cinder River was found to be entirely covered by volcanic ejecta, but the source of the material was not discovered until the summit of Elephant Mountain was occupied by Mr. Sargent and the writer. Prior to the view of the crater, however, locations on the topographic map of a group of peaks in the distance had assumed the form of a circle, but the reason of the arrangement was not suspected.

Aniakhak Crater is 24 miles northwest of Aniakhak Bay, approximately in the central part of the Alaska Peninsula, on the divide between the Pacific Ocean and Bering Sea. Although there are no trails, the district is easily accessible from either side of the peninsula. Aniakhak River, the largest stream on the peninsula flowing into the Pacific Ocean, rises within the crater and breaks through the east side of the rim in a narrow and picturesque castled canyon which has been named "The Gates" (Pl. XLIII, C), from two long, nearly symmetrical mountain spurs that diverge from the canyon, one on each side of the river, and inclose the upper valley. The crater is nearly circular in outline and has a maximum diameter of 6¾ miles and a minimum diameter of a little over 5¾ miles. The lowest part of the crater floor is 1,100 feet above sea level and contains Surprise Lake, a body of water with an area of 2 square miles. The walls are well preserved and rise almost vertically in places to altitudes of 1,200 to 3,000 feet above the bottom of the crater. A large truncated cinder cone, the summit of which is 2,200 feet above Surprise Lake, occupies the south-central part of the area inclosed within the crater rim.

¹ Waring, G. A., Mineral springs of Alaska: U. S. Geol. Survey Water-Supply Paper 418, pl. 1, 1917.

² It has recently been reported that W. W. French, an engineer, and party visited the crater in July, 1921, but no report of the discovery was made at that time.

The crater was first mentioned without a name by R. F. Griggs³ in a book that was printed shortly after the return of Mr. Sargent's party. Under the name "Old Crater"

report⁴ on the geology of the district. In a chronologic account of observed volcanic activity in Alaska, Grewingk⁵ gives the geographic position of Veniaminof volcano as




FIGURE 11.—Sketch map showing location of Aniakhak Crater, Alaska Peninsula.

its position is indicated on the map of Alaska published by the Survey in 1923. A very brief description of the crater was included in a

⁴ Smith, W. R., and Baker, A. A., The Cold Bay-Chignik district, U. S. Geol. Survey Bull. 755, p. 157, 1924.

⁵ Grewingk, Constantin, Beiträge zur Kenntniss der orographischen und geognostischen Beschaffenheit der Nord-west Kuste Amerikas, mit den anliegenden Inseln (with geologic and other maps), St. Petersburg, 1850.

³ The Valley of Ten Thousand Smokes, p. 65, footnote, 1922.

latitude 56° and longitude 158° to 159° . This position is rather close to that of Aniachak Crater, which is latitude $56^{\circ} 45'$ and longitude $158^{\circ} 9'$, but from the short account given it is quite evident that Grewingk meant the volcano known at present as Veniaminof, latitude $56^{\circ} 17'$ and longitude $159^{\circ} 15'$, and not Aniachak Crater. Although the Russian scientists and travelers recorded many interesting and valuable observations regarding the volcanoes of Alaska Peninsula, they made no mention, so far as known to the writer, of the immense crater discovered in 1922. It is quite probable that some of the Russian fur hunters, or perhaps a prospector or trapper of more recent days, came across the crater but either failed to recognize it as such or failed to report it. Inquiries were made of the inhabitants of Chignik, the nearest settlement, and of local trappers, all of whom knew nothing about the crater. The present paper is based on observations made by the writer during a few hours. Time was not available to determine the age of the crater or the exact manner in which it was formed.

The observer at a distance would not suspect the existence of a huge crater within the cluster of rather gentle peaks which are separated from the other mountains of the district by broad level areas. The valleys of Meshik and Aniachak rivers west of the crater are in general less than 200 feet above sea level. The boldest portion of the outer rim of the crater is along the east side of the mountains, where cliffs rise abruptly from the tributary valleys of Aniachak and Meshik rivers. Over these precipitous cliffs many streams cascade from the glaciers near the summit of the rim. One of these streams has a vertical fall of at least 1,000 feet. The west and north sides of the rim slope more gently away from the crater and are covered to a great depth by ash and cinders. Gullies 30 feet deep on these slopes do not expose the solid rock beneath but show rudely stratified layers of ash and large lava and scoria boulders 4 feet and less in diameter.

The panorama of the crater (Pl. XLI) conveys a rather poor conception of its magnitude, owing to the large size of the area inclosed within the proportionally low rim and to the absence of an object of known dimensions. However, the greatest length of Surprise Lake visible in the photograph is $2\frac{1}{2}$ miles, and the

small cones near the upper end of the lake rise about 200 feet above the floor of the crater. The summit of the large cinder cone is $4\frac{1}{2}$ miles from the point at which the picture was taken; the mountains beyond the cone are 6 miles from the observer. The inner wall is abrupt—at places nearly vertical—and so far as known it can not be easily descended except just north of The Gates. A view of Black Nose (Pl. XLIV, A), the peak immediately south of The Gates, taken from a cinder ridge within the crater, gives an idea of the steepness of parts of the inner wall. The south side of the crater rim affords a gathering ground for the snow, and here vigorous alpine glaciers have been formed on the steep slopes. Evidence of slumping along the inner wall was not noted except for several talus slopes in the canyon of Aniachak River and near the base of Black Nose. The circumference of the crater is nearly 19 miles. The rim is broken in two places—the sharp notch through which Aniachak River flows and a depression on the west side which was not seen in 1922 but has been reported by M. W. Taylor, of Seattle, who visited the crater in 1923. On the south side the highest point on the rim is 4,200 feet above sea level and 3,000 feet above Surprise Lake. This peak and several others close by were obscured by clouds at the time the photograph reproduced in Plate XLIV was taken. The greater part of the crest of the rim is not jagged, although Black Nose and several other sharp peaks rise above the average altitude, which is approximately 3,000 feet.

Along the north and northeast sides the inner wall is partly covered by detritus, but in the few exposures seen the rocks appeared to be layers of pink and black lava, probably obsidian, several hundred feet thick, overlying a light-colored quartz diorite. Most of these exposures are inaccessible and were not closely examined, but a specimen of the quartz diorite was taken on the outer wall northeast of the crater. Exposures in the cliffs along the canyon and near the bottom of Black Nose consist of nearly horizontal sedimentary rocks, chiefly very massive gray sandstone. In the lower 500 feet the rocks exposed in the canyon are abundantly fossiliferous; the greater part of the fossils belong to several species of *Aucella* that are characteristic of the Naknek formation, of Upper Jurassic age. From a distance the summit of Black Nose apparently consists of

lava, but the upper portion may be in part sedimentary rock, presumably of Tertiary age. The unconformity and the contrast in lithology between the rocks of the two kinds can be seen in Plate XLIV, A. Although the high mountains forming the south side of the rim were not closely examined, they probably consist entirely of sedimentary rocks, the greater part of which are Upper Jurassic. A rugged spur north of the north wing of The Gates is composed of very coarse agglomerate or tuff similar to that overlying large areas in the vicinity of Aniakhak and Kejulik bays. Most of the large and small boulders that make up the agglomerate are angular, but some are rounded. Individual blocks attain 16 feet in diameter and display the darker shades of red, green, and gray. Many of the boulders are black, and all of them show a scoriaceous texture.

The area of the bottom of the crater is approximately 30 square miles. The entire floor is covered to an unknown depth by black and gray scoria ranging in coarseness from very fine material to pieces several feet in diameter. About one-third of the floor is level; around the base of the large cone ridges of cinders, 200 to 800 feet high, radiate toward the crater wall. The formation of the cones and the position of the material within the rim are undoubtedly the results of activity subsequent to the major eruptions that produced the great crater. The later activity probably decreased the original depth of the crater by partly filling it with cinders. The upper 800 feet of the large cone is remarkably circular and well preserved. It is thought to be formed entirely of cinders, as no lava was observed on the ridges around the base. The summit is truncated and may be slightly depressed. An attempt was made to reach the top, but traveling over the loose material is wearisome, and the attempt was given up late in the evening with the realization that a trip of 10 miles to camp was yet to be made. The small cones near the head of the lake were not examined. Toward the west side of the crater, 5 miles from the writer's nearest point of observation, another small cone, which was not caught by the camera, could be seen by aid of a field glass. Near this cone a curious bowl-shaped deposit of white material, probably formed by hot mineral waters, stands out in contrast to the surrounding black scoria, which appeared to be very

coarse at that locality. Only a rough estimate of the dimensions of the bowl can be given, but it is at least 200 feet in diameter.

Surprise Lake, in the northeastern part of the crater, is irregular in outline and has a maximum length of $2\frac{1}{2}$ miles and average width of three-fourths of a mile. It covers an area of nearly 2 square miles, but the bluish-green color of the water indicates that the depth is not great. The lake may have formerly covered a much larger area, before the river had deepened its channel through the canyon, but terraces or high-water marks could not be detected on the wall at the few places examined. An ill-defined bench occurs well up on the sides of the cinder cone, however, but the bench is not continuous nor sufficiently well preserved to consider it an ancient lake shore. A small circular lake or pool several hundred feet higher than Surprise Lake is situated at the base of Black Nose. Although a stream of considerable size flows from the lake and unites with Aniakhak River just inside The Gates, the lake has no affluents and must derive its supply of water from the snow fields and glaciers by seepage through the cinders.

The narrow notch in the crater's rim through which Aniakhak River flows has undoubtedly been deepened by the erosive power of the stream, which is very turbulent as it emerges through The Gates. The break in the wall, however, is thought to be caused by a rift, which may extend across the crater and account for the depression on the west side. No evidence of relative vertical displacement of the opposite sides of the canyon was noted. The photograph of The Gates (Pl. XLIII, C), taken through the haze from a point $5\frac{1}{2}$ miles down the valley, does not do justice to their profoundly impressive beauty. Several terraces formed by the river in the ash and pumice can be seen in the foreground. The crater can be entered through The Gates without difficulty on the south side of the river. The stream was too swift to be waded within the canyon when it was visited in August, and an attempt to pass through on the north side of the river terminated by climbing a precipitous cliff 1,600 feet high.

Only the east side of the crater was visited, but the country to the north could be seen from Jaw Mountain and the southeast outer

wall from Pinnacle Mountain. All the valleys of the surrounding country for many miles are covered by thick deposits of ejecta from the crater. The material consists of fine ash, fragments of black and pink lava, and gray, black, and red pumice. The greatest observed thickness is exposed in the steep banks of Lava Creek. The stream has developed many beautiful terraces and its present bed, west of Jaw Mountain, is cut, perhaps 50 feet, into a deposit of black volcanic glass, as reported by Mr. Sargent. The glass or obsidian is 10 miles from the nearest point on the crater's rim, and whatever its source it was extruded before the great eruptions of the volcano, as it is overlain by at least 150 feet of loose material, the greater part of which is black pumice. This is the only lava seen in place except on the inner wall of the crater, but many fragments of black obsidian, containing a few glassy lath-shaped crystals of feldspar, were found on the mountains many miles from the crater. Indications of extinct fumaroles were sought in the ash-covered area east of the crater, but much of the volcanic material has been worked over by streams, and no evidence of fumarolic activity was found. The upper valley of Meshik River and also the valley of Aniachak River in the vicinity of Albert Johnson Creek have been filled with volcanic debris to such an extent that the present valley floors are very broad and nearly level. A few low isolated hills near the margins of the level area appear to have been partly buried, with only their summits left exposed, but this is probably an illusion. Several high mountains southeast of the crater are sharp-peaked, especially Pinnacle Mountain, and are composed of hornblende andesite. A stream of considerable size, Hidden Creek, a tributary of Aniachak River, flows through a subterranean channel beneath the ash and cinders for a distance of 5 or 6 miles and emerges on the west bank of the river. The Aniachak is about 100 feet wide and 4 feet deep below the North Fork. The current is swift enough to transport small pieces of pumice along the bottom. At 4 miles from its mouth the river has eroded a picturesque gorge through a thick deposit of volcanic agglomerate. The southwest side of the gorge is formed by a rugged spur that projects across the valley from the hills to the southwest and is locally known as Cape Horn. (See fig. 11.)

The river is navigable by small boats as far as the meanders below the mouth of Mystery Creek.

A broad marshy lowland lies between the shore line of Aniachak Bay and the hills toward the northwest. In this area a remarkable series of nine crescent-shaped ancient beaches can be seen from the mountains north of the river in the form of low ridges conforming in direction to the present shore line. A large part of the material in these ridges consists of volcanic ash and pumice washed ashore by the waves after having been transported into the bay by the river. The bay is uncharted and is not a protected harbor, although deep water is reported by masters of fishing schooners who have entered the bay. Hook Bay, 25 miles directly south of the crater, affords an excellent harbor for boats of moderate size. An old trail leads from this bay over the mountains toward Meshik River valley and is probably the best route from the Pacific coast to the crater, although in fair weather a landing can be made at the mouth of the river at Aniachak Bay. From this point the distance to the crater is not so great as from Hook Bay, and the route is over level country. Chignik is the nearest settlement and is 40 miles southwest in a direct line from the crater but at least 60 miles by trail. A trapper's cabin has been built near the lagoon at Aniachak Bay and one on the south side of the valley below Albert Johnson Creek. The trappers occupying the cabins during the winter had traveled over the cinders as far west as Meshik Lake for many years, but they did not know of the existence of Aniachak Crater.

Vegetation has gained a footing in places throughout the area covered by ash, and even within the crater several tufts of grass and low flowering plants were seen. Large areas, however, within a radius of 20 miles of the crater, are entirely barren of plant life and have the appearance of arid plains. Alder bushes have grown to a height of 6 feet in the valleys of High and Lava creeks and furnish sufficient fuel for camp purposes, but they are not straight enough to be used for tent poles. Foxes and Kodiak bears inhabit the region, Small herds of caribou were seen near the valley of Pumice Creek.

The wide general use of the term "crater" includes several varieties or types, which have

been differentiated as impact craters, subsidence craters, upbuilt craters, and explosion craters. An impact crater may be formed by the fall of a body, such as a large meteorite or a projectile from a cannon. Such a bowl-shaped hollow or pit may have nothing in common with volcanic activity. A subsidence crater is a depression below the general level of the country or a concave area on a mountain, made by the collapse of the roof of a cavity. In limestone countries such depressions are common, but the use of the term is restricted by many geographers to pits occurring in volcanic regions. These pits are usually caused by the collapse of cavities formed in volcanoes by the eruption of lava and other material. Craters formed in this manner are also known as caldrons. The largest and some of the best-known craters on the earth are of this variety, to which belong Kilauea, in the Hawaiian Islands (2.93 miles in diameter); the pit of Crater Lake, Oregon (5½ miles in diameter); Ngorongoro, East Africa (12 miles in diameter); and Aso-san, in Japan (dimensions, 9 by 14 miles). An upbuilt crater has the form of a more or less circular mound surrounding an aperture, relatively small in diameter but usually deep, which serves as a vent for the lava, ash, and fumes thrown out by a volcano. This is the normal and most common type of craters and occurs on the summits of Lassen, Etna, Cotopaxi, Fuji, and scores of other volcanoes. Stromboli, in the Mediterranean Sea, is taken as the type of the explosion craters. Better examples, however, are known, such as the crater formed by the ancient explosion of Vesuvius, A. D. 79, of which Mount Somma is a remnant, and the ruins of Krakatoa in 1883 and of Katmai in 1912. Excavations made by erosive agents have been referred to and classified as craters, but they have nothing to do with volcanic action.

All the evidence now available points to the origin of Aniakhak Crater by explosive activity and not by subsidence. Besides the vast quantities of rather fine material thrown from the crater and concentrated in the valleys of the surrounding country, huge projectiles were hurled many miles. On the north slope of Elephant Mountain, 1,000 feet above the valley of Meloy Creek and 23½ miles from the center of the crater, a block of black obsidian was found which measured at least 2 cubic feet in volume. About 14 miles southeast of the crater an angular mass of sandstone, 5 feet in

diameter and containing fossils similar to those found near Black Nose, occurs near the summit of a small hill composed of agglomerate. The large boulder is not waterworn, and there is slight probability that it was carried to its present position by a glacier. The ash contains smaller fragments of sandstone that have the same lithologic character as that exposed in the southeast wall of the crater and were not derived from rock in the immediate vicinity in which they were found. That tremendous explosive forces were active at the time the present crater was formed is quite evident. All the known craters of the world as large as Aniakhak Crater or larger are thought to have been formed by subsidence. Katmai Crater, 3 miles in diameter, 151 miles to the northeast, was formerly considered the largest crater of the explosive type known, but it is greatly surpassed by Aniakhak Crater, 6¾ miles in diameter. In diameter and general outline the caldron of Crater Lake, Oreg., and Aniakhak Crater are remarkably similar, although the two are of entirely different types.

The form of the mountain mass that occupied the present site of the crater prior to the eruptions that resulted in the formation of the depression can of course only be conjectured. Before the existence of the volcano the area now within the crater was probably occupied by one or more mountains of moderate altitude—that is, by analogy with other mountains in the district, between 3,000 and 4,000 feet—which were composed of very slightly folded sedimentary rocks, intruded on the north by a large mass of quartz diorite. Why volcanic vents should be formed near the summits of high mountains composed of nearly horizontal strata is difficult to explain, but on the Alaska Peninsula many of the active and extinct volcanoes originated in mountains of this type. This is especially true of the volcanoes toward the northwest, of which Mounts Katmai, Magick, Peulik, and Chiginagak are examples. These volcanoes and many others on the Alaska Peninsula have formed cones and increased their altitudes to heights ranging from 5,000 to nearly 10,000 feet by successive lava flows over the sedimentary rocks. Mount Chiginagak (Pl. XLIV, B), an active volcano 43 miles northwest of Aniakhak Crater, is an impressive and typical example of an upbuilt cone. That a similar cone once towered above the site of Aniakhak Crater is suggested by beheaded lava flows,

remnants of which are exposed on the east wall of the crater, on the summit of Black Nose, and possibly in the bed of Lava Creek. An imaginary cone reconstructed a mile in height above the present floor of the crater would be slightly over 6,000 feet above sea level, an altitude which is very low in proportion to the diameter of the base and also relatively low if compared with the altitude of other volcanic cones. But even if a low cone is postulated the amount of material to be accounted for is enormous and the force necessary to remove it in one or several explosions is beyond imagination. The capacity of the crater alone, if 1,600 feet is taken as the average height of the rim, is about 9 cubic miles. Subtracting 1 cubic mile for the large cinder cone still leaves space for 8 cubic miles within the crater. To this must be added at least 7 cubic miles if the low cone, suggested above, ever existed. Between 300 and 400 square miles of adjacent country on the east and south is known to be covered to various depths by volcanic ash and cinders, and if an equally large area north and west of the crater is also overlain by ejecta, certainly an appreciable part of the supposed total of 15 cubic miles is still to be seen.

Ordinarily the Bering Sea coastal plain on the northwest side of the peninsula is dotted with thousands of small lakes, but in the area north of the crater no lakes could be seen from Jaw Mountain. That all depressions in this part of the plain were filled by material from the crater is a fair deduction. The volume of material ejected from Katmai is estimated at 4.75 cubic miles, but Katmai Crater has a capacity of only 2 cubic miles. The ejection of vast quantities of material from a single source and perhaps during a single period of eruption is not unheard of in volcanism. Capps⁶ has estimated a volume of 10 cubic miles of volcanic ash thrown from a vent in the upper White River basin, and other large estimates have been made. Most of these computations, however, were made from measurements of the material ejected and not from the capacity of the crater.

Few data are now available for estimating the age of Aniakchak Crater. However, it is believed to be one of the oldest of the series of volcanoes along the Aleutian Range. That

the eruption antedates historic record is suggested by the amount of stream erosion, especially in Lava Creek, and the vegetation of the area, including alders. If the absence of evidence of glaciation since the eruption is taken as a criterion it would appear that the eruption is of post-Pleistocene age. Although the ash is more or less concentrated in the valleys it is not in the form of glacial moraines, and unless an ice sheet in which there was little or no movement covered the entire area, moraines of the loose pyroclastic material would be expected had this material been deposited before glaciation. A few of the valleys, such as that of Ray Creek, in which thick deposits of ash occur, are rather narrow and probably have been formed almost entirely by stream erosion. Most of the valleys are broad and suggest extensive glaciation before they were partly filled by ash. A detailed study is necessary for a more definite age determination.

An examination of the relief map (Pl. XLII) will perhaps convey a better idea of the giant crater than a description. Nearly all distances were scaled from Mr. Sargent's skillfully drawn topographic map, on which the relief map by Mr. Renshaw is based. The topographic map is published separately as the Cold Bay-Chignik map.

The writer wishes to acknowledge his indebtedness to Mr. Sargent for cooperation during the field work and for valuable assistance and criticism in the preparation of this paper. Thanks are also expressed to Mr. Sidney Old, who accompanied the writer into the crater and led the way across Aniakchak River.

The scenery of the Aniakchak district, with its broad prairie-like valleys, margined by majestic, sharp-peaked mountains, is impressive and in many respects unique. The once active scene of terrific earth convulsions is now almost oppressively silent. The coloration of the country is somber and together with the fretfully driven clouds tends to create a rather pleasing weirdness. Aniakchak Crater is one of the great natural curiosities of North America and is certainly worthy of further investigation. Although small in comparison, it is probably the nearest counterpart on earth to the craters of the moon in regard to the manner of its formation. Were it not so remote from the usual paths of travel the setting apart of this crater as a national monument would be justified.

⁶ Capps, S. R., An ancient volcanic eruption in the upper Yukon basin: U. S. Geol. Survey Prof. Paper 95, p. 62, 1915.

DEPARTMENT OF THE INTERIOR
Hubert Work, Secretary

U. S. GEOLOGICAL SURVEY
George Otis Smith, Director

Professional Paper 132

SHORTER CONTRIBUTIONS TO GENERAL GEOLOGY

1923-1924

W. C. MENDENHALL, CHIEF GEOLOGIST




WASHINGTON
GOVERNMENT PRINTING OFFICE
1925

ILLUSTRATIONS.

PLATE XLI. Aniakhak Crater, Alaska Peninsula.....	Page. 140
XLII. Relief map of Aniakhak Crater and vicinity, Alaska Peninsula.....	140
XLIII. A, B, Exposures of volcanic ash on Ray Creek, Alaska Peninsula; C, "The Gates" of Aniakhak Crater, Alaska Peninsula.....	140
XLIV. A, Black Nose, Alaska Peninsula; B, Mount Chiginagak, Alaska Peninsula.....	140
FIGURE 11. Sketch map showing location of Aniakhak Crater, Alaska Peninsula.....	140

U. S. GEOLOGICAL SURVEY


PROFESSIONAL PAPER, 132 PLATE XII




ANIACHAK CRATER, ALASKA PENINSULA



RELIEF MAP OF ANIAKCHAK CRATER AND VICINITY, ALASKA PENINSULA



A




B


A, B. EXPOSURES OF VOLCANIC ASH ON RAY CREEK, ALASKA PENINSULA



C. "THE GATES" OF ANIARCHAK CRATER, ALASKA PENINSULA



A. BLACK NOSE, ALASKA PENINSULA



B. MOUNT CHIGINAGAK, ALASKA PENINSULA

	Page.
PLATE X. A, View of Waterpocket Fold and Cretaceous plateaus between the Circle Cliffs and the Henry Mountains, Utah; B, Sketch of the country between the Circle Cliffs and the Henry Mountains, Utah, showing the surface distribution of the rock formations.....	13
XI-XXI. Fossils of the Colorado group of southern Montana.....	33
XXII. The curve of atomic volumes.....	58
XXIII. Fossil plants from the early Eocene at Sayersville, Tex.....	92
XXIV. Geologic map showing the distribution of the Wasatch and Green River formations in northwestern Colorado and southern Wyoming.....	94
XXV. A, Conglomerate in the lower part of the Wasatch formation at Sugarloaf Butte, Colo.; B, Oil shale in the Green River formation, showing characteristic thin, regular bedding; C, Papery oil shale, showing flexibility.....	100
XXVI. Fossils of the Idaho formation.....	112
XXVII. Microphotographs of <i>Elacophyton</i>	138
XXVIII. Microphotographs of <i>Elacophyton</i>	138
XXIX. Microphotographs of <i>Elacophyton</i>	138
XXX. Microphotographs of <i>Elacophyton</i>	138
XXXI. Microphotographs of <i>Elacophyton</i>	138
XXXII. Microphotographs of Australian boghead (kerosene shale).....	138
XXXIII. Microphotographs of Australian boghead (kerosene shale).....	138
XXXIV. A, Microphotograph of Australian boghead (kerosene shale); B, Microsection of torbanite from Scotland.....	138
XXXV. Microsections of torbanite from Scotland.....	138
XXXVI. Microsections of torbanite from Scotland.....	138
XXXVII. Microsections of torbanite from Scotland.....	138
XXXVIII. A, Microsection of torbanite from Scotland; B, Microsection of boghead from Alaska.....	138
XXXIX. A, Microsection of boghead from Alaska; B, Microsection of cannel coal from Pennsylvania.....	138
XL. Microsections of cannel coal from Pennsylvania.....	140
XLI. Aniakhak Crater, Alaska Peninsula.....	140
XLII. Relief map of Aniakhak Crater and vicinity, Alaska Peninsula.....	140
XLIII. A, B, Exposures of volcanic ash on Ray Creek, Alaska Peninsula; C, "The Gates" of Aniakhak Crater, Alaska Peninsula.....	140
XLIV. A, Black Nose, Alaska Peninsula; B, Mount Chiginagak, Alaska Peninsula.....	140
FIGURE 1. Index map of parts of Utah and Arizona showing location of columnar sections given in Plates I and II.....	6
2. Sketch map of Green River between Green River, Wyo., and Green River, Utah.....	36
3. Map showing Green River between the mouth of Henrys Fork and the mouth of Skull Canyon, Daggett County, Utah.....	37
4. Section of the north slope of the Boars Tusk ridge, Daggett County, Utah.....	38
5. Cross section of the ridge east of the Boars Tusk ridge, Daggett County, Utah.....	39
6. Diagrammatic sketch of the bluff northwest of Neilsons Flat, Daggett County, Utah.....	40
7. Diagram showing correlation of sections near Flaming Gorge, Daggett County, and in Island Park and near the Carnegie Museum dinosaur quarry, Uinta County, Utah.....	45
8. Sketch map showing areal relations of the Wilcox floras.....	90
9. Index map showing location of area in northwestern Colorado and southern Wyoming.....	94
10. Sections from northwest to southeast in the Green River Basin, Wyo., showing interfingering of the Wasatch and Green River formations.....	99
11. Sketch map showing location of Aniakhak Crater, Alaska Peninsula.....	140



CONTENTS

The letters in parentheses preceding the titles are those used to designate the papers for advance publication.]

	Page
(A) Rock formations in the Colorado Plateau of southeastern Utah and northern Arizona, by C. R. Longwell, H. D. Miser, R. C. Moore, Kirk Bryan, and Sidney Paige (published July 27, 1923).....	1
(B) A new fauna from the Colorado group of southern Montana, by J. B. Reeside, jr. (published November 5, 1923).....	25
(C) Notes on the geology of Green River valley between Green River, Wyo., and Green River, Utah, by J. B. Reeside, jr. (published November 30, 1923).....	35
(D) The evolution and disintegration of matter, by F. W. Clarke (published April 9, 1924).....	51
(E) An early Eocene florule from central Texas, by E. W. Berry (published April 10, 1924).....	87
(F) Relations of the Wasatch and Green River formations in northwestern Colorado and southern Wyoming, with notes on oil shale in the Green River formation, by J. D. Sears and W. H. Bradley (published November 6, 1924).....	93
(G) Discovery of a Balkan fresh-water fauna in the Idaho formation of Snake River valley, Idaho, by W. H. Dall (published November 10, 1924).....	109
(H) The resuscitation of the term Bryn Mawr gravel, by F. Bascom (published November 12, 1924).....	117
(I) Origin of the boghead coals, by Reinhardt Thiessen (published May 14, 1925).....	121
(J) Aniakchak Crater, Alaska Peninsula, by W. R. Smith (published May 15, 1925).....	139
Index.....	147

ILLUSTRATIONS

	Page
PLATE I. Generalized columnar sections of the Paleozoic and Mesozoic rocks of northern Arizona and southeastern Utah.....	6
II. Generalized columnar sections of the Paleozoic and Mesozoic rocks of southeastern Utah.....	6
III. A, View looking down San Juan Canyon, Utah, at a point 9 miles by stream above Honaker trail; B, View looking down Green River toward its junction with Colorado River, Utah; C, View in Cataract Canyon of the Colorado, Utah.....	6
IV. A, View looking down San Juan Canyon, Utah, at a point between Clay Hill Crossing and the mouth of Moonlight Creek; B, View near Colorado River 2 miles below mouth of Trachyte Creek, Utah; C, View looking down San Juan Canyon at the mouth of Slickhorn Gulch, Utah.....	6
V. A, View looking southwest down head of Marble Canyon from point near Lees Ferry, Ariz.; B, View looking up Colorado River from a high point near the mouth of Crescent Creek, Utah; C, Erosion remnants standing on platform of Coconino sandstone near mouth of Fremont River, Utah.....	6
VI. A, View looking up San Juan Canyon from point near mouth of Clay Gulch, Utah; B, View looking west toward high mesa at Clay Hill Crossing on San Juan River, Utah.....	7
VII. A, Castle Butte, near mouth of Red Canyon, Utah; B, View looking north across Colorado River at Lees Ferry, Ariz.....	12
VIII. A, View looking northwest up Colorado River toward proposed dam site near Lees Ferry, Ariz.; B, View from point half a mile south of Spencer Camp looking south down San Juan Canyon toward Navajo Mountain, Utah.....	12
IX. A, View looking northeast toward Wilson Mesa from point on Wilson Creek near San Juan Canyon, Utah; B, View looking across Colorado River opposite Smiths Fork, Utah; C, Sheer cliff of Navajo sandstone at Warm Spring Creek on Colorado River, Utah.....	12

INDEX.

A

Page.

Acknowledgments for aid.....	94, 135
Adams, L. H., and Williamson, E. D., determinations of compressibility of rocks by.....	79
Anelylus undulatus Meek.....	112
Aniakchak Crater, Alaska Peninsula, age of.....	145
description of.....	139-145
discovery of.....	139
origin of.....	139
plate showing.....	144
volcanic deposits near.....	140
and vicinity, relief map of.....	139, 143
Aniakchak River, Alaska Peninsula, features of.....	142, 143
Animal substances, complexity of.....	68
Aphanotylus Brusina.....	115
white Dall, n. sp.....	115, Pl. XXVI
Arizona, northern. See Colorado Plateau.	
Asplenium ? primo Knowlton.....	91, Pl. XXXIII
Aston, F. W., "mass spectra" of.....	60
work of, on isotopes.....	83-84
Atomic numbers, features of.....	59
importance of.....	59
table showing.....	56
Atomic volumes, curve of, described.....	57-58
curve of, plate showing.....	58
Atomic weights, divergences of, from whole numbers.....	60
numerical relations of.....	59
three standards for.....	55
usefulness of.....	64-65
Atoms, complexity of.....	59

B

Bascom, F., The resuscitation of the term Bryn Mawr gravel.....	117-119
Berry, E. W., An early Eocene florule from central Texas.....	87-92
Berzelius, system of, for atomic weights.....	55
"Black layer," Shinarump conglomerate so called.....	11
Black Nose, Alaska Peninsula, features of.....	141-142
plate showing.....	140
Blue Gate sandstone, features of.....	15-16
"Blue Gate shale," features of.....	15
Bears Tusk Ridge, Utah, rocks exposed in.....	37-39
Boghead coals, algal theory of origin of.....	121-126, 130-131
discovery of, in Australia.....	123
in Scotland.....	122
from Alaska, description of.....	132-133
microsections of.....	138
from Australia, description of.....	131-132
photomicrographs of.....	138
from Bathgate, Scotland, description of.....	132
from Pennsylvania, description of.....	133
from Torbane Hill, Scotland, description of.....	132
microsections of.....	138
nature and origin of.....	123-126
organism supplying carbonaceous matter to.....	126-127
origin of, summary on.....	134-135
other names for.....	122
use of term.....	122
yellow bodies in.....	122, 124
source of.....	130, 131
Boodle, L. A., cited.....	128-129
Bradley, W. H. See Sears and Bradley,	
Brandywine formation, history of the term.....	117
Browns Park, Utah-Colo., rocks exposed in and near.....	41
Bryn Mawr gravel, history of the term.....	118
resuscitation of the term proposed.....	117-119

C

Campbell, W. W., cited.....	53
Cannel coal from Pennsylvania and Kentucky, description of.....	133-134
microsections of.....	138

Page.

Cannizzaro, scheme of atomic weights revised by.....	50
Carbon atom, tetrahedral, theory of.....	65
Carbon compounds, possible similarity of elements to.....	64
Carinifex, basis for name.....	112
(Vorticifex) tryoni Meek.....	111
Carnegie Museum dinosaur quarry, above Jensen, Utah, section at.....	44
Caspi Dybowskii.....	114
Castle Butte, Utah, plate showing.....	12
Cerium, plural valency of.....	58
Chisle formation, nature of.....	1
Chlorine, possible isotopes of.....	84, 85
Chlorophyll, production of.....	68
Chollat, Paul, cited.....	28
Circle Cliffs, Utah, sections in.....	20-21, 22-23
Clarke, Frank Wigglesworth, The evolution and disintegration of matter.....	51-86
Coal, materials of.....	121
Cocconino sandstone, nature and occurrence of.....	8
Colorado, northwestern, sections of Green River formation in.....	101-107
topography of.....	94
Colorado group of southern Montana, new fauna from.....	25-33
Colorado Plateau, Cretaceous rocks of.....	14-16
erosion interval in.....	9
Jurassic rocks in, terminology of.....	11-12
previous work on.....	1-2
sandstones and shales, unnamed, in.....	14
sections of rocks in.....	16-23
plates showing.....	6
stratigraphy of.....	2-5, 11-12, 14-15, 16
Tertiary rocks of.....	16
topography of.....	2
Colorado River, plates showing.....	6, 12
Compounds, general nature of.....	65
solid, first formed in a cooling celestial mass.....	66
syntheses of, in the laboratory.....	69
Compression within the earth.....	78
Corongite, chemical properties of.....	129-130
formation of.....	126-127
occurrence and physical properties of.....	127-128
Cratium, existence and possible relations of.....	53, 61
Craters, types of.....	143-144
Crescent Creek, Utah, sections on Colorado River near.....	18-19
Crow Indian Reservation, Mont., section in.....	25-26

D

Dakota(?) sandstone, features of.....	15
Dall, W. H., Discovery of a Balkan fresh-water fauna in the Idaho formation of Snake River valley, Idaho.....	109-115
De Hautplek, E., cited.....	128
Desolation Canyon, Utah, rocks exposed.....	45-47
Devolution, evidence obtainable from.....	64
Digestion of food, breaking down and regeneration of compounds in.....	69-70

E

Earle, Tex., fossil plants found at.....	87
fossil plants found at, age of.....	88-89
distribution of.....	90
Earth, average composition of.....	76-77
average composition of known materials of.....	71
chemical development of the crust of.....	60-67
compounds formed in the cooling of.....	66
compression within.....	78
diameter of the nucleus of.....	77-78
distribution of matter in.....	66, 77
temperatures in the interior of.....	78-79
volume of atmosphere, ocean, and crust of.....	70
Echo Park and Mitten Park, Colo., rocks exposed in.....	42

	Page.		Page.
Elaeophyton coorongiana, material of boghead coals derived from.....	130-131	Island Park, Utah, rocks exposed in.....	42-43, 45
coorongiana, name proposed.....	126	Isotopes, production and nature of.....	83-84
photomicrographs of.....	138	of lead, features of.....	82-83
Electronic structure of atoms, theory of.....	59		K
Electrons, planetary, nature of.....	59	Kalbar limestone, occurrence and thickness of.....	8-9
Elements, chemical, decay of.....	51-52	Kaiparowits Plateau, Utah, section in front of.....	21
chemical, dissociated condition of, in the hotter stars postulated by Lockyer.....	51	Kerogen, nature and variations of.....	100-101
evolution of, suggested by Clarke.....	51	Kerosene, introduction of.....	122-123
Rutherford's scheme for.....	59	Kerosene shale. See Boghead coals.	
normal and abnormal, differences between.....	80, 82	Kingfisher Canyon and Flaming Gorge, Utah, rocks exposed in and between.....	39-40, 45
order of development of.....	83	Knowlton, F. H., cited.....	91
perfect and imperfect.....	60-61		L
speculations regarding a common origin of.....	51	Latn dalli White.....	112
Emery, W. B., naming of Jurassic formations by.....	12	Lead, uncertain atomic weight of.....	83
Erosion, remnants of, near Fremont River, Utah, plate showing.....	6	Lees Ferry, Ariz., section near.....	16
Evolution of substances, increasing complexity and instability produced by.....	52	Lewis, H. C., cited.....	118
	F	Lithasia antiqua Gabb.....	114
Farrington, O. C., on the average composition of the Earth.....	76	campbelli Dall, n. sp.....	115, Pl. XXVI
on the origin of meteorites.....	74	Lithoglyphus antiquus (Gabb).....	114-115, Pl. XXVI
Field work on the Colorado Plateau, record of.....	1	Lithosphere, approximate thickness of.....	77-78
Flaming Gorge and Kingfisher Canyon, Utah, rocks exposed in and between.....	39-40, 45	average composition of.....	75
Fremont River, Utah, section on Colorado River near.....	17-18	Lodore Canyon, Colo., rocks exposed in.....	41-42
	G	Longwell, C. R., and others, Rock formations in the Colorado Plateau of southeastern Utah and northern Arizona.....	1-23
Girty, G. H., cited.....	7	Lynnaea? sp.....	111
Goniobasis taylori (Gabb).....	114, Pl. XXVI		M
taylori (Gabb) var. calkinsi.....	114, Pl. XXVI	McElmo formation, features of, in the Colorado Plateau.....	14
Goodridge formation, nature and age of.....	7	"Mass spectra," process for producing.....	83
Gray Canyon, Utah, rocks exposed in.....	47	work of Aston on.....	60
Green River, plate showing.....	6	Masuk sandstone, features of.....	16
Green River, Wyo., rocks along Green River near.....	35	"Masuk shale," features of.....	16
Green River formation in Colorado and Wyoming, examinations of.....	93	Megasytrophia binneyi Meek.....	112
history of.....	99-100	Melania taylori Gabb.....	114
kerogen in oil shale of.....	100-101	Mendeleef, existence and properties of undiscovered elements deduced by.....	51, 57
map showing distribution of.....	94	periodic law discovered by.....	56
nomenclature of.....	98-99	Mercury, isotopic varieties of.....	84-85
oil shale in, chemical properties of.....	100-101	Merriam, John C., cited.....	109
distillation yields of.....	102-107	Merrill, G. P., average composition of meteoric stones computed by.....	71-72
physical properties of.....	100	Mespidolaphne precoushatta Berry, n. sp.....	91, Pl. XXIII
plates showing.....	100	Meteorite iron, composition of.....	72-74
sections of.....	101-107	former molten state of.....	73-74
source of material of.....	96-97	Meteorites, origin of.....	74
stratigraphic features of.....	95	stony, composition of.....	72
structure of.....	95	Meteors, disruption of.....	74-75
variations in material of.....	97-98, 101	Meyer, Lothar, curve of atomic volumes devised by.....	57-58
Green River valley, Wyo.-Utah, lithologic succession in, compared with those of southwestern Utah and central Wyoming.....	47-50	Micromelania Brusina.....	114
Gregory, H. E., naming of Jurassic formations by.....	11-12	Moenkopi formation, nature and divisions of.....	9-10
	H	Molecules, polyatomic, formation of.....	54
Halls Creek, Utah, sections on.....	21-22	Montana, fauna from Colorado group of.....	25-33
Halogens, relations of, to inert gases and alkaline metals.....	58	Moon, mean density of.....	75
Hancock, J. Gething, cited.....	128	Moseley, H. G. J., law of relation of the elements proposed by.....	59-60
Harkins, W. D., scheme of, for evolution of the elements.....	61	Mount Chignagak, Alaska Peninsula, origin of.....	144
Heat, generation of, in the earth.....	78-79	plate showing.....	140
produced by union of atoms.....	54		N
Helicoceras parietense White?.....	31, Pl. XX	Navajo sandstone, deposition and thickness of.....	13-14
Helium, elemental nature of.....	61	rock carving in.....	13
presence of, in nebulae and cooling stars.....	53, 54	sheer cliff of, plate showing.....	12
Henry's Fork, Utah, rocks along Green River near.....	36-37	Nebulas, development of, into stars and planets.....	52-54
Hydrocarbons, origin of.....	67	Nebulium, disappearance of, from cooling stars.....	51
presence of, in meteoric stones.....	72	existence and properties of.....	53
Hydrogen, liberation of, from elements of higher atomic weight.....	85-86	possible relations of.....	58, 61, 62
	I	Nematurella Sandberger.....	114
Inoceramus labiatus (Schlotheim).....	31, Pl. XIV	N'hangellite, microscopic features of.....	128-129
sp.....	31, Pl. XIV	occurrence of.....	128
Ionium, admixture of, with thorium, effect of.....	82		O
Iron, meteoric, composition of.....	72-74	Ocean, a possible source of the chlorine in.....	73
		Oil, occurrence of, in the Colorado Plateau.....	2, 7
		Oil shales. See Boghead coals and Green River formation.....	000
		Optum. See Poppy.	
		Organic compounds, complexity of.....	67-68

	Page.		Page.
Organisms, living, material for the substance of.....	67	Supai (?) formation, occurrence and correlation of	7-8
Orygoceras (Bovillina) tuba Dall, n. sp.	113	Surprise Lake, Alaska Peninsula, features of.....	141, 142
(Ibiceornu) areolatum Dall, n. sp.	113, Pl. XXVI		
crenulatum Dall, n. sp.	113, Pl. XXVI		
idahoense Dall, n. sp.	113, Pl. XXVI		
		T	
P		Temperatures within the earth	78-79
Payettia dalli (White)	112	Terminalia lesleyana (Lesquereux) Berry	91-92, Pl. XXIII
Periodic classification of the elements, description of.....	56-57	The Gates, Alaska Peninsula, features of	139, 142
table showing	56	plate showing	140
Pervinquier, Leon, cited	28	Theories, scientific, value of	63, 64
Pila scotica, occurrence of, in boghead coals from Scotland	132	Thiessen, Reinhardt, Origin of the boghead coals.....	121-138
Plute Farms, Utah, section near	17	Thorium, atomic weight of, not known.....	82
Planorbis? sp.	111	products of disintegration of	80, 81
Plants, different compounds produced by.....	67-68	Todillo(?) formation, nature of	13
reproduction of, by seeds	68	Trachyte Creek, Utah, section on Colorado River near	19
Poppy, hydrocarbons produced by	67-68	view near, plate showing	6
Prout, hydrogen basis of atomic weights proposed by	55, 59	Tumuk sandstone, features of	15
Pseudotissotia (Choffaticeras) sp.?	30-31, Pl. XII	"Tumuk shale," features of	15
		Twomile Canyon, Utah, section on	19-20
R			
Radioactivity, discovery of.....	51, 79	U	
products of	80-83	Upper Desolation Canyon, Utah, rocks exposed in.....	45-46
Radium, atomic weight of	82	Uranium, atomic weight of, not known.....	82
Rare-earth metals, origin of	58	products of disintegration of	80, 81
place of, in the periodic table	56-58	Utah, southeastern. See Colorado Plateau.	
relation of, to Moseley's "law"	59		
Ray Creek, Alaska Peninsula, volcanic ash on, plate showing	140	V	
Red Canyon, Utah, rocks exposed in	40-41	Valvata O. F. Müller?	115
Reeside, John B., jr., A new fauna from the Colorado group of	25-33	Vascones, Chiofat, features of	28-29
southern Montana	25-33	moultoni Reeside, n. sp.	30, Pls. XVII, XXIII
Notes on the geology of Green River valley between Green	35-50	stantoni Reeside, n. sp.	30, Pls. XIX-XXI
River, Wyo., and Green River, Utah	35-50	thomi Reeside, n. sp.	29-30, Pls. XI-XXI
Reactions, chemical, what atomic adjustments are involved in?	63	sp.	30, Pl. XX
Reinschia australis, occurrence of, in Australian boghead coal.....	131-132	Volcanoes, series of, along Alaska Peninsula and the Aleutian	139
Rocks, igneous, average composition of	71	Islands	139
Rutherford, Sir Ernest, hypothesis of, for evolution of the elements	59-63	Vorticifex troni Meek	111
work of, on the disintegration of elements.....	85-86		
S		W	
San Juan Canyon, Utah, plates showing	6, 7, 12	Wasatch formation in Colorado and Wyoming, conglomerate in, ...	100
San Juan River, Utah, sections on	17	plate showing	100
Sandria Brusina	114	examinations of	93
Sayersville, Tex., fossil plants found near	87	history of	99-100
fossil plants found near, age of	88-89	map showing distribution of	94
distribution of	90	nomenclature of	98-99
Sears, J. D., and Bradley, W. H., Relations of the Wasatch and	93-107	source of material of	96-97
Green River formations in northwestern Colorado and	93-107	stratigraphic features of	91-95
southern Wyoming	93-107	structure of	95
Shale, bituminous, use of term	121	variations in material of	97-98
bituminous, varieties of	121	Washington, H. S., analyses of igneous rocks assembled by.....	71
carbonaceous, use of term	121	Waterpocket Fold, Utah, plate showing	13
oil, use of term	121	section across	21-22
Shinarump conglomerate, occurrence and deposition of	10-11	surface distribution of rock formations near, plate showing	13
Silver Falls Creek, south fork of, Utah, section near	20	Whirlpool Canyon, Colo.-Utah, rocks exposed in	42
Smith, Walter R., Aniakchak Crater, Alaska Peninsula	139-145	Wlebert, E., on the average composition of the lithosphere	75-76
Snake River valley, Idaho, Balkan fresh-water fauna in	109-115	Williamson, E. D., with Adams, L. H., determinations of com-	79
Solar system, composition of members of	77	pressibility of rocks by	79
Solar systems, differences in composition among	64	Wilson Mesa, Utah, view near, plate showing	12
Spencer Camp, Utah, section near	14	Wingate sandstone, nature of	12-13
Sphaerium idahoense Meek	111	Wonsits Valley, Utah, rocks exposed in	44-45
meeki Dall, n. sp.	111, Pl. XXVI	Wyoming, southern, sections of Green River formation in	101-107
rugosum Meek	110	southern, topography of	94
Split Mountain Canyon, Utah, rocks exposed in	43-44		
Stability of substances, conditions affecting	52	Z	
Stars, "new," phenomena of	86	Zahns Camp, Utah, section at	17
		Zinc, possible complexity of	85

Properties of the
crystals of Alun