

Fire and Vegetation History of the Jemez Mountains

by Craig D. Allen, U.S. Geological Survey, Jemez Mountains Field Station,
Midcontinent Ecological Science Center, Los Alamos

Day One

Historic patterns of fire occurrence and vegetation change in the Jemez Mountains of northern New Mexico have been described in detail by using multiple lines of evidence. Data sources include old aerial and ground-based photographs, historic records, charcoal deposits from bogs, fire-scarred trees (Fig. 1), tree-ring reconstructions of precipitation, and field sampling of vegetation and soils. The forests and woodlands that cloak the southwestern uplands provide the most extensive and detailed regional-scale network of fire history data available in the world (Swetnam and Baisan, 1996; Swetnam et al., 1999; Allen, in press).

Modern climate/vegetation patterns basically developed in the Southwest about 11,000–8,000 years before present.

Substantial fire activity apparently emerged in the Southwest during that time, as evidenced by the contemporaneous and rapid spread of fire-adapted ponderosa pine forests across the region (Anderson, 1989), and by the abundant charcoal deposits found in lake and bog sediments (Brunner-Jass, 1999; Weng and Jackson, 1999). Charcoal sediments from Alamo Bog in the central Jemez Mountains indicate essentially continuous fire activity extending back almost 9,000 years (Brunner-Jass, 1999).

About 5,200 historic fires have been mapped in the Jemez Mountains for the period 1909–1996 from administrative records of local land-management agencies (Fig. 2). Lightning caused fully 75% of the recorded fires, with acreage burned




FIGURE 1—Map of fire scar sample site locations in the Jemez Mountains.




FIGURE 2—Point locations of more than 5,000 historic wildfires in the Jemez Mountains, 1909–1996, compiled from the administrative records of land-management agencies (Snyderman and Allen, 1997). Person-caused fires cluster near major roadways, campgrounds, habitations, and other human use areas.

peaking in the dry months of May and June before the onset of summer monsoon rains. High levels of lightning activity naturally foster fire ignitions here. For example, 62 thunderstorm-days/year are observed at Los Alamos, generating large numbers of lightning strikes. An automated lightning detection system recorded 165,117 cloud-to-ground lightning strikes over a 2,994 mi² area centered on the Jemez Mountains during the period 1985–1994 (Fig. 3). The annual number of recorded lightning strikes varied between 9,410 and 23,317. Particularly important for fire ignitions is the substantial lightning activity during the warm, dry, foreshummer months of April through June. Lightning strikes during this period are the most significant sources of fire ignition because lightning is much more likely to start a spreading fire if it strikes dry fuels. Because lightning ignitions are so frequent and ubiquitous in the Southwest, climate and fuel conditions are the main drivers of fire regime dynamics in this region.

Fire scars were sampled from over 600 trees, snags, and logs at 42 sites around the Jemez Mountains in northern New Mexico (Fig. 1), resulting in over 4,000 dendrochronologically dated fire scars. Fire scar dates extend back to 1422 A.D. These data have been used to develop fire histories at multiple spatial scales, from individual trees to watersheds and finally the entire mountain range. Fire histories were reconstructed for vegetation types ranging from piñon-juniper woodlands up through ponderosa pine forests and mixed conifer forests into high-elevation spruce-fir forests (Touchan et al., 1996; Allen et

al., 1996). These fire histories show that frequent, low-intensity surface fires naturally characterized most southwestern forests. These fires spread widely through grassy understory fuels, maintaining relatively open forest conditions (Fig. 4).

Pre-1900 mean fire intervals ranged from 5 to 25 years across the Jemez Mountains (Fig. 5). Significant spatial variation in past fire regimes is evident, depending upon such local factors as vegetation/fuel type, topography, and land-use history. Fire frequencies and area burned have been greatest in mid-elevation ponderosa pine forests. Fire activity commonly occurred over extensive areas (Allen et al., 1998); for example, watershed-wide fires occurred about every 16 years across the 9-mi-long Frijoles watershed in Bandelier before 1900 (Allen, 1989). In some years fires apparently burned across most of the Jemez Mountains (Allen et al., 1998), and indeed even across the Southwest (Swetnam et al., 1999; see graphics at: <http://biology.usgs.gov/luhna/chap9.html>).

Climate variability acted to regionally synchronize prehistoric fire activity, as major fire years were clearly associated with drought conditions, while wet periods recorded little fire activity (Touchan et al., 1996; Swetnam and Baisan, 1996; Swetnam and Betancourt, 1998). The most extensive fire activity in ponderosa pine forests occurred in dry years that followed within 1–3 years of wet conditions. This pattern of major fire years suggests the importance of both fuel production and fuel moisture in these fire regimes, with antecedent wet conditions stimulating the buildup of continuous fuels and subsequent drought conditions enabling the fuels to burn widely (Swetnam and Baisan, 1996). The common occurrence of persistent drought conditions in the Southwest likely allowed some fires to burn for months.

In most cases the seasonality of fire occurrence can be inferred by the relative position of a fire scar within the annual growth rings. The patterns of fire seasonality developed from prehistoric fire scars and modern fire records are generally




FIGURE 3—Map of 23,317 lightning strikes recorded across 2,994 mi² in the Jemez Mountains area during 1986 by the national automated lightning detection system. The nominal resolution of the locational data is ± 2 km.



FIGURE 4—Open ponderosa pine forest representing “typical” pre-1900 conditions, with grassy understory and surface fire activity.

indistinguishable, indicating that prehistoric fires occurred during the same seasons as modern lightning-ignited fires—predominantly in the spring and summer. Fall fires were rare.

Spatial patterns of consecutive-year fire events indicate the importance of herbaceous fuels in supporting fire spread in pre-settlement forests. Railroads linked northern New Mexico to external markets by about 1880, leading to a local boom in livestock numbers. Abrupt declines in fire frequency throughout the Jemez Mountains in the late 1800s (Fig. 5), decades before active fire suppression, support the hypothesis that overgrazing induced suppression of surface fires as livestock (particularly sheep in mountain forests) literally ate the grassy fuels through which fires previously had spread. Fires would

have repeatedly burned across widespread parts of the Southwest during the 20th century if the many natural and human-caused fires had not been vigorously suppressed after 1910 (Swetnam et al., 1999).

This history of livestock grazing and fire suppression in the Jemez Mountains has driven such landscape-wide vegetation changes as: increased density of woody species and accelerated erosion rates in piñon-juniper woodlands; conversion of ponderosa pine forests into thickets (or crown-fire-created grasslands and shrublands); changes in species composition and structure in mixed conifer forests (Fig. 6); and invasion of grasslands and meadows by trees and shrubs (Allen, 1989). Similar changes have occurred throughout the Southwest (Bogan et al., 1998; see graphics at <http://biology.usgs.gov/s+t/SNT/noframe/sw152.htm>). The increased densities of forests over the past century (often 10-fold increases) have markedly changed many ecosystem processes, including patterns of runoff and water yield from regional watersheds. For example, increased forest densities lead to decreases in total streamflow, peak flow, and base flow (Ffolliott et al., 1989), important concerns in the water-limited Southwest.

Fire behavior has also greatly changed due to the landscape-wide build ups of woody fuels associated with a century of fire suppression. As a result the frequency and severity of wildfire activity (including lightning-ignited fires) has been escalating despite increasing human suppression efforts, as the mean number of lightning fires/year in the Southwest grew by over 50% from 1940 to 1975 (Barrows, 1978) and the mean annual acreage burned has increased continuously since about 1960 (Swetnam and Betancourt, 1998). Unnatural stand-replacing conflagrations like the 1977 La Mesa fire (Allen, 1996), the 1996 Dome fire (Fig. 7), and the 2000 Cerro Grande fire are occurring more often in over-dense ponderosa pine forests




FIGURE 5—Fire scar chronology for Quemazon locality, western edge of Los Alamos townsite. Horizontal lines represent the life spans of individual trees, while fire scar events are shown by short vertical bars. Fire years are listed along the lower axis.




FIGURE 6—Altered ponderosa pine stand in need of restoration, showing changes in both stand structure and species composition. The dense midstory of mixed conifer trees provides ladder fuels that favor crown-fire development.



FIGURE 7—Dome fire, Day 2, April 26, 1996, near headwaters of Capulin Canyon.

(Covington and Moore, 1994). Extensive (>0.5 mi^2) stand-replacing fires rarely (if ever) occurred in pure, southwestern ponderosa pine forests before the middle of the 20th century. Severe crown fires typically cause major watershed impacts, including accelerated flooding and erosion. Twentieth century landscape scars created by stand-replacing fires in ponderosa pine and lower elevation mixed conifer are long-lasting lega-




FIGURE 8—Changes in vegetation cover between 1954 and 1963 on Frijolito Mesa in Bandelier National Monument, showing persistent ponderosa pine forest (1.4 mi^2), persistent piñon-juniper woodland (5.9 mi^2), and the ecotone shift zone (1.9 mi^2) where forest changed to woodland due to the death of the overstory ponderosa pine trees.

cies of human error in managing these ecosystems. Recovery of forest communities within such burned and eroded landscapes may not occur for centuries. Fire history data and evidence of extreme geomorphic responses following extensive crown fires provide strong justification for management programs aimed at preventing the future occurrence of these ecological and societal disasters (Covington et al., 1997; Allen et al., in revision).

It is interesting to note that droughts can also cause extensive ecosystem changes by rapidly killing vegetation. For example, a severe, regional drought occurred during the 1950s in the Southwest. Associated tree mortality in the Jemez Mountains caused the ecotone between ponderosa pine forests and piñon-juniper woodlands to shift upslope by as much as 1.2 mi in less than 5 years (Fig. 8), while mixed piñon-juniper woodlands were converted to overstories of only juniper at many sites (Allen and Breshears, 1998). The 1950s drought may have also reduced herbaceous ground cover in these ecotone zones, contributing to current high erosion rates (discussed in a separate minipaper). Projected global climate changes may render over-dense southwestern forests increasingly susceptible to rapid decline through drought-induced mortality, associated insect outbreaks, and crown fires (Swetnam and Betancourt, 1998).

References

- Allen, C. D., 1989, Changes in the landscape of the Jemez Mountains, New Mexico: Unpublished PhD dissertation, University of California, Berkeley, 346 p.

- Allen, C. D. (tech. ed.), 1996, Fire effects in southwestern forests—proceedings of the Second La Mesa Fire Symposium, Los Alamos, New Mexico: USDA Forest Service, Gen. Technical Report RM-GTR-286, Fort Collins, Colorado, 216 p.
- Allen, C. D., in press, An ecological history of fire in the American Southwest; in Vale, T. R. (ed.), Western wilderness—fire, native peoples, and the natural landscape: Island Press, Covelo, California.
- Allen, C. D., and Breshears, D. D., 1998, Drought-induced shift of a forest/woodland ecotone—rapid landscape response to climate variation: Proceedings of the National Academy of Sciences of the United States of America, v. 95, p. 14,839–14,842.
- Allen, C. D., Betancourt, J. L., and Swetnam, T. W., 1998, Landscape changes in the southwestern United States—techniques, long-term datasets, and trends; in Sisk, T. D. (ed.), Perspectives on the land use history of North America—a context for understanding our changing environment: U.S. Geological Survey, Biological Science Report USGS/BRD/BSR-1998-0003, p. 71–84.
- Allen, C. D., Touchan, R., and Swetnam, T. W., 1996, Overview of fire history in the Jemez Mountains, New Mexico; in Goff, F., Kues, B. S., Rogers, M. A., McFadden, L. D., and Gardner, J. N. (eds.), Jemez Mountains region: New Mexico Geological Society, Guidebook 47, p. 35–36.
- Allen, C., Falk, D., Hoffman, M., Klingel, J., Morgan, P., Savage, M., Schulke, T., Stacey, P., Suckling, K., and Swetnam, T., in revision, Ecological restoration of southwestern ponderosa pine ecosystems—a broad framework: Restoration Ecology.
- Anderson, R. S., 1989, Development of the southwestern ponderosa pine forests—what do we really know? USDA Forest Service, Gen. Technical Report RM-185, Fort Collins, Colorado, p. 15–22.
- Barrows, J. S., 1978, Lightning fires in southwestern forests: Final Report to USDA Forest Service, Intermountain Forest and Range Experiment Station, 154 p.
- Bogan, M. A., Allen, C. D., Muldavin, E. H., Platania, S. P., Stuart, J. N., Farley, G. H., Melhop, P., and Belnap, J., 1998, Southwest; in Mac, M. J., Opler, P. A., and Doran, P. D. (eds.), National status and trends report: U.S. Geological Survey, Washington, D.C., p. 543–592.
- Brunner-Jass, R., 1999, Fire occurrence and paleoecology at Alamo Bog and Chihuahuénos Bog, Jemez Mountains, New Mexico: Unpublished MS thesis, Northern Arizona University, Flagstaff, 140 p.
- Covington, W. W., and Moore, M. M., 1994, Southwestern ponderosa pine forest structure—changes since Euro-American settlement: Journal of Forestry, v. 92, p. 39–47.
- Covington, W. W., Fulé, P. Z., Moore, M. M., Hart, S. C., Kolb, T. E., Mast, J. N., Sackett, S. S., and Wagner, M. R., 1997, Restoring ecosystem health in ponderosa pine forests of the Southwest: Journal of Forestry, v. 95, p. 23–29.
- Ffolliott, P. F., Gottfried, G. S., and Baker, M. B., Jr., 1989, Water yield from forest snowpack management—research findings in Arizona and New Mexico: Water Resources Research, v. 25, p. 1999–2007.
- Snyderman, D., and Allen, C. D., 1997, Fire in the mountains—analysis of historical fires for Bandelier National Monument, Santa Fe National Forest, and surrounding areas, 1909–1996: Unpublished report on file at USGS Jemez Mountains Field Station, 27 p., 18 figs.
- Swetnam, T. W., and Baisan, C. H., 1996, Historical fire regime patterns in the southwestern United States since A.D. 1700; in Allen, C. D. (tech. ed.), Fire effects in southwestern forests—proceedings of the Second La Mesa Fire Symposium: USDA Forest Service, Gen. Technical Report RM-GTR-286, Fort Collins, Colorado, p. 11–32.
- Swetnam, T. W., and Betancourt, J. L., 1998, Mesoscale disturbance and ecological response to decadal climatic variability in the American Southwest: Journal of Climate, v. 11, p. 3128–3147.
- Swetnam, T. W., Allen, C. D., and Betancourt, J. L., 1999, Applied historical ecology—using the past to manage for the future: Ecological Applications, v. 9, p. 1189–1206.
- Touchan, R., Allen, C. D., and Swetnam, T. W., 1996, Fire history and climatic patterns in ponderosa pine and mixed-conifer forests of the Jemez Mountains, northern New Mexico; in Allen, C. D. (tech. ed.), Fire effects in southwestern forests—proceedings of the Second La Mesa Fire Symposium: USDA Forest Service, Gen. Technical Report RM-GTR-286, Fort Collins, Colorado, p. 33–46.
- Weng, C., and Jackson, S. T., 1999, Late glacial and Holocene vegetation history and paleoclimate of the Kaibab Plateau, Arizona: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 153, p. 179–201.

Craig D. Allen

Research Ecologist

USGS Midcontinent Ecological Science Center, Jemez Mountains Field Station

HCR 1, Box 1, #15 Los Alamos, NM 87544

505-672-3861 ext. 541

Fax: 505-672-9607

craig_allen@usgs.gov

Education: BS and MS, University of Wisconsin at Madison; PhD, University of California at Berkeley

Craig Allen is a research ecologist with the U.S. Geological Survey. He received degrees in geography from the University of Wisconsin (Madison), and a PhD focused on forest and landscape ecology from the University of California (Berkeley). He has worked as an ecologist with the Department of Interior since 1989, beginning with the National Park Service through his current post with USGS. Allen conducts research on the ecology and environmental history of southwestern landscapes, oversees the USGS Jemez Mountains Field Station at Bandelier National Monument (New Mexico), and provides technical support to land management agencies in the region. Ongoing research activities with multiple collaborators include: development of vegetation and fire histories in the Southwest; fire effects on Mexican spotted owls, Jemez Mountains salamanders, and nitrogen cycling; responses of semiarid forests and woodlands to drought (and links to global change issues); development of long-term ecological monitoring networks across landscape gradients at Bandelier; dynamics of runoff, erosion, and restoration of piñon-juniper watersheds; and determining elk movements and habitat effects in the Jemez Mountains. Allen lives with his wife, Sharon, and children Kiyana, Benjamin, and Nikolas in Los Alamos.