



Invasive Non-native Plant Inventory for Montezuma Castle and Tuzigoot National Monuments

Natural Resource Technical Report NPS/SODN/NRTR—2009/268



ON THE COVER

Redstem storksbill (*Erodium cicutarium*). Photograph by Sheri Hagwood @ USDA-NRCS PLANTS Database.

Invasive Non-native Plant Inventory for Montezuma Castle and Tuzigoot National Monuments

Natural Resource Technical Report NPS/SODN/NRTR—2009/268

Authors

Theresa Mau-Crimmins
Melissa Mauzy
Sarah Studd
Sonoran Desert Network
7660 E. Broadway Blvd., Suite 303
Tucson, AZ 85710

Guy R. McPherson
School of Natural Resources and the Environment
Biological Sciences East
University of Arizona
Tucson, AZ 85721

Editing and Design

Alice Wondrak Biel
Sonoran Desert Network
7660 E. Broadway Blvd., Suite 303
Tucson, AZ 85710

November 2009

U.S. Department of the Interior
National Park Service
Natural Resource Program Center
Fort Collins, Colorado

The National Park Service, Natural Resource Program Center publishes a range of reports that address natural resource topics of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Technical Report Series is used to disseminate results of scientific studies in the physical, biological, and social sciences for both the advancement of science and the achievement of the National Park Service mission. The series provides contributors with a forum for displaying comprehensive data that are often deleted from journals because of page limitations.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner. Data in this report were collected and analyzed using methods based on established, peer-reviewed protocols and were analyzed and interpreted within the guidelines of the protocols.

Views, statements, findings, conclusions, recommendations, and data in this report are those of the author(s) and do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the National Park Service.

This report is available from Sonoran Desert Network website, <http://science.nature.nps.gov/im/units/sodn/index.cfm>, and the Natural Resource Publications Management website, <http://www.nature.nps.gov/publications/NRPM>.

Please cite this publication as:

Mau-Crimmins, T., M. Mauzy, S. Studd, and G. McPherson. 2009. Invasive non-native plant inventory for Montezuma Castle and Tuzigoot national monuments. Natural Resource Technical Report NPS/SODN/NRTR—2009/268. National Park Service, Fort Collins, Colorado.

Contents

Figures v

Tables v

Executive Summary vii

Acronyms ix

Acknowledgements xi

1 Introduction 1

 1.1 Project goals 1

 1.2 Study area 2

2 Methods..... 3

 2.1 Roaming surveys..... 3

 2.2 Circular-plot sampling..... 3

3 Results 7

 3.1 Roaming surveys..... 7

 3.1.1 2003 surveys 7

 3.1.2 2004 surveys 7

 3.2 Circular plots..... 8

 3.2.1 2003 surveys 8

 3.2.2 2004 surveys 8

 3.3 Alien Plants Ranking System 8

4 Discussion 29

 4.1 Comparison of methods 29

 4.2 Alien Plants Ranking System 30

5 Literature Cited 33

 Appendix A. Sample Circular Plot Field Data Collection Sheet..... 35

 Appendix B. Maps of Species Detected During Roaming Surveys, 2003. 37

 Appendix C. Maps of Species Detected During Roaming Surveys, 2004. 101

Figures

Figure 1-1. Location of Montezuma Castle and Tuzigoot national monuments, Arizona.....	2
Figure 2-1. Circular-plot layout used for spatial bias and accuracy assessment.....	3
Figure 2-2. Location of circular plots in three park units, 2003.	4
Figure 2-3. Location of circular plots in three park units, 2004.	5
Figure 3-1. Percent land area affected by non-native species, calculated from roaming surveys, 2003.....	11
Figure 3-2. Area infested by non-native species, calculated from roaming surveys, 2003.....	13
Figure 3-3. Percent land area affected by non-native species, calculated from roaming surveys, 2004.....	19
Figure 3-4. Area infested by non-native species, calculated from roaming surveys, 2004.....	22
Figure 3-5. Plot of species ranked using the APRS, showing ease of control versus level of impact.	25
Figure 3-6. Plot of species ranked using the APRS, showing ease of control versus ability to become a pest.....	27
Figure 4-1. Relative abundance of target species collected via roaming surveys and circular plots, 2003... 31	
Figure 4-2. Relative abundance of target species collected via roaming surveys and circular plots, 2004... 32	

Tables

Table 3-1. Target species occurrences mapped by park unit during roaming surveys, summer 2003.	9
Table 3-2. Area and percent affected by target species measured through roaming surveys, 2003.	10
Table 3-3. Area and percent infested by target species measured by roaming surveys, 2003.....	12
Table 3-4. Most densely concentrated species, Montezuma Castle National Monument, 2003.....	14
Table 3-5. Target species point and polygon occurrences mapped by park unit during roaming surveys, summer 2004.....	15
Table 3-6. Area and percent affected by target species measured through roaming surveys, 2004.	17
Table 3-7. Area and percent infested by target species measured by roaming surveys, 2004.....	20
Table 3-8. Small, concentrated patches of target species in Montezuma Castle and Tuzigoot national monuments, 2004.	22
Table 3-9. Target species occurrences and relative abundances within circular plots, 2003.....	23
Table 3-10. Target species occurrences and relative abundances within circular plots, 2004.....	24
Table 3-11. On a 100-point scale, ranks of target species for significance of impact, innate ability to become a pest, and feasibility of control or management.	26
Table 4-1. Target species detected by roaming surveys and circular plots, 2003.	29

Executive Summary

Early detection and trend detection of invasive non-native plant species comprise two of the Sonoran Desert Network Inventory and Monitoring Program's vital signs. Baseline information on the presence, location, and abundance of invasive non-native plant species is necessary to accomplish two goals: (1) to identify which invasive non-native species are not yet present in the park (early detection monitoring); and (2) to map the spatial location, distribution, and abundance of those species that already occur in the park as the basis for future trend assessment.

The goals of this project were to: (1) Map the spatial location, distribution, and abundance of 50 non-native plants in Montezuma Castle (MOCA) and Tuzigoot (TUZI) national monuments, in central Arizona; (2) Map the spatial location, distribution, and abundance of 50 non-native plants in the proposed expansion lands adjacent to Tuzigoot National Monument; (3) Test the efficacy of roaming surveys for non-native plant mapping using a modified version of the "Beyond NAWMA" weed mapping standards suggested by the North American Weed Management Association in these park units (Stohlgren et al. no date); and (4) Implement an analytical prioritization technique, such as the Alien Plants Ranking System (Hiebert and Stubbendieck 1993), for management planning.

Through a 2003 Cooperative Ecosystems Studies Unit agreement between the SODN and the University of Arizona School of Natural Resources, the spatial location, distribution, and abundance of target non-native plant species were mapped at the two units of MOCA (Montezuma Castle and Montezuma Well), at TUZI, and (in 2004 only) on proposed expansion lands at TUZI. Fieldwork took place in 2003 and 2004. Roaming surveys were performed with the intent of comprehensively covering each park. Additional, randomly located, plot-based surveys were performed in order to assess any observer bias and the accuracy of the roaming surveys. The data collected from both field seasons were fed directly into an analytical prioritization system, to facilitate management planning relative to invasive non-natives. All of this information enhances efforts toward an exotic plant management plan for MOCA and TUZI.

Of the 50 target species for this project, 34 species were encountered during roaming surveys in at least one of the three main park units and mapped as points or polygons during 2003 (see Table 3-1). In 2004, 44 species were encountered during roaming surveys in at least one of the four park units and mapped as points or polygons (see Table 3-5). Redstem storksbill (*Erodium cicutarium*) and red brome (*Bromus madritensis* ssp. *rubens*) were found to be the most dominant species across the units.

Area affected is a measure of plant distribution calculated directly from mapped polygon boundaries; this measure does not reflect the intensity of invasion. For example, a 1.5-ha polygon estimated at 25–50% cover would represent 1.5 ha affected. Affected-area value varied substantially by species, with several species being widespread throughout the units. Red brome was the most widely distributed species across the park units. Redstem storksbill, barley (*Hordeum* spp.), and London rocket (*Sisymbrium irio*) also occurred widely in the three park units (see Tables 3-2 and 3-6).

Area infested was calculated using polygon data by multiplying the cover class midpoint by the polygon's area. For example, a polygon 1.5 ha in area, estimated at 25–50% cover, would represent 0.5625 ha ($1.5 \times .375 = 0.5625$ ha) infested. Levels of infestation varied by species, with 36 species infesting less than 1% of the total park areas, 8 species infesting 1–5% of the total areas, and 5 species infesting 5–24% of the total areas in either 2003 or 2004 (see Tables 3-3 and 3-7). Bermuda grass (*Cynodon dactylon*) and johnsongrass (*Sorghum halepense*) were found to occupy 50–95% of small, concentrated patches in MOCA's Well unit in both 2003 and 2004 (see Table 3-8).

It is recommended that the Alien Plants Ranking System (Hiebert and Stubbendieck 1993) be used to identify species that are easy to control but have large impacts, and that management efforts should be concentrated on these plants first. No species fell within this quadrant of the graph for the units surveyed (see Figure 3-5). Several target species fall within the quadrant, "easy to control, lesser impact." Efforts could be placed on this group of species to minimize the overall impact of the weed communities at the parks. Alternatively, the most widespread species, red brome, redstem storksbill, barley, London

rocket, and cheatgrass (*Bromus tectorum*), are also the most difficult to control of the target species, and have the greatest impact. Ripgut brome (*Bromus rigidus*), red brome, cheatgrass, and barley also exhibit high potential to become pests. Careful consideration of these plants' potential for control, likelihood of spread, and current level of infestation should be weighed in planning for management efforts.

Acronyms

ft	feet
GPS	global positioning system
m	meters
MOCC	Montezuma Castle National Monument (Castle unit)
MOWE	Montezuma Castle National Monument (Well unit)
MOCA	Montezuma Castle National Monument
NAIPMS	North American Invasive Plant Mapping Standards
NAWMA	North American Weed Management Association
NM	national monument
NPS	National Park Service
SODN	Sonoran Desert Network
TUZI	Tuzigoot National Monument
USDA	U.S. Department of Agriculture
UTM	Universal Transverse Mercator

Acknowledgements

Thanks to Kathy Davis and the staff at Montezuma Castle and Tuzigoot national monuments for their support of this project. Much appreciation also goes to Melissa Mauzy and Meg Quinn for field data collection and plant identification. Kristen Beaupré performed invaluable data management and manipulation tasks, and Sarah Studd played a large role in final map production. Thanks also go to the Sonoran Desert Inventory and Monitoring Program for project support and facilities.

1 Introduction

The Sonoran Desert Inventory & Monitoring Network (SODN), whose purpose is to inventory and monitor natural resources in 11 national parks in southern Arizona and New Mexico, has identified invasive exotic plant species as a significant threat to park resources. The SODN is currently developing methods for both early detection and long-term monitoring of invasive non-native plant species in SODN park units.

The first step in invasive non-native plant monitoring is establishing baseline data on the presence, distribution, and abundance of target species in park units. Through the project documented in this report, the spatial location, distribution, and abundance of 50 key non-native plant species were mapped at Montezuma Castle and Tuzigoot national monuments, in central Arizona.

Non-native invasive species are a major threat to native species diversity and ecosystem function, causing economic impacts within the U.S. estimated at more than \$100 billion annually (Pimentel et al. 1999). In addition to competing with and displacing native species, these introduced species can hybridize with natives and alter conditions to promote the establishment and spread of other non-native species. Invasive species have been called the “single most formidable threat of natural disaster of the twenty-first century” (Schnase et al. 2002).

The National Park Service (NPS) defines exotic species as those species that occupy or could occupy park lands directly or indirectly as the result of deliberate or accidental human activities (NPS 2006). Exotic species are also commonly referred to as non-native, alien, or invasive species. Because an exotic species did not evolve in concert with the species native to an area, the exotic species is not a natural component of the natural ecosystem at that location. According to the *NPS Management Policies* (NPS 2006), “(e)xotic species will not be allowed to displace native species if displacement can be prevented.” In addition, these policies state that “high priority will be given to managing exotic species that have, or potentially could have, a substantial impact on park resources, and that can reasonably be expected to be successfully controllable.”

The initial fieldwork for this project, made possible through a 2003 Cooperative Ecosystem Stud-

ies Unit agreement between the SODN and the University of Arizona School of Natural Resources, took place in late July and early August 2003.

Though many of the target species were mapped sufficiently in 2003, numerous dead target species individuals were observed, suggesting that sampling may have missed the peak in abundance of the target species and some may have been missed altogether. To supplement these data, a second field season was undertaken in 2004, occurring earlier in the growing season.

Also, after the 2003 field season, it was recommended that we implement an analytical prioritization technique, such as Hiebert and Stubbendieck (1993), to facilitate management planning relative to invasive non-natives. This system uses species location, distribution, and abundance information, in conjunction with phenological information, to determine which non-native species should receive first priority for control measures. The data collected through both field seasons were fed directly into such a system, resulting in a prioritization for management efforts. All of this information enhances efforts toward an exotic plant management plan for Montezuma Castle and Tuzigoot national monuments.

The information collected through this project will:

1. Increase the ability of resource managers to analyze and prioritize invasive plant management needs, enhancing the time- and cost-effectiveness of management actions;
2. Serve as a baseline for long-term monitoring and assist with the evaluation of changes in alien plant populations over time as well as detecting new infestations;
3. Serve as a critical tool for increasing public and political awareness and education on invasive plant issues; and
4. Implement the Alien Plants Ranking System (Hiebert and Stubbendieck 1993) to prioritize the target species for management efforts.

1.1 Project goals

The goals of this project (followed by the years to which they apply) were to:

1. Map the spatial location, distribution, and abundance of 50 non-native plants in

Montezuma Castle and Tuzigoot national monuments (2003, 2004);

2. Map the spatial location, distribution, and abundance of 50 non-native plants in the expansion lands adjacent to Tuzigoot National Monument (2004);
3. Test the efficacy of roaming surveys for non-native plant mapping using a modified version of the “Beyond NAWMA” (Stohlgren et al. no date) weed mapping standards suggested by the North American Weed Management Association (NAWMA) in these national parks (2003, 2004); and
4. Implement an analytical prioritization technique, such as Hiebert and Stubbendieck (1993), for management planning (2004).


1.2 Study area

Located midway between the cities of Flagstaff and Phoenix along Interstate Highway 17, Montezuma Castle National Monument (MOCA) occupies 671 hectares of the upper Verde Valley

in central Arizona (Figure 1-1). MOCA comprises two comparably sized units located approximately seven kilometers apart: the Castle unit (MOCC), containing the primary structures and the park visitor center, and the Well unit (MOWE). Both are located along Beaver Creek. The Castle unit preserves a five-story, 20-room prehistoric cliff dwelling, built some 600 years ago by people of the Sinagua culture. The Well unit preserves a unique limestone sinkhole (or “well”) located on a mesa overlooking Wet Beaver Creek, a perennial stream. The well is ringed with additional structures, likely because of the nearby perennial water source.

Tuzigoot National Monument (TUZI), located 35 kilometers west of Montezuma Castle NM (Figure 1-1), was established in 1939 to protect Tuzigoot Pueblo, a 110-room multistory structure also built by people of the Sinagua culture. Tavasci Marsh, an adjacent 130-hectare wetland, has been proposed for incorporation into park boundaries and was included in the surveys during 2004.

Figure 1-1. Location of Montezuma Castle and Tuzigoot national monuments, Arizona.



2 Methods

The location, abundance, and distribution of non-native species were mapped at Montezuma Castle and Tuzigoot national monuments following the North American Invasive Plant Mapping Standards (NAWMA 2002) and the “Beyond NAWMA” guidelines suggested by researchers at the U.S. Geological Survey and Colorado State University (Stohlgren et al. no date). Because the parks to be mapped were small (less than 500 ha), an intensive mapping method was undertaken. Using handheld computers and GPS units, field crews mapped individual target plants mapped as points, and mapped invaded regions as polygons. The methodology employed was the same for both seasons, to ensure comparability.

2.1 Roaming surveys

Fifty weed species identified by the National Park Service as problematic within Arizona were the targets for this mapping effort. Spatial location, distribution, and abundance of these plants were recorded as ArcView shapefiles (.shp) using pocket PCs equipped with HGIS software (www.starpal.com) attached to GPS units. Two University of Arizona researchers identified locations of target species via walking surveys. Target-species individuals were mapped as points; patches larger than three meters in diameter (spatial resolution of GPS units) were mapped as polygons. Up to five species were recorded for each polygon, in descending order of invasion by target species. Abundance of target species was recorded for polygons in the following cover classes: <1%, 1–5%, 6–25%, 26–50%, 51–75%, 76–95%, and 96–100%.

In 2003, data-collection surveys took place in July (21–25, 28) and August (11–15, 18–22) 2003. In 2004, data were collected in May (24–28) and June (2–6, 21–26). Data were collected following the North American Invasive Plant Mapping Standards (NAIPMS) (NAWMA 2002), including the following fields: collection date, examiner, plant name (genus, species), canopy cover, location (UTM coordinates).

Data files were manipulated in the office to incorporate several additional data fields suggested by the NAIPMS (NAWMA 2002), including:

- NPS region (Intermountain Region, abbreviated as IMR)
- NPS Inventory & Monitoring Network (Sonoran Desert Network, abbreviated as SODN)
- Park unit (MOCC, MOWE, or TUZI)
- State (AZ)
- County (Yavapai)
- Ownership (NPS)
- Type of survey (observed-mapped)
- Authority (following Kearney and Peebles 1951)
- Plant code (following the USDA Natural Resources Conservation Service’s PLANTS database, <http://plants.usda.gov/plants/index.html>)
- Area of polygons

Data files were split and merged to generate individual ArcView shapefiles by species and park unit, with point and polygons appearing in separate files.

2.2 Circular-plot sampling

In the “Beyond NAWMA” guidelines, Stohlgren et al. (no date) recommend supplementing roaming surveys with randomly located circular-plot sampling to estimate spatial bias and accuracy. Approximately 10% of field resources were used to perform circular-plot sampling.

The circular plots were 7.32 m (24 ft) in diameter, encompassing three 1-m² quadrats (Figure 2-1). Percent cover of all species was recorded within the 1-m² quadrats to the nearest 1%. Percent cover of all ground variables (rock, leaf litter, dead wood, dung) was also recorded to the nearest 1%. All additional species observed within the entire circular plot were also recorded. A sample




Figure 2-1. Circular-plot layout used for spatial bias and accuracy assessment.

data sheet is provided in Appendix A. Data from all circular plot data sheets were entered into an MS Access database.

In 2003, 29 circular plots were sampled within Tuzigoot National Monument, providing excellent coverage of this small park, at 17.5 ha (Figure 2-2). Equal coverage was initially planned for each unit of Montezuma Castle National Monument; however, due to time constraints, 12 points were sampled in each unit (Figure 2-2).

To better align with the recommendation by Stohlgren and others (no date) that no more than 10% of field resources be allocated to the supplemental circular-plot method, only 10 circular plots were sampled within TUZI in 2004. Twelve plots were sampled in the Castle unit of MOCA that year, and 8 plots were sampled in the Well unit (Figure 2-3).




Figure 2-2. Location of circular plots in three park units, 2003.




Figure 2-3. Location of circular plots in three park units, 2004.

3 Results

3.1 Roaming surveys

3.1.1 2003 surveys

Of the 50 target species for this project, 34 species were encountered during roaming surveys in at least one of the three park units and mapped as points or polygons during 2003 (Table 3-1; tables and figures begin on page 7). Maps of each species by park unit appear in Appendix B.

The area of land affected by each target species is provided in Table 3-2. These values represent all land area within park units invaded by these plants, measured through the roaming surveys. Therefore, these values represent distribution of non-native plant cover, but do not reflect the intensity of invasion. Points were estimated to represent 0.5 m². Figure 3-1 displays these data, highlighting red brome (*Bromus madritensis* ssp. *rubens*) as the most widely distributed species across the three park units. Redstem storksbill (*Erodium cicutarium*), cheatgrass (*Bromus tectorum*), common Mediterranean grass (*Schismus barbatus*), barley (*Hordeum* spp.), and wild oats (*Avena fatua*) also occur widely in MOCC and MOWE. London rocket (*Sisymbrium irio*) is rather widely distributed in MOWE and TUZI.

Figure 3-1 depicts species that are widely distributed, but gives no indication of the level of infestation. In contrast, Figure 3-2 displays actual land area occupied by target species. Land area infested was calculated using polygon data by multiplying the cover class midpoint by the polygon's area. For example, a polygon 1.5 ha in area, estimated at 25–50% cover, would represent 0.5625 ha ($1.5 \times .375 = 0.5625$ ha) infested. Points were estimated to represent 0.5 m².

Figure 3-2 highlights redstem storksbill and red brome as the most dominant species across the three park units. These species are also the two most widely dispersed (Figure 3-1), as well as the most ubiquitous, infesting more than 50 hectares each across the three park units. As shown in Figure 3-2, cheatgrass, barley, and Mediterranean grass (*Schismus barbatus*) also infest sizeable land areas (<10 ha total), primarily in MOCC. These target plants are also widely distributed (Figure 3-1).

Species totals by park unit appear in Table 3-3. These values are much smaller than those in Table 3-2 because they represent total hectares completely infested by the target species.

In addition to recognizing species that are widely distributed across the park units, it is important to recognize target plants that occur in small but concentrated patches. These species may have the potential to become more widely dispersed across the study area in the future. The densest patches for these species are listed in Table 3-4, where johnsongrass (*Sorghum halepense*) and common Mediterranean grass are shown to have infested 50–75% of certain small-scale patches at MOCC and MOWE.

3.1.2 2004 surveys

Of the 50 target species for this project, 44 species were encountered during roaming surveys in at least one of the four park units surveyed and mapped as points or polygons in 2004 (Table 3-5). Maps of each species by park unit appear in Appendix C.

The area of land affected by each target species is provided in Table 3-6. These values represent all land area within park units invaded by these plants, measured through the roaming surveys. Therefore, these values represent distribution of non-native plant cover, but do not reflect the intensity of invasion. Points were estimated to represent 0.5 m². Figure 3-3 displays these data, highlighting red brome as the most widely distributed species across the park units. Redstem storksbill, barley, and London rocket also occur widely in the park units surveyed. Ripgut brome (*Bromus rigidus*) and cheatgrass are widespread in MOCC and MOWE, and Bermuda grass (*Cynodon dactylon*) and rabbitsfoot grass (*Polypogon monspeliensis*) are common on the TUZI expansion lands.

Figure 3-3 depicts species that are widely distributed, but gives no indication of the level of infestation. In contrast, Figure 3-4 displays actual land area occupied by target species. Land area infested was calculated using polygon data by multiplying the cover class midpoint by the polygon's area. For example, a polygon 1.5 ha in area estimated at 25–50% cover would represent 0.5625 ha ($1.5 \times .375 = 0.5625$ ha) infested. Points were estimated to represent 0.5 m².

Figure 3-4 highlights red brome and redstem storksbill as the most dominant species across the four park units. These species are the two most widely dispersed (Figure 3-3) as well as the most ubiquitous; red brome was measured to infest over 80 ha, and redstem storksbill was calculated at over 30 ha across the four park units (see Table 3-7). As shown in Figure 3-4, ripgut brome, barley, and cheatgrass also infest sizeable land areas (<10 ha total), primarily in MOCC; these four target plants are also widely distributed (Figure 3-3).

Species totals by park unit appear in Table 3-7. These values are much smaller than those in Table 3-6 because these represent total hectares completely infested by the target species.

In addition to recognizing species that are widely distributed across the park units, it is important to recognize target plants that are occurring in small but concentrated patches. These species may have the potential to become more widely dispersed across the study area in the future. The densest patches for these species appear in Table 3-8, where johnsongrass and common Mediterranean grass are shown to have infested 50–75% of certain small-scale patches at MOWE.

3.2 Circular plots

3.2.1 2003 surveys

Of the 50 target species, a total of 17 species were recorded within the circular plots in 2003: 10 of these in MOCC, 13 in MOWE, and 5 in TUZI (Table 3-9). Three 1-m² quadrats were located within each circular plot, in which percent cover of all species was recorded to the nearest 1%. Using these data, the relative abundance of the target species was calculated (Table 3-9).

3.2.2 2004 surveys

Of the 50 target species, 10 species were recorded within the circular plots in 2004: 6 of these in MOCC, 7 in MOWE, and 4 in TUZI (Table 3-10). Three 1-m² quadrats were located within each circular plot, in which percent cover of all species were recorded to the nearest 1%. Using these data, the relative abundance of the target species was calculated (Table 3-10).

3.3 Alien Plants Ranking System

Of the 40 species ranked using the APRS, all fell within the low- and medium-impact categories, and ranged from easy-to-control to difficult-to-control (Figure 3-5). On the 100-point scale, plants ranged from 33 to 79 (tamarisk) for significance of impact; from 7 to 51 (red brome) for innate ability to become a pest; and from 17 to 73 (red brome, ripgut brome) for difficulty to control or manage (Table 3-11). The species determined to be hardest to control, with the most serious impact, were red brome and barley.

Figure 3-6 displays the difficulty of control and ability to be a pest for the target species. The target species are rather evenly distributed along the y-axis, ability to become a pest. Several of the target species are very problematic; others appear to have less direct impact on the ecosystem they invade. The species determined to be hardest to control, with the greatest ability to become pests, were ripgut brome, red brome, and cheatgrass.

Note: Some species mapped were not ranked using the APRS because insufficient data was available on their life history strategies or ecological impacts. Also, some species ranked were only mapped during either the 2003 or 2004 season, but may have been observed during both seasons.

Table 3-1. Target species occurrences mapped by park unit during roaming surveys, summer 2003.

Species	Common name	MOCA				TUZI		Total	
		Castle unit		Well unit		Points	Polygons	Points	Polygons
		Points	Polygons	Points	Polygons				
<i>Avena fatua</i>	wild oats	9	7	0	2	0	0	9	9
<i>Boerhavia coccinea</i>	scarlet spiderling	1	0	10	6	0	0	11	6
<i>Brassica tournefortii</i>	Sahara mustard	1	0	0	1	0	0	1	1
<i>Bromus madritensis</i> ssp. <i>rubens</i>	red brome	52	49	46	18	387	29	485	96
<i>Bromus rigidus</i>	ripgut brome	65	36	52	14	0	0	117	50
<i>Bromus tectorum</i>	cheatgrass	72	23	4	12	0	0	76	35
<i>Centaurea melitensis</i>	Malta starthistle	3	3	0	3	0	0	3	6
<i>Centaurea solstitialis</i>	yellow starthistle	0	0	0	1	0	0	0	1
<i>Conyza canadensis</i>	Canadian horseweed	0	0	5	2	0	0	5	2
<i>Cynodon dactylon</i>	Bermuda grass	70	3	42	4	0	0	112	7
<i>Cyperus esculentus</i>	yellow nutsedge	3	0	3	0	0	0	6	0
<i>Digitaria sanguinalis</i>	hairy crabgrass	68	0	33	0	0	0	101	0
<i>Echinochloa crus-galli</i>	barnyard grass	6	0	4	0	0	0	10	0
<i>Eragrostis curvula</i>	weeping lovegrass	0	0	1	0	0	0	1	0
<i>Erodium cicutarium</i>	redstem storksbill	0	18	1	8	0	0	1	26
<i>Helianthus annuus</i>	common sunflower	2	0	5	1	0	0	7	1
<i>Heterotheca subaxillaris</i>	camphorweed	2	0	1	0	0	0	3	0
<i>Hordeum</i> ssp.	common barley	8	28	0	9	0	0	8	37
<i>Lactuca serriola</i>	prickly lettuce	17	0	26	2	0	0	43	2
<i>Linaria dalmatica</i>	Dalmatian toadflax	55	0	0	0	0	0	0	0
<i>Marrubium vulgare</i>	horehound	4	5	27	12	0	1	31	18
<i>Melilotus albus</i>	white sweetclover	81	0	28	0	0	0	109	0
<i>Polygonum aviculare</i>	prostrate knotweed	6	0	0	0	0	0	6	0
<i>Polypogon monspeliensis</i>	rabbitsfoot grass	116	5	22	0	0	0	138	5
<i>Rumex crispus</i>	curly dock	0	0	2	0	0	0	2	0
<i>Salsola</i> ssp.	Russian thistle	3	2	1	2	0	0	4	4
<i>Schismus barbatus</i>	common Mediterranean grass	2	13	0	4	0	0	2	17
<i>Setaria viridis</i>	green bristlegrass	4	0	4	0	0	0	8	0
<i>Sisymbrium irio</i>	London rocket	18	45	39	16	248	32	305	93
<i>Sorghum halepense</i>	johnsongrass	72	3	17	2	0	0	89	5
<i>Tamarix</i> ssp.	tamarisk	56	1	16	0	0	0	72	1
<i>Tribulus terrestris</i>	puncture vine	0	0	0	2	0	0	0	2
<i>Verbascum thapsus</i>	woolly mullein	0	0	1	0	0	0	1	0
<i>Xanthium strumarium</i>	rough cocklebur	0	3	35	0	0	0	35	3

Table 3-2. Area and percent affected by target species measured through roaming surveys, 2003.

Species	Common name	MOCA				TUZI		Total ha affected
		Castle unit		Well unit		Ha affected	Percent affected	
		Ha affected	Percent affected	Ha affected	Percent affected			
<i>Bromus madritensis</i> ssp. <i>rubens</i>	red brome	221.79	95.8%	86.31	83.7%	2.33	13.3%	310.43
<i>Erodium cicutarium</i>	redstem storksbill	175.05	75.6%	64.04	62.1%	-	-	239.09
<i>Bromus tectorum</i>	cheatgrass	169.29	73.1%	54.42	52.8%	-	-	223.71
<i>Schismus barbatus</i>	common Mediterranean grass	109.47	47.3%	39.21	38.0%	-	-	148.68
<i>Hordeum</i> spp.	common barley	47.53	20.5%	43.50	42.2%	-	-	91.03
<i>Sisymbrium irio</i>	London rocket	T	T	45.32	44.0%	2.80	15.9%	48.12
<i>Avena fatua</i>	wild oats	21.71	9.4%	17.75	17.2%	-	-	39.46
<i>Bromus rigidus</i>	ripgut brome	T	T	31.22	30.3%	-	-	31.22
<i>Marrubium vulgare</i>	horehound	0.82	0.4%	17.81	17.3%	0.07	0.4%	18.70
<i>Boerhavia coccinea</i>	scarlet spiderling	T	T	11.01	10.7%	-	-	11.01
<i>Conyza canadensis</i>	Canadian horseweed	-	-	8.90	8.6%	-	-	8.90
<i>Centaurea melitensis</i>	Malta starthistle	0.34	0.1%	7.66	7.4%	-	-	8.00
<i>Cynodon dactylon</i>	Bermuda grass	0.25	0.1%	5.97	5.8%	-	-	6.22
<i>Lactuca serriola</i>	prickly lettuce	T	T	3.93	3.8%	-	-	3.93
<i>Salsola</i> spp.	Russian thistle	0.69	0.3%	2.85	2.8%	-	-	3.54
<i>Helianthus annuus</i>	common sunflower	T	T	2.23	2.2%	-	-	2.23
<i>Sorghum halepense</i>	johnsongrass	0.13	0.1%	2.03	2.0%	-	-	2.16
<i>Brassica tournefortii</i>	Sahara mustard	T	T	1.36	1.3%	-	-	1.36
<i>Centaurea solstitialis</i>	yellow starthistle	-	-	1.29	1.3%	-	-	1.29
<i>Tribulus terrestris</i>	puncture vine	-	-	0.92	0.9%	-	-	0.92
<i>Xanthium strumarium</i>	rough cocklebur	0.11	T	T	T	-	-	0.11
<i>Tamarix</i> spp.	tamarisk	0.07	T	T	T	-	-	0.07
<i>Polypogon monspeliensis</i>	rabbitsfoot grass	0.01	T	T	T	-	-	0.01
<i>Cyperus esculentus</i>	yellow nutsedge	T	T	T	T	-	-	T
<i>Digitaria sanguinalis</i>	hairy crabgrass	T	T	T	T	-	-	T
<i>Echinochloa crus-galli</i>	barnyard grass	T	T	T	T	-	-	T
<i>Eragrostis curvula</i>	weeping lovegrass	-	-	T	T	-	-	T
<i>Heterotheca subaxillaris</i>	camphorweed	T	T	T	T	-	-	T
<i>Linaria dalmatica</i>	Dalmatian toadflax	T	T	-	-	-	-	T
<i>Melilotus albus</i>	white sweetclover	T	T	T	T	-	-	T
<i>Polygonum aviculare</i>	prostrate knotweed	T	T	-	-	-	-	T
<i>Rumex crispus</i>	curly dock	-	-	T	T	-	-	T
<i>Setaria viridus</i>	green bristlegrass	T	T	T	T	-	-	T
<i>Verbascum thapsus</i>	woolly mullein	T	T	T	T	-	-	T

T denotes area <0.01 ha or percent <0.1%.

- denotes species not observed.




Figure 3-1. Percent land area affected by non-native species, calculated from roaming surveys, 2003.

Table 3-3. Area and percent infested by target species measured by roaming surveys, 2003.

Species	Common name	MOCA				TUZI		Total ha infested
		Castle unit		Well unit				
		Ha infested	Percent infested	Ha infested	Percent infested	Ha infested	Percent infested	
<i>Erodium cicutarium</i>	redstem storksbill	49.49	21.4%	18.13	16.9%	-	-	67.62
<i>Bromus madritensis</i> ssp. <i>rubens</i>	red brome	35.24	15.2%	20.23	18.9%	0.68	3.9%	56.15
<i>Bromus tectorum</i>	cheatgrass	10.55	4.6%	1.16	1.1%	-	-	11.71
<i>Hordeum</i> spp.	common barley	9.97	4.3%	0.93	0.9%	-	-	10.90
<i>Schismus barbatus</i>	common Mediterranean grass	7.88	3.4%	0.52	0.5%	-	-	8.40
<i>Tamarix</i> spp.	tamarisk	7.20	3.1%	T	T	-	-	7.20
<i>Bromus rigidus</i>	ripgut brome	0.01	0.0%	5.07	4.7%	-	-	5.08
<i>Sisymbrium irio</i>	London rocket	0.85	0.4%	0.77	0.7%	0.86	4.9%	2.48
<i>Marrubium vulgare</i>	horehound	T	T	0.94	0.9%	T	T	0.94
<i>Conyza canadensis</i>	Canadian horseweed	-	-	0.37	0.3%	-	-	0.37
<i>Cynodon dactylon</i>	Bermuda grass	0.12	0.1%	0.24	0.2%	-	-	0.36
<i>Sorghum halepense</i>	johnsongrass	0.01	0.0%	0.32	0.3%	-	-	0.33
<i>Avena fatua</i>	wild oats	0.13	0.1%	0.09	0.1%	-	-	0.22
<i>Centaurea solstitialis</i>	yellow starthistle	-	-	0.16	0.1%	-	-	0.16
<i>Boerhavia coccinea</i>	scarlet spiderling	T	T	0.13	0.1%	-	-	0.13
<i>Lactuca serriola</i>	prickly lettuce	T	T	0.12	0.1%	-	-	0.12
<i>Salsola</i> spp.	Russian thistle	0.01	0.0%	0.09	0.1%	-	-	0.10
<i>Digitaria sanguinalis</i>	hairy crabgrass	0.01	0.0%	T	-	-	-	0.01
<i>Helianthus annuus</i>	common sunflower	T	T	0.01	0.0%	-	-	0.01
<i>Melilotus albus</i>	white sweetclover	0.01	0.0%	T	T	-	-	0.01
<i>Polypogon monspeliensis</i>	rabbitsfoot grass	0.01	0.0%	T	T	-	-	0.01
<i>Brassica tournefortii</i>	Sahara mustard	T	T	-	-	-	-	T
<i>Centaurea melitensis</i>	Malta starthistle	T	T	-	-	-	-	T
<i>Cyperus esculentus</i>	yellow nutsedge	T	T	T	T	-	-	T
<i>Echinochloa crus-galli</i>	barnyard grass	T	T	T	T	-	-	T
<i>Eragrostis curvula</i>	weeping lovegrass	-	-	T	T	-	-	T
<i>Heterotheca subaxillaris</i>	camphorweed	T	T	T	T	-	-	T
<i>Linaria dalmatica</i>	Dalmatian toadflax	T	T	-	-	-	-	T
<i>Polygonum aviculare</i>	prostrate knotweed	T	T	-	-	-	-	T
<i>Rumex crispus</i>	curly dock	-	-	T	T	-	-	T
<i>Setaria viridus</i>	green bristlegrass	T	T	T	T	-	-	T
<i>Tribulus terrestris</i>	puncture vine	-	-	-	-	-	-	T
<i>Verbascum thapsus</i>	woolly mullein	T	T	T	T	-	-	T
<i>Xanthium strumarium</i>	rough cocklebur	-	-	T	T	-	-	T

T denotes area <0.01 ha.
 - denotes species not observed.




Figure 3-2. Area infested by non-native species, calculated from roaming surveys, 2003.

Table 3-4. Most densely concentrated species, Montezuma Castle National Monument, 2003.

Species	Common name	Unit	Area occupied (ha)	% cover
<i>Cynodon dactylon</i>	Bermuda grass	Castle	0.087	50–75%
<i>Cynodon dactylon</i>	Bermuda grass	Well	0.551	50–75%
<i>Sorghum halepense</i>	johnsongrass	Well	0.740	50–75%
<i>Conyza canadensis</i>	Canadian horseweed	Well	2.226	5–25%
<i>Hordeum</i> spp.	common barley	Well	1.641	5–25%
<i>Marrubium vulgare</i>	horehound	Well	4.811	5–25%
<i>Tamarix</i> spp.	tamarisk	Castle	0.069	5–25%

Table 3-5. Target species point and polygon occurrences mapped by park unit during roaming surveys, summer 2004.

		MOCA				TUZI				Total	
		Castle unit		Well unit		Main unit		Expansion unit			
		Points	Polygons	Points	Polygons	Points	Polygons	Points	Polygons	Points	Polygons
Scientific name	Common name										
<i>Acroptilon repens</i>	Russian knapweed	0	0	0	0	0	0	11	0	11	0
<i>Ailanthus altissima</i>	tree of heaven	2	0	0	0	0	0	1	0	3	0
<i>Arundo donax</i>	giant reed	0	0	0	0	1	0	0	0	1	0
<i>Avena fatua</i>	wild oats	9	2	0	0	0	0	0	0	9	2
<i>Boerhavia coccinea</i>	scarlet spiderling	3	0	1	2	0	0	2	0	6	2
<i>Bromus japonicus</i>	Japanese brome	0	0	0	0	0	0	9	0	9	0
<i>Bromus madritensis</i> ssp. <i>rubens</i>	red brome	15	28	33	16	485	24	15	7	548	75
<i>Bromus rigidus</i>	ripgut brome	42	17	18	10	0	0	6	2	66	29
<i>Bromus tectorum</i>	cheatgrass	28	13	0	9	0	0	9	0	37	22
<i>Centaurea melitensis</i>	Malta starthistle	3	3	1	1	0	0	1	0	5	4
<i>Conyza canadensis</i>	Canadian horseweed	0	0	3	0	0	0	0	0	3	0
<i>Cynodon dactylon</i>	Bermuda grass	48	3	24	3	3	1	32	3	104	10
<i>Digitaria sanguinalis</i>	hairy crabgrass	49	0	15	0	0	0	3	0	67	0
<i>Elaeagnus angustifolia</i>	Russian olive	0	0	0	0	0	0	16	0	16	0
<i>Eragrostis cilianensis</i>	stinkgrass	0	0	0	0	2	0	0	0	2	0
<i>Eragrostis curvula</i>	weeping lovegrass	0	0	14	0	0	0	0	0	14	0
<i>Eragrostis lehmannia</i>	Lehmann lovegrass	12	1	0	0	1	1	0	0	12	2
<i>Erodium cicutarium</i>	redstem storksbill	8	16	7	11	22	27	3	5	40	59
<i>Heterotheca subaxillaris</i>	camphorweed	11	2	7	1	0	0	6	0	24	3
<i>Hordeum leporinum</i>	hare barley	11	17	0	7	0	2	1	5	12	31
<i>Kochia scoparia</i>	fireweed	0	0	0	0	0	0	54	0	54	0
<i>Lactuca serriola</i>	prickly lettuce	9	0	11	1	0	0	2	0	22	1
<i>Lamium amplexicaule</i>	henbit	0	0	0	0	2	0	0	0	2	0
<i>Linaria dalmatica</i>	Dalmatian toadflax	25	2	1	2	0	0	0	0	26	4
<i>Lolium perenne</i>	perennial ryegrass	3	0	0	0	0	0	0	0	3	0
<i>Marrubium vulgare</i>	horehound	9	4	7	8	10	3	4	0	30	15
<i>Medicago</i> sp.	alfalfa	0	0	0	0	1	0	0	0	1	0
<i>Melilotus albus</i>	white sweetclover	7	0	6	0	0	0	6	0	19	0

Table 3-5. Target species point and polygon occurrences mapped by park unit, summer 2004, cont.

		MOCA				TUZI				Total	
		Castle unit		Well unit		Main unit		Expansion unit			
		Points	Polygons	Points	Polygons	Points	Polygons	Points	Polygons	Points	Polygons
Scientific name	Common name										
Melilotus indicus	yellow sweetclover	18	0	16	0	0	0	2	0	36	0
Polygonum aviculare	prostrate knotweed	10	0	0	0	0	0	1	0	11	0
Polypogon monspeliensis	rabbitsfoot grass	35	0	33	0	0	0	27	2	95	2
Rumex crispus	curly dock	12	0	5	0	0	0	8	0	25	0
Salsosa kali	Russian thistle	8	4	0	2	0	0	6	1	14	7
Schismus barbatus	common Mediterranean grass	1	7	0	1	4	0	1	1	6	9
Sisymbrium irio	London rocket	6	7	2	0	116	24	3	3	127	34
Solanum elaeagnifolium	silverleaf nightshade	0	0	0	0	0	1	0	0	0	1
Sonchus asper	spiny sowthistle	0	0	13	1	0	0	4	0	17	1
Sonchus oleraceus	common sowthistle	0	0	0	0	0	0	1	0	1	0
Sorghum halepense	johnsongrass	32	0	12	3	0	0	0	0	44	3
Tamarix ramosissima	five-stamen tamarisk	48	0	12	0	0	0	15	0	75	0
Torilis arvensis	spreading hedgeparsley	0	0	0	0	0	0	2	0	2	0
Tribulus terrestris	puncture vine	0	0	0	1	0	0	1	0	1	1
Verbascum thapsus	woolly mullein	12	0	8	0	0	0	2	0	22	0
Xanthium strumarium	rough cocklebur	43	0	14	0	0	0	9	0	66	0

Table 3-6. Area and percent affected by target species measured through roaming surveys, 2004.

Species	Common name	MOCA			TUZI			Total area affected (ha)		
		Castle unit		Well unit	Main unit		Expansion unit			
		Ha affected	Percent affected	Ha affected	Percent affected	Ha affected	Percent affected			
<i>Bromus madritensis</i> spp. <i>rubens</i>	red brome	195.00	84.2	83.41	80.9	9.06	51.6	34.83	36.9	322.30
<i>Erodium cicutarium</i>	redstem storksbill	143.13	61.8	73.87	71.7	15.55	88.6	28.45	30.2	261.00
<i>Bromus rigidus</i>	ripgut brome	65.60	28.3	27.92	27.1	0.00	-	3.14	3.3	96.66
<i>Bromus tectorum</i>	cheatgrass	58.05	25.1	36.59	35.5	0.00	-	0.00	T	94.64
<i>Hordeum leporinum</i>	hare barley	66.48	28.7	14.18	13.8	0.34	1.9	14.03	14.9	95.04
<i>Sisymbrium irio</i>	London rocket	55.59	24.0	0.00	T	8.92	50.8	7.65	8.1	72.16
<i>Schismus barbatus</i>	common Mediterranean grass	69.13	29.9	0.16	0.2	0.00	T	1.94	2.1	71.23
<i>Eragrostis lehmannia</i>	Lehmann lovegrass	49.36	21.3	0.00	0.0	0.28	1.6	0.00	-	49.64
<i>Avena fatua</i>	wild oats	49.50	21.4	0.00	0.0	0.00		0.00	-	49.50
<i>Cynodon dactylon</i>	Bermuda grass	0.70	0.3	1.15	T	0.28	1.6	34.70	36.8	36.83
<i>Marrubium vulgare</i>	horehound	20.17	8.7	13.46	13.1	1.53	8.7	0.00	T	35.16
<i>Salsosa kali</i>	Russian thistle	26.70	11.5	4.14	0.0	0.00	-	5.16	5.5	36.01
<i>Polypogon monspeliensis</i>	rabbitsfoot grass	0.00	T	0.00	T	0.00	-	30.94	32.8	30.94
<i>Boerhavia coccinea</i>	scarlet spiderling	0.00	T	21.37	20.7	0.00	-	0.00	T	21.37
<i>Heterotheca subaxillaris</i>	camphorweed	12.87	5.6	3.49	3.4	0.00	-	0.00	T	16.36
<i>Linaria dalmatica</i>	Dalmatian toadflax	5.08	2.2	3.75	3.6	0.00	-	0.00	-	8.83
<i>Sorghum halepense</i>	johnsongrass	0.00	T	4.85	4.7	0.00	-	0.00	-	4.85
<i>Sonchus asper</i>	spiny sowthistle	0.00	0.0	2.00	1.9	0.00	-	0.00	T	2.00
<i>Centaurea melitensis</i>	Malta starthistle	0.55	0.2	0.23	0.2	0.00	-	0.00	T	0.78
<i>Lactuca serriola</i>	prickly lettuce	0.00	T	0.46	0.4	0.00	-	0.00	T	0.46
<i>Tribulus terrestris</i>	puncture vine	0.00	0.0	0.16	0.2	0.00	-	0.00	T	0.16
<i>Tamarix ramosissima</i>	five-stamen tamarisk	0.02	T	0.01	T	0.00	-	0.01	T	0.04
<i>Solanum elaeagnifolium</i>	silverleaf nightshade	0.00	0.0	0.00	0.0	0.03	T	-	-	0.03
<i>Kochia scoparia</i>	fireweed	0.00	0.0	0.00	0.0	0.00	-	0.03	T	0.03
<i>Elaeagnus angustifolia</i>	Russian olive	0.00	0.0	0.00	0.0	0.00	-	0.01	T	0.01
<i>Acroptilon repens</i>	Russian knapweed	0.00	0.0	0.00	0.0	0.00	-	0.00	T	T
<i>Ailanthus altissima</i>	tree of heaven	0.00	T	0.00	0.0	0.00	-	0.00	T	T
<i>Arundo donax</i>	giant reed	0.00	0.0	0.00	0.0	0.00	T	0.00	-	T
<i>Bromus japonicus</i>	Japanese brome	0.00	0.0	0.00	0.0	0.00	-	0.00	T	T

Table 3-6. Area and percent affected by target species measured through roaming surveys, 2004, cont.										
Species	Common name	MOCA				TUZI				Total area affected (ha)
		Castle unit		Well unit		Main unit		Expansion unit		
		Ha affected	Percent affected	Ha affected	Percent affected	Ha affected	Percent affected	Ha affected	Percent affected	
<i>Conyza canadensis</i>	Canadian horseweed	0.00	0.0	0.00	T	0.00	-	0.00	-	T
<i>Digitaria sanguinalis</i>	hairy crabgrass	0.00	T	0.00	T	0.00	-	0.00	T	T
<i>Eragrostis cilianensis</i>	stinkgrass	0.00	0.0	0.00	0.0	0.00	T	0.00	-	T
<i>Eragrostis curvula</i>	weeping lovegrass	0.00	0.0	0.00	T	0.00	-	0.00	-	T
<i>Lamium amplexicaule</i>	henbit	0.00	0.0	0.00	0.0	0.00	-	0.00	T	T
<i>Lolium perenne</i>	perennial ryegrass	0.00	T	0.00	0.0	0.00	-	0.00	-	T
<i>Medicago</i> sp.	alfalfa	0.00	0.0	0.00	0.0	0.00	-	0.00	T	T
<i>Melilotus albus</i>	white sweetclover	0.00	T	0.00	T	0.00	-	0.00	T	T
<i>Melilotus indicus</i>	yellow sweetclover	0.00	T	0.00	T	0.00	-	0.00	T	T
<i>Polygonum aviculare</i>	prostrate knotweed	0.00	T	0.00	0.0	0.00	-	0.00	T	T
<i>Rumex crispus</i>	curly dock	0.00	T	0.00	T	0.00	-	0.00	T	T
<i>Sonchus oleraceus</i>	common sowthistle	0.00	0.0	0.00	0.0	0.00	-	0.00	T	T
<i>Torilis arvensis</i>	spreading hedgeparsley	0.00	0.0	0.00	0.0	0.00	-	0.00	T	T
<i>Verbascum thapsus</i>	woolly mullein	0.00	T	0.00	T	0.00	-	T	T	T
<i>Xanthium strumarium</i>	rough cocklebur	0.00	T	0.00	T	0.00	-	T	T	T

T denotes area of <0.01 ha or <0.1 %.

- denotes species not observed.




Figure 3-3. Percent land area affected by non-native species, calculated from roaming surveys, 2004.

Table 3-7. Area and percent infested by target species measured by roaming surveys, 2004.

Scientific name	Common name	MOCA				TUZI			
		Castle unit		Well unit		Main unit		Expansion	
		Ha affected	Percent affected	Ha affected	Percent affected	Ha affected	Percent affected	Ha affected	Percent affected
<i>Bromus madritensis</i> ssp. <i>rubens</i>	red brome	53.96	23.3%	21.34	19.9%	1.89	10.7%	6.03	6.4%
<i>Erodium cicutarium</i>	redstem storksbill	14.94	6.5%	4.15	3.9%	8.47	48.2%	2.81	3.0%
<i>Bromus rigidus</i>	ripgut brome	20.82	9.0%	5.97	5.6%	-		0.08	0.1%
<i>Hordeum leporinum</i>	hare barley	14.32	6.2%	2.28	2.1%	T		2.19	2.3%
<i>Bromus tectorum</i>	cheatgrass	9.63	4.2%	1.45	1.4%	-		T	
<i>Salsola kali</i>	Russian thistle	2.85	1.2%	0.09	0.1%	-		1.94	2.1%
<i>Schismus barbatus</i>	common Mediterranean grass	2.83	1.2%	T		T		0.05	0.1%
<i>Sisymbrium irio</i>	London rocket	0.30	0.1%	T		0.48	2.7%	0.79	0.8%
<i>Cynodon dactylon</i>	Bermuda grass	0.03	0.0%	0.47	0.4%	0.04	0.2%	0.18	0.2%
<i>Heterotheca subaxillaris</i>	camphorweed	0.32	0.1%	0.09	0.1%	-		T	
<i>Sorghum halepense</i>	johnsongrass	T		0.31	0.3%	-		-	
<i>Marrubium vulgare</i>	horehound	0.10	0.0%	0.19	0.2%	0.01	0.1%	T	
<i>Linaria dalmatica</i>	Dalmatian toadflax	0.13	0.1%	0.13	0.1%	-		-	
<i>Avena fatua</i>	wild oats	0.25	0.1%	-		-		-	
<i>Eragrostis lehmannia</i>	Lehmann lovegrass	0.25	0.1%	-		T		-	
<i>Polypogon monspeliensis</i>	rabbitsfoot grass	T		T		-		0.16	0.2%
<i>Boerhavia coccinea</i>	scarlet spiderling	T		0.11	0.1%			T	
<i>Kochia scoparia</i>	fireweed	-		-		-		0.02	0.0%
<i>Tamarix ramosissima</i>	five-stamen tamarisk	0.02	0.0%	T		-		T	
<i>Xanthium strumarium</i>	rough cocklebur	0.02	0.0%	T		-		-	
<i>Sonchus asper</i>	spiny sowthistel	-		0.01	0.0%	-		T	
<i>Eragrostis curvula</i>	weeping lovegrass	-		-		-		-	
<i>Acrotilon repens</i>	Russian knapweed	-		-		-		T	
<i>Ailanthus altissima</i>	tree of heaven	T		-		-		T	
<i>Arundo donax</i>	giant reed	-		-		T		-	
<i>Bromus japonicus</i>	Japanese brome	-		-		-		T	
<i>Centaurea melitensis</i>	Malta starthistle	T		T		-		T	
<i>Conyza canadensis</i>	Canadian horseweed	-		T		-		-	
<i>Digitaria sanguinalis</i>	hairy crabgrass	T		T		-		T	
<i>Elaeagnus angustifolia</i>	Russian olive	-		-		-		T	

Table 3-7. Area and percent infested by target species measured by roaming surveys, 2004, cont.

Scientific name	Common name	MOCA				TUZI				Total ha infested
		Castle unit		Well unit		Main unit		Expansion		
		Ha affected	Percent affected	Ha affected	Percent affected	Ha affected	Percent affected	Ha affected	Percent affected	
<i>Elaeagnus cilianensis</i>	stinkgrass	-	-	-	-	T	-	-	-	T
<i>Lactuca serriola</i>	prickly lettuce	T	-	T	-	-	-	T	-	T
<i>Lamium amplexicaule</i>	henbit	-	-	-	-	T	-	-	-	T
<i>Lolium perenne</i>	perennial ryegrass	T	-	-	-	-	-	-	-	T
<i>Medicago</i> sp.	alfalfa	-	-	-	-	T	-	-	-	T
<i>Melilotus albus</i>	white sweetclover	T	-	T	-	-	-	T	-	T
<i>Melilotus indicus</i>	yellow sweetclover	T	-	T	-	-	-	T	-	T
<i>Polygonum aviculare</i>	prostrate knotweed	T	-	-	-	-	-	T	-	T
<i>Rumex crispus</i>	curly dock	T	-	T	-	-	-	T	-	T
<i>Solanum elaeagnifolium</i>	silverleaf nightshade	-	-	-	-	T	-	-	-	T
<i>Sonchus oleraceus</i>	common sowthistle	-	-	-	-	-	-	T	-	T
<i>Torilis arvensis</i>	spreading hedgeparsley	-	-	-	-	-	-	T	-	T
<i>Tribulus terrestris</i>	puncture vine	-	-	T	-	-	-	T	-	T
<i>Verbascum thapsus</i>	woolly mullein	T	-	T	-	-	-	-	-	T

T denotes area of <0.01 ha.
 - denotes species not observed.




Figure 3-4. Area infested by non-native species, calculated from roaming surveys, 2004.

Table 3-8. Small, concentrated patches of target species in Montezuma Castle and Tuzigoot national monuments, 2004.

Scientific name	Common name	Park unit	Area occupied (ha)	Percent cover
<i>Cynodon dactylon</i>	Bermuda grass	MOWE	0.53	75–95%
<i>Sorghum halepense</i>	johnsongrass	MOWE	2.32	75–95%
<i>Sisymbrium irio</i>	London rocket	TUZI	5.16	5–25%
<i>Schismus barbatus</i>	common Mediterranean grass	MOCC	175.06	5–25%
<i>Salsola kali</i>	Russian thistle	MOCC	175.06	5–25%

Table 3-9. Target species occurrences and relative abundances within circular plots, 2003.

Species	Common name	MOCA				TUZI	
		Castle unit		Well unit		Occurrences	Foliar cover
		Occurrences	Foliar cover	Occurrences	Foliar cover		
<i>Avena fatua</i>	wild oats	-	-	1	0.03%	-	-
<i>Boerhavia coccinea</i>	scarlet spiderling	-	-	2	0.11%	-	-
<i>Brassica tournefortii</i>	Sahara mustard	-	-	-	-	-	-
<i>Bromus rigidus</i>	ripgut brome	6	2.78%	3	2.72%	-	-
<i>Bromus madritensis</i> ssp. <i>rubens</i>	red brome	20	8.78%	21	5.39%	48	4.02%
<i>Bromus tectorum</i>	cheatgrass	12	2.50%	1	0.03%	-	-
<i>Centaurea melitensis</i>	Malta starthistle	-	-	1	0.03%	-	-
<i>Centaurea solstitialis</i>	yellow starthistle	-	-	-	-	-	-
<i>Conyza canadensis</i>	Canadian horseweed	-	-	-	-	-	-
<i>Cynodon dactylon</i>	Bermuda grass	-	-	1	0.69%	-	-
<i>Cyperus esculentus</i>	yellow nutsedge	-	-	-	-	-	-
<i>Digitaria sanguinalis</i>	hairy crabgrass	-	-	-	-	-	-
<i>Echinochloa crus-galli</i>	barnyard grass	-	-	-	-	-	-
<i>Eragrostis curvula</i>	weeping lovegrass	-	-	-	-	-	-
<i>Erodium cicutarium</i>	redstem storksbill	3	0.08%	8	3.61%	67	1.30%
<i>Helianthus annuus</i>	common sunflower	-	-	-	-	-	-
<i>Heterotheca subaxillaris</i>	camphorweed	-	-	-	-	-	-
<i>Hordeum</i> spp.	common barley	4	1.31%	-	-	-	-
<i>Lactuca serriola</i>	prickly lettuce	-	-	-	-	-	-
<i>Marrubium vulgare</i>	horehound	-	-	1	0.28%	1	0.01%
<i>Melilotus albus</i>	white sweetclover	-	-	-	-	-	-
<i>Polygonum aviculare</i>	prostrate knotweed	-	-	1	T	5	T
<i>Polypogon monspeliensis</i>	rabbitsfoot grass	1	T	-	-	-	-
<i>Rumex crispus</i>	curly dock	1	0.03%	-	-	-	-
<i>Salsola</i> spp.	Russian thistle	-	-	1	0.03%	-	-
<i>Schismus barbatus</i>	common Mediterranean grass	7	0.56%	5	0.14%	-	-
<i>Setaria viridis</i>	green bristlegrass	-	-	-	-	-	-
<i>Sisymbrium irio</i>	London rocket	3	0.61%	4	0.11%	23	0.77%
<i>Sorghum halepense</i>	johnsongrass	1	0.03%	-	-	-	-
<i>Tamarix</i> spp.	tamarisk	-	-	-	-	-	-
<i>Tribulus terrestris</i>	puncture vine	-	-	-	-	-	-
<i>Verbascum thapsus</i>	woolly mullein	-	-	-	-	-	-
<i>Xanthium strumarium</i>	rough cocklebur	-	-	-	-	-	-

T denotes relative abundance <0.01%.

- denotes species not observed.

Table 3-10. Target species occurrences and relative abundances within circular plots, 2004.

Scientific name	Common name	MOCA				TUZI	
		Castle unit		Well unit		Occurrences	Foliar cover
		Occurrences	Foliar cover	Occurrences	Foliar cover		
<i>Acroptilon repens</i>	Russian knapweed	-	-	-	-	-	-
<i>Ailanthus altissima</i>	tree of heaven	-	-	-	-	-	-
<i>Avena fatua</i>	wild oats	-	-	-	-	-	-
<i>Boerhavia coccinea</i>	scarlet spiderling	-	-	-	-	-	-
<i>Bromus japonicus</i>	Japanese brome	-	-	2	2.3%	-	-
<i>Bromus madritensis</i> ssp. <i>rubens</i>	red brome	10	1.3%	8	1.3%	10	4.7%
<i>Bromus rigidus</i>	ripgut brome	-	-	-	-	-	-
<i>Bromus tectorum</i>	cheatgrass	-	-	2	0.1%	-	-
<i>Centaurea melitensis</i>	Malta starthistle	-	-	-	-	-	-
<i>Conyza canadensis</i>	Canadian horseweed	-	-	-	-	-	-
<i>Cynodon dactylon</i>	Bermuda grass	-	-	-	-	-	-
<i>Digitaria sanguinalis</i>	hairy crabgrass	-	-	-	-	-	-
<i>Elaeagnus angustifolia</i>	Russian olive	-	-	-	-	-	-
<i>Eragrostis lehmannia</i>	Lehmann lovegrass	-	-	-	-	-	-
<i>Erodium cicutarium</i>	redstem storksbill	5	2.7%	6	1.5%	10	25.6%
<i>Heterotheca subaxillaris</i>	camphorweed	-	-	-	-	-	-
<i>Hordeum leporinum</i>	hare barley	1	1.1%	3	T	-	-
<i>Kochia scoparia</i>	fireweed	-	-	-	-	-	-
<i>Lactuca serriola</i>	prickly lettuce	1	T	-	-	-	-
<i>Linaria dalmatica</i>	Dalmatian toadflax	-	-	-	-	-	-
<i>Lolium perenne</i>	perennial ryegrass	-	-	-	-	-	-
<i>Marrubium vulgare</i>	horehound	-	-	1	1.7%	1	T
<i>Melilotus albus</i>	white sweetclover	-	-	-	-	-	-
<i>Melilotus indicus</i>	yellow sweetclover	-	-	-	-	-	-
<i>Polygonum aviculare</i>	prostrate knotweed	-	-	-	-	-	-
<i>Polypogon monspeliensis</i>	rabbitsfoot grass	-	-	-	-	-	-
<i>Rumex crispus</i>	curly dock	-	-	-	-	-	-
<i>Salsola kali</i>	Russian thistle	2	T	3	0.2%	-	-
<i>Schismus barbatus</i>	common Mediterranean grass	6	0.1%	-	-	-	-
<i>Sisymbrium irio</i>	London rocket	-	-	-	-	8	0.2%
<i>Sonchus asper</i>	spiny sowthistle	-	-	-	-	-	-
<i>Sorghum halepense</i>	johnsongrass	-	-	-	-	-	-
<i>Tamarix ramosissima</i>	five-stamen tamarisk	-	-	-	-	-	-
<i>Torilis arvensis</i>	spreading hedgeparsley	-	-	-	-	-	-
<i>Tribulus terrestris</i>	puncture vine	-	-	-	-	-	-
<i>Verbascum thapsus</i>	woolly mullein	-	-	-	-	-	-
<i>Xanthium strumarium</i>	rough cocklebur	-	-	-	-	-	-

T denotes relative abundance <0.01%.

- denotes species not observed.

Table 3-11. On a 100-point scale, ranks of target species for significance of impact, innate ability to become a pest, and feasibility of control or management.

Scientific name	Common name	Impact	Pest	Control
<i>Bromus tectorum</i>	cheatgrass	78	38	72
<i>Bromus madritensis</i> ssp. <i>rubens</i>	red brome	59	51	73
<i>Erodium cicutarium</i>	redstem storksbill	70	49	59
<i>Tamarix</i> spp.	tamarisk	79	42	52
<i>Hordeum</i> spp.	common barley	56	47	68
<i>Bromus rigidus</i>	ripgut brome	59	31	73
<i>Sorghum halepense</i>	johnsongrass	75	35	49
<i>Polypogon monspeliensis</i>	rabbitsfoot grass	65	45	49
<i>Sisymbrium irio</i>	London rocket	59	44	52
<i>Cynodon dactylon</i>	Bermuda grass	65	27	59
<i>Linaria dalmatica</i>	Dalmatian toadflax	60	38	53
<i>Digitaria sanguinalis</i>	hairy crabgrass	63	16	65
<i>Salsola kali</i>	Russian thistle	60	33	43
<i>Ailanthus altissima</i>	tree of heaven	57	31	45
<i>Avena fatua</i>	wild oats	65	15	52
<i>Schismus barbatus</i>	common Mediterranean grass	59	38	32
<i>Xanthium strumarium</i>	rough cocklebur	52	25	52
<i>Centaurea melitensis</i>	Malta starthistle	67	15	45
<i>Elaeagnus angustifolia</i>	Russian olive	56	15	56
<i>Conyza canadensis</i>	Canadian horseweed	63	9	52
<i>Rumex crispus</i>	curly dock	44	18	60
<i>Acroptilon repens</i>	Russian knapweed	62	22	32
<i>Bromus japonicus</i>	Japanese brome	59	13	44
<i>Setaria viridis</i>	green bristlegrass	56	11	48
<i>Eragrostis curvula</i>	weeping lovegrass	48	7	60
<i>Brassica tournefortii</i>	Sahara mustard	60	11	40
<i>Tribulus terrestris</i>	puncture vine	51	13	47
<i>Boerhavia coccinea</i>	scarlet spiderling	35	35	41
<i>Kochia scoparia</i>	fireweed	52	29	29
<i>Centaurea solstitialis</i>	yellow starthistle	57	11	39
<i>Cyperus esculentus</i>	yellow nutsedge	46	7	49
<i>Lactuca serriola</i>	prickly lettuce	46	22	29
<i>Melilotus albus</i> or <i>indicus</i>	sweetclover	44	15	37
<i>Echinochloa crus-galli</i>	barnyard grass	49	7	35
<i>Verbascum thapsus</i>	woolly mullein	40	9	33
<i>Polygonum aviculare</i>	prostrate knotweed	49	9	23
<i>Heterotheca subaxillaris</i>	camphorweed	33	15	33
<i>Sonchus asper</i>	spiny sowthistle	41	13	21
<i>Marrubium vulgare</i>	horehound	38	15	20
<i>Helianthus annuus</i>	common sunflower	37	16	17




Figure 3-6. Plot of species ranked using the APRS, showing ease of control versus ability to become a pest. Size of dot represents the impact score derived using the APRS.

4 Discussion

To our knowledge, this represents one of the first applications of roaming surveys to non-native species mapping in Arizona. As a result, this effort served as a test of the method as well as a data-collection exercise. Some adjustments were made as data collection proceeded. For example, the number of circular plots collected for MOCC and MOWE was drastically reduced (from 100 to 19 per park unit) in 2003. In addition, we believe that several species were underestimated in 2003 due to the time of year of the surveys. By July, several of the annual species were already dormant or dead, making them difficult to identify, so additional fieldwork was conducted earlier in the growing season in 2004.

This method has yielded detailed maps of the location, distribution, and abundance of the most problematic plants within Arizona national parks, which will enhance decisions made by land managers. In addition, the data collected will serve as a baseline for long-term monitoring, allowing for evaluation in changes of weed populations over time. It will also feed directly into the Exotic Plant Management Plan under development for Montezuma Castle and Tuzigoot national monuments.

4.1 Comparison of methods

The circular plots were implemented to estimate spatial bias and accuracy of the roaming surveys. Stohlgren et al. (no date) suggest randomly locating these plots and comparing the foliar cover of the target species within to the foliar cover of target species collected during the roaming surveys. This comparison identifies observer bias to particular regions of the study area (i.e., near roads or riparian areas).

As a whole, the circular plots detected fewer numbers and lower abundances of target species (Table 4-1). In 2003, the circular-plot method added one target species each at MOCC and MOWE, and two target species at TUZI. The stark difference in target species detected by the two methods in 2003 suggests that MOCC and MOWE were not sufficiently sampled using the circular-plot method. Per unit area, TUZI received many more circular plots (1 plot/0.9 ha, compared to 1 plot/19.3 ha for MOCC and 1 plot/8.6 ha for MOWE). However, Stohlgren et al. (no date) suggest allocating only 10% of field resources to the supplemental circular-plot method. Nineteen

Table 4-1. Target species detected by roaming surveys and circular plots, 2003.

Method	MOCA				TUZI	
	Castle unit		Well unit			
	2003	2004	2003	2004	2003	2004
Roaming surveys	26	28	32	27	3	8
Circular plots	10	6	13	7	5	4
Species in common	9	6	12	6	3	4

circular plots for TUZI were sampled in 2½ days, and roaming surveys for TUZI were completed in three days, allocating 45% of field resources to sufficient circular plots. Based on these numbers, circular plots were drastically reduced for the remaining two park units. As a result, fewer species were detected by circular plots within the 10%-of-resources rule.

In 2004, approximately 10% of field resources were allocated to collecting circular-plot data. However, it appears, from Table 4-1, that more time spent with the circular-plot method would have yielded many more target species. The target species detected with the circular-plot method repeated those species located using the roaming surveys for all three units.

The circular-plot method described by Stohlgren et al. (no date) for assessing the roaming surveys performed similarly to previous efforts (Mau-Crimmins et al. 2003), but seemed to require much more than the 10% of resources specified in these guidelines. The small sample sizes of circular plots employed for MOCC and MOWE in 2003 performed much worse than roaming surveys for detecting target species; however, foliar cover for detected species was consistent among the methods. In 2004, the small sample sizes of circular plots yielded much smaller calculated relative abundances for the target species, suggesting the need for many more circular plots. When implemented in small numbers, the circular-plot method appears to be most useful for checking the consistency in measurements made by field crew members. Larger numbers of circular plots are required to assess consistency in estimating foliar cover across the two methods, as in previous efforts (Mau-Crimmins et al. 2003).

Overall, roaming surveys consistently estimated higher percent cover than circular plots. This may be due to the cover classes used in roaming surveys, or due to the small number of circular

plots that can be estimated using the 10% of field resources suggested by Stohlgren et al. (no date). From these two seasons of surveys, we conclude that the roaming surveys are performing well on these small park units, and that the best use of the circular plots is to check the consistency of estimates made by the field observers.

Figures 4-1 and 4-2 depict foliar cover of target species collected via circular plots against that collected via roaming surveys in 2003 and 2004, respectively. Both datasets were transformed using a cube-root transformation. The relative abundance of target species was somewhat consistent between the methods: $R^2 = 0.77$ (2003) and $R^2 = 0.53$ (2004). However, abundance calculated from the circular plots in 2004 was two orders of magnitude lower than that calculated from the roaming surveys. The positive correlation observed in this study is consistent with that of a previous study (Mau-Crimmins et al. 2003). However, in the previous study, much more time in the field was allocated to the circular-plot method.

The maps and weeds rankings produced in this report reflect approximately 700 hours of field-work and 360 hours of work in the office. Based on the accuracy assessment performed and the effort expended to yield these products, we feel that roaming surveys are a cost-effective method for producing maps of non-native species for these park units.

4.2 Alien Plants Ranking System

The Alien Plants Ranking System (Hiebert and Stubbendieck 1993) is a sensible approach to prioritizing non-native species for management efforts. The data collected and ranking performed for this project creates a foundation for the recommendations to be made in the Exotic Plant Management Plan being drafted for Montezuma Castle and Tuzigoot national monuments. A prioritized list of the target species, based on their impacts, ability to become a pest, and ease of control is a strong complement to the distribution and abundance data collected in the field.

Hiebert and Stubbendieck (1993) recommend implementing the APRS to identify species that are easy to control but have large impacts and to concentrate management efforts on these plants first. No species fell within this quadrant of the graph for Montezuma Castle and Tuzigoot NMs (see Figure 3-5). Several target species fall within the quadrant labeled “easy to control, lesser impact.” Efforts could be placed on this group of species to minimize the overall impact of the weed communities at the parks. Alternatively, some of the most widespread species, red brome, redstem storksbill, barley, London rocket, and cheatgrass (see Tables 3-2 and 3-6), are also the most difficult to control of the target species, and have the greatest impact (see Figure 3-5).

Additionally, several of these (ripgut brome, red brome, cheatgrass, and barley) also exhibit the greatest ability to become pests (see Figure 3-6). Careful consideration of these plants’ potential for control, likelihood of spread, and current level of infestation should be weighed as recommendations are made for the Exotic Plant Management Plan.




Figure 4-1. Relative abundance of target species collected via roaming surveys and circular plots, 2003.




Figure 4-2. Relative abundance of target species collected via roaming surveys and circular plots, 2004.

5 Literature Cited

- Hiebert, R. D., and J. Stubbendieck. 1993. Handbook for ranking exotic plants for management and control. Natural Resources Report NPS/NRMWRO/NRR-93/08. National Park Service, Fort Collins, Colorado.
- Kearney, T. H., and R. H. Peebles. 1951. Arizona flora. Berkeley: University of California Press.
- Mau-Crimmins, T., J. A. Hubbard, and G. R. McPherson. 2003. Non-native plant mapping at Montezuma Castle and Tuzigoot national monuments. Sonoran Desert Network I&M Program, Tucson, Az.
- National Park Service (NPS). 2006. Management Policies 2006. <http://www.nps.gov/policy/MP2006.pdf>.
- North American Weed Management Association (NAWMA). 2002. North American invasive plant mapping standards. <http://www.nawma.org/documents/Mapping%20Standards/MAIN%20DOCUMENT.pdf>.
- Pimentel, D., L. Lach, R. Zuniga, and D. Morrison. 1999. Environmental and economic costs associated with non-indigenous species in the United States. *BioScience* 50:53–65.
- Schnase, J. L., J. A. Smith, T. J. Stohlgren, S. Graves, and C. Trees. 2002. Biological invasions: A challenge in ecological forecasting. Pages 122–124 in IEEE International Geoscience and Remote Sensing Symposium, 24th Canadian Symposium.
- Stohlgren, T. J., D. T. Barnett, S. E. Simonson. No date. Beyond North American Weed Management Association standards. <http://www.nawma.org/documents/Mapping%20Standards/BEYOND%20NAWMA%20STANDARDS.pdf>.
- Vitousek, P. M., and L. R. Walker. 1989. Biological invasion by *Myrica faya* in Hawai'i: Plant demography, nitrogen fixation, ecosystem effects. *Ecological Monographs* 59:247–265.

Appendix A. Sample Circular Plot Field Data Collection Sheet

Sample data sheet

Botanist:

Date:

UTM:

Plot Number

Location:

Q = QUAD

SS = SUBPLOT SEARCH (1 for presence)

TRAMPLING: 1 undisturbed, 2 moderate, 3 heavy

.01 indicates less than 1 percent cover


1 indicates species presence in the subplot search column


All cover estimates are to the closest 1%


N/I Native or introduced

[illegible]


Appendix B. Maps of Species Detected During Roaming Surveys, 2003.











Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
 Data projection: UTM NAD83
 Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.















Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
 Data projection: UTM NAD83
 Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.













Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
 Data projection: UTM NAD83
 Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.












Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
 Data projection: UTM NAD83
 Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.












Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.


Data projection: UTM NAD83


Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.
















Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
 Data projection: UTM NAD83
 Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Avena fatua Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Boerhavia coccinea Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Brassica tournefortii Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Bromus madritensis spp. *rubens* Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Bromus rigidus
Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Bromus tectorum
Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Centaurea melitensis
Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Centaurea solstitialis Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Conyza canadensis
Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Cynodon dactylon Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Cyperus esculentus Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Digitaria sanguinalis Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Echinochloa crus-galli Montezuma Castle National Monument, Well Unit, AZ



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.


Eragrostis curvula Montezuma Castle National Monument, Well Unit, AZ



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.



Erodium cicutarium Montezuma Castle National Monument, Well Unit, AZ

- Legend**
- Erodium cicutarium points
 - ▭ Park boundary
 - ▭ Buildings
 - ▭ Roads
- Erodium cicutarium percent cover**
- <1%
 - 1-5%
 - 5-25%
 - 25-50%
 - 50-75%
 - 75-95%
 - 95-100%





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Helianthus annuus Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Heterotheca subaxillaris Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Hordeum spp.
Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Lactuca serriola
Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Marrubium vulgare
Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Melilotus albus
Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Polypogon monspeliensis
Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Rumex crispus
Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Salsola spp.
Montezuma Castle National Monument, Well Unit, AZ



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.


Schismus barbatus Montezuma Castle National Monument, Well Unit, AZ



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Montezuma Castle National Monument, Well Unit, AZ *Setaria viridis*


- Legend**
- *Setaria viridis* points
 - Park boundary
 - Buildings
 - Roads
- Setaria viridis* percent cover**
- | | |
|---|---------|
| | <1% |
| | 1-5% |
| | 5-25% |
| | 25-50% |
| | 50-75% |
| | 75-95% |
| | 95-100% |



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
 Data projection: UTM NAD83
 Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.



Sisymbrium irio
Montezuma Castle National Monument, Well Unit, AZ

- Legend**
- *Sisymbrium irio* points
 - Park boundary
 - Buildings
 - Roads
- Sisymbrium irio* percent cover**
- | | |
|--|---------|
| | <1% |
| | 1-5% |
| | 5-25% |
| | 25-50% |
| | 50-75% |
| | 75-95% |
| | 95-100% |





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
 Data projection: UTM NAD83
 Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Sorghum halepense Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Tamarix spp.
Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Tribulus terrestris
Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Verbascum thapsus
Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Xanthium strumarium Montezuma Castle National Monument, Well Unit, AZ



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Bromus madritensis spp. *rubens*
Tuzigoot National Monument, AZ




Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins,
NPS Sonoran Desert Network I&M, 2004.

Legend

- *B. madritensis* spp. *rubens* points
- Roads
- Buildings
- Park boundary
- B. madritensis* spp. *rubens* percent cover
- 1
- 1-5%
- 5-25%
- 25-50%
- 50-75%
- 75-95%
- 95-100%

Marrubium vulgare
Tuzigoot National Monument, AZ



Legend

- Marrubium vulgare points
- Roads
- Buildings
- Park boundary
- Marrubium vulgare percent cover
- 1
- 1-5%
- 5-25%
- 25-50%
- 50-75%
- 75-95%
- 95-100%

Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Sisymbrium irio
Tuzigoot National Monument, AZ





Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, summer 2003.
Data projection: UTM NAD83
Map created by: A. Tendick and T. Mau-Crimmins,
NPS Sonoran Desert Network I&M, 2004.

Legend


- *Sisymbrium irio* points
- Roads
- Buildings
- Park boundary
- Sisymbrium irio* percent cover
- 1
- 1-5%
- 5-25%
- 25-50%
- 50-75%
- 75-95%
- 95-100%

Appendix C. Maps of Species Detected During Roaming Surveys, 2004.







Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.







Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.













Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.







Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.







Data source: collected using Garmin III GPS units and handheld computers. M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.







Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.







Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.






Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.







Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.







Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.







Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Boerhavia coccinea Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Bromus madritensis ssp. *rubens* Montezuma Castle National Monument, Well Unit, AZ

Legend

- *Bromus madritensis* ssp. *rubens* points
- ▭ Park boundary
- ▭ Buildings
- ▭ Roads

Bromus madritensis ssp. *rubens*
percent cover




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Bromus rigidus Montezuma Castle National Monument, Well Unit, AZ



Legend

- Bromus rigidus points
 - Park boundary
 - Buildings
 - Roads
- Bromus rigidus percent cover**
- | | |
|---|---------|
| | <1% |
| | 1-5% |
| | 5-25% |
| | 25-50% |
| | 50-75% |
| | 75-95% |
| | 95-100% |



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Bromus tectorum Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Centaurea melitensis Montezuma Castle National Monument, Well Unit, AZ

Legend

- *Centaurea melitensis* points
 - Park boundary
 - Buildings
 - Roads
- Centaurea melitensis*
percent cover
- | | |
|--|---------|
| | <1% |
| | 1-5% |
| | 5-25% |
| | 25-50% |
| | 50-75% |
| | 75-95% |
| | 95-100% |




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Montezuma Castle National Monument, Well Unit, AZ

Conyza canadensis



Legend

- Conyza canadensis points
 - Park boundary
 - Buildings
 - Roads
- Conyza canadensis percent cover
- | | |
|---|---------|
| | <1% |
| | 1-5% |
| | 5-25% |
| | 25-50% |
| | 50-75% |
| | 75-95% |
| | 95-100% |





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Cynodon dactylon Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Digitaria sanguinalis Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Eragrostis curvula
Montezuma Castle National Monument, Well Unit, AZ





Data Source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, summer 2004.
Data projection: UTM NAD 83
Map created by: A. Tendick and T. Mau-Crimmins, NPS Sonoran Desert Network I&M, 2004.

Erodium cicutarium Montezuma Castle National Monument, Well Unit, AZ



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Heterotheca subaxillaris Montezuma Castle National Monument, Well Unit, AZ



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.



Montezuma Castle National Monument, Well Unit, AZ

Hordeum spp.

Legend

- Hordeum spp. points
- ▬ Park boundary
- ▬ Buildings
- ▬ Roads


Hordeum spp.
percent cover



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Lactuca serriola Montezuma Castle National Monument, Well Unit, AZ


- Legend**
- Lactuca serriola points
 - Park boundary
 - Buildings
 - Roads
- Lactuca serriola percent cover
- | | |
|--|---------|
| | <1% |
| | 1-5% |
| | 5-25% |
| | 25-50% |
| | 50-75% |
| | 75-95% |
| | 95-100% |



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Montezuma Castle National Monument, Well Unit, AZ *Linaria dalmatica*

- Legend**
- Linaria dalmatica points
 - ▭ Park boundary
 - ▭ Buildings
 - ▭ Roads
- Linaria dalmatica percent cover
- | |
|---------|
| <1% |
| 1-5% |
| 5-25% |
| 25-50% |
| 50-75% |
| 75-95% |
| 95-100% |




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Marrubium vulgare Montezuma Castle National Monument, Well Unit, AZ

Legend

- *Marrubium vulgare* points
 - Park boundary
 - Buildings
 - Roads
- Marrubium vulgare*
percent cover
- | | |
|--|---------|
| | <1% |
| | 1-5% |
| | 5-25% |
| | 25-50% |
| | 50-75% |
| | 75-95% |
| | 95-100% |



Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.



Montezuma Castle National Monument, Well Unit, AZ

Melilotus albus

Legend


- Melilotus albus points
- ▭ Park boundary
- ▭ Buildings
- ▭ Roads

Melilotus albus percent cover




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Montezuma Castle National Monument, Well Unit, AZ *Melilotus indicus*




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Polypogon monspeliensis Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Schismus barbatus Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Montezuma Castle National Monument, Well Unit, AZ *Sisymbrium irio*




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Sonchus asper Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Sorghum halepense Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
 Data projection: UTM NAD83
 Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Tribulus terrestris Montezuma Castle National Monument, Well Unit, AZ




Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.

Verbascum thapsus Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.


Xanthium strumarium Montezuma Castle National Monument, Well Unit, AZ





Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring and summer 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy, NPS Sonoran Desert Network I&M, 2004.




























































































Arundo donax
Tuzigoot National Monument, AZ



Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Legend

- *Arundo donax* points
-  Roads
-  Buildings
-  Park boundary

Arundo donax percent cover

-  1
-  1-5%
-  5-25%
-  25-50%
-  50-75%
-  75-95%
-  95-100%

Bromus madritensis ssp. rubens
Tuzigoot National Monument, AZ



Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Legend

- *Bromus madritensis ssp. rubens* points
- Roads
- Buildings
- Park boundary
- Bromus madritensis ssp. rubens* percent cover
 - 1
 - 1-5%
 - 5-25%
 - 25-50%
 - 50-75%
 - 75-95%
 - 95-100%

Cynodon dactylon
Tuzigoot National Monument, AZ




Legend

- *Cynodon dactylon* points
 - Roads
 - Buildings
 - Park boundary
- Cynodon dactylon* percent cover**
- 1
 - 1-5%
 - 5-25%
 - 25-50%
 - 50-75%
 - 75-95%
 - 95-100%

Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Eragrostis cilianensis
Tuzigoot National Monument, AZ



0.05 0 0.05 Kilometers

Legend


- *Eragrostis cilianensis* points
-  Roads
-  Buildings
-  Park boundary

***Eragrostis cilianensis* percent cover**

-  1
-  1-5%
-  5-25%
-  25-50%
-  50-75%
-  75-95%
-  95-100%

Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Eragrostis lehmanniana
Tuzigoot National Monument, AZ




Legend

- *Eragrostis lehmanniana* points
- Roads
- Buildings
- Park boundary
- Eragrostis lehmanniana*
percent cover
- 1
- 1-5%
- 5-25%
- 25-50%
- 50-75%
- 75-95%
- 95-100%

Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Erodium cicutarium
Tuzigoot National Monument, AZ



Legend

- Erodium cicutarium points
- Roads
- ▬ Buildings
- ▭ Park boundary

Erodium cicutarium percent cover

- 1
- 1-5%
- 5-25%
- 25-50%
- 50-75%
- 75-95%
- 95-100%

Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Hordeum spp.
Tuzigoot National Monument, AZ



Legend


- *Hordeum spp.* points
- Roads
- Buildings
- Park boundary

***Hordeum spp.* percent cover**

- 1
- 1-5%
- 5-25%
- 25-50%
- 50-75%
- 75-95%
- 95-100%

Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Lamium amplexicaule
Tuzigoot National Monument, AZ



0.05 0 0.05 Kilometers

Legend

- *Lamium amplexicaule* points
 -  Roads
 -  Buildings
 -  Park boundary
- Lamium amplexicaule* percent cover
-  1
 -  1-5%
 -  5-25%
 -  25-50%
 -  50-75%
 -  75-95%
 -  95-100%

Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Marrubium vulgare
Tuzigoot National Monument, AZ



Legend


- *Marrubium vulgare* points
-  Roads
-  Buildings
-  Park boundary

***Marrubium vulgare* percent cover**

-  1
-  1-5%
-  5-25%
-  25-50%
-  50-75%
-  75-95%
-  95-100%

Data source: collected using Garmin III GPS units and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Medicago spp.
Tuzigoot National Monument, AZ



0.05 0 0.05 Kilometers

Legend


- *Medicago spp.* points
-  Roads
-  Buildings
-  Park boundary

***Medicago spp.* percent cover**

-  1
-  1-5%
-  5-25%
-  25-50%
-  50-75%
-  75-95%
-  95-100%

Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Schismus barbatus
Tuzigoot National Monument, AZ



Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.


Legend

- *Schismus barbatus* points
- Roads
- Buildings
- Park boundary

***Schismus barbatus* percent cover**

- 1
- 1-5%
- 5-25%
- 25-50%
- 50-75%
- 75-95%
- 95-100%

Sisymbrium irio
Tuzigoot National Monument, AZ



Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.


Legend

- *Sisymbrium irio* points
- Roads
- Buildings
- Park boundary

Sisymbrium irio percent cover

- 1
- 1-5%
- 5-25%
- 25-50%
- 50-75%
- 75-95%
- 95-100%

Solanum elaeagnifolium
Tuzigoot National Monument, AZ



Data source: collected using Garmin III GPS units
and handheld computers, M. Mauzy and M. Quinn, spring 2004.
Data projection: UTM NAD83
Map created by: M. Mauzy,
NPS Sonoran Desert Network I&M, 2004.

Legend

- *Solanum elaeagnifolium* points
- Roads
- Buildings
- Park boundary
- Solanum elaeagnifolium*
percent cover
- 1
- 1-5%
- 5-25%
- 25-50%
- 50-75%
- 75-95%
- 95-100%

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS 309/100671, November 2009

National Park Service
U.S. Department of the Interior



Natural Resource Program Center

1201 Oak Ridge Drive, Suite 150
Fort Collins, Colorado 80525

www.nature.nps.gov